

A NOTE ON REPRESENTATIONS OF SOME AFFINE VERTEX ALGEBRAS OF TYPE D

OZREN PERŠE

University of Zagreb, Croatia

ABSTRACT. In this note we construct a series of singular vectors in universal affine vertex operator algebras associated to $D_\ell^{(1)}$ of levels $n-\ell+1$, for $n \in \mathbb{Z}_{>0}$. For $n = 1$, we study the representation theory of the quotient vertex operator algebra modulo the ideal generated by that singular vector. In the case $\ell = 4$, we show that the adjoint module is the unique irreducible ordinary module for simple vertex operator algebra $L_{D_4}(-2, 0)$. We also show that the maximal ideal in associated universal affine vertex algebra is generated by three singular vectors.

1. INTRODUCTION

The classification of irreducible modules for simple vertex operator algebra $L_{\mathfrak{g}}(k, 0)$ associated to affine Lie algebra $\hat{\mathfrak{g}}$ of level k is still an open problem for general $k \in \mathbb{C}$ ($k \neq -h^\vee$). This problem is connected with the description of the maximal ideal in the universal affine vertex algebra $N_{\mathfrak{g}}(k, 0)$. One approach to this classification problem is through construction of singular vectors in $N_{\mathfrak{g}}(k, 0)$.

The known (non-generic) cases include positive integer levels (cf. [14, 19, 20]) and some special cases of rational admissible levels, in the sense of Kac and Wakimoto ([17]) (cf. [1, 3, 4, 6, 9, 21, 22]). It turns out that negative integer levels also have some interesting properties. They appeared in bosonic realizations in [10], and also recently in the context of conformal embeddings (cf. [5]).

In this note we study a vertex operator algebra associated to affine Lie algebra of type $D_\ell^{(1)}$ and negative integer level $-\ell + 2$. This level appeared in

2010 *Mathematics Subject Classification.* 17B69, 17B67, 81R10.

Key words and phrases. Vertex operator algebra, affine Kac-Moody algebra, Zhu's algebra.

[5] in the context of conformal embedding of $L_{B_{\ell-1}}(-\ell+2, 0)$ into $L_{D_\ell}(-\ell+2, 0)$. This conformal embedding is in some sense similar to the conformal embedding of $L_{D_\ell}(-\ell+\frac{3}{2}, 0)$ into $L_{B_\ell}(-\ell+\frac{3}{2}, 0)$.

We will show that there are also similarities in singular vectors in universal affine vertex algebras $N_{B_\ell}(-\ell+\frac{3}{2}, 0)$ (studied in [21]) and $N_{D_\ell}(-\ell+2, 0)$. More generally, we construct a series of singular vectors

$$v_n = \left(\sum_{i=2}^{\ell} e_{\epsilon_1 - \epsilon_i}(-1) e_{\epsilon_1 + \epsilon_i}(-1) \right)^n \mathbf{1}$$

in $N_{D_\ell}(n-\ell+1, 0)$, for any $n \in \mathbb{Z}_{>0}$. For $n=1$, we study the representation theory of the quotient $N_{D_\ell}(-\ell+2, 0)$ modulo the ideal generated by v_1 . Using the methods from [1, 2, 4, 20], we obtain the classification of irreducible weak modules in the category \mathcal{O} for that vertex algebra. It turns out that there are infinitely many of these modules.

In the special case $\ell=4$, we obtain the classification of irreducible weak modules from the category \mathcal{O} for simple vertex operator algebra $L_{D_4}(-2, 0)$. This vertex algebra also appeared in [5] in the context of conformal embedding of $L_{G_2}(-2, 0)$ into $L_{D_4}(-2, 0)$. It follows that there are finitely many irreducible weak $L_{D_4}(-2, 0)$ -modules from the category \mathcal{O} , that the adjoint module is the unique irreducible ordinary $L_{D_4}(-2, 0)$ -module, and that every ordinary $L_{D_4}(-2, 0)$ -module is completely reducible. We also show that the maximal ideal in $N_{D_4}(-2, 0)$ is generated by three singular vectors.

2. PRELIMINARIES

We assume that the reader is familiar with the notion of vertex operator algebra (cf. [7, 11–14, 16, 18, 19]) and Kac-Moody algebra (cf. [15]).

Let V be a vertex operator algebra. Denote by $A(V)$ the associative algebra introduced in [23], called the Zhu's algebra of V . As a vector space, $A(V)$ is a quotient of V , and we denote by $[a]$ the image of $a \in V$ under the projection of V onto $A(V)$. We recall the following fundamental result from [23]:

PROPOSITION 2.1. *The equivalence classes of the irreducible $A(V)$ -modules and the equivalence classes of the irreducible \mathbb{Z}_+ -graded weak V -modules are in one-to-one correspondence.*

Let \mathfrak{g} be a simple Lie algebra with a triangular decomposition $\mathfrak{g} = \mathfrak{n}_- \oplus \mathfrak{h} \oplus \mathfrak{n}_+$, and $\hat{\mathfrak{g}}$ the (untwisted) affine Lie algebra associated to \mathfrak{g} . Denote by $V(\mu)$ the irreducible highest weight \mathfrak{g} -module with highest weight μ , and by $L(k, \mu)$ the irreducible highest weight $\hat{\mathfrak{g}}$ -module with highest weight $k\Lambda_0 + \mu$.

Furthermore, denote by $N(k, 0)$ (or $N_{\mathfrak{g}}(k, 0)$) the universal affine vertex algebra associated to $\hat{\mathfrak{g}}$ of level $k \in \mathbb{C}$. For $k \neq -h^\vee$, $N(k, 0)$ is a vertex operator algebra with Segal-Sugawara conformal vector, and $L(k, 0)$ is a

simple vertex operator algebra. The Zhu's algebra of $N(k, 0)$ was determined in [14]:

PROPOSITION 2.2. *The associative algebra $A(N(k, 0))$ is canonically isomorphic to $U(\mathfrak{g})$. The isomorphism is given by $F : A(N(k, 0)) \rightarrow U(\mathfrak{g})$*

$$F([x_1(-n_1 - 1) \cdots x_m(-n_m - 1)\mathbf{1}]) = (-1)^{n_1 + \cdots + n_m} x_m \cdots x_1,$$

for any $x_1, \dots, x_m \in \mathfrak{g}$ and any $n_1, \dots, n_m \in \mathbb{Z}_+$.

We have:

PROPOSITION 2.3. *Assume that a $\hat{\mathfrak{g}}$ -submodule J of $N(k, 0)$ is generated by m singular vectors ($m \in \mathbb{Z}_{>0}$), i.e. $J = U(\hat{\mathfrak{g}})v^{(1)} + \dots + U(\hat{\mathfrak{g}})v^{(m)}$. Then*

$$A(N(k, 0)/J) \cong U(\mathfrak{g})/I,$$

where I is the two-sided ideal of $U(\mathfrak{g})$ generated by $u^{(1)} = F([v^{(1)}]), \dots, u^{(m)} = F([v^{(m)}])$.

Let $J = U(\hat{\mathfrak{g}})v^{(1)} + \dots + U(\hat{\mathfrak{g}})v^{(m)}$ be a $\hat{\mathfrak{g}}$ -submodule of $N(k, 0)$ generated by singular vectors $v^{(1)}, \dots, v^{(m)}$. Now we recall the method from [1, 2, 4, 20] for the classification of irreducible $A(N(k, 0)/J)$ -modules from the category \mathcal{O} by solving certain systems of polynomial equations.

Denote by ${}_L$ the adjoint action of $U(\mathfrak{g})$ on $U(\mathfrak{g})$ defined by $X_L f = [X, f]$ for $X \in \mathfrak{g}$ and $f \in U(\mathfrak{g})$. Let $R^{(j)}$ be a $U(\mathfrak{g})$ -submodule of $U(\mathfrak{g})$ generated by the vector $u^{(j)} = F([v^{(j)}])$ under the adjoint action, for $j = 1, \dots, m$. Clearly, $R^{(j)}$ is an irreducible highest weight $U(\mathfrak{g})$ -module. Let $R_0^{(j)}$ be the zero-weight subspace of $R^{(j)}$.

The next proposition follows from [1, 4, 20]:

PROPOSITION 2.4. *Let $V(\mu)$ be an irreducible highest weight $U(\mathfrak{g})$ -module with the highest weight vector v_μ , for $\mu \in \mathfrak{h}^*$. The following statements are equivalent:*

- (1) $V(\mu)$ is an $A(N(k, 0)/J)$ -module,
- (2) $R^{(j)}V(\mu) = 0$, for every $j = 1, \dots, m$,
- (3) $R_0^{(j)}v_\mu = 0$, for every $j = 1, \dots, m$.

Let $r \in R_0^{(j)}$. Clearly there exists the unique polynomial $p_r \in S(\mathfrak{h})$ such that

$$rv_\mu = p_r(\mu)v_\mu.$$

Set $\mathcal{P}_0^{(j)} = \{p_r \mid r \in R_0^{(j)}\}$, for $j = 1, \dots, m$. We have:

COROLLARY 2.5. *There is one-to-one correspondence between*

- (1) irreducible $A(N(k, 0)/J)$ -modules from the category \mathcal{O} ,
- (2) weights $\mu \in \mathfrak{h}^*$ such that $p(\mu) = 0$ for all $p \in \mathcal{P}_0^{(j)}$, for every $j = 1, \dots, m$.

In the case $m = 1$, we use the notation R , R_0 and \mathcal{P}_0 for $R^{(1)}$, $R_0^{(1)}$ and $\mathcal{P}_0^{(1)}$, respectively.

3. VERTEX OPERATOR ALGEBRA ASSOCIATED TO $D_\ell^{(1)}$ OF LEVEL $-\ell + 2$

In this section we study the representation theory of the quotient of universal affine vertex operator algebra associated to $D_\ell^{(1)}$ of level $-\ell + 2$, modulo the ideal generated by a singular vector of conformal weight two.

Denote by \mathfrak{g} the simple Lie algebra of type D_ℓ . We fix the root vectors for \mathfrak{g} as in [8, 10]. We have:

THEOREM 3.1. *Vector*

$$v_n = \left(\sum_{i=2}^{\ell} e_{\epsilon_1 - \epsilon_i}(-1) e_{\epsilon_1 + \epsilon_i}(-1) \right)^n \mathbf{1}$$

is a singular vector in $N_{D_\ell}(n - \ell + 1, 0)$, for any $n \in \mathbb{Z}_{>0}$.

PROOF. Direct verification of relations $e_{\epsilon_k - \epsilon_{k+1}}(0)v_n = 0$, for $k = 1, \dots, \ell - 1$, $e_{\epsilon_{\ell-1} + \epsilon_\ell}(0)v_n = 0$ and $f_{\epsilon_1 + \epsilon_2}(1)v_n = 0$. \square

In the case $n = 1$, we obtain the singular vector

$$(3.1) \quad v = \sum_{i=2}^{\ell} e_{\epsilon_1 - \epsilon_i}(-1) e_{\epsilon_1 + \epsilon_i}(-1) \mathbf{1}$$

in $N_{D_\ell}(-\ell + 2, 0)$.

REMARK 3.2. Vector v from relation (3.1) has a similar formula as singular vector

$$-\frac{1}{4} e_{\epsilon_1}(-1)^2 \mathbf{1} + \sum_{i=2}^{\ell} e_{\epsilon_1 - \epsilon_i}(-1) e_{\epsilon_1 + \epsilon_i}(-1) \mathbf{1}$$

for $B_\ell^{(1)}$ in $N_{B_\ell}(-\ell + \frac{3}{2}, 0)$. The representation theory of the quotient of $N_{B_\ell}(-\ell + \frac{3}{2}, 0)$ modulo the ideal generated by that vector was studied in [21].

We will consider representations of the vertex operator algebra

$$\mathcal{V}_{D_\ell}(-\ell + 2, 0) = \frac{N_{D_\ell}(-\ell + 2, 0)}{U(\hat{\mathfrak{g}})v}.$$

Proposition 2.3 gives:

PROPOSITION 3.3. *The associative algebra $A(\mathcal{V}_{D_\ell}(-\ell + 2, 0))$ is isomorphic to the algebra $U(\mathfrak{g})/I$, where I is the two-sided ideal of $U(\mathfrak{g})$ generated by*

$$u = \sum_{i=2}^{\ell} e_{\epsilon_1 - \epsilon_i} e_{\epsilon_1 + \epsilon_i}.$$

We have the following classification:

THEOREM 3.4. *For any subset $S = \{i_1, \dots, i_k\} \subseteq \{1, 2, \dots, \ell - 2\}$, $i_1 < \dots < i_k$, and $t \in \mathbb{C}$, we define weights*

$$\mu_{S,t} = \sum_{j=1}^k \left(i_j + 2 \sum_{s=j+1}^k (-1)^{s-j} i_s + (-1)^{k-j+1} (t + \ell - 1) \right) \omega_{i_j} + t\omega_{\ell-1},$$

$$\mu'_{S,t} = \sum_{j=1}^k \left(i_j + 2 \sum_{s=j+1}^k (-1)^{s-j} i_s + (-1)^{k-j+1} (t + \ell - 1) \right) \omega_{i_j} + t\omega_{\ell},$$

where $\omega_1, \dots, \omega_{\ell}$ are fundamental weights for \mathfrak{g} . Then the set

$$\{L_{D_{\ell}}(-\ell + 2, \mu_{S,t}), L_{D_{\ell}}(-\ell + 2, \mu'_{S,t}) \mid S \subseteq \{1, 2, \dots, \ell - 2\}, t \in \mathbb{C}\}$$

provides the complete list of irreducible weak $\mathcal{V}_{D_{\ell}}(-\ell + 2, 0)$ -modules from the category \mathcal{O} .

PROOF. We use the method for classification of irreducible $A(\mathcal{V}_{D_{\ell}}(-\ell + 2, 0))$ -modules in the category \mathcal{O} from Corollary 2.5. In this case $R \cong V_{D_{\ell}}(2\omega_1)$, and similarly as in [21, Lemma 28] one obtains that

$$\dim R_0 = \ell - 1.$$

Furthermore, one obtains by direct calculation that

$$(f_{\epsilon_1 - \epsilon_2} f_{\epsilon_1 + \epsilon_2})_L u \in p_1(h) + U(\mathfrak{g})\mathfrak{n}_+,$$

$$(f_{\epsilon_1 - \epsilon_{i+1}} f_{\epsilon_1 + \epsilon_{i+1}} - f_{\epsilon_1 - \epsilon_i} f_{\epsilon_1 + \epsilon_i})_L u \in p_i(h) + U(\mathfrak{g})\mathfrak{n}_+, \quad i = 2, \dots, \ell - 1,$$

where

$$(3.2) \quad p_i(h) = h_i(h_{\epsilon_i + \epsilon_{i+1}} + \ell - i - 1), \quad \text{for } i = 1, \dots, \ell - 1$$

are linearly independent polynomials in \mathcal{P}_0 . Here h_i ($i = 1, \dots, \ell$) denote the simple coroots for \mathfrak{g} and

$$h_{\epsilon_i + \epsilon_{i+1}} = h_i + 2h_{i+1} + \dots + 2h_{\ell-2} + h_{\ell-1} + h_{\ell}, \quad \text{for } i < \ell - 1.$$

Corollary 2.5 now implies that the highest weights of irreducible $A(\mathcal{V}_{D_{\ell}}(-\ell + 2, 0))$ -modules from the category \mathcal{O} are given as solutions of polynomial equations

$$(3.3) \quad p_i(h) = 0, \quad i = 1, \dots, \ell - 1.$$

First we note that for $i = \ell - 1$, we obtain the equation

$$h_{\ell-1} h_{\ell} = 0.$$

Thus, either $h_{\ell-1} = 0$ or $h_{\ell} = 0$. Assume first that $h_{\ell-1} = 0$, and let $S = \{i_1, \dots, i_k\}$, $i_1 < \dots < i_k$ be the subset of $\{1, 2, \dots, \ell - 2\}$ such that

$h_i = 0$ for $i \notin S$ and $h_i \neq 0$ for $i \in S$. Then we have the system

$$(3.4) \quad \begin{aligned} h_{i_1} + 2h_{i_2} + \dots + 2h_{i_k} + h_\ell + \ell - i_1 - 1 &= 0, \\ h_{i_2} + 2h_{i_3} + \dots + 2h_{i_k} + h_\ell + \ell - i_2 - 1 &= 0, \\ &\vdots \\ h_{i_{k-1}} + 2h_{i_k} + h_\ell + \ell - i_{k-1} - 1 &= 0, \\ h_{i_k} + h_\ell + \ell - i_k - 1 &= 0. \end{aligned}$$

The solution of this system is given by

$$\begin{aligned} h_{i_j} &= i_j + 2 \sum_{s=j+1}^k (-1)^{s-j} i_s + (-1)^{k-j+1} (t + \ell - 1), \quad \text{for } j = 1, \dots, k; \\ h_\ell &= t \quad (t \in \mathbb{C}). \end{aligned}$$

It follows that $V_{D_\ell}(\mu'_{S,t})$ is an irreducible $A(\mathcal{V}_{D_\ell}(-\ell+2, 0))$ -module. Similarly, the case $h_\ell = 0$ gives that $V_{D_\ell}(\mu_{S,t})$ is irreducible $A(\mathcal{V}_{D_\ell}(-\ell+2, 0))$ -module. We conclude that the set

$$\{V_{D_\ell}(\mu_{S,t}), V_{D_\ell}(\mu'_{S,t}) \mid S \subseteq \{1, 2, \dots, \ell-2\}, t \in \mathbb{C}\}$$

provides the complete list of irreducible $A(\mathcal{V}_{D_\ell}(-\ell+2, 0))$ -modules from the category \mathcal{O} . The claim of theorem now follows from Zhu's theory. \square

EXAMPLE 3.5. For $\ell = 4$, we have subsets $S = \emptyset, \{1\}, \{2\}, \{1, 2\}$ of the set $\{1, 2\}$, so we obtain that the set

$$(3.5) \quad \begin{aligned} &\{L_{D_\ell}(-\ell+2, t\omega_3), L_{D_\ell}(-\ell+2, t\omega_4), L_{D_\ell}(-\ell+2, (-2-t)\omega_1 + t\omega_3), \\ &L_{D_\ell}(-\ell+2, (-2-t)\omega_1 + t\omega_4), L_{D_\ell}(-\ell+2, (-1-t)\omega_2 + t\omega_3), \\ &L_{D_\ell}(-\ell+2, (-1-t)\omega_2 + t\omega_4), L_{D_\ell}(-\ell+2, t\omega_1 + (-1-t)\omega_2 + t\omega_3), \\ &L_{D_\ell}(-\ell+2, t\omega_1 + (-1-t)\omega_2 + t\omega_4) \mid t \in \mathbb{C}\} \end{aligned}$$

provides the complete list of irreducible weak $\mathcal{V}_{D_4}(-2, 0)$ -modules from the category \mathcal{O} .

Recall that a module for vertex operator algebra is called ordinary if $L(0)$ acts semisimply with finite-dimensional weight spaces. We have:

COROLLARY 3.6. *The set*

$$\{L_{D_\ell}(-\ell+2, t\omega_{\ell-1}), L_{D_\ell}(-\ell+2, t\omega_\ell) \mid t \in \mathbb{Z}_{\geq 0}\}$$

provides the complete list of irreducible ordinary $\mathcal{V}_{D_\ell}(-\ell+2, 0)$ -modules.

PROOF. If $L_{D_\ell}(-\ell+2, \mu)$ is an ordinary $\mathcal{V}_{D_\ell}(-\ell+2, 0)$ -module, then μ is a dominant integral weight. Then $\mu(h_{\epsilon_i + \epsilon_{i+1}}) \in \mathbb{Z}_{\geq 0}$, for $i = 1, \dots, \ell-1$. Relations (3.2) and (3.3) then give that

$$\mu(h_i) = 0, \quad \text{for } i = 1, \dots, \ell-2,$$

and $\mu(h_{\ell-1}) = 0$ or $\mu(h_\ell) = 0$. Thus, $\mu = t\omega_{\ell-1}$ or $\mu = t\omega_\ell$, and $t \in \mathbb{Z}_{\geq 0}$ since μ is a dominant integral weight. \square

It follows that:

COROLLARY 3.7. *The set of irreducible ordinary $L_{D_\ell}(-\ell + 2, 0)$ -modules is a subset of the set*

$$\{L_{D_\ell}(-\ell + 2, t\omega_{\ell-1}), L_{D_\ell}(-\ell + 2, t\omega_\ell) \mid t \in \mathbb{Z}_{\geq 0}\}.$$

4. CASE $\ell = 4$

In this section we study the case $\ell = 4$. We determine the classification of irreducible weak $L_{D_4}(-2, 0)$ -modules from the category \mathcal{O} . It turns out that there are finitely many of these modules and that the adjoint module is the unique irreducible ordinary $L_{D_4}(-2, 0)$ -module. We also show that the maximal ideal in $N_{D_4}(-2, 0)$ is generated by three singular vectors.

Denote by θ the automorphism of $N_{D_4}(-2, 0)$ induced by the automorphism of the Dynkin diagram of D_4 of order three such that

$$\begin{aligned} \theta(\epsilon_1 - \epsilon_2) &= \epsilon_3 - \epsilon_4, & \theta(\epsilon_2 - \epsilon_3) &= \epsilon_2 - \epsilon_3, \\ \theta(\epsilon_3 - \epsilon_4) &= \epsilon_3 + \epsilon_4, & \theta(\epsilon_3 + \epsilon_4) &= \epsilon_1 - \epsilon_2. \end{aligned}$$

Relation (3.1) implies that

$v = (e_{\epsilon_1 - \epsilon_2}(-1)e_{\epsilon_1 + \epsilon_2}(-1) + e_{\epsilon_1 - \epsilon_3}(-1)e_{\epsilon_1 + \epsilon_3}(-1) + e_{\epsilon_1 - \epsilon_4}(-1)e_{\epsilon_1 + \epsilon_4}(-1))\mathbf{1}$ is a singular vector in $N_{D_4}(-2, 0)$. Furthermore,

$$\begin{aligned} \theta(v) &= (e_{\epsilon_3 - \epsilon_4}(-1)e_{\epsilon_1 + \epsilon_2}(-1) - e_{\epsilon_2 - \epsilon_4}(-1)e_{\epsilon_1 + \epsilon_3}(-1) \\ &\quad + e_{\epsilon_2 + \epsilon_3}(-1)e_{\epsilon_1 - \epsilon_4}(-1))\mathbf{1}, \end{aligned}$$

and

$$\begin{aligned} \theta^2(v) &= (e_{\epsilon_3 + \epsilon_4}(-1)e_{\epsilon_1 + \epsilon_2}(-1) - e_{\epsilon_2 + \epsilon_4}(-1)e_{\epsilon_1 + \epsilon_3}(-1) \\ &\quad + e_{\epsilon_1 + \epsilon_4}(-1)e_{\epsilon_2 + \epsilon_3}(-1))\mathbf{1} \end{aligned}$$

are also singular vectors in $N_{D_4}(-2, 0)$. We consider the vertex operator algebra

$$\tilde{L}_{D_4}(-2, 0) = \frac{N_{D_4}(-2, 0)}{J},$$

where J is the ideal in $N_{D_4}(-2, 0)$ generated by vectors v , $\theta(v)$ and $\theta^2(v)$.

Proposition 2.3 gives that the associative algebra $A(\tilde{L}_{D_4}(-2, 0))$ is isomorphic to the algebra $U(\mathfrak{g})/I$, where I is the two-sided ideal of $U(\mathfrak{g})$ generated by u , $\theta(u)$ and $\theta^2(u)$, and

$$u = e_{\epsilon_1 - \epsilon_2}e_{\epsilon_1 + \epsilon_2} + e_{\epsilon_1 - \epsilon_3}e_{\epsilon_1 + \epsilon_3} + e_{\epsilon_1 - \epsilon_4}e_{\epsilon_1 + \epsilon_4}.$$

PROPOSITION 4.1. *We have:*

(i) *The set*

$$\{L_{D_4}(-2, 0), L_{D_4}(-2, -2\omega_1), L_{D_4}(-2, -2\omega_3), L_{D_4}(-2, -2\omega_4), L_{D_4}(-2, -\omega_2)\}$$

provides a complete list of irreducible weak $\tilde{L}_{D_4}(-2, 0)$ -modules from the category \mathcal{O} .

(ii) *$L_{D_4}(-2, 0)$ is the unique irreducible ordinary module for $\tilde{L}_{D_4}(-2, 0)$.*

PROOF. (i) We use the method for classification from Corollary 2.5. In this case $R^{(1)} \cong V_{D_4}(2\omega_1)$, $R^{(2)} \cong V_{D_4}(2\omega_3)$, $R^{(3)} \cong V_{D_4}(2\omega_4)$ and

$$\dim R_0^{(1)} = \dim R_0^{(2)} = \dim R_0^{(3)} = 3.$$

Using polynomials from relation (3.2) and automorphisms θ and θ^2 , one obtains that the highest weights μ of $A(\tilde{L}_{D_4}(-2, 0))$ -modules $V_{D_4}(\mu)$ are obtained as solutions of these 9 polynomial equations:

$$\begin{aligned} h_{\epsilon_1 - \epsilon_2}(h_{\epsilon_1 + \epsilon_2} + 2) &= 0, \\ h_{\epsilon_2 - \epsilon_3}(h_{\epsilon_2 + \epsilon_3} + 1) &= 0, \\ h_{\epsilon_3 - \epsilon_4}h_{\epsilon_3 + \epsilon_4} &= 0, \\ h_{\epsilon_3 - \epsilon_4}(h_{\epsilon_1 + \epsilon_2} + 2) &= 0, \\ h_{\epsilon_2 - \epsilon_3}(h_{\epsilon_1 + \epsilon_4} + 1) &= 0, \\ h_{\epsilon_3 + \epsilon_4}h_{\epsilon_1 - \epsilon_2} &= 0, \\ h_{\epsilon_3 + \epsilon_4}(h_{\epsilon_1 + \epsilon_2} + 2) &= 0, \\ h_{\epsilon_2 - \epsilon_3}(h_{\epsilon_1 - \epsilon_4} + 1) &= 0, \\ h_{\epsilon_1 - \epsilon_2}h_{\epsilon_3 - \epsilon_4} &= 0. \end{aligned}$$

This easily gives that $\mu = 0, -2\omega_1, -2\omega_3, -2\omega_4$ or $-\omega_2$, and the claim follows from Zhu's theory.

Claim (ii) follows from the fact that $\mu = 0$ is the only dominant integral weight such that $L_{D_4}(-2, \mu)$ is in the set given in the claim (i). \square

We have:

THEOREM 4.2. *Vertex operator algebra $\tilde{L}_{D_4}(-2, 0)$ is simple, i.e.,*

$$L_{D_4}(-2, 0) = \frac{N_{D_4}(-2, 0)}{U(\hat{\mathfrak{g}}).v + U(\hat{\mathfrak{g}}).\theta(v) + U(\hat{\mathfrak{g}}).\theta^2(v)}.$$

PROOF. Let w be a singular vector for $\hat{\mathfrak{g}}$ in $\tilde{L}_{D_4}(-2, 0)$. The classification result from Proposition 4.1 (ii) implies that $U(\hat{\mathfrak{g}}).w$ is a highest weight $\hat{\mathfrak{g}}$ -module with highest weight $-2\Lambda_0$ and that w is proportional to $\mathbf{1}$. The claim follows. \square

We conclude:

THEOREM 4.3. (i) *The set*

$$\{L_{D_4}(-2, 0), L_{D_4}(-2, -2\omega_1), L_{D_4}(-2, -2\omega_3), L_{D_4}(-2, -2\omega_4), L_{D_4}(-2, -\omega_2)\}$$

provides a complete list of irreducible weak $L_{D_4}(-2, 0)$ -modules from the category \mathcal{O} .

- (ii) *$L_{D_4}(-2, 0)$ is the unique irreducible ordinary module for $L_{D_4}(-2, 0)$.*
 (iii) *Every ordinary $L_{D_4}(-2, 0)$ -module is completely reducible.*

PROOF. Proposition 4.1 and Theorem 4.2 imply claims (i) and (ii).

(iii) Let M be an ordinary $L_{D_4}(-2, 0)$ -module, and let w be a singular vector for $\hat{\mathfrak{g}}$ in M . The classification result from (ii) implies that $U(\hat{\mathfrak{g}}).w$ is a highest weight $\hat{\mathfrak{g}}$ -module with highest weight $-2\Lambda_0$. Claim (ii) also implies that any singular vector in $U(\hat{\mathfrak{g}}).w$ has highest weight $-2\Lambda_0$ and it is proportional to w . Thus, $U(\hat{\mathfrak{g}}).w$ is an irreducible $\hat{\mathfrak{g}}$ -module and the claim follows. \square

ACKNOWLEDGEMENTS.

The author thanks Dražen Adamović for his helpful advice and valuable suggestions.

REFERENCES

- [1] D. Adamović, *Some rational vertex algebras*, Glas. Mat. Ser. III **29(49)** (1994), 25–40.
- [2] D. Adamović, *A construction of some ideals in affine vertex algebras*, Int. J. Math. Math. Sci. **2003** (2003), 971–980.
- [3] D. Adamović, *A construction of admissible $A_1^{(1)}$ -modules of level $-\frac{4}{3}$* , J. Pure Appl. Algebra **196** (2005), 119–134.
- [4] D. Adamović and A. Milas, *Vertex operator algebras associated to modular invariant representations for $A_1^{(1)}$* , Math. Res. Lett. **2** (1995), 563–575.
- [5] D. Adamović and O. Perše, *Some general results on conformal embeddings of affine vertex operator algebras*, Algebr. Represent. Theory **16** (2013), 51–64.
- [6] J. D. Axtell and K.-H. Lee, *Vertex operator algebras associated to type G affine Lie algebras*, J. Algebra **337** (2011), 195–223.
- [7] R. E. Borcherds, *Vertex algebras, Kac-Moody algebras, and the Monster*, Proc. Nat. Acad. Sci. U.S.A. **83** (1986), 3068–3071.
- [8] N. Bourbaki, *Éléments de mathématique*. (French) Fasc. XXXVIII: Groupes et algèbres de Lie. Chapitre VII: Sous-algèbres de Cartan, éléments réguliers. Chapitre VIII: Algèbres de Lie semi-simples déployées. Actualités Scientifiques et Industrielles, No. 1364. Hermann, Paris, 1975.
- [9] C. Dong, H. Li and G. Mason, *Vertex operator algebras associated to admissible representations of $\widehat{\mathfrak{sl}}_2$* , Comm. Math. Phys. **184** (1997), 65–93.
- [10] A. J. Feingold and I. B. Frenkel, *Classical affine algebras*, Adv. in Math. **56** (1985), 117–172.
- [11] E. Frenkel and D. Ben-Zvi, *Vertex algebras and algebraic curves*, American Mathematical Society, Providence, 2001.
- [12] I. Frenkel, Y.-Z. Huang and J. Lepowsky, *On axiomatic approaches to vertex operator algebras and modules*, Mem. Amer. Math. Soc. **104**, 1993.

- [13] I. Frenkel, J. Lepowsky and A. Meurman, *Vertex operator algebras and the Monster*, Academic Press, Boston, 1988.
- [14] I. Frenkel and Y.-C. Zhu, *Vertex operator algebras associated to representations of affine and Virasoro algebras*, *Duke Math. J.* **66** (1992), 123–168.
- [15] V. G. Kac, *Infinite dimensional Lie algebras*, 3rd ed., Cambridge Univ. Press, Cambridge, 1990.
- [16] V. G. Kac, *Vertex algebras for beginners*, AMS, Providence, 1998.
- [17] V. Kac and M. Wakimoto, *Modular invariant representations of infinite dimensional Lie algebras and superalgebras*, *Proc. Nat. Acad. Sci. U.S.A.* **85** (1988), 4956–4960.
- [18] J. Lepowsky and H. Li, *Introduction to vertex operator algebras and their representations*, Birkhäuser, Boston, 2004.
- [19] H.-S. Li, *Local systems of vertex operators, vertex superalgebras and modules*, *J. Pure Appl. Algebra* **109** (1996), 143–195.
- [20] A. Meurman and M. Primc, *Annihilating fields of standard modules of $sl(2, \mathbb{C})$ and combinatorial identities*, *Mem. Amer. Math. Soc.* **137**, AMS, Providence RI, 1999.
- [21] O. Perše, *Vertex operator algebras associated to type B affine Lie algebras on admissible half-integer levels*, *J. Algebra* **307** (2007), 215–248.
- [22] O. Perše, *Vertex operator algebras associated to certain admissible modules for affine Lie algebras of type A*, *Glas. Mat. Ser. III* **43(63)** (2008), 41–57.
- [23] Y.-C. Zhu, *Modular invariance of characters of vertex operator algebras*, *J. Amer. Math. Soc.* **9** (1996), 237–302.

O. Perše

Department of Mathematics

University of Zagreb

10 000 Zagreb

Croatia

E-mail: perse@math.hr

Received: 15.5.2012.