

ISPRAVAK

In the paper Loew SJ, Watson, K. (2013): The prevalence of symptoms of scotopic sensitivity / Meares-Irlen syndrome in subjects diagnosed with ADHD: - Does misdiagnosis play a significant role?, Croatian Review of Rehabilitation Research, 49, Supplement, 64-72., the table with incorrect data is published on the page 68. We apologize to the authors and publish the correct table.

Editorial

Table 2. Mean incidences of SSS symptoms in the ADHD-diagnosed group compared to SSS-diagnosed group means with results of one-way ANOVA.

DIAGNOSTIC SYMPTOM OF SSS:	ADHD group	SSS group	Variation		
			d.f.	F	p
1. Slow reading:	50.00%	72.22%	1	1.51	0.230
2. Strain & fatigue with extended reading:	83.33%	83.33%	1	0.00	1.000
3. Print distortions, esp. black print on white paper:	50.00%	88.89%	1	6.38	0.017*
4. Prefer reading text on coloured or off-white paper:	41.67%	66.67%	1	1.82	0.188
5. Lack of depth perception, difficulty catching a ball:	58.33%	61.11%	1	0.02	0.884
6. Clumsiness when walking on uneven ground:	58.33%	50.00%	1	0.19	0.667
7. Dislike of bright light or glare (photophobia):	41.67%	61.11%	1	1.06	0.312
8. Dislike of reading/writing under fluorescent lighting:	16.67%	61,11%	1	6.70	0.015*
9. Difficulty keeping train of thought in a conversation:	58.33%	55,56%	1	0.02	0.885

d.f. = degrees of freedom; F = variance ratio; p = probability