

Spectral and Thermal Studies of Light Lanthanide Complexes with 4-Methoxy-3-nitrobenzoic Acid

Wiesława Ferenc, Beata Bocian, and Daria Mazur

Department of Inorganic and General Chemistry, Faculty of Chemistry
Maria Curie-Skłodowska University, PL 20-031, Lublin, Poland

Received June 25, 1998; revised October 26, 1998; accepted February 24, 1999

4-Methoxy-3-nitrobenzoates of light lanthanides with a metal to ligand ratio of 1:3 have been obtained as hydrated salts. They have colours typical of Ln^{3+} ions (Pr-green, Nd-pink, Sm-cream, and La, Ce, Eu and Gd-white). The carboxylate group appears to be a bidentate chelating ligand. 4-Methoxy-3-nitrobenzoates of light lanthanides are crystalline compounds. During heating in air, they decompose explosively. Therefore, their thermal stability was studied in the temperature range 273–523 K. The complexes dehydrate in one or three steps. The solubilities of complexes were measured in water at 293 K. They are in the order of 10^{-4} mol dm^{-3} .

Key words: light lanthanides, 4-methoxy-3-nitrobenzoates, rare earth complexes, spectra, synthesis, thermal studies

INTRODUCTION

4-Methoxy-3-nitrobenzoic acid, $\text{C}_8\text{H}_7\text{O}_5\text{N}$, is a cream-yellow, crystalline solid, sparingly soluble in water and readily soluble in hot alcohol and ether.¹ Being crystallized from water, it forms needles. Its melting point is equal to 186–189 °C. The compounds of 4-methoxy-3-nitrobenzoic acid are not well-known. It is known from literature that its complexes have been obtained only with the following cations:¹ Na^+ , K^+ , Ag^+ , Ca^{2+} , Sr^{2+} , Ba^{2+} and Pb^{2+} . 4-Methoxy-3-nitrobenzoates of Na and K are monohydrates, those of Ca and Sr are tetrahydrates while those of Ag, Ba and Pb are anhydrous salts. There is no information in the literature about the complexes of 4-methoxy-3-nitrobenzoic acid with light lanthanide elements. Therefore, we decided to obtain them as solids and to examine some of their physico-

chemical properties, including their thermal characterization during heating in air to 523 K and also their solubility in water at 293 K. The properties of 4-methoxy-3-nitrobenzoates of light lanthanides were compared with those of 3-nitro- and 4-methoxybenzoates of lanthanide elements.²⁻⁴ The comparison allows us to estimate the influence of the position of substituents in aromatic ring on the properties of the complexes. Thermal stability investigations enable to evaluate the assumed position of water molecules in the outer or inner spheres of coordination, to know the mechanism of complex decomposition and to determine the *endo*- or *exo*- effects. Determination of the solubility is valuable because it gives information about the practical use of the acids for the separation of rare earth elements by extraction or ion-exchange chromatography methods.

EXPERIMENTAL

4-Methoxy-3-nitrobenzoates of light lanthanides were prepared by adding equivalent quantities of 0.1 M ammonium 4-methoxy-3-nitrobenzoate (pH ~ 5) to a hot solution containing light lanthanide chlorides. The solids formed were filtered off, washed with hot water to remove ammonium ions and dried at 303 K to constant weight. The carbon, hydrogen and nitrogen contents were determined by elemental analysis using a Perkin Elmer CHN 2400 analyzer. The contents of light lanthanides were analyzed by oxalic acid method (Table I).

In order to confirm the chemical composition of the salts, the IR spectra of the hydrated and dehydrated complexes, the spectra of 4-methoxy-3-nitrobenzoic acid, its sodium salt and sodium nitrite were run in the range 4000–400 cm⁻¹ using a SPECORD M-80 spectrophotometer. Samples were prepared as KBr discs Table II.

TABLE I

Elemental analysis of 4-methoxy-3-nitrobenzoates of light lanthanides

Complex	% H		% C		% N		% Ln	
	Calcd.	Found	Calcd.	Found	Calcd.	Found	Calcd.	Found
LaL ₃ · 6H ₂ O	3.62	3.48	34.50	34.03	5.03	5.04	16.64	16.15
CeL ₃ · 0.5H ₂ O	2.59	2.52	39.08	38.94	5.69	5.60	18.99	18.74
PrL ₃ · H ₂ O	2.69	2.54	38.57	38.54	5.62	5.50	18.85	18.51
NdL ₃ · H ₂ O	2.68	2.53	38.40	38.40	5.59	5.54	19.21	19.06
SmL ₃ · 2H ₂ O	2.86	2.58	37.20	37.20	5.42	5.42	19.41	19.36
EuL ₃ · H ₂ O	2.66	2.52	38.01	38.08	5.54	5.48	20.04	19.63
GdL ₃ · 4H ₂ O	3.19	3.02	35.12	35.39	5.12	5.54	19.23	19.16

L = C₈H₆O₅N

TABLE II

Frequencies of the absorption bands of COO⁻ and NO₂ group vibrations for 4-methoxy-3-nitrobenzoates of light lanthanides and sodium, and of CO and NO₂ for 4-methoxy-3-nitrobenzoic acid (ν / cm⁻¹)

Complex	$\nu(\text{C}=\text{O})$	$\nu_{\text{as}}(\text{OCO})$	$\nu_{\text{s}}(\text{OCO})$	$\Delta\nu(\text{OCO})$	$\nu_{\text{as}}(\text{NO}_2)$	$\nu_{\text{s}}(\text{NO}_2)$	$\Delta\nu(\text{NO}_2)$	$\nu(\text{M}-\text{O})$
LaL ₃ · 6H ₂ O	–	1595	1415	180	1545	1365	180	525
CeL ₃ · 0.5H ₂ O	–	1590	1410	180	1540	1365	175	535
PrL ₃ · H ₂ O	–	1590	1410	180	1540	1370	170	540
NdL ₃ · H ₂ O	–	1590	1410	180	1540	1365	175	540
SmL ₃ · 2H ₂ O	–	1600	1420	180	1540	1365	175	545
EuL ₃ · H ₂ O	–	1595	1415	180	1540	1370	170	545
GdL ₃ · 4H ₂ O	–	1595	1420	175	1545	1365	180	550
NaL	–	1610	1425	185	1545	1365	180	520
HL	1700	–	–	–	1540	1360	180	–

L = C₈H₆O₅N

X-ray diffraction patterns of the complexes were measured on a DRON-2 diffractometer using CuK_α radiation. The measurements were performed within the range $2\theta = 5\text{--}80^\circ$ according to the Deby-Scherrer-Hull method.

TABLE III

Temperature range of the dehydration process of 4-methoxy-3-nitrobenzoates of light lanthanides in air atmosphere and their solubilities in water at 293 K

Complex	ΔT / K	Loss of mass / %		<i>n</i>	Dehydration product	<i>E</i> _a / kJ mol ⁻¹	Solubility / mol dm ⁻³
		Calcd.	Found				
LaL ₃ · 6H ₂ O	320–405	12.94	12.71	6	LaL ₃	40.0	2.710 ⁻⁴
CeL ₃ · 0.5H ₂ O	410–450	1.22	1.35	0.5	CeL ₃	83.6	3.110 ⁻⁴
PrL ₃ · H ₂ O	420–450	2.41	2.25	1	PrL ₃	71.5	3.510 ⁻⁴
NdL ₃ · H ₂ O	420–458	2.40	2.10	1	NdL ₃	98.4	4.210 ⁻⁴
SmL ₃ · 2H ₂ O (I stage)	348–383	1.66	1.20	0.5	SmL ₃ · 1.5H ₂ O	37.5	4.810 ⁻⁴
SmL ₃ · 1.5H ₂ O (II stage)	410–450	3.48	3.33	1.0	SmL ₃ · 0.5H ₂ O	88.9	
SmL ₃ · 0.5H ₂ O (III stage)	454–510	4.65	4.53	0.5	SmL ₃	63.4	
EuL ₃ · H ₂ O	410–448	2.37	2.52	1	EuL ₃	89.8	5.210 ⁻⁴
GdL ₃ · 4H ₂ O	355–425	8.78	9.04	4	GdL ₃	54.4	5.510 ⁻⁴

L = C₈H₆O₅N

ΔT = temperature range of dehydration process,

n = number of crystallization water molecules being lost in one or three endothermic steps,

*E*_a = energy activation of dehydration reaction

The thermal decomposition of the complexes in air in the range 273–523 K was determined with the Paulik-Paulik-Erday Q-1500D derivatograph with Derill converter of the MOM Budapest Firm, recording TG, DTG and DTA curves. The measurements were made at a heating rate of 10 K min⁻¹. The 100 mg samples were heated in platinum crucibles in static air to 523 K with a sensitivity TG = 50 mg. DTG and DTA sensitivities were regulated by the computer Derill programme worked out by MOM the Budapest Firm (Table III, Figure 1). The products of dehydration were calculated from TG curves and verified by the diffraction pattern registration and IR spectra.

The solubilities of complexes in water at 293 K were determined by measuring the concentration of Ln³⁺ ions in a saturated solution by the oxalic acid method (Table III).

RESULTS AND DISCUSSION

4-Methoxy-3-nitrobenzoates of light lanthanides were obtained as crystalline products of the general formula: Ln(C₈H₆O₅N)₃ · nH₂O (where n = 0.5; 1; 2; 4 and 6) (Table I). 4-Methoxy-3-nitrobenzoates of Pr, Nd and Eu are monohydrates, those of Sm, Gd and La are dihydrate, tetrahydrate and hexahydrate, respectively. Only the complex of cerium 4-methoxy-3-nitrobenzoate forms hemihydrate.




Figure 1. TG, DTG and DTA curves of praseodymium 4-methoxy-3-nitrobenzoate.

3-Nitrobenzoates of Y, Pr-Gd contain two water molecules while those of La and Ce two and a half.² The 4-methoxybenzoates of light lanthanides form hydrates (La, Ce) or anhydrous salts (Pr-Gd).^{3,4}

The colours of 4-methoxy-, 3-nitro- and 4-methoxy-3-nitrobenzoates of light lanthanides are typical of Ln^{3+} ions²⁻⁴ (it means: Pr-green, Nd-pink, Sm-cream and La, Ce, Eu and Gd-white), which is connected with the similar electron density in the complexes. Probably, in these molecules, the $f \rightarrow f$ electronic transitions of the central ions are those of the lowest energy and absorption occurs at relatively high wave-lengths, depending on the nature, of the metal ion.⁵⁻¹¹ 4-Methoxy-3-nitrobenzoates of light lanthanides show similar IR spectra. However, the characteristic frequencies related to the carbonyl group are altered markedly in going from acid to salts. Table II presents the frequencies of typical absorption bands arising from asymmetric and symmetric vibrations of COO^- and $\text{C}=\text{O}$ groups and metal-oxygen bond for 4-methoxy-3-nitrobenzoates of light lanthanides and sodium, and of CO group for 4-methoxy-3-nitrobenzoic acid.¹²⁻¹⁷ In the IR spectra of light lanthanide 4-methoxy-3-nitrobenzoates, the band at 1700 cm^{-1} completely disappears, which confirms that no COOH group is present in the complexes. In these spectra, the intense broad absorption bands at $3580\text{--}3110 \text{ cm}^{-1}$ and strong bands at $1630\text{--}1620 \text{ cm}^{-1}$ confirm the presence of crystallization water. The bands resulting from asymmetric and symmetric vibrations of $-\text{O}-\text{CH}_3$ group occur at $2960\text{--}2955 \text{ cm}^{-1}$ and $2870\text{--}2860 \text{ cm}^{-1}$, respectively.¹²⁻¹⁷ The bands due to asymmetric and symmetric NO_2 group vibrations are not significantly shifted with respect to those of the parent acid, thus one may conclude that the group does not coordinate with metal ions, similarly as in the case of 3-nitrobenzoates of light lanthanide elements.² In 4-methoxy-3-nitrobenzoates, the bands confirming M-O vibrations are being shifted to higher frequencies with increased atomic number of the element. It may suggest that 4-methoxy-3-nitrobenzoic acid forms with light lanthanides complexes of increasingly greater stability. For 3-nitrobenzoates of rare earth elements, the bands of M-O appear in the range $425\text{--}410 \text{ cm}^{-1}$.² The frequencies are lower compared to those of 4-methoxy-3-nitrobenzoates of rare earth elements (Table II). This difference is connected with the presence of $-\text{O}-\text{CH}_3$ group in *para* position and with the stronger influence of inductive and mesomeric effects of nitro group on electron density in 4-methoxy-3-nitrobenzoate molecules than in 3-nitrobenzoates. In 4-methoxy-3-nitrobenzoates of light lanthanides, the M-O bands are stronger than in the case of 3-nitrobenzoates. The bands of skeleton and of $-\text{O}-\text{CH}_3$ group vibrations in the IR spectra of light lanthanide 4-methoxy-3-nitrobenzoates are shifted to lower frequencies compared to those in the parent acid. The changes in their position are probably caused by the interaction between aromatic ring vibrations and $-\text{O}-\text{CH}_3$ group with groups of atoms or ions.

In the IR spectra of 4-methoxy-3-nitrobenzoates, two bands due to asymmetric, ν_{as} , and symmetric, ν_s , vibrations of COO^- group are shifted to lower frequencies than those for sodium salt. Accordingly, it is possible to assume that the carboxylate ion is a bidentate, chelating ligand.^{12-14, 18} In 3-nitro- and 4-methoxybenzoates of rare earth elements, the carboxylate ion is also a bidentate, chelating ligand.²⁻⁴ The magnitude of separation, $\Delta\nu_{\text{OCO}^-}$, between the frequencies due to $\nu_{as\text{OCO}^-}$ and $\nu_{s\text{OCO}^-}$ in the light lanthanide 4-methoxy-3-nitrobenzoates indicates a lower degree of ionic bond in these complexes compared to the sodium salt.

In order to verify whether the obtained light lanthanide 4-methoxy-3-nitrobenzoates convert to the nitrito isomers (during precipitation or on heating) the IR spectra for hydrates of 4-methoxy-3-nitrobenzoates of light lanthanides, of dehydrated complexes and of sodium nitrite were recorded. A survey of the literature shows that, under influence of sunlight or, with an increase of temperature, the transformation process of nitro complexes to nitrito ones may occur.^{19,20} It was found that isomerization of nitro group may accompany the dehydration process of the complex or it may be connected with the transformation process in its inner sphere.²⁰ As a monodentate the nitrite ion may bond either through a nitrogen atom forming the nitro complex or through one of the oxygen atoms yielding a nitrito one. The general effect of coordination via nitrogen is to rise the frequencies of both $\nu_{as(\text{NO}_2)}$ and $\nu_{s(\text{NO}_2)}$ from the free ion values $\sim 1328 \text{ cm}^{-1}$ and 1260 cm^{-1} to higher values.²¹ The IR spectra of 4-methoxy-3-nitrobenzoates of light lanthanides isolated as solids show these frequencies at $1545\text{--}1540 \text{ cm}^{-1}$ and $1370\text{--}1365 \text{ cm}^{-1}$, respectively (Table II).

The IR spectra recorded also for 3-nitrobenzoates of rare earth elements show that no transformation process of nitro group to nitrito takes place with rising temperature.^{2,20,21}

From the X-ray powder diffraction of 4-methoxy-3-nitrobenzoates of light lanthanides it follows that the complexes are polycrystalline compounds. However, their structures have not been determined since their single crystals have not been obtained.

On heating in air, 4-methoxy-3-nitro- and 3-nitrobenzoates of light lanthanides decompose explosively above 523 K .² Their instability is caused by the presence of the NO_2 group in benzene ring.^{2,9} Contrary to them, 4-methoxybenzoates of rare earth elements are not decomposed in this way.^{3,4} The thermal stability of 4-methoxy-3-nitrobenzoates of light lanthanides was studied in the range $273\text{--}523 \text{ K}$. The results obtained from their thermal decomposition reveal them to be hydrates containing various molecules of crystallization water, as it was established on the basis of elemental analysis (Tables I, III). All complexes are stable in air at room temperature and do not

change their mass after storage. When heated at 273–523 K, they dehydrate in various ways. 4-Methoxy-3-nitrobenzoates of La-Nd, Eu and Gd lose crystallization water molecules in one step (320–458 K) and form anhydrous complexes. The dehydration process is connected with an endothermic effect shown in DTA curves.^{22,23} The values change according to the appropriate amounts of water molecules in these complexes. The complex of samarium is decomposed in three steps. At first, it loses only half a molecule of water and then gradually one and finally a half to form the anhydrous complex. In the case of the decomposition of praseodymium 4-methoxy-3-nitrobenzoate (Figure 1), the weight loss starting at 420 K corresponds to the beginning of the dehydration process that takes place in the range 420–450 K. The strong DTG peak connected with weight loss is observed and endothermic effect also appears in the DTA curve. Next, above 450 K, the TG curve gains a constant level, which is indicative of the anhydrous complex formation. Considering the temperatures at which the dehydration process takes place and the way in which it proceeds, it is possible to assume that the molecules of crystallization water are of the outer-sphere water. According to Nikolaev *et al.*²⁴ and Singh *et al.*²⁵, water eliminated below 423 K can be considered as crystallization water whereas the water eliminated above 423 K may be that coordinated to the central ion. In our investigations, water is assumed to be crystallization water (with the exception of hemihydrate of samarium complex). This was also confirmed by IR studies.

From TG and DTA curves, the activation energies of dehydration reactions were calculated by the Fatieev and Pletniev method²⁶ using equation:

$$E_a = \frac{RT_{\max}^2}{m_0} \cdot \frac{\Delta m}{\Delta T}$$

where: R – gas constant; T_{\max} – temperature of maximum weight loss; m_0 – weight loss at T_{\max} ; m – weight loss at T , T – given temperature, $\Delta m = m_0 - m$, $\Delta T = T_{\max} - T$.

The different values of activation energy may suggest that molecules of outer-sphere water are coordinated by different forces, depending on their position. The smallest value for samarium 4-methoxy-3-nitrobenzoate indicates that the water molecules are the most weakly coordinated whereas the highest value for neodymium complex shows that the water molecule is presumably the most strongly bound in complex coordination. From preliminary investigations on thermal properties of the complexes during their heating to 1173 K, it appears that they are ultimately decomposed to the oxides of respective metals. Oxides of lanthanides were identified by X-ray powder diffraction^{27,28} and the measurements revealed them to be for La, Nd-Gd and CeO₂ and Pr₆O₁₁. In the case of 3-nitrobenzoates of rare earth

elements, the initial dehydration temperature values are lower than those for 4-methoxy-3-nitrobenzoates.² Hydrates of 4-methoxy-3-nitrobenzoates are more stable than those of 3-nitrobenzoates, which is connected with the presence of methoxy- group in 3-nitrobenzoate ring that causes a decrease of electron density in aromatic ring.

The solubilities of 4-methoxy-3-nitrobenzoates of light lanthanides in water at 293 K were measured (Table III) and they are in the order of 10^{-4} mol dm⁻³. The values of solubility increase with increasing atomic number of elements. The solubility order of light lanthanide 4-methoxy-3-nitrobenzoates is less than that determined for 4-methoxy- and 3-nitrobenzoates of those elements (10^{-3} mol dm⁻³). The difference in the solubility values of the complexes is connected with the different electron densities established by the influence of the inductive and mesomeric effects in the molecules of the compounds, causing their different degrees of dissociation in water. The presence of $-O-CH_3$ group in 3-nitrobenzoate ring causes a decrease of solubility compared to 3-nitrobenzoates and also the presence of NO_2 group in 4-methoxybenzoate ring decreases the solubilities, compared to 4-methoxybenzoates. Taking into account the values of 4-methoxy-3-nitrobenzoate solubilities it is possible to state that 4-methoxy-3-nitrobenzoic acid cannot be used for the separation of rare earth elements by ion-exchange chromatography or extraction methods because it forms with rare earth elements complexes that are not easily soluble.

REFERENCES

1. Beilsteins Handbuch der organischen Chemie, Bd. 10, Springer-Verlag, Berlin, 1927, pp. 181–182
2. W. Ferenc, *Mh. Chem.* **119** (1988) 1345–1355.
3. W. Brzyska and W. Ferenc, *Zesz. Nauk. Politech. Śląsk.* **113** (1985) 5–14.
4. W. Brzyska and W. Blaszczyk, *Ann. Univ. Mariae Curie-Skłodowska*, Sect. AA, Vol. 34, **6** (1979) 45–48.
5. P. Karrer, *Chemia organiczna*, **T.2**, PWN, Warszawa, 1985, pp. 98–102.
6. R. Q. Brewster, E. William and Mc Ewen, *Podstawy chemii organicznej*, PWN, Warszawa, 1968, pp. 507–509.
7. H. A Staab., *Einführung in die theoretische organische Chemie*, Verlag Chemie, Weinheim, 1962, pp. 343–364.
8. C. N. R. Rao, *Ultra-Violet and Visible Spectroscopy. Chemical Applications*, Butterworths, London, 1975, pp. 76–96.
9. B. I. Stiepanow, *Podstawy chemii technologii barwników organicznych*, Warszawa, PWN, 1980, pp. 157–165.
10. A. Bartecki, *Spektroskopia elektronowa związków nieorganicznych i kompleksowych*, PWN, Warszawa, 1971, pp. 60–65.
11. J. Shorter, *Correlation Analysis in Organic Chemistry. An Introduction to Linear Free-Energy Relationships*, Clarendon Press, Oxford, 1973, pp. 27–33.

12. K. Nakamoto, *Infrared and Raman Spectra of Inorganic and Coordination Compounds*, John Wiley and Sons, Toronto, 1968, pp. 191–268.
13. K. Nakamoto K. and P. J. McCarthy, *Spectroscopy and Structure of Metal Chelate Compounds*, John Wiley and Sons, New York, 1968, pp. 217–285.
14. R. C. Mehrotra and R. Bohra, *Metal Carboxylates*, Academic Press, London, 1983, pp. 48–89.
15. L. J. Bellamy, *The Infrared Spectra of Complex Molecules*, Chapman and Hall, London, 1975, pp. 72–231.
16. K. Burger, *Coordination Chemistry: Experimental Methods*, Akadémiai Kiadó, Budapest, 1973, pp. 53–130.
17. A. D. Cross and A. R. Jones, *An Introduction to Practical Infrared Spectroscopy*, Butterworths, London, 1969, pp. 43–100.
18. B. S. Manhas and A. K. Trikha, *J. Indian Chem. Soc.* **59** (1982) 315–316.
19. I. N. Kukuszkin, *Koord. Chim.*, **4** (1972) 1170–1173.
20. D. L. Goodgame and M. A. Hitchman, *Inorg. Chem.* **6** (1987) 813–819.
21. W. Fee, C. S. Gamer and I. N. Harrowfield, *Inorg. Chem.* **6** (1987) 1–5.
22. D. N. Todor, *Thermal Analysis of Minerals*, Abacus Press, Tunbridge Wells, Kent 1976, pp. 55–96.
23. F. Paulik, *Special Trends in Thermal Analysis*, John Wiley and Sons, Chichester, 1995, pp. 76–173.
24. A. V. Nikolaev, V. A. Logvienko and L. I. Myachina, *Thermal Analysis*, **Vol 2**, Academic Press, New York, 1989, pp. 72–80.
25. B. Singh, B. V. Agarwala, P. L. Mourya and A. K. Dey, *J. Indian Chem. Soc.* **59** (1992) 1130–1134.
26. W. Brzyska, *Wstęp do chemii koordynacyjnej*, UMCS, Lublin 1996, pp. 70–71.
27. M. Von Meersche and J. Feneau-Dupont, *Introduction à la cristallographie et à la chimie structurale*, OYEZ, Bruxelles, 1976, pp. 394–624.
28. P. Pascal, *Nouveau traité de chimie minérale*, Masan et c^{ie}, Paris, 1959, pp. 777–831.

SAŽETAK

Spektralna i termalna studija kompleksa lakih lantanoida s 4-metoksi-3-nitrobenzojevom kiselinom

Wiesława Ferenc, Beata Bocian i Daria Mazur

4-Metoksi-3-nitrobenzoati lakih lantanoida dobiveni su kao hidratne soli s omjerom metal : ligand = 1 : 3. Boje su im tipične za ione Ln³⁺ (Pr: zelen, Na: ružičast, Sm: žućkasto-bijel te La, Ce, Eu i Gd: bijeli). Karboksilna skupina javlja se kao bidentatan kelatni ligand. 4-Metoksi-3-nitrobenzoati lakih lantanoida kristalni su spojevi. Tijekom zagrijavanja na zraku, eksplozivno se raspadaju, pa je zato proučavana njihova toplinska stabilnost u rasponu od 273 K do 523 K. Kompleksi se dehidratiziraju u jednomu ili tri stupnja. Topljivosti kompleksa izmjerene u vodi pri 293 K, iznose oko 10⁻⁴ mol dm⁻³.