

“Double fire” — rijetka, a još češće neprepoznata aritmija

“Double fire” — a rare and commonly unrecognized arrhythmia

Zoran Baković*, Ante Anić, Marin Bištić, Albino Jović

Opća bolnica Zadar, Zadar, Hrvatska
Zadar General Hospital, Zadar, Croatia

SAŽETAK: Dvojna fiziologija provođenja atrioventrikularnim (AV) čvorom, odnosno prisutnost tzv. sporog puta, prema raznim studijama, elektrofiziološkim ispitivanjem se može dokazati u do čak 35% ljudi. Međutim, kod samo manjeg broja on ima i klinički značaj.

Prikazujemo slučaj pacijentice s vrlo rijetkom elektrofiziološkom manifestacijom aktivnog sporog puta, dvostrukog odgovora ventrikula na jedan atrijski kompleks. Problem je uspješno rješen radiofrekventnom ablacijom. Ovakav način provođenja AV čvorom vjerovatno je i znatno češći nego što se opisuje u literaturi, ali se na žalost rijetko prepozna te je uglavnom refraktoran na medikamentoznu terapiju.

KLJUČNE RIJEČI: dualna fiziologija atrioventrikularnog čvora, “double fire” fenomen, radiofrekventna ablacija.

SUMMARY: Dual atrioventricular node (AV) pathway physiology or the presence of so-called slow conduction pathway is according to various studies demonstrable in up to 35% of normal people during electrophysiology study. In only a small number of them, it has a clinical significance.

We present a case of a patient with a very rare electrophysiological manifestation of active slow pathway, double ventricular response to one atrial complex. The problem was successfully treated with radiofrequency ablation.

This form of conduction via AV node is probably much more common than it was previously described in the literature, but unfortunately it is rarely recognized and is generally refractory to medical therapy.

KEYWORDS: dual atrioventricular node physiology, “double fire” phenomenon, radiofrequency ablation.

CITATION: Cardiol Croat. 2014;9(1-2):44-47.

Prikaz slučaja

Prikazujemo slučaj 62 godišnje pacijentice naručene na invazivnu kardiološku obradu zbog učestalih palpitacija te intolerancije napora praćene opresijama u prekordiju. Od ranije se liječi zbog arterijske hipertenzije i hiperlipidemije, a zbog palpitacija u terapiji ima i propafenon 2x150 mg te verapamil 180 mg.

Već na rutinskom 12-kanalnom elektrokardiogramu kod prijema prepoznat je potencijalni uzrok tegoba. Naime, prisutni su brojni ventrikularni kompleksi koji po morfološkoj odgovaraju supraventrikularnim ekstrasistolama (SVES), bez vidljive atrijske aktivnosti, a u uskoj vezi s prethodno uredno provedenim sinusnim kompleksom (**Slika 1**). U 24-satnom kontinuiranom snimanju EKG zabilježeno je više od 10.000 takvih SVES. Ehokardiografski se radi o strukturno zdravom srcu očuvane sistoličke funkcije, a koronarografijom je isključena okluzivna bolest epikardijalnih arterija.

Case study

We present a case of a 62 year-old woman admitted for further evaluation because of frequent palpitations and effort intolerance accompanied by precordial oppressions. She has been treated for arterial hypertension and hyperlipidemia since earlier, and takes propafenone 2x150 mg, and verapamil 180 mg in her therapy for palpitations.

A potential cause of problems was recognized already on a routine 12-lead electrocardiogram at the time of admission. In fact, there are numerous ventricular complexes that according to their morphology correspond to supraventricular extrasystoles (SVES), with no visible atrial activity, and are closely related to the previously duly conducted sinus complex (**Figure 1**). More than 10,000 such SVES were recorded in the 24-hour continuous ECG recording. Echocardiography showed that it is a structurally normal heart with preserved systolic function, and coronary angiography excluded coronary artery disease.




Figure 1. Sinus rhythm with lot of narrow QRS extra beats, without preceding P wave. It is difficult to differentiate whether it is ectopic activity from the AV junction (His region) or dual conduction of one P wave through both fast and slow pathway to ventricle.

Odlučili smo se za elektrofiziološko ispitivanje (EPS). Bazičnim intrakardijalnim elektrogramima potvrđena je sumnja da se radi o tzv. "double fire" fenomenu, odnosno dvostrukom atrioventrikularno (AV) provođenju. Na jedan atrijski kompleks dolaze dva ventrikularna od kojih je prvi proveden brzim, a drugi sporim putem (**Slika 2a i 2b**). Standardnim EPS protokolom dokazana je dualna fiziologija AV čvora, isključeno je retrogradno provođenje, a nije inducirana tachikardija niti ev. "echo" udari. U regiju sporog puta postavljen je ablacijski kateter te se nakon nekoliko kraćih aplikacija energije u potpunosti eliminira dvojna AV fiziologija s isključivim provođenjem kroz brzi put (**Slika 3a i 3b**). Aritmija se ne javlja niti nakon primjene izoproterenola. Otpuštena je kući bez antiaritmika.

Electrophysiology study (EPS) was performed. Basic intracardiac elektograms confirmed the suspicion that this is the so-called "double fire" phenomenon, or dual atrioventricular (AV) conduction. Two ventricular complexes come to one atrial complex, of which the first is conducted by the fast and the second by the slow pathway (**Figure 2a and 2b**). Standard EPS protocol proved dual AV node pathway physiology, excluded retrograde conduction, and induced no tachycardia or "echo" beats. After a few short energy applications in the region of the slow pathway, dual AV node physiology was completely eliminated, only with conduction via the fast pathway (**Figure 3a and 3b**). Arrhythmia was not inducible even after the application of isoproterenol. She was discharged home without antiarrhythmics.




Figure 2a.
Intracardiac recording during electrophysiology study.



Figure 2b. Schematic presentation of pulse propagation from atrium to ventricle: First atrial (sinus) impulse is conducted through both fast and slow pathway to the region of His, and down to the ventricle (1:2 conduction). Second atrial stimulus is blocked in the atrioventricular node which is still refractory from the previous depolarization by slow pathway. Third atrial impulse is conducted only by fast pathway. (HRAd — high right atrium; HISd — region of His with its potential; RVd — apex of the right ventricle).




Figure 3a.

ECG after successful ablation in the slow pathway region — intracardiac recording.

Zaključak

Dvojna fiziologija provođenja AV čvorom, odnosno prisutnost tzv. sporog puta, prema raznim studijama, elektrofiziolškim ispitivanjem se može dokazati u do čak 35% ljudi^{1,2}.

Fenomen dvostrukog ventrikularnog odgovora na jedan atrijski kompleks prvi je opisao Csapo 1979. godine i nazvao

Conclusion

Dual AV node pathway physiology or the presence of so-called slow conduction pathway is according to various studies present in up to 35% of people during EPS^{1,2}.

The phenomenon of dual ventricular response to a single atrial complex was first described by Csapo in 1979 and cal-



Figure 3b. ECG after successful ablation in the slow pathway region — standard 12-lead ECG.

ga je "double fire" ili "non-reentrant" tahikardija³. Rijedak je jer zahtjeva posebne karakteristike oba puta — anterogradnu provodljivost i retrogradni blok, a spori put mora biti dovoljno spor da dopusti His-Purkinjeovom tkivu da oporavi podražljivost nakon prethodne stimulacije⁴⁻⁶. Zbog toga fenomen provodljivosti 1:2 nije konstantan, AV čvor se neujednačeno depolarizira, pa su često prisutni i različiti oblici funkcionalnog AV bloka (PR prolongacija ili Wenckebach). Na vanjskom elektrokardiogramu to se manifestira nepravilnim ventrikularnim ritmom, a odnos s P valom je ponekad teško pratiti. Zato i nije neobično da se ova aritmija teško prepozna, a kod brže frekvencije i niže volatžne P vala lako zamjeni za npr. fibrilaciju atrija. Dominantan simptom su palpitacije, a u literaturi su opisani slučajevi tahikardiomiopatiye uzrokovani ovom aritmijom, uspješno rješeni ablacijskom sporog puta⁷⁻⁹. Ovi pacijenti u pravilu nemaju kružne tahikardije koje su tipične za aktivni spori put, kao što je atrioventrikularna nodalna kružna tahikardija.

Loš odgovor na medikamentoznu terapiju (zbog rezistencije sporog puta na većinu klasičnih antiaritmika) i mogućnost izlječenja radiofrekventnom ablacijskom, nameće potrebu da razmišljamo o ovom tipu aritmije koja se može prepoznati već iz standardnog 12-kanalnog elektrokardiograma. Nаравно, za dokaz je ipak potrebna elektrofiziološka studija.

Received: 12th Jan 2014; Updated 19th Jan 2014; Accepted 30th Jan 2014

*Address for correspondence: Opća bolnica Zadar, Bože Perićića 5, HR-23000 Zadar, Croatia.

Phone: +385-23-505-505

E-mail: zbakotic@gmail.com

Literature

- Lee KW, Badhwar N, Scheinman MM. Supraventricular Tachycardia - Part I. Curr Probl Cardiol. 2008;33:467-546.
- Brooks R, Goldberger J, Kadish A. Extended protocol for demonstration of dual AV nodal physiology. Pacing Clin Electrophysiol. 1993;16(2):277-84.
- Csapo G. Paroxysmal nonreentrant tachycardias due to simultaneous conduction in dual atrioventricular nodal pathways. Am J Cardiol. 1979;43(5):1033-45.
- Francis J, Krishnan M. Dual ventricular response or 1:2 atrioventricular conduction in dual atrioventricular nodal physiology. Indian Pacing Electrophysiol J. 2008;8(2):77-9.
- Germano JJ, Essebag V, Papageorgiou P, Josephson ME. Concealed and manifest 1:2 tachycardia and atrioventricular nodal reentrant tachycardia: manifestations of dual atrioventricular nodal physiology. Heart Rhythm. 2005;2:536-9.
- Fraticelli A, Saccomanno G, Pappone C, Oreto G. Paroxysmal supraventricular tachycardia caused by 1:2 atrioventricular conduction in the presence of dual atrioventricular nodal pathways. J Electrocardiol. 1999;32:347-54.
- Josephson ME. Tachycardia-mediated cardiomyopathy. Card Electrophysiol Clin. 2010; 2:191-6.
- Clementy N, Casset-Senon D, Giraudeau C, Cosnay P. Tachycardiomyopathy secondary to nonreentrant atrioventricular nodal tachycardia: recovery after slow pathway ablation. Pacing Clin Electrophysiol. 2007;30:925-8.
- Anselm F, Frederiks J, Boyle NG, Papageorgiou P, Josephson ME. An unusual cause of tachycardia-induced myopathy. Pacing Clin Electrophysiol. 1996;19:115-9.