

Synthesis and potential antimycotic activity of 4-substituted-3-(thiophene-2-yl-methyl)- Δ^2 -1,2,4-triazoline-5-thiones

MONIKA WUJEC¹
MONIKA PITUCHA¹
MARIA DOBOSZ^{1*}
URSZULA KOSIKOWSKA²
ANNA MALM²

¹ Department of Organic Chemistry
Faculty of Pharmacy, Medical University
Lublin, Poland

² Department of Pharmaceutical
Microbiology, Faculty of Pharmacy
Medical University, Lublin, Poland

In the reaction of hydrazide of thiophene-2-acetic acid (1) with isothiocyanates, the respective thiosemicarbazides **2a-g** were obtained. Further cyclization with 2% NaOH led to formation of 4-substituted-3-(thiophene-2-yl-methyl)- Δ^2 -1,2,4-triazoline-5-thiones (**3a-g**). These compounds showed promising antimycotic activity.

Keywords: 4-substituted-3-(thiophene-2-yl-methyl)- Δ^2 -1,2,4-triazoline-5-thiones, synthesis, spectral analysis, antimycotic activity

Received March 12, 2004

Accepted September 3, 2004

1,2,4-Triazole system is a structural element of many drugs that have antimycotic activity such as fluconazol, itraconazol, voriconazol (1, 2). One of the methods of preparing derivatives of 1,2,4-triazole is the cyclization reaction of acyl derivatives of thiosemicarbazide in alkaline medium (3–13). Using the cyclization reactions of thiosemicarbazide derivatives of 1,2,4-triazole-1-acetic acid and 1,2,4-triazole-4-acetic acid compounds composed of two 1,2,4-triazole systems connected by a methylene group were prepared (3, 4).

As a continuation of investigations of compounds of a similar structure, a series of new 1,2,4-triazoline-5-thione derivatives with potential pharmacological activity on the central nervous system were prepared (14). When investigated on mice, these compounds did not produce neurotoxic activity, but acted on the central nervous system calmly, and prolonged the hexobarbital sleeping time significantly and depressed the locomotor activity (14). Derivatives of 1,2,4-triazoline-5-thione show different biological activities: analgesis (15), antiinflammatory (16), bacteriostatic (17, 18), antifungal (19).

* Correspondence, e-mail: wujec@panaceum.am.lublin.pl

This work is a continuation of our search for 1,2,4-triazole derivatives with antimycotic and antibacterial activity.

EXPERIMENTAL

Melting points were determined in Fisher-Johns blocs (Sanyo, Japan) and are presented without corrections. IR spectra were recorded in KBr using a Specord IR-75 spectrophotometer (Perkin Elmer, UK). The ^1H NMR spectra were recorded on a Bruker Avance 300 (Bruker, Germany) in $\text{DMSO-}d_6$ with TMS as internal standard.

Chemicals were purchased from Merck (Germany) or Fluka (Switzerland) and were used without purification. Purity of all compounds was checked by TLC on aluminium oxide 60 F_{254} plates (Merck), in a $\text{CHCl}_3/\text{C}_2\text{H}_5\text{OH}$ (10:2) solvent system with UV visualization ($\lambda = 254 \text{ nm}$).

Syntheses

Thiophene-2-acetic acid hydrazide (1). – Ethyl thiophene-2-acetate (0.01 mol, 1.7 g), dry ethanol (5 mL) and hydrazine hydrate (0.02 mol) were boiled for 3 h under reflux. After cooling, the precipitate was filtered off, dried and recrystallized from 95% ethanol. m.p. 89–90 °C; Yield 1.28 g (89%). Analysis for $\text{C}_5\text{H}_8\text{N}_2\text{OS}$ (M_r 144.31): calcd. C 41.65%, H 5.59%, N 19.43%; found: C 41.52%, H 5.09%, N 19.37%. Spectral data were as follows: IR (ν_{max}): 3240 (NH), 3047 (CH arom.), 1427 (CH aliph.), 1678 (C=O) cm^{-1} .

^1H NMR (δ ppm): 3.57 (s, 2H, CH_2), 4.28 (s, 2H, NH_2), 6.89–6.95 (m, 2H, 2CH), 7.32–7.34 (m, 1H, CH), 9.23 (s, 1H, NH).

Thiosemicarbazide derivatives of thiophene-2-acetic acids (2a-g). – Hydrazide **1** (0.01 mol) and isothiocyanates (0.01 mol) were mixed carefully and then placed in a round-bottom flask, and heated on oil bath at 90 °C for 10 h. The reaction product was washed with diethyl ether in order to remove the unreacted isothiocyanate and then with water to remove unreacted hydrazide. Next, the product was filtered, dried and crystallized from 95% ethanol.



4-Substituted-3-(thiophene-2-yl-methyl)- Δ^2 -1,2,4-triazoline-5-thiones (3a-g). – Thiosemicarbazide **2a-g** (0.01 mol) and sodium hydroxide (2%, 40 mL) were boiled for 2 h. After cooling, the solution was neutralized with diluted hydrochloric acid. The precipitate was filtered off and then crystallized from 95% ethanol.

The data relating to compounds **2a-g** and **3a-g** are listed in Table I.

Microbiological tests

Compounds **2a**, **2c-e**, **3a** and **3c-e** were assayed for their antimicrobial activity *in vitro* against 11 strains of aerobic bacteria – 3 strains of Gram-positive bacteria (*Staphylococcus aureus*, *Staphylococcus epidermidis*, *Enterococcus faecalis*), 8 strains of Gram-negative bacteria (*Escherichia coli*, *Klebsiella pneumoniae*, *Proteus vulgaris*, *Proteus mirabilis*, *Salmonella enteritidis*, *Salmonella agona*, *Yersinia enterocolitica*, *Pseudomonas aeruginosa*) and 12

Table I. Physical and analytical data for compounds 2a-g and 3a-g



Compd. No.	R	Molecular formula (M_r)	Yield (%) M.p. (°C)	Elemental analysis Calc./found (%)			IR (KBr, cm^{-1})	^1H NMR (DMSO- d_6 , δ ppm)
				C	H	N		
2a	C_2H_5	$\text{C}_9\text{H}_{13}\text{N}_3\text{OS}_2$ (243.34)	80 138–140	44.41 44.52	5.38 5.43	17.26 17.30	3312 (NH) 3142 (CH arom.) 2971, 1432 (CH aliph.) 1695 (C=O) 1375 (C=S)	1.03–1.08 (t, 3H, CH_3) $J = 7.1$ Hz 3.41–3.48 (q, 2H, CH_2) $J = 6.9$ Hz 3.68 (s, 2H, CH_2) 6.88–6.97 (m, 2H, 2CH) 7.36–7.38 (m, 1H, CH) 7.93, 9.24, 9.92 (3s, 3H, 3NH)
2b	$\text{CH}_2\text{C}_6\text{H}_5$	$\text{C}_{14}\text{H}_{15}\text{N}_3\text{OS}_2$ (305.41)	85 154–156	55.04 54.96	4.95 5.02	13.75 13.67	3285 (NH) 3128 (CH arom.) 2931, 1423 (CH aliph.) 1690 (C=O) 1342 (C=S)	3.72 (s, 2H, CH_2) 4.73 (s, 2H, CH_2) 6.94–6.97 (m, 2H, 2CH) 7.22–7.31 (m, 5H, arom. benz.) 7.33–7.38 (m, 1H, CH) 8.52, 9.46, 10.04 (3s, 3H, 3NH)
2c	C_6H_5	$\text{C}_{13}\text{H}_{13}\text{N}_3\text{OS}_2$ (291.38)	82 145–147	53.58 53.60	4.50 4.62	14.42 14.57	3243 (NH) 3051 (CH arom.) 2921, 1496 (CH aliph.) 1687 (C=O) 1358 (C=S)	3.76 (s, 2H, CH_2) 6.94–6.97 (m, 2H, 2CH) 7.14–7.19 (m, 1H, CH) 7.31–7.44 (m, 5H, arom. benz.) 9.63 (s, 2H, 2NH) 10.19 (s, 1H, NH)
2d	4- CH_3 - C_6H_4	$\text{C}_{14}\text{H}_{15}\text{N}_3\text{OS}_2$ (305.41)	84 140–142	55.04 55.21	4.95 4.94	13.75 13.82	3232 (NH) 3130 (CH arom.) 2956, 1406 (CH aliph.) 1680 (C=O) 1336 (C=S)	2.28 (s, 3H, CH_3) 3.75 (s, 2H, CH_2) 6.93–6.97 (m, 2H, 2CH) 7.12–7.36 (m, 4H, arom. benz.) 7.37–7.39 (m, 1H, CH) 9.56 (s, 2H, 2NH) 10.15 (s, 1H, NH)

2e	4-OCH ₃ -C ₆ H ₄	C ₁₄ H ₁₅ N ₃ O ₂ S ₂ (321.41)	87 155–157	52.30	4.70	13.07	3133 (NH) 3053 (CH arom.) 2958, 1435 (CH aliph.) 1668 (C=O) 1336 (C=S)	3.32 (s, 3H, CH ₃) 3.75 (s, 2H, CH ₂) 6.88–6.93 (m, 2H, 2CH)
				52.42	4.81	13.06		6.95–7.27 (m, 4H, arom. benz.) 7.37–7.39 (m, 1H, CH) 9.52 (s, 2H, 2NH) 10.14 (s, 1H, NH)
2f	4-Br-C ₆ H ₄	C ₁₃ H ₁₂ BrN ₃ OS ₂ (370.28)	85 168–170	42.16	3.27	11.35	3211 (NH) 3028 (CH arom.) 1699 (C=O) 1487 (CH aliph.) 1353 (C=S)	3.77 (s, 2H, CH ₂) 6.94–6.98 (m, 2H, 2CH)
				42.15	3.28	11.34		7.35–7.51 (m, 4H, arom. benz.) 7.53–7.54 (m, 1H, CH) 9.74 (s, 2H, 2NH) 10.20 (s, 1H, NH)
2g	4-Cl-C ₆ H ₄	C ₁₃ H ₁₂ ClN ₃ OS ₂ (325.83)	88 150–152	48.07	3.73	12.94	3219 (NH) 3027 (CH arom.) 1678 (C=O) 1492 (CH aliph.) 1341 (C=S)	3.78 (s, 2H, CH ₂) 6.94–6.98 (m, 2H, 2CH)
				48.10	3.86	12.89		7.32–7.37, 7.47–7.50 (m, 4H, arom. benz.) 7.38–7.41 (m, 1H, CH) 9.73 (s, 2H, 2NH) 10.21 (s, 1H, NH)
3a	C ₂ H ₅	C ₉ H ₁₁ N ₃ S ₂ (225.33)	70 146–147	47.95	4.92	18.64	3094 (CH arom.) 2953,1418 (CH aliph.) 1568 (C=N) 1496 (C-N) 1354 (C=S)	0.97–1.02 (t, 3H, CH ₃) <i>J</i> = 7.1 Hz 3.89–3.96 (q, 2H, CH ₂) <i>J</i> = 7.1 Hz 4.37 (s, 2H, CH ₂) 6.98–7.05 (m, 2H, 2CH)
				47.86	4.90	18.62		7.44–7.46 (m, 1H, CH) 13.62 (s, 1H, NH)
3b	CH ₂ C ₆ H ₅	C ₁₄ H ₁₃ N ₃ S ₂ (287.39)	69 158–160	58.50	4.56	14.62	3087 (CH arom.) 2919,1437 (CH aliph.) 1574 (C=N) 1499 (C-N) 1358 (C=S)	4.17 (s, 2H, CH ₂) 5.23 (s, 2H, CH ₂) 6.81–6.82 (m, 1H, CH)
				58.57	4.59	14.61		6.87–6.89, 7.30–7.37 (m, 5H, arom. benz.) 7.18–7.21 (m, 2H, 2CH) 13.83 (s, 1H, NH)

3c	C ₆ H ₅	C ₁₃ H ₁₁ N ₃ S ₂ (273.37)	68 160–162	57.11 57.17	4.06 4.09	15.37 15.34	3079 (CH arom.) 2907,1494 (CH aliph.) 1590 (C=N) 1494 (C-N) 1338 (C=S)	4.10 (s, 2H, CH ₂) 6.57–6.58 (m, 1H, CH) 6.82–6.85, 7.30–7.52 (m, 5H, arom. benz.) 7.27–7.29 (m, 2H, 2CH) 13.84 (s, 1H, NH)
3d	4-CH ₃ -C ₆ H ₄	C ₁₄ H ₁₃ N ₃ S ₂ (287.39)	71 154–156	58.50 58.62	4.56 4.59	14.62 14.54	3075 (CH arom.) 2918,1492 (CH aliph.) 1564 (C=N) 1492 (C-N) 1338 (C=S)	2.37 (s, 3H, CH ₃) 4.07 (s, 2H, CH ₂) 6.61–6.63 (m, 1H, CH) 6.85–6.87, 7.30–7.40 (m, 4H, arom. benz.) 7.14–7.17 (m, 2H, 2CH) 13.80 (s, 1H, NH)
3e	4-OCH ₃ -C ₆ H ₄	C ₁₄ H ₁₃ N ₃ OS ₂ (303.39)	73 164–165	55.42 55.49	4.32 4.38	13.85 13.74	3040 (CH arom.) 2928,1438 (CH aliph.) 1569 (C=N) 1468 (C-N) 1335 (C=S)	3.81 (s, 3H, CH ₃) 4.05 (s, 2H, CH ₂) 6.81–6.82 (m, 1H, CH) 6.87–6.89, 7.30–7.37 (m, 4H, arom. benz.) 7.18–7.21 (m, 2H, 2CH) 13.83 (s, 1H, NH)
3f	4-Br-C ₆ H ₄	C ₁₃ H ₁₀ BrN ₃ S ₂ (352.26)	67 134–136	44.32 44.38	2.86 2.83	11.93 11.91	3040 (CH arom.) 2921,1491 (CH aliph.) 1572 (C=N) 1413 (C-N) 1332 (C=S)	3.77 (s, 2H, CH ₂) 6.63–6.65 (m, 1H, CH) 6.84–6.87 (m, 1H, CH) 7.25–7.30, 7.69–7.74 (m, 4H, arom. benz.) 7.34–7.36 (m, 1H, CH) 13.88 (s, 1H, NH)
3g	4-Cl-C ₆ H ₄	C ₁₃ H ₁₀ ClN ₃ S ₂ (307.81)	65 143–145	50.72 50.78	3.27 3.29	13.65 13.69	3042 (CH arom.) 2921,1493 (CH aliph.) 1574 (C=N) 1438 (C-N) 1333 (C=S)	4.13 (s, 2H, CH ₂) 6.63–6.64 (m, 1H, CH) 6.84–7.59 (m, 4H, arom. benz.) 7.50–7.59 (m, 2H, 2CH) 13.89 (s, 1H, NH)



strains of fungi – yeast-like fungi (*Candida albicans*, *Candida famata*, *Geotrichum candidum*), moulds (*Aspergillus fumigatus*, *Aspergillus niger*, *Penicillium* spp., *Scopulariopsis* spp.) and dermatophytes (*Trichophyton granulosum*, *Trichophyton galinae*, *Trichophyton interdigitale*, *Trichophyton mentagrophytes*, *Trichophyton rubrum*). Tested microorganisms were isolated from clinical specimens. In addition, two reference strains of dermatophytes – *Trichophyton mentagrophytes* ATCC9533 and *Trichophyton rubrum* ATCC28188 were included in the studies.

The minimal inhibitory concentration (MIC) defined as the lowest concentration of the compound preventing visible growth, was estimated for all compounds, by agar dilution method, using Mueller-Hinton medium without or with 4% glucose for the growth of bacteria and fungi, respectively (20).

The compounds were dissolved in mixture of acetone and DMSO (1:1). The final concentration of each solvent in the medium (2%) had no influence on the growth of tested microorganisms. Microbial cultures (optical density of 0.5 McFarland standard, 0.025 mL) were put into Petri dishes with 20 mL medium containing several concentrations of the compounds (500, 250, 125, 62.5, 31.25 mg L⁻¹). The media were incubated for 18 h at 37 °C for bacteria or for 5 days at 30 °C for fungi. The medium with no synthesized compounds added served as the control.

RESULTS AND DISCUSSION

Hydrazide of thiophene-2-acetate (**1**) was obtained in the reaction of ethyl thiophene-2-acetate with hydrazine hydrate. The new thiosemicarbazide derivatives (**2**) were obtained in the reaction of hydrazide with isothiocyanates. These compounds were subjected to cyclization in a 2% solution of sodium hydroxide, giving suitable, 4-substituted-3-(thiophene-2-yl-methyl)- Δ^2 -1,2,4-triazoline-5-thiones (**3**). The reaction was performed according to Scheme 1.



Scheme 1

The structure of the new compounds was confirmed by elemental analysis as well as the IR and ^1H NMR spectra. ^1H NMR spectra of thiosemicarbazide derivatives (2) gave proton signals typical of the NH group in the range δ (ppm) 7.93–10.21. In ^1H NMR spectra of 4-substituted-3-(thiophene-2-yl-methyl)- Δ^2 -1,2,4-triazoline-5-thiones (3), proton signals for –NH group in the δ (ppm) 13.62–13.89 range were observed. The above observations suggest that the compounds exist mainly in the thione form.

According to the results from Table II, none of the compounds tested (2a, 2c-e, 3a, 3c-e) inhibited the growth of Gram-positive and Gram-negative bacterial strains, not

Table II. Antibacterial activity of the tested compounds

	Species	MIC (mg L ⁻¹)							
		2a	2c	2d	2e	3a	3c	3d	3e
Gram-positive bacteria	<i>Staphylococcus aureus</i>			> 500 ^a			> 500	> 500 ^a	
	<i>Staphylococcus epidermidis</i>	> 500	> 500	> 500	> 500	> 500	> 500 ^a	> 500	> 500
	<i>Enterococcus faecalis</i>			> 500			> 500	> 500	
	<i>Escherichia coli</i>								
Gram-negative bacteria	<i>Klebsiella pneumoniae</i>								
	<i>Proteus vulgaris</i>								
	<i>Proteus mirabilis</i>	> 500	> 500	> 500	> 500	> 500	> 500	> 500	> 500
	<i>Salmonella enteritidis</i>								
	<i>Salmonella agona</i>								
	<i>Yersinia enterocolitica</i>								
	<i>Pseudomonas aeruginosa</i>								
Yeast-like fungi	<i>Candida albicans</i>								
	<i>Candida famata</i>	> 500	> 500	> 500	> 500	> 500	> 500	> 500	> 500
	<i>Geotrichum candidum</i>								
Moulds	<i>Aspergillus fumigatus</i>			> 500					
	<i>Aspergillus niger</i>	> 500	> 500 ^a	> 500	> 500	> 500	> 500	> 500	> 500
	<i>Penicillium</i> spp.		> 500						
	<i>Scopulariopsis</i> spp.		> 500						
Dermato-Phytes	<i>Trichophyton granulosum</i>			> 500 ^b	> 500 ^a	> 500 ^a		> 500 ^b	> 500 ^a
	<i>Trichophyton galinae</i>			500	500	> 500		250	500
	<i>Trichophyton interdigitale</i>			500	> 500 ^b	> 500		> 500	> 500 ^a
	<i>Trichophyton mentagrophytes</i>								
	<i>Trichophyton mentagrophytes</i> ATCC9533	> 500	> 500	> 500 ^b	> 500 ^a	> 500 ^a	> 500	> 500 ^b	500
	<i>Trichophyton rubrum</i>			> 500 ^b	> 500 ^a	> 500		> 500 ^b	500
	<i>Trichophyton rubrum</i> ATCC28188			> 500 ^b	500	250		250	500
				> 500 ^b	500	250		250	500
				> 500 ^b	500	250		250	500
				> 500 ^b	500	250		250	500

^a Growth inhibition by about 50–70% at a concentration of 500 mg L⁻¹ compared to growth intensity on the medium without the tested compound (visual assessment).

^b Growth inhibition by about 80–90% at a concentration of 125, 250 or 500 mg L⁻¹ compared to growth intensity on the medium without the tested compound (visual assessment).

even at a concentration $> 500 \text{ mg L}^{-1}$. However, compounds **2d**, **3c**, **3d** affected the growth of some *Staphylococci* by about 50–70%.

As shown in Table II, none of the compounds except compound **2d**, inhibits the growth of yeast-like fungi and moulds even at a high concentration of 500 mg L^{-1} . Only compound **2d** affected the growth of *Aspergillus niger* by 50–70% at 500 mg L^{-1} .

Likewise, compounds **2a**, **2c** and **3c** showed no inhibitory effect on the growth of dermatophytes ($MIC > 500 \text{ mg L}^{-1}$). In contrast, compounds **2d**, **2e**, **3a**, **3d**, **3e** showed total or partial inhibitory effect on the growth of some clinical and reference species of *Trichophyton* spp. at concentrations of 250 or 500 mg L^{-1} . Besides, some of them affected the growth of dermatophytes by 80–90% at a concentration of 125 mg L^{-1} . It is worth noting that even $MICs$ of some triazoles currently used in the treatment of mycoses, including dermatophytoses, showed high variability in activity against different species of *Trichophyton* (*e.g.*, for itraconazole MIC range from 0.015 to 8 mg L^{-1} , for fluconazole from 0.5 to 64 mg L^{-1}).

CONCLUSIONS

In this report, an easy method of synthesizing 4-substituted-3-(thiophene-2-yl-methyl)- Δ^2 -1,2,4-triazoline-5-thione in good yield is presented. Eight new compounds were tested for their antibacterial and antifungal activity. Our preliminary data indicate that the assessed, newly synthesized compounds had no significant antibacterial activity but some of them (compounds **2d**, **2e**, **3a**, **3d**, **3e**) showed promising antifungal activity against some species belonging to *Trichophyton* spp. known as the causative agents of superficial mycoses. The most effective derivatives have the carbon ring substituted with a methyl or methoxy group. Although $MICs$ values for the newly synthesized compounds were higher than those for currently available triazoles (*e.g.*, fluconazole, itraconazole), we found them as a good basis for further research.

REFERENCES

1. *The Merck Index*, 12th ed., Merck Co. Inc. Whitehouse Station 1996.
2. J. Haber, Present status and perspectives on antimycotics with systematic effects, *Cas. Lek. Cesk.* **140** (2001) 596–604.
3. M. Dobosz and M. Sikorska, The reaction of cyclization of thiosemicarbazide derivatives with 1,2,4-triazole system. I. The reaction of cyclization of thiosemicarbazide derivatives of 1,2,4-triazole-1-acetic acid, *Acta Polon. Pharm.* **51** (1994) 369–376.
4. M. Dobosz and M. Sikorska, The reaction of cyclization of thiosemicarbazide derivatives with 1,2,4-triazole system. II. The reaction of cyclization of thiosemicarbazide derivatives of 3-methyl-5-mercapto-1,2,4-triazole-4-acetic acid, *Acta Polon. Pharm.* **51** (1994) 377–385.
5. M. Dobosz and A. Pachuta-Stec, Synthesis of new thiourea derivatives of Δ^2 -1,2,4-triazoline-5-thione and 1,3,4-thiadiazole, *Acta Polon. Pharm.* **51** (1994) 457–465.
6. M. Dobosz and A. Pachuta-Stec, Cyclization of 1-cyanoacetyl-4-substituted thiosemicarbazides to 1,2,4-triazole or 1,3,4-thiadiazole derivatives, *Acta Polon. Pharm.* **52** (1995) 103–111.

7. M. Dobosz and A. Maliszewska-Guz, Synthesis of thiosemicarbazide derivatives with 1,2,4-triazole system, *Ann. UMCS, Lublin, Sectio AA* 46/47 (1991/1992) 51–59.
8. M. Dobosz, M. Pitucha and M. Wujec, The reaction of cyclization of thiosemicarbazide derivatives to 1,2,4-triazole or 1,3,4-thiadiazole system, *Acta Polon. Pharm.* **53** (1996) 31–38.
9. M. Dobosz and A. Pachuta-Stec, Synthesis of new derivatives of 3-benzyl- Δ^2 -1,2,4-triazoline-5-thione and 5-benzyl-1,3,4-thiadiazole, *Acta Polon. Pharm.* **53** (1996) 123–131.
10. Fujisawa Pharmaceutical, *Triazoles*, Jpn Kokai Tokkyo Koho JP 59 10,574, 20 Jan 1984; ref. *Chem. Abstr.* **101** (1984) 23485e.
11. F. Kurzer and J. Canelle, Cyclization of 4-substituted 1-amidinothio- semicarbazides to 1,2,4-triazole and 1,3,4-thiadiazole derivatives, *Tetrahedron* **19** (1963) 1603–1610.
12. M. Dobosz, M. Wujec and M. Pitucha, Synthesis of *trans*-1,2-bis(2-R-amino-1,3,4-thiadiazol-5-yl)ethene and *trans*-1,2-bis[3-(4-substituted- Δ^2 -1,2,4-triazoline-5-thione)]ethane, *Ann. UMCS, Lublin, Sectio AA* **50/51** (1995/1996) 67–79.
13. M. Dobosz and M. Wujec, Synthesis of 3-amino-1,2,4-triazole derivatives, *Ann. UMCS, Lublin, Sectio AA* **52/53** (1997/1998) 99–109.
14. M. Dobosz, Studies on products of transformations of 4-carbethoxymethyl-1,2,4-triazoline-thiones, *Acta Polon. Pharm.* **41** (1984) 43–48.
15. G. Mekuskiene, P. Gaidelis and P. Vainilavicius, Synthesis and properties of 5-(4,6-diphenyl-2-pyrimidin-2-yl)-1,2,4-triazoline-5-thione and its derivatives, *Pharmazie* **53** (1998) 94–96.
16. G. Sahin, E. Palaska, P. Kelicen, R. Demirdamar and G. Altmok, Synthesis of some new 1-acyl-thiosemicarbazides, 1,3,4-oxadiazoles, 1,3,4-thiadiazoles and 1,2,4-triazole-3-thiones and their anti-inflammatory activities, *Arzneim. Forsch.*, **51** (2001) 478–484.
17. N. F. Eweiss, A. A. Bahajaj and E. A. Elsherbini, Synthesis and antimicrobial activity of some 4-amino-5-aryl-1,2,4-triazole-3-thiones and their derivatives, *J. Heterocycl. Chem.* **23** (1986) 1451–1458.
18. G. Mazzone, F. Bonina, R. Arrigo Reina and G. Blandino, Synthesis of 1-aryl-4H (R)-thiosemicarbazides, the corresponding 5-aryl 4H (R)-1,2,4-triazoline-3-thiones and some derivatives of pharmaceutical interest, *Farmaco* **36** (1981) 181–196.
19. P. D. Knight, R. A. Demauriac and P. A. Graham, *Silver Salts of 3-amino-5-mercapto-1,2,4-triazole Derivatives*, Ger. Offen. 2,811,025, 05 Oct 1978; ref. *Chem. Abstr.* **90** (1979) 79146s.
20. J. A. Hindler, B. J. Howard and J. F. Keiser, *Clinical and Pathogenic Microbiology*, Mosby-Year Book Inc., St. Louis 1994, pp. 145–195.
21. A. Kucers, S. M. Crowe, M. L. Grayson and J. F. Hoy, *Use of Antibiotics. A clinical Review of Antibacterial, Antifungal and Antiviral Drugs*, Butterworth-Heinmann, Oxford 1997.
22. S. Perea, A. W. Fothergill, D. A. Sutton and M. G. Rinaldi, Comparison of in vitro activities of voriconazol and five established antifungal agents against different species of dermatophytes using a broth macrodilution method, *J. Clin. Microb.* **1** (2001) 385–388.

S A Ž E T A K

**Sinteza i antimikotsko djelovanje 4-supstituiranih
3-(tiofen-2-il-metil)- Δ^2 -1,2,4-triazolin-5-tiona**

MONIKA WUJEC, MONIKA PITUCHA, MARIA DOBOSZ, URSZULA KOSIKOWSKA i ANNA MALM

Rekacijom hidrazida tiofen-2-otene kiseline (**1**) s izotiocijanatima sintetizirani su odgovarajući tiosemikarbazidi (**2a-g**), a njihovom ciklizacijom u 2% NaOH 4-supstituirani 3-(tiofen-2-il-metil)- Δ^2 -1,2,4-triazolin-5-tioni (**3a-g**). Sintetizirani spojevi su potencijalni antimikotici.

Ključne riječi: 4-supstituirani 3-(tiofen-2-il-metil)- Δ^2 -1,2,4-triazolin-5-tioni, sinteza, spektralna analiza, antimikotsko djelovanje

Department of Organic Chemistry, Faculty of Pharmacy, Medical University, Lublin, Poland

Department of Pharmaceutical Microbiology, Faculty of Pharmacy, Medical University, Lublin, Poland