

original scientific paper / izvorni znanstveni rad

## PRELIMINARY CHECK-LIST OF INVASIVE ALIEN PLANT SPECIES (IAS) IN CROATIA

IGOR BORŠIĆ<sup>1</sup>, MILENKO MILOVIĆ<sup>2</sup>, IVA DUJMOVIĆ<sup>3</sup>,  
SANDRO BOGDANOVIĆ<sup>1</sup>, PETRA CIGIĆ<sup>1</sup>, IVANA REŠETNIK<sup>1</sup>,  
TONI NIKOLIĆ<sup>1</sup> & BOŽENA MITIĆ<sup>1</sup>

<sup>1</sup>Department of Botany and Botanical garden, Faculty of Science, University  
of Zagreb, Marulićev trg 9<sup>a</sup>, HR-10000 Zagreb, Croatia  
(E-mail: bozena@botanic.hr)

<sup>2</sup>Medicinsko-kemijska škola, Ante Šupuka bb, HR-22000 Šibenik, Croatia

<sup>3</sup>»Svijet biljaka d.o.o.«, Antonio Smareglia 65, HR-52215 Vodnjan, Croatia

Boršić, I., Milović, M., Dujmović, I., Bogdanović, S., Cigić, P., Rešetnik, I., Nikolić, T. & Mitić,  
B.: Preliminary check-list of invasive alien plant species (IAS) in Croatia. Nat. Croat., Vol. 17,  
No. 2., 55–71, 2008, Zagreb.

In this paper a preliminary check-list of invasive alien plant species (IAS) in Croatia is presented. It has been created on the basis of the literature and of field observations. The list consists of 64 taxa, with family, life-form and geographic origin assigned to each IAS. Analyses according to family affiliation, life-form and origin were made. Out of 27 families the majority (24 families) belong to the dicotyledons. The most numerous family is *Asteraceae*, and genera with the highest number of IAS are *Conyza*, *Erigeron* and *Impatiens*. Life-form analysis showed the predominance of therophytes but also presence of hemicryptophytes, phanaerophytes and geophytes. In the origin analysis, IAS from the Americas predominated, followed by those from Asia and Africa.

**Key words:** invasive alien plant species (plant IAS), alien flora, introduced flora, Croatia

Boršić, I., Milović, M., Dujmović, I., Bogdanović, S., Cigić, P., Rešetnik, I., Nikolić, T. & Mitić,  
B.: Preliminarni popis invazivnih stranih biljnih vrsta (IAS) u Hrvatskoj. Nat. Croat., Vol. 17,  
No. 2., 55–71, 2008, Zagreb.

U radu je predstavljen preliminarni popis invazivnih stranih biljnih vrsta (IAS) u Hrvatskoj. Izrađen je na temelju literaturnih podataka i opažanja na terenu. Popis se sastoji od 64 svojte, kojima je pridružena pripadajuća porodica, životni oblik i geografsko podrijetlo. Izvršene su analize prema pripadnosti porodicama, životnom obliku i podrijetlu. Od 27 porodica većina (24 porodice) pripada dvosupnicama. Najbrojnija porodica je porodica *Asteraceae* dok su rodovi *Conyza*, *Erigeron* i *Impatiens* zastupljeni s najvećim brojem svojti. Analiza životnih oblika pokazala je dominaciju terofita, ali i prisutnost hemikriptofita, fanerofita i geofita. Po podrijetlu dominiraju invazivne svoje iz obiju Amerika, a slijede ih one iz Azije te Afrike.

**Ključne riječi:** invazivne strane biljne vrste, strana flora, alohtona flora, Hrvatska

## INTRODUCTION

Alien species are considered to be »species introduced and growing outside of their natural distribution« (MITIĆ *et al.*, 2008). Their spread was enabled and in the last decades considerably enhanced by long-distance trade, travel and tourism. Although Europe has mostly been a source of taxa in these processes (CROSBY, 1986), it has also received a great number of alien plants from other parts of the world. Until recently, just a few European countries had exhaustive information regarding their alien floras while for most of them at least some data on alien plant species existed (cf. PYŠEK *et al.*, 2002). This situation has positively improved with the DAISIE project which included 49 countries/regions in its inventory of alien plant species at the European level (LAMBDON *et al.*, 2008).

According to the proposed categorisation of alien (non-native) flora in Croatia, plant invasive alien species (IAS) are defined as »naturalized plants that produce reproductive offspring, often in very large numbers and at considerable distances from the parent plants and thus have the potential to spread over large areas (produce reproductive offspring more than 100 m in less than 50 years through generative reproduction and/or more than 6 m in three years through vegetative reproduction)« (RICHARDSON *et al.*, 2000; MITIĆ *et al.*, 2008). Thus, it is necessary for the naturalized species to overcome environmental barriers as well as to spread successfully into new areas in order to become effective invaders (cf. RICHARDSON *et al.*, 2000). The habitats they mostly invade are disturbed, human-made or semi-natural.

The problem of IAS has become one of the most important issues in conservation biology during the last decades. Invasive species are nowadays recognized as the second most important cause of biodiversity loss, just after direct habitat destruction (GENOVESI & SHINE, 2003). For example, VILA *et al.* (2006) have shown that invasion of *Carpobrotus* species, *Ailanthus altissima* (Mill.) Swingle and *Oxalis pes-caprae* L. considerably decreased species richness on Mediterranean islands and altered environmental conditions. Apart from being deleterious from the conservational and environmental point of view, IAS have a strong negative economic impact (PIMENTEL, 2002). In addition to that, some species pose a great problem to public health (e. g. allergenic plant *Ambrosia artemisiifolia* L.; PETERNEL *et al.*, 2005).

Compared to the generally rich history of floristic research in the territory of Croatia, the bibliography on alien plant taxa in Croatia is not abundant. While, for example, alien plant taxa were systematically recorded from the beginning of the 20<sup>th</sup> century in what is now the Czech Republic (PYŠEK *et al.*, 2002), in Croatia only sporadic and in many cases accidental records concerning adventive/neophyte plant taxa exist. They mostly concern the first records of new alien plant taxa for Croatia (e.g. ILIJANIĆ, 1990; BOGDANOVIĆ *et al.*, 2006). Only a minority of articles deal with the distribution and spread of neophyte species in Croatia (e.g. FRANJIĆ *et al.*, 1998; SMITAL *et al.*, 1998).

The first more or less comprehensive list of alien species in Croatia with analyses regarding their life forms and origins was presented in DOBROVIĆ *et al.* (2005). The list comprised casuals, naturalized non-invasive and naturalized invasive alien spe-

cies while cultivated species and archaeophytes were not included. Out of 206 plant taxa on the list, 48 were considered invasive. Apart from this data, there is also an unpublished document of Starmühler that mentions 157 alien taxa for Croatia. This list was used in the inventory of alien species in the framework of the DAISIE project, where Croatia was represented only by this total number of alien plants. As Croatian botanists were not involved in that project, other data on alien and/or neophyte species were not included (LAMBDON *et al.*, 2008). With regard to invasive alien species only, DOBROVIĆ *et al.* (2006) reported 58 taxa on their preliminary list.

As a result of few national projects carried out during last years in Croatia (STANČIĆ, 2007; NIKOLIĆ, 2008), the module »Allochthonous plants« was especially developed and integrated in the already existing Flora Croatica Database (NIKOLIĆ, 2008) as a national standard, and a list of neophytes and the corresponding habitats has been produced (STANČIĆ, 2007).

As can be seen from the above, there has been no systematic research into either alien or invasive plant taxa in Croatia and no comprehensive check-list of alien flora exists. Categorisation has only recently been proposed for alien flora (MITIĆ *et al.*, 2008). Based on this categorisation, this is one of the first attempts to determine which alien plant taxa could be considered a threat to the native flora and should therefore be treated as invasive alien plant species in Croatia.

## MATERIAL AND METHODS

A preliminary check-list of invasive alien plant species in Croatia was created on the basis of two different procedures. Primarily, a survey of the literature was made. All relevant literature concerning non-native species, their first records and their spread on the territory of Croatia was examined. For the sake of completeness, the whole list of references is given, even though some of them are not cited in the text (see References). The articles were mostly from *Acta Botanica Croatica* (e.g. ILIJA-NIĆ & TOPIĆ, 1986), *Natura Croatica* (e.g. BOGDANOVIĆ *et al.*, 2003) and *Fragmenta phytomedica et herbologica* (e.g. GAŽI-BASKOVA & ŠEGULJA, 1978b). Especial consideration was given to the weeds from HULINA (1998a). As a second source, our own field observations were used.

The nomenclature of plant invasive alien species follows the Croatian Flora Check-list (NIKOLIĆ, 2008).

Life-form and origin were attributed to each IAS given on the check-list. The life-forms were interpreted according to Raunkier's system (RAUNKIER, 1934) as presented by ELLENBERG *et al.* (1991) and OBERDORFER (2001) and were denoted with the following abbreviations: **P** – Phanaerophytes, **Ch** – Chamaephytes, **H** – Hemicyclopediae, **T** – Therophytes, **G** – Geophytes, **Hy** – Hydrophytes. For the IAS not listed in ELLENBERG *et al.* (1991) and OBERDORFER (2001), available literature (see References) was mostly used.

Geographic origins of IAS were attributed only in general terms (at the continental level except for Mediterranean species) and were assigned with the following

abbreviations: **Am** – the Americas (North and South), **As** – Asia, **Af** – Africa, **EA** – Eurasia, **M** – Mediterranean. Data on origin were taken mostly from the available literature (see References).

Basic statistic analyses were carried out to show the relative abundance of IAS according to the family affiliation, life-form and geographic origin. Taxa that originate from two different continents were considered as representatives for each of them in the origin analysis.

## RESULTS AND DISCUSSION

The preliminary check-list of invasive alien plant species in Croatia consists of 64 taxa and is presented in Table 1. There are neither pteridophytes nor gymnosperms on the list although alien representatives of these two groups exist in the flora of Croatia (cf. NIKOLIĆ, 2008). The species *Pinus nigra* Arnold is considered by SUŠIĆ & RADEK (2008) to be invasive on the island of Cres. It was often used for forestation because of its resistance and fast-growth, as was the case on the island of Cres. We did not include it on our list because it is a native species in Croatia, naturally distributed on Grobničko polje, near Senj, on Mala Kapela, Velebit (Paklenica) and Biokovo mountains, on the islands of Brač, Hvar and Korčula as well as on the Pelješac peninsula (LIBER, 2000). However, this species is also considered to be a transformer on rocky grasslands in Hungary (TÖRÖK et al., 2003) and should therefore be monitored outside its native range.

The check-list presented here contains six more taxa, added on the basis of newer field observations, than the previous list of IAS produced by DOBROVIĆ et al. (2006). These added taxa are: *Cenchrus incertus* M. A. Curtis, *Duchesnea indica* (Andrews) Focke, *Nicotiana glauca* Graham and *Parthenocissus quinquefolia* (L.) Planchon. We have also included three subspecies in the checklist: *Erigeron annuus* (L.) Pers. subsp. *annuus*, *E. annuus* subsp. *septentrionalis* (Fernald et Wiegand) Wagenitz and *E. annuus* subsp. *strigosus* (Mühlensb. ex Willd.) Wagenitz, instead of the species *Erigeron annuus* (L.) Pers. which was on the list of DOBROVIĆ et al. (2006).

The list consists of invasive alien plant species that belong to 27 families (Tab. 2) of angiosperms, the majority of which (24 families with 55 taxa) belong to dicotyledons. Monocotyledons are represented by only three families with nine taxa. It is approximately the same ratio of dicotyledon and monocotyledon families as in the Croatian vascular flora in general (cf. NIKOLIĆ, 2001). The family with the highest number of IAS is *Asteraceae*, which is the second biggest family of the entire Croatian flora (with 448 taxa; NIKOLIĆ, 2001). It includes 22 invasive taxa, which account for more than one third of all IAS in Croatia. Other families with a considerable number of invasive taxa are *Poaceae*, *Solanaceae* and *Balsaminaceae*. Out of them, only *Poaceae* is among the 15 biggest families of Croatian vascular flora (6% of all taxa; NIKOLIĆ, 2001). The family *Solanaceae* with 41 taxa (NIKOLIĆ, 2008) accounts for less than 0.8% of Croatian vascular flora, while four invasive taxa belonging to this family total 6.3% of all IAS. The family *Balsaminaceae* is represented in the flora of Croatia by only five species (NIKOLIĆ, 2008), out of which a full three are consid-

**Tab. 1.** List of plant invasive alien species in Croatia.

Taxon	Family	Life form	Origin
<i>Abutilon theophrasti</i> Medik.	Malvaceae	T	EA Af
<i>Acer negundo</i> L.	Aceraceae	P	Am
<i>Ailanthus altissima</i> (Mill.) Swingle	Simaroubaceae	P	As
<i>Amaranthus retroflexus</i> L.	Amaranthaceae	T	Am
<i>Ambrosia artemisiifolia</i> L.	Asteraceae	T	Am
<i>Amorpha fruticosa</i> L.	Fabaceae	P	Am
<i>Angelica archangelica</i> L. subsp. <i>archangelica</i>	Apiaceae	H	EA
<i>Artemisia annua</i> L.	Asteraceae	T	EA
<i>Artemisia verlotiorum</i> Lamotte	Asteraceae	G	As
<i>Asclepias syriaca</i> L.	Asclepiadaceae	G	Am
<i>Aster squamatus</i> (Spreng.) Hieron.	Asteraceae	T	Am
<i>Bidens frondosa</i> L.	Asteraceae	T	Am
<i>Bidens subalternans</i> DC.	Asteraceae	T	Am
<i>Broussonetia papyrifera</i> (L.) Vent.	Moraceae	P	As
<i>Carpobrotus edulis</i> (L.) N.E.Br. in Phillips	Aizoaceae	Ch	Af
<i>Cenchrus incertus</i> M.A.Curtis	Poaceae	T	Am
<i>Chamomilla suaveolens</i> (Pursh) Rydb.	Asteraceae	T	As Am
<i>Chenopodium ambrosioides</i> L.	Chenopodiaceae	T	Am
<i>Conyza bonariensis</i> (L.) Cronquist	Asteraceae	T	Am
<i>Conyza canadensis</i> (L.) Cronquist	Asteraceae	T	Am
<i>Conyza sumatrensis</i> (Retz.) E.Walker	Asteraceae	T	Am
<i>Cuscuta campestris</i> Yuncker	Cuscutaceae	T	Am
<i>Datura innoxia</i> Mill.	Solanaceae	T	Am
<i>Datura stramonium</i> L.	Solanaceae	T	Am
<i>Diplotaxis erucoides</i> (L.) DC.	Brassicaceae	T	M
<i>Duchesnea indica</i> (Andrews) Focke	Rosaceae	H	As
<i>Echinocystis lobata</i> (Michx.) Torr. et Gray	Cucurbitaceae	T	Am
<i>Eleusine indica</i> (L.) Gaertn.	Poaceae	T	As
<i>Elodea canadensis</i> Michx.	Hydrocharitaceae	Hy	Am
<i>Epilobium ciliatum</i> Raf.	Onagraceae	H	Am
<i>Erigeron annuus</i> (L.) Pers. subsp. <i>annuus</i>	Asteraceae	T	Am
<i>Erigeron annuus</i> (L.) Pers. subsp. <i>septentrionalis</i> (Fernald et Wiegand) Wagenitz	Asteraceae	T	Am
<i>Erigeron annuus</i> (L.) Pers. subsp. <i>strigosus</i> (Mühlb. ex Willd.) Wagenitz	Asteraceae	T	Am
<i>Euphorbia maculata</i> L.	Euphorbiaceae	T	Am
<i>Euphorbia prostrata</i> Aiton	Euphorbiaceae	T	Am
<i>Galinsoga ciliata</i> (Raf.) S.F.Blake	Asteraceae	T	Am

Tab. 1. continued

Taxon	Family	Life form	Origin
<i>Galinsoga parviflora</i> Cav.	Asteraceae	T	Am
<i>Helianthus tuberosus</i> L.	Asteraceae	G	Am
<i>Impatiens balfourii</i> Hooker f.	Balsaminaceae	T	As
<i>Impatiens glandulifera</i> Royle	Balsaminaceae	T	As
<i>Impatiens parviflora</i> DC.	Balsaminaceae	T	As
<i>Juncus tenuis</i> Willd.	Juncaceae	H	Am
<i>Lepidium virginicum</i> L.	Brassicaceae	T	Am
<i>Nicotiana glauca</i> Graham	Solanaceae	P	Am
<i>Oenothera biennis</i> L.	Onagraceae	H	Am
<i>Oxalis pes-caprae</i> L.	Oxalidaceae	G	Af
<i>Panicum capillare</i> L.	Poaceae	T	Am
<i>Panicum dichotomiflorum</i> Michx.	Poaceae	T	Am
<i>Parthenocissus quinquefolia</i> (L.) Planchon	Vitaceae	P	Am
<i>Paspalum dilatatum</i> Poir.	Poaceae	H	Am
<i>Paspalum paspalodes</i> (Michx.) Scribn.	Poaceae	H	Am
<i>Phytolacca americana</i> L.	Phytolaccaceae	H	Am
<i>Reynoutria japonica</i> Houtt.	Polygonaceae	G	As
<i>Reynoutria sachalinensis</i> (F.S.Petrop.) Nakai in T. Mori	Polygonaceae	G	As
<i>Robinia pseudoacacia</i> L.	Fabaceae	P	Am
<i>Rudbeckia laciniata</i> L.	Asteraceae	G	Am
<i>Solanum eleagnifolium</i> Cav.	Solanaceae	H	Am
<i>Solidago canadensis</i> L.y	Asteraceae	H	Am
<i>Solidago gigantea</i> Aiton	Asteraceae	H	Am
<i>Sorghum halepense</i> (L.) Pers.	Poaceae	H	Af As
<i>Tagetes minuta</i> L.	Asteraceae	T	Am
<i>Veronica persica</i> Poir.	Scrophulariaceae	T	As
<i>Xanthium spinosum</i> L.	Asteraceae	T	Am
<i>Xanthium strumarium</i> L. subsp. <i>italicum</i> (Moretti) D.Löve	Asteraceae	T	Am


ered invasive. There are no representatives of the families *Phytolaccaceae* and *Simaroubaceae* in the Croatian flora other than the invasive *Phytolacca americana* L. and *Ailanthus altissima* (Mill.) Swingle, respectively. Genera with the highest number of IAS are *Conyza*, *Erigeron* and *Impatiens*.

The list of neophytes compiled by STANČIĆ (2007) contains among other data information on the status (naturalized/casual) and abundance (from rare to very frequent) of the taxa. The invasive status of the species is not indicated but most of the naturalized and at the same time frequent neophytes are included in our list as well.

**Tab. 2.** Family analysis of plant invasive alien species in Croatia.

Family	Number of taxa	%
Asteraceae	22	34.4
Poaceae	7	10.9
Solanaceae	4	6.3
Balsaminaceae	3	4.7
Brassicaceae	2	3.1
Euphorbiaceae	2	3.1
Fabaceae	2	3.1
Onagraceae	2	3.1
Polygonaceae	2	3.1
Aceraceae	1	1.6
Aizoaceae	1	1.6
Amaranthaceae	1	1.6
Apiaceae	1	1.6
Asclepiadaceae	1	1.6
Chenopodiaceae	1	1.6
Cucurbitaceae	1	1.6
Cuscutaceae	1	1.6
Hydrocharitaceae	1	1.6
Juncaceae	1	1.6
Malvaceae	1	1.6
Moraceae	1	1.6
Oxalidaceae	1	1.6
Phytolaccaceae	1	1.6
Rosaceae	1	1.6
Scrophulariaceae	1	1.6
Simaroubaceae	1	1.6
Vitaceae	1	1.6


Life-form analysis showed the predominance of therophytes (Fig. 1). They include 36 taxa which account for 56.3% of all IAS. The second share goes to hemicyclopediae with 12 taxa (18.7%). Phanaerophytes and geophytes are represented by seven taxa (10.9%) each. There is just one hydrophyte (*Elodea canadensis* Michx.) and one chamaephyte (*Carpobrotus edulis* (L.) N. E. Br. in Phillips). These results and especially the high percentage of therophytes were expected because therophytes (annual plants), with their short life cycle and production of many, easily dispersed seeds, are very effective as invasive plants. Moreover, weeds are generally also therophytes.



**Fig. 1.** Life-form analysis of invasive alien plant species in Croatia.

According to the results of geographic origin analysis (Fig. 2), most of the IAS in Croatia originate from the Americas (71.9%). They contain aliens from both North and South America. As much as 20.3% of invasive plant taxa in Croatia are native to Asia and 6.3% originate from Africa. Eurasian taxa account for 4.7% of all the IAS while there is only one species (1.6%) that originates in the Mediterranean (*Diplotaxis erucoides* (L.) DC.). The small percentage of invasive plants of Eurasian and Mediterranean origin could be explained by the fact that these taxa are likely archaeophytes and are by now a constitutive part of the natural vegetation. On the other hand, plants from the Americas have been spreading more vigorously during the last decades.

Comparing different countries regarding their plant invasive alien species in a quantitative and a qualitative sense is difficult because of the different terminology used. Also, different alien plants invade different habitat types. On the other hand, invasion history, data on invasiveness and effects of alien species observed in other (neighbouring) countries and areas are quite informative and therefore should be used for considering potentially invasive species. In comparison with the Slovenian list of IAS which does not include weeds, out of 42 plant taxa (JOGAN, 2006), 21 are also found on our list. These are mainly some of the most widespread alien species in Europe, such as *Acer negundo* L., *Ailanthus altissima* Desf., *Elodea canadensis* Michx., *Helianthus tuberosus* L., *Reynoutria japonica* Houtt., *Solidago* species etc. (cf. LAMBDON *et al.*, 2008). A list of invasive and transformer (invasive plants that change the character, condition, form or nature of plant communities and/or ecosystems; MITIĆ *et al.*, 2008) plant species occurring in major habitat types in Hungary also contains



**Fig. 2.** Origin analysis of invasive alien plant species in Croatia.

most of the species that are believed to be invasive in Croatia (cf. TÖRÖK *et al.*, 2003). According to ESSL & RABITSCH (2002), 17 vascular plant species are considered invasive and 18 potentially invasive in Austria. Most of them are included in our list as well. STEŠEVIĆ (2005) cites eight major plant invaders in Montenegro, out of which only *Chenopodium album* L. and *Cardaria draba* (L.) Desv. are not included in our list. No data regarding invasive alien plant species were available to us for Serbia and Bosnia and Herzegovina.

The list presented in this article (Tab. 1) is one of our first attempts to define which alien plant species are invasive in Croatia. Strict nomination of some taxa as invasive according to the definition (see above) is hindered by the fact that the real data on their dispersal, accurate distribution, spreading dynamics etc., are mostly unknown. This list should not be considered either definite or complete, hence the attribute preliminary. The taxa from the list should constantly be monitored and the list should continuously be revised and updated depending on newly obtained data and field observations. It is recommended that the updates are done through the new module »Allochthonous plants« of the Flora Croatica Database (NIKOLIĆ, 2008; MITIĆ *et al.*, 2008) which is officially recognized as the national database by the responsible government department.

Although we have compiled much information regarding alien and especially invasive plant species in Croatia and are still working on it, there are yet many important data that need to be obtained, such as mode and year of introduction, accurate distribution data, velocity of spreading and invaded habitat types. In spite of that, we have decided to present our current results and thereby draw attention to the problem of invasive alien species in Croatia as well as contribute to this concerning issue.

## ACKNOWLEDGEMENTS

We would like to thank Dr Jasenka Topić, Dr Renata Šoštarić and Dr Nejc Jogan for their helpful comments while preparing the list, and Igor Rešetnik for improving our English. This work has been partly financed by the State Institute for Nature Protection, Republic of Croatia, through project 1228/05-2.

Received May 15, 2008

## REFERENCES

- BOGDANOVIĆ, S., I. DOBROVIĆ, A. OSTOJIĆ & I. BORŠIĆ, 2003: *Oxalis pes-caprae* L. (Oxalidaceae) a new species in the flora of Croatia. Nat. Croat. **12**(1), 31–37.
- BOGDANOVIĆ, S. & B. MITIĆ, 2004: *Nicotiana glauca* Graham (Solanaceae) a new adventive species in the flora of Croatia. In: MITIĆ, B. & R. ŠOŠTARIĆ (eds.), 1<sup>st</sup> Croatian Botanical Symposium 2004, Book of Abstracts. Croatian Botanical Society, Zagreb, 17–18.
- BOGDANOVIĆ, S., B. MITIĆ, M. RUŠČIĆ & K. DOLINA, 2006: *Nicotiana glauca* Graham (Solanaceae), a new invasive plant in Croatia. Acta Bot. Croat. **65**(2), 203–209.

- BRITVEC, M. & I. TRINAJSTIĆ, 1999: New localities of the *Sisyrinchium bermudiana* L. (Iridaceae) species in Croatia. Nat. Croat. 8(1), 49–54.
- CIGIĆ, P., T. NIKOLIĆ, M. PLAZIBAT, V. HRŠAK & S. JELASKA, 2003: The distribution of the genus *Impatiens* L. (Balsaminaceae) in Medvednica Nature Park, Croatia. Nat. Croat. 12(1), 19–29.
- CROSBY, A.W., 1986: Ecological Imperialism: The biological expansion of Europe, 900–1900. Cambridge University Press, Cambridge.
- DEVIDÉ, Z., 1956: Nova adventivna biljka hrvatske flore: *Echinocystis lobata* (Michx) Torr. & Gray. Izv. Bot. instituta Univ. 14–15, 186–187.
- DOBROVIĆ, I., S. BOGDANOVIC, I. BORŠIĆ & P. CIGIĆ, 2005: Analisi delle specie esotiche della flora croata. In: Longo, N. (ed.), Inform. Bot. Ital. 37(1, Parte A). Società Botanica Italiana Onlus, Firenze, 330–331.
- DOBROVIĆ, I., I. BORŠIĆ, M. MILOVIĆ, S. BOGDANOVIC, P. CIGIĆ, I. REŠETNIK, T. NIKOLIĆ & B. MITIĆ, 2006: Invasive alien species in Croatia – preliminary report. In: BESENDORFER, V. & G.I.V. KLOBUČAR (eds.), Proceeding of abstracts of the 9<sup>th</sup> Croatian Biological Congress with International Participation. Rovinj, September 23–29, 2006. Croatian Biological Society 1885, Zagreb, 146–147.
- DUBRAVEC, K., 1972: New sites of *Echinocystis lobata* (Michaux) Torr. et Gray in Croatia and Slovenia. Fragm. Herbol. Croat. 5, 1–2.
- DUBRAVEC, K., 1994: O raširenosti vrste *Echinocystis lobata* (Michx.) Torr. & Gray (Cucurbitaceae) u Hrvatskoj. In: TRINAJSTIĆ, I. (ed.), Simpozij – Pevalek. Flora i vegetacija Hrvatske, zbornik radova sa skupa održanog u povodu stote obljetnice rođenja hrvatskog botaničara akademika Ive Pevaleka (1893.–1967.) u Koprivnici 20.–22. svibnja 1993. Šumarski fakultet Sveučilišta u Zagrebu, Koprivnica-Zagreb, 19–23.
- DUBRAVEC, K., N. PLAVŠIĆ-GOJKOVIĆ & M. BRITVEC, 1989: Nova nalazišta adventivne vrste *Eleusine indica* (L.) Gaertn. u Hrvatskoj. Fragm. Herbol. Jugosl. 18(1), 95–97.
- ELLENBERG, H., H. E. WEBER, R. DÜLL, V. WIRTH, W. WERNER & D. PAULSEN, 1991: Zeigwerte von Pflanzen in Mitteleuropa. Scripta Geobot. 18, Lehrstuhl für Geobotanik der Universität Göttingen.
- ESSL, F. & W. RABITSCH, 2002: Neobiota in Österreich. Umweltbundesamt, Wien.
- FRANJIĆ, J., 1993: Nova nalazišta vrste *Datura inoxia* Miller (Solanaceae) u Hrvatskoj. Acta Bot. Croat. 52, 97–100.
- FRANJIĆ, J. & I. TRINAJSTIĆ, 1996: Sadašnje stanje rasprostranjenosti vrste *Datura inoxia* Miller (Solanaceae) u Hrvatskoj. Fragm. Phytom. Herbol. 24(2), 5–9.
- FRANJIĆ, J., I. TRINAJSTIĆ & Ž. ŠKVORC, 1998: Prilog poznavanju širenja nekih neofita u Hrvatskoj. Fragm. Phytom. Herbol. 26(1–2), 5–17.
- GAßMANN, A. & E. WEBER, 2005: Plants-Planta. In: WITTENBERG, R. (ed.), An inventory of alien species and their threat to biodiversity and economy in Switzerland. CABI Bioscience Switzerland Centre report to the Swiss Agency for Environment, Forests and Landscape.
- GAŽI-BASKOVA, V., 1978: Širenje vrste *Reynoutria japonica* Houtt. u ruderalnu vegetaciju SR Hrvatske (in Cyrillic). Fragm. Herbol. Jugosl. 5, 41–44.
- GAŽI-BASKOVA, V. & N. ŠEGULJA, 1977: Nove vrste roda *Solanum* (Solanaceae) u flori Jugoslavije. In: ANONYMOUS, Sadržaji refer. III Simpoz. biosistematičara Jugoslavije. Novi Sad, 89.
- GAŽI-BASKOVA, V. & N. ŠEGULJA, 1978a: Nove vrste roda *Solanum* (Solanaceae) u flori Jugoslavije. Acta Biol. Jugosl. Biosistemática ser. G 4(1), 67–74.
- GAŽI-BASKOVA, V. & N. ŠEGULJA, 1978b: Pojava opasnih korova iz roda *Solanum* na kvarnerskom otoku Plavnik. Fragm. Herbol. Jugosl. 6(106–115), 55–59.
- GENOVESI, P. & C. SHINE, 2003: European strategy on invasive alien species. Council of Europe T-PVS 2003/7.

- HODAK, N., 1956: Rasprostranjenost trave *Paspalum distichum* L. ssp. *paspalodes* (Michx.) Thell. u Jugoslaviji. Biol. glas. Period. biol. **9**, 81–85.
- HODAK, N., 1959/1960: Nalazišta tropske vrste *Eleusine indica* Gaertn. u flori Jugoslavije. Acta Bot. Croat. **18/19**, 65–67.
- HORVATIĆ, S., 1947: Nekoliko novih pridošlica u flori grada Zagreba. Glas. biol. sek. Period. biol. ser. II/B **1**, 69–76.
- HORVATIĆ, S., 1949: *Paspalum distichum* L. ssp. *paspalodes* (Michx.) Thell. na području donje Neretve. Izv. Bot. Inst. Univ. Zagrebu **12–13**, 231–238.
- HORVATIĆ, S. & Lj. GOSPODARIĆ, 1960: *Sporobolus vaginæflorus* (Torr.) Wood u bilnjom pokrovu Hrvatske. Acta Bot. Croat. **18–19**, 79–103.
- HRUŠEVAR, D. & V. POSAVEC VUKELIĆ, 2008: *Ailanthus altissima* (Mill.) Swingle. In: NIKOLIĆ, T., Flora. Priručnik za inventarizaciju i praćenje stanja – Svojte. State Institute for Nature Protection, Zagreb.
- HRUŠEVAR, D. & V. POSAVEC VUKELIĆ, 2008: *Ambrosia artemisiifolia* L. In: NIKOLIĆ, T., Flora. Priručnik za inventarizaciju i praćenje stanja – Svojte. State Institute for Nature Protection, Zagreb.
- HULINA, N., 1971: *Eleusine indica* (L.) Gaertn. u turopoljskom čretu. In: ANONYMOUS, Zborn. sa I Simpoz. biosistematičara Jugoslavije, Sarajevo, 11.–14.10.1971, Biološko društvo SR Bosne i Hercegovine, Sarajevo, 17–19.
- HULINA, N., 1985: Vrsta *Panicum dichotomiflorum* Michx. – novi korov u Jugoslaviji. Fragm. Herbol. Jugosl. **14**(1–2), 113–120.
- HULINA, N., 1998a: Korovi. Školska knjiga, Zagreb.
- HULINA, N., 1998b: Nova nalazišta i rasprostranjenost nekih rijetkih biljnih vrsta u flori Hrvatske. Acta Bot. Croat. **55/56**, 41–51.
- HULINA, N., 1998c: Rare, endangered or vulnerable plants and neophytes in a drainage system in Croatia. Nat. Croat. **7**(4), 279–289.
- HULINA, N., 2008: Wild Marigold – *Tagetes minuta* L., new weed on the island of Hvar, and new contribution to the knowledge of its distribution in Dalmatia (Croatia). Agriculturae Conspectus Scientificus **73**(1), 23–26.
- ILIJANIĆ, Lj., 1957: Nova nalazišta dviju adventivnih mlječika (*Euphorbia maculata* L. i *Euphorbia nutans* Lag.) u Hrvatskoj. Acta Bot. Croat. **16**, 105–108.
- ILIJANIĆ, Lj., 1989: Die Stadt Split, neuer Fundort von *Eleusine indica* (L.) Gaertner (Poaceae). In: Anonymous, Bios. Tessaloniki, 117.
- ILIJANIĆ, Lj., 1990: *Paspalum dilatatum* Poiret, eine neue Adventivpflanze in der Flora Kroatiens. Razprave IV. razreda SAZU **5**(31), 89–96.
- ILIJANIĆ, Lj. & S. HEĆIMOVIĆ, 1983: Nova nalazišta adventivne vrste *Bidens bipinnata* L. u istočnojadranskom primorju. Acta Bot. Croat. **42**, 123–126.
- ILIJANIĆ, Lj. & Lj. MARKOVIĆ, 1986: *Panicum dichotomiflorum* Michaux in the surroundings of Zagreb. Acta Bot. Croat. **45**, 137–139.
- ILIJANIĆ, Lj., Lj. MARKOVIĆ & Z. STANČIĆ, 1994: *Impatiens balfourii* Hooker fil. in Kroatien. Acta Bot. Croat. **53**, 115–119.
- ILIJANIĆ, Lj., M. RADIĆ & Ž. ROKOV, 1991: Prilog adventivnoj flori Splita i okolice. Acta Bot. Croat. **50**, 59–65.
- ILIJANIĆ, Lj. & J. TOPIĆ, 1986: *Paspalum dilatatum* Poirer, a new adventitious plant in the flora of Yugoslavia. Acta Bot. Croat. **45**, 141–144.

- IVKOVIĆ, O., 1981a: Novi nalazi adventivne vrste *Eleusine indica* (L.) Gaertn. 1788 (*Poales, Poaceae*) u Jugoslaviji. In: SPRINGER, O. (ed.), Zb. saž. priopćenja I. kongr. biologa Hrv., Poreč, 21.-26.9.1981. Hrvatsko biološko društvo, Zagreb, 96.
- IVKOVIĆ, O., 1981b: Rasprostranjeње adventivne vrste *Lepidium virginicum* L. 1753 (*Capparidales, Brassicaceae*) u Jugoslaviji. In: SPRINGER, O. (ed.), Zb. saž. priopćenja I. kongr. biologa Hrv., Poreč, 21.-26.9.1981. Hrvatsko biološko društvo, Zagreb, 97–98.
- IVKOVIĆ, O., 1982: Novi nalazi adventivne vrste *Eleusine indica* (L.) Gaertn. 1788 (*Poales, Poaceae*) u Jugoslaviji (in Cyrillic). Zb. prir. nauke Matice srp. **63**, 77–81.
- JOGAN, N., 2000: Some new or interesting records of grasses (*Poaceae*) in Croatia. In: LJUBEŠIĆ, N. (ed.), Proceeding of abstracts of the 7<sup>th</sup> Croatian Biological Congress with International Participation. Hvar, September 24–29, 2000. Croatian Biological Society 1885, Zagreb, 291–292.
- JOGAN, N., 2006: Plant invaders in coastal Slovenia. In: BRUNEL, S. (ed.), Invasive plants in Mediterranean type regions of the World. Mèze (France), 25–27 May 2005. Council of Europe Publishing, 332.
- KOŠČEC, F., 1913: *Helodea (Elodea) canadensis* Rich. u varaždinskoj okolici. Glas. hrv. prir. druš. **25**, 30–39.
- KREMER, D., R. PERNAR & M. ANČIĆ, 2006: Distribution of North American ash species in the Drava River basin and Danube basin (Croatia). Acta Bot. Croat. **65**(1), 57–66.
- LAMBDON, P.W., P. PYŠEK, C. BASNOU, M. ARIANOUTSOU, F. ESSL, M. HEJDA, V. JAROŠÍK, J. PERGL, M. WINTER, P. ANASTASIU, P. ANDRIOPoulos, I. BAZOS, G. BRUNDU, L. CELESTI-GRAPOW, P. CHASSOT, P. DELIPETROU, M. JOSEFSSON, S. KARK, S. KLOTZ, Y. KOKKORIS, I. KÜHN, H. MARCHANTE, I. PERGLOVA, J. PINO, M. VILLA, A. ZIKOS, D. ROY & P.E. HULME, 2008: Alien flora of Europe: species diversity, temporal trends, geographical patterns and research needs. Preslia **80**: 101–149.
- LIBER, Z., 2000: Filogenetski i taksonomski odnosi populacija crnoga bora (*Pinus nigra* Arnold) u Hrvatskoj. PhD Thesis. University of Zagreb, Zagreb.
- LUKAČ, G., 1988: Neke značajke strukture sastojina *Solidago gigantea* i *Helianthus tuberosus* i njihove ornitocene u sjeverozapadnoj Hrvatskoj. Acta Bot. Croat. **47**, 63–75.
- LUKAČ, G., 1989: Nova nalazišta vrste *Impatiens glandulifera* Royle u Hrvatskoj. Acta Bot. Croat. **48**, 165–168.
- LUKAČ, G. & Lj. MARKOVIĆ, 1987: O rasprostranjenosti adventivne vrste kamilice (*Matricaria discoidea* DC.) u Hrvatskoj. In: GOMERČIĆ, H. (ed.), Proceedings of abstracts of the papers presented at the third congress of Croatian biologists, Mali Lošinj, 5–10.10.1987. Croatian Biological Society, Zagreb, 111.
- LUKAČ, G. & S. VUJIĆIĆ-KARLO, 2000: Habitat characteristics and the importance of some plant species as singing places for Marsh Warblers (*Acrocephalus palustris*, Aves) in Croatian neophyte structures. Nat. Croat. **9**(3), 169–177.
- MARKOVIĆ, Lj., 1970: Prilozi neofitskoj flori savskih obala u Hrvatskoj. Acta Bot. Croat. **29**, 203–211.
- MARKOVIĆ, Lj., 1973: *Sporobolus neglectus* Nash, nova adventivna vrsta Jugoslavije. Acta Bot. Croat. **32**, 237–242.
- MARKOVIĆ, Lj., 1978: *Iva xanthifolia* Nutt. i *Amaranthus gracilis* Desf. – dvije nove pridošlice u flori Hrvatske. Acta Bot. Croat. **37**, 207–210.
- MARKOVIĆ, Lj., 1979: Neofiti kao šumski korovi u poplavnim područjima kontinentalne Hrvatske. In: ANONYMOUS, I Jugoslavensko savjetovanje o suzbijanju korova u šumarstvu. Sarajevo, 21–29.

- MARKOVIĆ, Lj., 1981: *Chenopodium integrifolium* Worosch. in der Pflanzendecke Kroatiens. Acta Bot. Croat. **40**, 229–232.
- MARKOVIĆ, Lj., 1981: Zur Syntaxonomie der *Xanthium italicum*-Bestände in Kroatien. Berichte der Internationalen Symposien der Internationalen Vereinigung für Vegetationskunde. Gantner, Vaduz, 279–288.
- MARKOVIĆ, Lj., 1984: Zur Verbreitung und Vergesellschaftung von *Impatiens glandulifera* in Kroatien. Acta Bot. Slov., S.A, Taxon., Geobot., Suppl. (1.1984) **1**, 209–215.
- MARKOVIĆ, Lj., 1987: Američki neofiti u flori Hrvatske. In: GOMERČIĆ, H. (ed.), Proceedings of abstracts of the papers presented at the third congress of Croatian biologists, Mali Lošinj, 5–10.10.1987. Croatian Biological Society, Zagreb, 110.
- MARKOVIĆ, Lj. & N. HULINA, 1970: *Eleusine indica* (L.) Gaertn. u širem području Zagreba. Acta Bot. Croat. **29**, 213–215.
- MARKOVIĆ, Lj. & G. LUKAČ, 1993: Verbreitung und Standorte von *Chamomilla suaveolens* in Kroatien. Nat. Croat. **2**(1), 89–98.
- MARKOVIĆ, Lj. & G. LUKAČ, 1994: Fitocenološka pripadnost sastojina čičoke (*Helianthus tuberosus* L. s.l.) u zapadnoj Hrvatskoj. In: TRINAJSTIĆ, I. (ed.), Simpozij – Pevalek. Flora i vegetacija Hrvatske, zbornik radova sa skupa održanog u povodu stote obljetnice rođenja hrvatskog botaničara akademika Ive Pevaleka (1893.–1967.) u Koprivnici 20.–22. svibnja 1993. Šumarski fakultet Sveučilišta u Zagrebu, Koprivnica-Zagreb, 63–69.
- MARKOVIĆ, Lj. & M. RUŠČIĆ, 1999: *Chenopodium multifidum* L. in the Croatian flora. Nat. Croat. **8**(4), 459–463.
- MARKOVIĆ-GOSPODARIĆ, Lj., 1963: Dvije zanimljive ruderalne biljke u flori Hrvatske. Acta Bot. Croat. **22**, 233–237.
- MARTINI, F., 1989: *Salpichroa organifolia* (Lam.) Baillon, eine neue Adventivpflanze für die jugoslawische Flora. Acta Bot. Croat. **48**, 173–176.
- MELZER, H., 1996: Neues zur Flora von Slowenien und Kroatien. Hladnikia **7**, 5–10.
- MILOVIĆ, M., 2001: A contribution to the knowledge of the neophytic flora of the county of Šibenik and Knin (Dalmatia, Croatia). Nat. Croat. **10**(4), 277–292.
- MILOVIĆ, M., 2004: Naturalised species from the genus *Conyza* Less. (Asteraceae) in Croatia. Acta Bot. Croat. **63**(2), 147–170.
- MITIĆ, B., I. BORŠIĆ, M. MILOVIĆ, S. BOGDANOVIC, I. DOBROVIĆ, P. CIGIĆ, I. REŠETNIK, R. ŠOŠTARIĆ, N. VUKOVIĆ & T. NIKOLIĆ, 2007: Treatment of invasive alien plant species in Croatia – Present and Future. In: BRITVEC, M. & Ž. ŠKVORC (eds.), Book of Abstracts, 2<sup>nd</sup> Croatian Botanical Congress. Croatian Botanical Society, Zagreb, 32–33.
- MITIĆ, B., I. BORŠIĆ, I. DUJMOVIĆ, S. BOGDANOVIC, M. MILOVIĆ, P. CIGIĆ, I. REŠETNIK & T. NIKOLIĆ, 2008: Alien flora of Croatia: proposals for standards in terminology, criteria and related database. Nat. Croat. **17**(2), 73–90.
- MITIĆ, B., I. DOBROVIĆ, I. BORŠIĆ, M. MILOVIĆ, S. BOGDANOVIC, P. CIGIĆ, I. REŠETNIK & T. NIKOLIĆ, 2006a: Proposal for Croatian national standards and criteria for autochthonous and allochthonous flora treatment. In: BESENDORFER, V. & G.I.V. KLOBUČAR (eds.), Proceeding of abstracts of the 9<sup>th</sup> Croatian Biological Congress with International Participation, Rovinj, September 23–29, 2006. Croatian Biological Society 1885, Zagreb, 144–145.
- MITIĆ, B., I. DOBROVIĆ, I. BORŠIĆ, M. MILOVIĆ, S. BOGDANOVIC, P. CIGIĆ, I. REŠETNIK & T. NIKOLIĆ, 2006b: Croatian botanical standards for IAS prevention and monitoring. In: RABITSCH, W., F. KLINGENSTEIN & F. ESSL (eds.), Neobiota, from Ecology to Conservation. 4th European Conference on Biological Invasions, Vienna (Austria), September 27–29, 2006. Bundesamt für Naturschutz (BfN), Bonn, 194.

- MITIĆ, B. & T. NIKOLIĆ, 2006: Invasive alien plants in Croatia – Situation and vision. In: BRUNEL, S. (ed.), Invasive plants in Mediterranean type regions of the World. Mèze (France), 25–27 May 2005. Council of Europe Publishing, 325.
- NIKOLIĆ, T., 2001: The diversity of Croatian vascular flora based on the Checklist and CRO-Flora database. *Acta Bot. Croat.* **60**(1), 49–67.
- NIKOLIĆ, T. (ed.), 2008: Flora Croatica baza podataka. On-Line (<http://hirc.botanic.hr/fcd>). Department of Botany, Faculty of Science, University of Zagreb.
- OBERDORFER, E., 2001: Pflanzensoziologische Exkursionsflora. E. Ulmer Verlag, Stuttgart.
- PANDŽA, M., J. FRANJIĆ, I. TRINAJSTIĆ, Ž. ŠKVORC & Z. STANČIĆ, 2001: The most recent state of affairs in the distribution of some neophytes in Croatia. *Nat. Croat.* **10**(4), 259–275.
- PANDŽA, M. & Z. STANČIĆ, 1999: New localities of the species *Datura innoxia* Miller and *Solanum elaeagnifolium* Cav. (Solanaceae) in Croatia. *Nat. Croat.* **8**(2), 117–124.
- PAVLETIĆ, Z., 1980: *Chenopodium* L. u flori Jugoslavije s posebnim obzirom na primjenu u prehrani ljudi i životinja. In: ANONYMOUS, I Kongr. o korovima. Banja Koviljača, 91–96.
- PAVLETIĆ, Z., 1987a: *Diplotaxis erucoides* (L.) DC. nova adventivna vrsta u flori Hrvatske. *Bull. Mus. Hist. Nat. Beogr.*, Ser. B **42**, 15–18.
- PAVLETIĆ, Z., 1987b: Prilozi poznavanju Biokovske flore. *Acta Biokov. Radovi o prirodi biokovskog područja* **4**, 25–30.
- PAVLETIĆ, Z., 1993: Prilog rasprostranjenosti vrste *Impatiens glandulifera* Royle (Balsaminaceae) u Hrvatskoj. *Fragm. Phytom. Herbol.* **21**(1), 63–66.
- PAVLETIĆ, Z., Z. DEVETAK & I. TRINAJSTIĆ, 1978: Novo značajno nalazište neotofita *Solanum elaeagnifolium* Cav. u flori Hrvatskog primorja. *Fragm. Herbol. Jugosl.* **6**(106–115), 69–72.
- PAVLETIĆ, Z. & M. PANDŽA, 1994: *Diplotaxis erucoides* (L.) DC. (Brassicaceae) u hrvatskoj flori. *Fragm. Phytom. Herbol.* **22**(1–2), 25–28.
- PAVLETIĆ, Z. & I. TRINAJSTIĆ, 1990: *Datura inoxia* Miller (Solanaceae) u flori Hrvatske. *Fragm. Herbol. Jugosl.* **19**(2), 133–138.
- PAVLETIĆ, Z., I. TRINAJSTIĆ & I. ŠUGAR, 1980: O rasprostranjenosti vrste *Sisyrinchium angustifolium* Miller u Hrvatskoj. *Acta Bot. Croat.* **39**, 171–174.
- PETERNEL R., J. ČULIG, L. SRNEC, B. MITIĆ, I. VUKUŠIĆ & I. HRGA, 2005: Variation in Ragweed (*Ambrosia artemisiifolia* L.) pollen concentration in central Croatia, 2002–2003. *Ann. Agric. Environ. Med.* **12**, 11–16.
- PEVALEK, I., 1947: *Dichrocephala sonchifolia* nov pripadnik zagrebačke flore. *Glas. biol. sek. Period. biol. ser. II/B* **1**, 13–16.
- PEVALEK, I., 1953: Dvije adventivne araceje u južnoj Dalmaciji. *God. Biol. Inst. Sar. (Zb. rad. usp. K. Maly)* **5**(1–2), 331–334.
- PIMENTEL, D., 2002: Biological Invasions. Economic and Environmental Costs of Alien Plant, Animal and Microbe Species. CRS Press, New York.
- PYŠEK, P., J. SÁDLO & B. MANDÁK, 2002: Catalogue of alien plants of the Czech Republic. *Preslia* **74**, 97–186.
- RAUNKIER, C., 1934: Life forms of plants and statistical plant geography. Clarendon Press, Oxford.
- RICHARDSON, D.M., P. PYŠEK, M. REJMÁNEK, M.G. BARBOUR, F.D. PANETTA & C.J. WEST, 2000: Naturalization and invasion of alien plants: concepts and definitions. *Diversity Distrib.* **6**, 93–107.
- RUŠČIĆ, M., 2003: The most frequent adventive plants in the flora of the town of Split, Croatia. In: REDŽIĆ, S. & S. ĐUG (eds.), Third International Balkan Congress »Plant resources in the

- creation of new values». Book of Abstracts, Sarajevo, 18–24 May 2003. Faculty of Science of the University of Sarajevo (Center for Ecology and Natural Resources), Sarajevo, 76.
- RUŠČIĆ, M., 2003: Urbana flora Splita. Master thesis. University of Zagreb, Zagreb.
- SMITAL, A., Lj. MARKOVIĆ & M. RUŠČIĆ, 1998: O širenju vrste *Artemisia verlotiorum* Lamotte u Hrvatskoj. Acta Bot. Croat. 55/56, 53–63.
- STANČIĆ, Z., 2007: Final report of the projects 'Neophytic species of vascular flora in Croatia' and 'Habitats of neophytic species of vascular flora in Croatia'. State Institute for Nature Protection, Zagreb.
- STEŠEVIĆ, D., 2005: Contribution to the knowledge on the invasive species in the flora of Montenegro. In: BRUNEL, S. (ed.), Invasive plants in Mediterranean type regions of the World. Mèze (France), 25–27 May 2005. Council of Europe Publishing, 322.
- STRGULC KRAJŠEK, S. & N. JOGAN, 2004: *Epilobium ciliatum* Raf., a new plant invader in Slovenia and Croatia. Acta Bot. Croat. 63(1), 49–58.
- SUŠIĆ, G. & V. RADEK, 2008: Invazivne strane biljne i životinjske vrste otoka Cresa. Problemi i mјere uklanjanja. Eko-centar Caput Insulae Beli, Istraživačko-edukacijski centar za zaštitu prirode, Rijeka.
- ŠAJINović, B. & B. KOLJADZINSKI, 1966: Nova adventivna vrsta *Iva xanthifolia* Nutt. (*Cyclachaena xanthifolia* Fresen.) u našoj zemlji. Bull. Mus. Hist. Nat. Beogr., S.B (11.1958-) 21, 217–220.
- ŠEGULJA, N. & Lj. REGULA-BEVILACQUA, 1994: *Sedum sarmentosum* Bunge a newcomer in Croatian flora. Nat. Croat. 3(1), 91–97.
- ŠILIĆ, Č., 1973: *Tagetes minutus* L. – sve masovniji i sve opasniji korov na poljoprivrednim površinama Dalmacije, Hercegovine, Crnogorskog Primorja i južne Makedonije. In: ŠARIĆ, T. (ed.), Yugoslav symposium on weed control in hilly and mountainous areas, Sarajevo, 21–22 June 1973. Yugoslav Weed Control Committee, Academy of Sciences and Arts of Bosnia and Herzegovina, Faculty of Agriculture Sarajevo, Institute for Forestry Sarajevo, Sarajevo, 27–34.
- ŠILIĆ, Č. & M. ŠOLIĆ, 1999: Contribution to the knowledge of the neophytic flora in the Biokovo area (Dalmatia, Croatia). Nat. Croat. 8(2), 109–116.
- TOPIĆ, J. & Lj. ILIJANIĆ, 2003: *Veronica peregrina* L. and *Veronica scardica* Griseb. (Scrophulariaceae), new species in Croatian flora. Nat. Croat. 12(4), 253–258.
- TOPIĆ, J. & B. KUSULJA, 1987: Učestalost vrste *Eleusine indica* L. Gaertn. u Slavoniji. In: GOMERČIĆ, H. (ed.), Proceedings of abstracts of the papers presented at the third congress of Croatian biologists, Mali Lošinj, 5–10.10.1987. Croatian Biological Society, Zagreb, 111.
- TOPIĆ, J. & B. KUSULJA, 1989: Frequency and abundance of the species *Eleusine indica* (L.) Gaertn. in the Slavonia and Baranja regions. Period. biol. 91(1), 174.
- TOPIĆ, J. & N. ŠEGULJA, 1978a: Novo nalazište vrste *Eleusine indica* (L.) Gaertn. u Hrvatskoj. Acta Bot. Croat. 37, 229–230.
- TOPIĆ, J. & N. ŠEGULJA, 1978b: *Azolla filiculoides* Lam. u Sremu i Slavoniji. Zb. prir. nauke Matice srpske 55, 117–118.
- TÖRÖK, K., Z. BOTTA-DUKÁT, I. DANCZA, I. NÉMETH, J. KISS, B. MIHÁLY & D. MAGYAR, 2003: Invasion gateways and corridors in the Carpathian Basin: biological invasions in Hungary. Biological Invasions 5, 349–356.
- TRINAJSTIĆ, I., 1973: *Duchesnea indica* (Andr.) Focke (Rosaceae), nova adventivna vrsta u flori Jugoslavije. Acta Bot. Croat. 32, 261–266.
- TRINAJSTIĆ, I., 1974a: Ein neuer Fundort von *Impatiens glandulifera* Royle in Kroatien. Fragm. Herbol. Jugosl. 40, 1–6.
- TRINAJSTIĆ, I., 1974b: *Galinsoga ciliata* (Rafin) Blake in der Ruderal-Flora Kroatiens. Fragm. Herbol. Jugosl. 47, 1–10.

- TRINAJSTIĆ, I., 1974c: Prilog poznavanju horologije neofita *Tagetes minutus* L. na području Jugoslavije. Acta Bot. Croat. **33**, 231–235.
- TRINAJSTIĆ, I., 1975a: *Bidens bipinnata* L. – nova adventivna vrsta u flori Hrvatske. Acta Bot. Croat. **34**, 171–173.
- TRINAJSTIĆ, I., 1975b: Kronološka klasifikacija antropohora s osvrtom na helenopaleofite jadranskog primorja Jugoslavije. Biosistematika, Acta Biologica Jugoslavica, serija G **1**(1), 79–85.
- TRINAJSTIĆ, I., 1975c: Trepavičasta konica – nova pridošlica u hrvatskoj flori. Priroda **64**(2), 40–42.
- TRINAJSTIĆ, I., 1977: Chronological classification of the anthropochors. Fragm. Herbol. Jugosl. **2**, 27–31.
- TRINAJSTIĆ, I., 1978a: Novo nalazište vrste *Bidens bipinnata* L. u Hrvatskoj. Fragm. Herbol. Jugosl. **5**(96–105), 5–7.
- TRINAJSTIĆ, I., 1978b: O rasprostranjenosti adventivne vrste *Galinsoga ciliata* (Rafin) Blake u Jugoslaviji. Fragm. Herbol. Jugosl. **6**(106–115), 61–68.
- TRINAJSTIĆ, I., 1979a: Još jedno novo nalazište taksona *Bidens bipinnata* L. u Hrvatskoj. Fragm. Herbol. Jugosl. **8**(126–135), 5–6.
- TRINAJSTIĆ, I., 1979b: O ulozi antropohora u sastavu regionalne flore i vegetacije. In: RAUŠ, Đ. (ed.), Second congress of ecologists of Yugoslavia, Book II, Zadar-Plitvice, 1-7.10.1978. The association of ecological societies of Yugoslavia, Zagreb, 1143–1152.
- TRINAJSTIĆ, I., 1984: Novi podaci o rasprostranjenosti adventivne vrste *Galinsoga ciliata* (Rafin.) Blake u u Jugoslaviji. Fragm. Herbol. Jugosl. **13**(2), 63–66.
- TRINAJSTIĆ, I., 1990: Prilog poznavanju rasprostranjenosti vrste *Reynoutria japonica* Houtt. (Polygonaceae) u Jugoslaviji. Fragm. Herbol. Jugosl. **19**(2), 139–143.
- TRINAJSTIĆ, I., 1993: *Bidens subalternans* DC. u neofitskoj flori Hrvatske. Acta Bot. Croat. **52**, 107–112.
- TRINAJSTIĆ, I., 2001: Turkestanski brijest – *Ulmus pinnato-ramosa* Dieck ex Koehne u dendroflori Hrvatske. Šumarski list **125**(9–10), 533–537.
- TRINAJSTIĆ, I. & J. FRANJIĆ, 1994: Prilog poznavanju rasprostranjenosti vrste *Impatiens glandulifera* Royle (Balsaminaceae) u Hrvatskoj. Fragm. Phytom. Herbol. **22**(1–2), 21–24.
- TRINAJSTIĆ, I. & J. FRANJIĆ, 1995: As. *Impatiens-Solidaginetum* M. Moor 1958 (*Calistegion sepium*) u vegetaciji Republike Hrvatske. Fragm. Phytom. Herbol. **23**(1), 25–30.
- TRINAJSTIĆ, I., J. FRANJIĆ & D. KAJBA, 1991: Prilog poznavanju rasprostranjenosti vrste *Echinocystis lobata* (Michx.) Torr. et Gray (Cucurbitaceae) u Hrvatskoj. Fragm. Herbol. Croat. **20** (1–2), 69–74.
- TRINAJSTIĆ, I., J. FRANJIĆ & D. KAJBA, 1993: Dosadašnje stanje rasprostranjenosti vrste *Galinsoga quadriradiata* Ruiz et Pavon (Asteraceae) u Hrvatskoj. Fragm. Phytom. Herbol. **21**(2), 117–121.
- TRINAJSTIĆ, I., J. FRANJIĆ & D. KAJBA, 1994: Prilog poznavanju vrste *Reynoutria japonica* Houtt. (Polygonaceae) u Hrvatskoj. Acta Bot. Croat. **53**, 145–149.
- TRINAJSTIĆ, I., J. FRANJIĆ, D. KAJBA & J. SAMARDŽIĆ, 1991: Dosadašnje stanje rasprostranjenosti vrste *Reynoutria japonica* Houtt. (Polygonaceae) u Hrvatskoj. Fragm. Herbol. Croat. **20**(1–2), 63–67.
- TRINAJSTIĆ, I. & N. JASPRICA, 1998: *Eleusine tristachya* (Lam.) Lam. (Poaceae) newcomer in the flora of Croatia. Acta Bot. Croat. **57**, 89–94.
- TRINAJSTIĆ, I. & Z. PAVLETIĆ, 1978: *Azolla filiculoides* Lam. u vegetaciji vodenjara istočne Hrvatske. Acta Bot. Croat. **37**, 159–162.

- TRINAJSTIĆ, I. & Z. PAVLETIĆ, 1989: *Guizotia abyssinica* (L.F.) Cass. – nova pridošlica u flori Hrvatske. *Fragm. Herbol. Jugosl.* **18**(1), 99–101.
- TRINAJSTIĆ, I., Z. PAVLETIĆ, J. FRANJIĆ & Z. LIBER, 1993: Prilog poznavanju neofitske flore makarskog primorja (Dalmacija, Hrvatska). *Fragm. Phytom. Herbol.* **21**(1), 57–62.
- TRINAJSTIĆ, I. & Ž. ŠPANJOL, 1994: *Crytomium falcatum* (L. fil.) C. Presl (*Polystichaceae*) a new-comer in Croatian flora. *Nat. Croat.* **3**(1), 87–90.
- VILA, M., M. TESSIER, C.M. SUEHS, G. BRUNDU, L. CARTA, A. GALANIDIS, P. LAMBDON, M. MANCA, F. MEDAIL, E. MORAGUES, A. TRAVESSET, A.Y. TROUMBIS & P.E. HULME, 2006: Local and regional assessments of the impacts of plant invaders on vegetation structure and soil properties of Mediterranean islands. *J. Biogeogr.* **33**, 853–861.
- VREŠ, B., 1996: New localities of the species *Eleusine indica* (L.) Gaertn. (*Poaceae*) in Croatia and Slovenia. *Nat. Croat.* **5**(2), 155–160.