

short communication / kratko priopćenje

NEW DATA ON CROATIAN CRUSTACEAN PLANKTON FAUNA (COPEPODA & CLADOCERA) IN ČINGI-LINGI LAKE

IGOR STANKOVIĆ¹ & IVANČICA TERNJEJ²

¹Hrvatske vode, Department of Water Protection,
Central Water Management Laboratory, Ulica grada Vukovara 220,
10000 Zagreb, Croatia (igor.stankovic@voda.hr)

²Department of Zoology, Faculty of Science, University of Zagreb,
Rooseveltov trg 6, 10000 Zagreb, Croatia

Stanković, I. & Ternjej, I.: New data on Croatian crustacean plankton fauna (Copepoda & Cladocera) in the Čingi-Lungi Lake. *Nat. Croat.*, Vol. 18, No. 1, 181–184, 2009, Zagreb.

Čingi-Lungi Lake is a gravel pit produced 25 years ago. During investigation of zooplankton in the year 2004 we determined eight species. Three of them are copepods and five of them are cladocerans. Čingi-Lungi Lake is the only locality where zooplankton in the Drava Valley region has been investigated.

Key words: Copepoda, Cladocera, plankton, Čingi-Lungi, Croatia

Stanković, I. & Ternjej, I.: Novi podaci za faunu zooplanktona (Crustacea, Copepoda & Cladocera) u Hrvatskoj u jezeru Čingi-Lungi. *Nat. Croat.*, Vol. 18, No. 1, 181–184, 2009, Zagreb.

Jezero Čingi-Lungi je šljunčara nastala prije 25 godina. Tijekom istraživanja u 2004. godini u jezeru je utvrđeno osam vrsta planktonskih račića. Tri vrste spadaju u skupinu veslonožaca, a preostalih pet u skupinu rašljoticalaca. Jezero Čingi-Lungi je prvi i za sada jedini lokalitet na kojem su vršena istraživanja zooplanktona u Podravini.

Ključne riječi: Copepoda, Cladocera, plankton, Čingi-Lungi, Hrvatska

Zooplankton in Croatia has been systematically investigated since 1888 (ŠOŠTARIĆ, 1888). Many species have been found and there are still many to be found. Thus every attempt to take samples from new localities is very valuable for our knowledge of Croatian fauna.

The investigation was conducted on August 1, August 16 and September 2, 2004 in Čingi-Lungi Lake (Fig. 1). The lake is a man-made gravel pit; it was described in detail in STANKOVIĆ & TERNJEJ (2007).



Fig. 1. Čingi-Lingi Lake (photo: I. Stanković).

Specimens were collected in the 16 m deep part of the lake by means of vertical hauls using plankton net with a mesh size of 67 µm. Samples were preserved in 4% formaldehyde. For the purpose of determination, animals were dissected in glycerol and analysed under an Olympus microscope. Copepod determination was carried out according to EINSLE (1993, 1996) and KIEFER (1978), while cladoceran determination was carried out according to AMOROS (1984), MARGARITORIA (1983) and KOROVCHINSKY (1992).

We determined 8 species of zooplankton fauna in Čingi-Lingi Lake. There were 3 species of Copepoda and 5 species of Cladocera. Copepod nauplii and copepodite stages were also established. A detailed systematic and species list is presented:

Class Maxillopoda Dahl, 1956

Subclass Copepoda Milne-Edwards, 1830

Order Calanoida G. O. Sars, 1903

Family Diaptomidae Baird, 1850

1. *Eudiaptomus gracilis* (G. O. Sars, 1862)

Order Cyclopoida Burmeister, 1834

Family Cyclopidae Dana, 1846

2. *Cyclops bohater* Kozminski, 1933

3. *Thermocyclops oithonoides* (G. O. Sars, 1863)

Class Branchiopoda Latreille, 1817

Suborder Cladocera Latreille, 1829

Family Bosminidae Baird, 1845

4. *Bosmina longirostris* (O. F. Müller, 1776)

Family Daphniidae Straus, 1820

5. *Ceriodaphnia quadrangula* (O. F. Müller, 1785)
6. *Daphnia galeata* G. O. Sars, 1864
7. *Daphnia longispina* (O. F. Mueller, 1785)

Family Sididae Baird, 1850

8. *Diaphanosoma brachyurum* (Liévin, 1848)

Čingi-Lingi Lake is the habitat where *C. bohater* was found for the first time in Croatia and the Balkan Peninsula (STANKOVIĆ & TERNJEJ, 2007). According to TERNJEJ & STANKOVIĆ (2007) the lake is also the first place to be investigated in the Drava Valley region (C7) (NIKOLIĆ *et al.*, 1998). Apart from this location, there are no data for the following regions: Banija (C2), Zagorje (C5), Baranja (D1) and the Đakovo-Vinkovci loessial zone (D3) (Fig. 2). Accordingly, this presents a big contribution to Croatian faunal research.

Low species richness is a result of the brevity of the period of investigation and of the investigation being preliminary in nature. Further investigation is expected.

Received February 19, 2008




Fig. 2. Map of Croatian mesoregions (according to NIKOLIĆ *et al.*, 1998) with regions marked in grey that have never been investigated.

REFERENCES

- AMOROS, C., 1984: Crustaces Cladoceres. Introduction Pratique a la Systematique des Organismes des Eaux Continentales Francaises. Université Claude Bernard, **5**, 1–64.
- EINSLE, U., 1993: Crustacea: Copepoda: Calanoida und Cyclopoida. In: SCHWOERBEL, J. & P. ZWICK (eds.), Süsswasserfauna von Mitteleuropa **8(4–1)**, 1–209, Gustav Fischer Verlag.
- EINSLE, U., 1996: Copepoda: Cyclopoida. Genera *Cyclops*, *Megacyclops*, *Acanthocyclops*. Guides to the Identification of the Microinvertebrates of the Continental Waters of the World. H. J. F. DUMONT (ed.), SPB Academic Publishing bv, **10**, 1–82.
- KIEFER, F., 1978: Freilebende Copepoda. In: KIEFER, F. & G. FRYER (eds.), Das Zooplankton der Binnengewässer, Binnengewässer 2. Teil. **26** (2), 1–380.
- KOROVCHINSKY, N. M., 1992: Sididae & Holopediidae (Crustacea: Daphniiformes). Guides to the Identification of the Microinvertebrates of the Continental Waters of the World. SPB Academic Publishing bv, **3**, 1–82.
- MARGARITORIA, F., 1983: Cladoceri (Crustacea: Cladocera). Guide per il riconoscimento delle specie animali delle acque interne Italiane. Consiglio Nazionale delle Ricerche **22**, 1–169.
- NIKOLIĆ, T., BUKOVEC, D., ŠOPF, J. & JELASKA, S. D., 1998: Mapping of Croatian flora – possibilities and standards. Natura Croatica **7**(1), 1–62.
- STANKOVIĆ, I. & TERNEJ, I., 2007: First record of *Cyclops bohater* Kozminski (Copepoda, Cyclopoida) in Croatia and the Balkan Peninsula. Natura Croatica **16** (2), 189–199.
- ŠOŠTARIĆ, D., 1888: Prilog poznavanju faune slatkovodnih korepnjaka Hrvatske. Beitrag zur Kenntniss der Fauna der Süsswasser-Crustaceen. Rad jugoslavenske Akademije Znanosti i Umjetnosti, **92**, 103–214.
- TERNEJ, I. & STANKOVIĆ, I., 2007: Checklist of fresh and brackish water free-living copepods (Crustacea: Calanoida and Cyclopoida) from Croatia. Zootaxa **1585**, 45–57.

S A Ž E T A K

Novi podaci za faunu zooplanktona (Crustacea, Copepoda & Cladocera) u Hrvatskoj u jezeru Čingi-Lingi

I. Stanković & I. Ternjej

Zooplankton se u Hrvatskoj istražuje od 1888. godine, no unatoč tome nisu u potpunosti istražena sva područja. Jezero Čingi-Lingi, nastalo kao šljunčara u neposrednoj blizini rijeke Drave, za sada je jedini lokalitet na kojem su vršena istraživanja zooplanktona u Podravini. Do sada su uz Podravinu potpuno neistraženi Banija, Hrvatsko zagorje, Baranja te područje Đakovo-Vinkovci. U jezeru su tijekom istraživanja 2004. godine utvrđene tri vrste veslonožaca (*Eudiaptomus gracilis*, *Cyclops bohater*, *Thermocyclops oithonoides*) i 5 vrsta rašljoticalaca (*Bosmina longirostris*, *Ceriodaphnia quadrangula*, *Daphnia galeata*, *Daphnia longispina*, *Diaphanosoma brachyurum*).