

Intracellular Ca^{2+} Modulation during Short Exposure to Ischemia-Mimetic Factors in Isolated Rat Ventricular Myocytes

Danijel Pravdić^{1,2}, Nikolina Vladić² and Željko Josip Bošnjak^{2,3}

¹ Department of Physiology, School of Medicine, University of Mostar, Mostar, Bosnia and Herzegovina

² Department of Anesthesiology, Medical College of Wisconsin, Milwaukee, USA

³ Department of Physiology, Medical College of Wisconsin, Milwaukee, USA

ABSTRACT

We investigated the effects of different ischemia-mimetic factors on intracellular Ca^{2+} concentration ($[\text{Ca}^{2+}]_i$). Ventricular myocytes were isolated from adult Wistar rats, and $[\text{Ca}^{2+}]_i$ was measured using fluorescent indicator fluo-4 AM by confocal microscopy. Intracellular pH was measured using c5-(and-6)-carboxy SNARF-1 AM, a dual emission pH-sensitive ionophore. Myocytes were exposed to hypoxia, extracellular acidosis (pH_o 6.8), Na-lactate (10 mM), or to combination of those factors for 25 min. Monitoring of $[\text{Ca}^{2+}]_i$ using fluo-4 AM fluorescent indicator revealed that $[\text{Ca}^{2+}]_i$ accumulation increased immediately after exposing the cells to Na-lactate and extracellular acidosis, but not during cell exposure to moderate ischemia. Increase in $[\text{Ca}^{2+}]_i$ during Na-lactate exposure decreased to control levels at the end of exposure period at extracellular pH 7.4, but not at pH 6.8. When combined, Na-lactate and acidosis had an additive effect on $[\text{Ca}^{2+}]_i$ increase. After removal of solutions, $[\text{Ca}^{2+}]_i$ continued to rise only when acidosis, hypoxia, and Na-lactate were applied together. Analysis of intracellular pH revealed that treatment of cells by Na-lactate and acidosis caused intracellular acidification, while short ischemia did not significantly change intracellular pH. Our experiments suggest that increase in $[\text{Ca}^{2+}]_i$ during short hypoxia does not occur if pH_i does not fall, while extracellular acidosis is required for sustained rise in $[\text{Ca}^{2+}]_i$ induced by Na-lactate. Comparing to the effect of Na-lactate, extracellular acidosis induced slower $[\text{Ca}^{2+}]_i$ elevation, accompanied with slower decrease in intracellular pH. These multiple effects of hypoxia, extracellular acidosis, and Na-lactate are likely to cause $[\text{Ca}^{2+}]_i$ accumulation after the hypoxic stress.

Key words: intracellular Ca^{2+} , intracellular acidosis, ischemia

Introduction

Elevated intracellular Ca^{2+} concentration ($[\text{Ca}^{2+}]_i$) during ischemia and reperfusion is known to induce damage to the heart. During ischemia, interruption of the oxygen supply and substrates promotes ATP hydrolysis and glycolysis, which together induce intracellular acidosis. Intracellular acidosis during ischemia may cause elevation of $[\text{Ca}^{2+}]_i$ by stimulation Na^+-H^+ exchanger resulting in Na^+ influx, which increases $[\text{Ca}^{2+}]_i$ by activation of $\text{Na}^+-\text{Ca}^{2+}$ exchanger. On the other hand, ATP depletion decreases function of Ca-ATPases of sarcoplasmic reticulum and sarcolemma further leading to elevation of $[\text{Ca}^{2+}]_i$ ¹⁻⁴.

Ischemia-induced changes in intracellular environment like depletion of ATP, lactate accumulation, and intracellular acidosis are often used to simulate ischemic damage of the heart^{3,5,6}. Different cellular models, including single fresh cardiomyocyte preparations, require short-lasting ischemia to study the effects of ischemia and reperfusion on cellular function^{3,4}. Differences in ischemia-mimetic solutions can differently affect $[\text{Ca}^{2+}]_i$, and thus be crucial for the research of simulated cardiac ischemic injury. In this study, we compared the effects of 25 min-lasting simulated hypoxia with glucose deficiency,

lactate accumulation, and extracellular acidosis on $[Ca^{2+}]_i$ during hypoxia and reoxygenation in single ventricular cardiomyocytes. Additionally, we examined the role of intracellular pH (pH_i) on changes of $[Ca^{2+}]_i$ under the same conditions. The results of this study provide new insight into effects of simulated cellular ischemia that alter $[Ca^{2+}]_i$, which produce cellular damage.

Materials and Methods

Animals

Adult male Wistar rats (175–250 g) were used in this study. Animals were kept on 12-h cycles of light and dark, and received water and standard food *ad libitum*. The investigation was conducted in accordance with the Institutional Animal Use and Care Committee of the Medical College of Wisconsin (Milwaukee, Wisconsin), and in accordance with U.S. National Institutes of Health standards (NIH Publication 95–23, revised 1996).

Cell isolation

Ventricular myocytes were isolated by enzymatic dissociation^{5,6}. Animals were anesthetized with an intraperitoneal injection of thiobutabarbital (Inactin, 100–150 mg/kg, Sigma-Aldrich, St. Louis, MO) and heparin (1000 U, Abraxis, Schaumburg, IL) to prevent blood clotting. The heart was rapidly removed and mounted to Langendorff perfusion system. All perfusion solutions were continuously gassed with 95% O₂ and 5% CO₂, and temperature was held at 37 °C. Perfusion with a 0 mM Ca²⁺ solution for 5 min was followed by 11 min of recirculation with the same solution containing 0.5 mg/mL collagenase type II (Invitrogen, Carlsberg, CA) and 0.25 mg/mL protease XIV (Sigma-Aldrich, St. Louis, MO). The left ventricle was minced and shaken for 5 min and then filtered through a nylon mesh (200 µm; Spectrum Laboratories, Inc. Rancho Dominguez, CA). After isolation, the myocytes were stored in the HEPES-buffered Tyrode solution at 20°–22°C and used for experiments within 5h after isolation. All cells used in this study were rod shaped in appearance, had well-defined striations, and did not spontaneously contract.

Measurement of $[Ca^{2+}]_i$

Fluorescence measurements were made with an inverted laser-scanning confocal microscope (Nikon Eclipse TE2000-U, Nikon, Japan). $[Ca^{2+}]_i$ was monitored with Ca²⁺ sensitive indicator Fluo-4 AM (Fluo-4). Cells were incubated with Fluo-4 AM (1 µM; Invitrogen, Carlsberg, CA) at room temperature for 20 min. Following incubation, the myocytes were superfused with dye-free Tyrode solution for additional 15 min. Fluo-4 was excited at the 488 nm with an argon laser, and emitted fluorescence was recorded at 520 nm using a 40×/1.4 oil-immersion objective (Nikon). Changes in $[Ca^{2+}]_i$ were expressed as relative changes compared to the baseline. Relative $[Ca^{2+}]_i$ was calculated as the ratio $(F/F_0 \times 100)$, where F is the

measured fluorescence and F₀ is the baseline fluorescence at the beginning of experiment.

Measurement of pH_i

The pH_i was monitored in single myocytes using 5-(and-6)-carboxy SNARF-1, acetoxymethyl ester, acetate (SNARF-1) as the fluorescent indicator^{7,8}. Myocytes were loaded with 10 µM SNARF-1 (Molecular Probes, Eugene, OR) in Tyrode solution at room temperature for 15 min⁹. After dye loading, cells were superfused with the same solution, containing no dye, for additional 20 min before recordings started. SNARF-1 was excited at 543 nm with a green HeNe laser. Emitted fluorescence was simultaneously collected at 590 and 650 nm via a 40×/1.4 oil-immersion objective (Nikon). The 640 nm/590 nm emission ratio from each cell was calculated and converted to pH_i values using nigericin calibration technique^{7,10}. Nigericin, a H⁺/K⁺ exchanger ionophore sets $[K^+]_o = [K^+]_i$ and thus $pH_o = pH_i$. Calibration curve was made by exposing the cells to different pH (pH 6.0, pH 7.5, and pH 9.0) buffers in a depolarizing high K⁺ buffer (140 mM KCl, 1.0 mM MgCl₂, 5 mM D-glucose, 10 mM HEPES, in the presence of 10 µM nigericin)¹¹. To reduce contamination of the superfusion system with nigericin¹², calibration was not done after every experiment. Instead, the pH calibration data were obtained separately from individual cells and were averaged for more than 15 cells.

Solutions

For baseline measurements, myocytes were superfused with normoxic Tyrode solution containing (in mM): 132 NaCl, 10 HEPES, 5 D-glucose, 5 KCl, 1 CaCl₂, and 1.2 MgCl₂ equilibrated at room air; PO₂ 199.1±14.1 mmHg; pH 7.4 adjusted with NaOH. For simulated hypoxia, cells were placed into a closed hypoxic chamber (RC-21B, Warner Instruments, Hamden, CT) and perfused under controlled conditions. Hypoxia was simulated by perfusing the myocytes with hypoxic Tyrode solution containing (in mM): 132 NaCl, 10 HEPES, 0 D-glucose, 8 KCl, 1 CaCl₂, and 1.2 MgCl₂; pH 7.4; PO₂ 26.5±2.8 mmHg, bubbled with 100% N₂ for 20 min. Acidosis was simulated by perfusing the cells with Tyrode solution at pH 6.8. Lactate accumulation was induced by adding 10 mM Na-lactate to Tyrode solution without osmotic compensation. Hypoxia, acidosis, and lactate accumulation were simulated with HEPES-buffered glucose-free Tyrode solution containing (in mM): 118 NaCl, 10 HEPES, 8 KCl, 1 CaCl₂, 1.2 MgCl₂, 1 Na₂HPO₄, and 10 mM Na-L-lactate, pH 6.8; gassed with 100% N₂ for 20 min; PO₂ 26.5±2.8 mmHg. To diminish contamination by room air, all solutions were placed in glass syringes and delivered to recording chamber using an air-tight tubing system. The PO₂ of superfusing solutions was measured with a gas analyzer (Radiometer ABL 505, Radiometer, Copenhagen, Denmark).

Statistical analysis

Data are presented as means±SD, with the sample size (n). To test differences between groups, we used one-way analysis of variance with Bonferroni's post-hoc test. Analysis of variance for repeated measures with Tukey's post-hoc test was used for temporal comparisons between groups. A p value of 0.05 or less was considered significantly different.

Results

Following equilibration in normoxic HEPES-buffered Tyrode solution for 5 min, myocytes were exposed to simulated hypoxia ($PO_2 = 25$ mmHg) at extracellular pH of 7.4. Figure 1a shows the effect of hypoxia applied for 25 min on $[Ca^{2+}]_i$ in ventricular myocytes. During the exposure to hypoxia there was a slow increase of $[Ca^{2+}]_i$. At the end of this period $[Ca^{2+}]_i$ increased to $106 \pm 5\%$ ($p > 0.05$). When hypoxic solution was replaced with normoxic Tyrode solution, $[Ca^{2+}]_i$ returned to the preischemic levels.




Fig 1. Intracellular Ca^{2+} changes in response to hypoxia and acidosis. a) Summarized recordings of relative fluo-4 fluorescence in cardiomyocytes exposed to glucose-free Tyrode solution with decreased O_2 concentration (25 mmHg) for 25 min and in cells which were perfused with normoxic Tyrode solution (Time control). b) Time course of the relative fluo-4 fluorescence change in cells exposed to normoxic Tyrode solution with decreased pH to 6.8. * $p < 0.05$ versus time control, $n = 12$ from 3 rats, error bars represent SD.

Figure 1b shows the effect of acidotic Tyrode solution (pH 6.8) on $[Ca^{2+}]_i$. The rate of $[Ca^{2+}]_i$ increase was faster compared to hypoxia-induced $[Ca^{2+}]_i$ increase, and at the end of 25 min period it was $116 \pm 4\%$ ($p < 0.05$). Washout of acidotic Tyrode solution slowly returned $[Ca^{2+}]_i$ to control levels. In contrast to the effect of hypoxia, $[Ca^{2+}]_i$ recovery was prolonged. This suggests that the $[Ca^{2+}]_i$ regulating mechanisms are impaired by lowering extracellular pH.

Na-lactate (10 mM) was used to induce intracellular lactate accumulation and intracellular acidosis. Figure 2a shows the changes in $[Ca^{2+}]_i$ during the application of lactate at extracellular pH 7.4. Initially, there was rapid increase in $[Ca^{2+}]_i$, which was similar to that induced by acidotic Tyrode solution, $118 \pm 10\%$ ($p < 0.05$). After the initial rise in $[Ca^{2+}]_i$, there was a slow partial recovery of $[Ca^{2+}]_i$, which reached control levels after 25 min. This result suggests that $[Ca^{2+}]_i$ regulating process is active at normal extracellular pH.

When cardiac cells were exposed to hypoxia together with acidosis (pH 6.8) and Na-lactate, there was rapid in-




Fig 2. The effect of Na-lactate and hypoxia with Na-lactate and extracellular acidosis on intracellular Ca^{2+} . a) Fluorescence measurements of intracellular pH in cardiomyocytes exposed to 10 mM Na-lactate (pH 7.4) for 25 min. b) Relative fluo-4 fluorescence intensity during hypoxia together with Na-lactate (10 mM) and extracellular acidosis (pH 6.8). * $p < 0.05$ versus time control, $n = 12$ from 3 rats, error bars represent SD.




Fig 3. Changes in intracellular Ca^{2+} in response to hypoxia, Na-lactate, extracellular acidosis, and combination of those factors. Graph shows summarized data for relative fluo-4 fluorescence of cardiomyocytes exposed to different ischemia-mimetic factors. * $p < 0.05$ versus time control, # $p < 0.05$ versus pH 6.8 or Na-lactate, $n = 10$ from 3 rats, error bars represent SD.

crease in $[Ca^{2+}]_i$ ($149 \pm 15\%$), which did not recover during the hypoxic period. This implies that Na-lactate induces fast increase in $[Ca^{2+}]_i$, which persist when extracellular pH is decreased, indicating that regulating processes are less active at decreased extracellular pH. During early reperfusion, $[Ca^{2+}]_i$ decreased to the preischemic levels, but then significantly increased through reoxygenation, indicating that cells were damaged during simulated hypoxia (Figure 2b). Figure 3 summarized the change in $[Ca^{2+}]_i$ induced by hypoxia, Na-lactate, and extracellular acidosis or with combination of these factors.

The changes in pH_i resulting from exposure to hypoxia, Na-lactate, extracellular acidosis, and combination of these factors are shown in Figure 4. Hypoxia induced small decrease in pH_i (0.07 ± 0.02 pH units, $p > 0.05$).




Fig 4. Changes in pH_i in response to hypoxia, Na-lactate, extracellular acidosis, and combination of those factors. Figure shows representative examples of pH_i changes of cardiomyocytes following exposure to different ischemia-mimetic factors. * $p < 0.05$ versus time control, # $p < 0.05$ versus pH 6.8 or Na-lactate, $n = 12$ from 5 rats, error bars represent SD.

Na-lactate induced faster pH_i decrease (0.15 ± 0.03 pH units, $p < 0.05$), which remained constant. Exposure of the cells to extracellular acidosis caused the decrease in pH_i that was slower than pH_i decrease during the Na-lactate exposure, but it was more profound (0.22 ± 0.05 pH units, $p < 0.05$). This corresponds to an increase in $[Ca^{2+}]_i$, as seen when cells were exposed to acidotic Tyrode solution. Under the conditions of hypoxia with decreased extracellular pH and Na-lactate, the decrease in pH_i was the largest (0.42 ± 0.08 pH units, $p < 0.05$).

Discussion

In this study, we examined the effects of different ischemia-mimicking factors on $[Ca^{2+}]_i$ and pH_i in isolated ventricular myocytes. The main objective of this study was to compare the measurements with different types of hypoxia-mimicking solutions and to determine temporal relationship between the rise in $[Ca^{2+}]_i$ and pH_i . Our results showed that moderate ischemia ($PO_2 = 25$ mmHg) had no effect on the rise in $[Ca^{2+}]_i$ and decrease in pH_i during the tested period. Na-lactate at 10 mM concentration rapidly increased $[Ca^{2+}]_i$, which subsequently recovered at extracellular pH 7.4. When Na-lactate was combined with extracellular acidosis (pH 6.8) this effect was prevented. Extracellular acidosis induced slower rise in $[Ca^{2+}]_i$ that was accompanied with slow intracellular acidification. Combination of those factors also induced the rise in $[Ca^{2+}]_i$ after the removal of solutions.

Accumulation of $[Ca^{2+}]_i$ is one of the main factors causing cardiac damage during ischemia, and rise in $[Ca^{2+}]_i$ causes irreversible myocardial damage⁴. In myocardial ischemia pH_i decreases due to ATP hydrolysis, increased glycolysis, and increased lactate production. In addition, protons can not be removed due to impaired circulation. The lack of ATP impairs the function of Ca^{2+} pumps, while acidosis induces an additional rise in $[Ca^{2+}]_i$ by activation of $Na^+ - Ca^{2+}$ exchanger^{1,2}. When myocytes are exposed longer to an acidic solution, $Na^+ - H^+$ exchanger activity is stimulated resulting in a Na^+ influx, which subsequently induces $Na^+ - Ca^{2+}$ exchanger to operate in reverse mode resulting in rise in $[Ca^{2+}]_i$. Studies performed on animal models showed that $Na^+ - H^+$ exchanger inhibitors, such as amiloride derivatives, can reduce intracellular Na^+ accumulation and subsequent rise in $[Ca^{2+}]_i$ during ischemia². These inhibitors were also beneficial in reducing myocardial ischemia and reperfusion injury¹³.

To study the effects of ischemia and reoxygenation on cardiac myocytes, different ischemia-mimicking solutions are used. Those solutions can induce hypoxia, glucose deficiency, hyperkalemia, hypercapnia, acidosis, lactate accumulation, or substrate deprivation^{3,14}. For ischemia and reperfusion experiments, extended cellular viability is essential to study different cellular processes at reoxygenation. The long duration of hypoxia is limiting factor to study ischemia and reperfusion injury in fresh cellular preparations, like cardiac myocytes, which have to be used within several hours of isolation. On the other

hand, to induce ischemic damage of resting cardiomyocytes by sole hypoxia, usually several hours of hypoxia are necessary. Additionally, cardioprotective strategies like ischemic and anesthetic preconditioning are maximally potent when fresh cells are used³. For that reason the use of the short ischemia and reoxygenation-mimicking factors is necessary in many experimental models.

It has been reported that rat cardiomyocyte superfused with an anoxic medium lacking glucose does not increase $[Ca^{2+}]_i$ during prolonged hypoxia (more than 1 hour)¹⁵. This is in agreement with our experiments where 25 min of hypoxia without glucose did not significantly increase $[Ca^{2+}]_i$. Our experiments suggest that increase in $[Ca^{2+}]_i$ during short hypoxia does not happen if pH_i does not fall. In an earlier study, it was found that 20 mM Na-lactate induces intracellular acidosis and rise in $[Ca^{2+}]_i$. But when it was accompanied with extracellular acidosis (pH 6.4) at used concentration, Na-lactate induced severe twitch potentiations and myocytes went into an irreversible contracture at reoxygenation¹⁶. We observed similar effect when 20 mM Na-lactate was used. The high lactate concentrations can damage the myocytes through increased osmolarity, intracellular acidosis, and increase in $[Ca^{2+}]_i$. Myocyte hypercontracture leads to rupture of plasma membrane and rapid cell death (within 10 min), which is not suitable for studying the effect of reoxygenation on cellular function. Contrary, others have reported that Na-lactate is not responsible for anoxia and reoxygenation damage in cardiac myocytes¹⁷. Intracellular lactate increases up to 5–10 mM during anoxia in isolated ferret heart. In the present study, we used 10 mM Na-lactate to mimic those conditions present during ischemia in isolated hearts. Cells exposed to Na-lactate and acidosis had more severe increase in $[Ca^{2+}]_i$, and decrease in pH_i . This is in agreement with the finding that intracellular acidosis induced by CO_2 and Na-lactate can increase Ca^{2+} transients¹. Furthermore, there was a faster increase in $[Ca^{2+}]_i$ during Na-lactate exposure, than during extracellular acidosis. Increase in $[Ca^{2+}]_i$ has an early rapid phase and slower delayed phase dependent on Na^+-H^+ exchanger activity. There are several mechanisms responsible for the early increase in $[Ca^{2+}]_i$ during Na-lactate exposure. One of the factors that could induce rapid rise in $[Ca^{2+}]_i$ upon exposure to Na-lactate is acute intracellular acidosis, when intracellular buffering mechanisms become hypersaturated. Under the conditions of intracellular acidosis, binding capacity of intracellular proteins for Ca^{2+} is reduced, and acidosis reduces Ca^{2+} uptake by sarcoplasmic reticulum. In addition to this, protons directly inhibit Na^+-Ca^{2+} exchanger, which normally extrudes Ca^{2+} in resting myocytes^{2,3}.

The decrease in $[Ca^{2+}]_i$ observed under prolonged exposure to Na-lactate was prevented with extracellular ac-

idosis. During continued exposure to lactate, another acidosis buffering mechanisms like Na^+-H^+ exchanger might be activated and thus attenuate decrease in pH_i and, therefore, rise in $[Ca^{2+}]_i$. Indeed, it has been shown that pH_i recovery of cardiac myocytes exposed to ischemia-mimetic solution containing Na-lactate can be prevented with Na^+-H^+ exchange inhibitors. Part of this increase results from uncharged form of lactate under conditions of extracellular acidosis, which is more membrane permeable, and lactate can induce greater intracellular acidosis, as seen when Na-lactate and extracellular acidosis were combined. In addition, it is known that reducing extracellular pH leads to an intracellular acidosis in cardiac myocytes, which promotes $[Ca^{2+}]_i$ accumulation¹. Low extracellular pH inhibits Na^+-H^+ exchanger so that H^+ efflux is inhibited, and cellular ability to regulate pH_i is impaired¹.

Increase in $[Ca^{2+}]_i$ during reoxygenation of an ischemic myocardium correlates with the degree of cellular damage¹. Cells damaged by hypoxia are not able to maintain $[Ca^{2+}]_i$ within physiological values. Monitoring changes in $[Ca^{2+}]_i$ and pH_i after cells were exposed to hypoxia-mimicking solutions revealed that pH_i recovered with the similar kinetics in all tested solutions. Faster pH_i recovery in solutions containing Na-lactate is due to activation of lactate- H^+ symport, which has been shown to be the muscle membrane transport systems with the highest capacity for H^+ removal in rat^{18,19}. However, $[Ca^{2+}]_i$ increased only when cells were exposed to hypoxia, acidosis, and Na-lactate together suggesting that some other mechanisms including hypoxia, together with decreased pH_i , are responsible for the myocyte damage with hypoxia-mimicking solutions.

In summary, the results show that intracellular acidosis is the main trigger for acute rise in $[Ca^{2+}]_i$ during acute exposure to ischemia-mimicking solutions in cardiomyocytes. The magnitude and duration of increase in $[Ca^{2+}]_i$ produced by superfusing the myocytes with lactate containing solution is increased at reduced extracellular pH. The inability of cells to prevent $[Ca^{2+}]_i$ rise at reduced extracellular pH showed that other processes, dependent on extracellular pH, contribute to this process. Continued $[Ca^{2+}]_i$ elevation after hypoxia suggests that other mechanisms, independently from intracellular acidosis and lactate accumulation, are involved in ischemic damage of the cardiac myocytes.

Acknowledgements

This work was supported in part by the National Institutes of Health Grants P01GM066730 and R01HL034708 (to ZJB), Bethesda, Maryland.

REFERENCES

- MURPHY E, STEENBERGEN C, *Physiol Rev*, 88 (2008) 581. — 2. MURPHY E, STEENBERGEN C, *Physiology*, 23 (2008) 115. — 3. O'BRIEN JD, FERGUSON JH, HOWLETT SE, *Am J Physiol Heart Circ Physiol*, 294 (2008) H2174. — 4. O'ROURKE B, *Circ Res*, 102 (2008) 1301. — 5. AIZAWA K, TURNER LA, WEIHRAUCH D, BOSNJAK ZJ, KWOK WM, *Anesthesiology*, 101 (2004) 381. — 6. MARINOVIC J, BOSNJAK ZJ, STADNICKA A, *Anesthesiology*, 103 (2005) 540. — 7. BUCKLER KJ, VAUGHAN-JONES RD, *Pflugers Arch*, 417 (1990) 234. — 8. HOSHINO K, AVKIRAN M, *Br J Pharmacol*, 134 (2001) 1587. — 9. VAUGHAN-JONES RD, PEERCY BE, KEENER JP, SPITZER KW, *J Physiol*, 541 (2002) 139. — 10. SPITZER KW, BRIDGE JH, *Am J Physiol*, 262 (1992) C316. — 11. SALVI A, QUILLAN JM, SADEE W, *AAPS PharmSci*, 4 (2002) E21. — 12. RICHMOND PH, VAUGHAN-JONES RD, *Pflugers Arch*, 434 (1997) 429. — 13. NAMEKATA I, SHIMADA H, KAWANISHI T, TANAKA H, SHIGENOBU K, *Eur J Pharmacol*, 543 (2006) 108. — 14. LU J, ZANG WJ, YU XJ, CHEN LN, ZHANG CH, JIA B, *Exp Physiol*, 90 (2005) 497. — 15. ALLSHIRE A, PIPER HM, CUTHBERTSON KS, COBBOLD PH, *Biochem J*, 244 (1987) 381. — 16. CAIRNS SP, WESTERBLAD H, ALLEN DG, *J Physiol*, 464 (1993) 561. — 17. GEISBUHLER TP, ROVETTO MJ, *J Mol Cell Cardiol*, 22 (1990) 1325. — 18. COADY MJ, WALLENDORFF B, BOURGEOIS F, CHARRON F, LAPOINTE JY, *Biophys J*, 93 (2007) 2325. — 19. PILEGAARD H, DOMINO K, NOLAND T, JUEL C, HELLSTEN Y, HALESTRAP AP, BANGSBO J, *Am J Physiol*, 276 (1999) E255.

D. Pravdić

School of Medicine, University of Mostar, Bijeli Brijeg bb, 88000 Mostar, Bosnia and Herzegovina
e-mail: danijel_pravdic@yahoo.com

PROMJENE UNUTARSTANIČNOG Ca^{2+} TIJEKOM KRTAKOTRAJNOG IZLAGANJA ČIMBENICIMA KOJI OPONAŠAJU HIPOKSIJU U VENTRIKULARNIM MIOCITIMA ŠTAKORA

SAŽETAK

Istražili smo učinak različitih čimbenika koji oponašaju ishemiju na koncentraciju unutarstaničnog Ca^{2+} ($[Ca^{2+}]_i$). Ventrikularne srčane stanice smo izolirali iz srca odraslih Wistar štakora, a $[Ca^{2+}]_i$ smo mjerili korištenjem fluorescentnog indikatora fluo-4 AM uz uporabu konfokalnog mikroskopa. Unutarstanični pH smo mjerili koristeći c5-(and-6)-carboxy SNARF-1 AM. Srčane stanice smo izložili hipoksiji, izvanstaničnoj acidozi (pH 6,8), Na-laktatu (10 mM) ili kombinaciji tih čimbenika u trajanju od 25 minuta. Mjerenje $[Ca^{2+}]_i$ koristeći fluo-4 AM indikator, pokazalo je da se nakupljanje $[Ca^{2+}]_i$ povisilo neposredno nakon izlaganja miocita izvanstaničnoj acidozi i Na-laktatu, ali ne i hipoksiji. Porast u nakupljanju $[Ca^{2+}]_i$ tijekom izloženosti Na-laktatu smanjen je na razinu kontrole vrijednosti na kraju vremena izloženosti uz izvanstanični pH 7,4, ali ne i pri pH 6,8. Na-laktat i izvanstanična acidoza imali su aditivni učinak na porast $[Ca^{2+}]_i$. Nakon odstranjivanja otopina, $[Ca^{2+}]_i$ je se povećao samo kad su acidoza, hipoksija i Na-laktat bili primijenjeni zajedno. Analiza unutarstaničnog pH pokazala je da su Na-laktat i izvanstanična acidoza uzrokovali smanjenje unutarstaničnog pH, dok kratkotrajna ishemija nije značajno promijenila unutarstanični pH. Naši rezultati upućuju da se $[Ca^{2+}]_i$ ne povećava tijekom kratkotrajne hipoksija ako se ne smanji unutarstanični pH, dok je izvanstanična acidoza potrebna za trajno povećanje $[Ca^{2+}]_i$ koje je uzrokovano Na-laktatom. U usporedbi s Na-laktatom, izvanstanična acidoza uzrokovala je sporiji porast $[Ca^{2+}]_i$, praćen sporijim smanjivanjem unutarstaničnog pH. Udruženi učinci hipoksije, izvanstanične acidoze i Na-laktata najvjerojatnije uzrokuju nakupljanje $[Ca^{2+}]_i$ nakon hipoksijskog stresa.