



# Variation and Characteristics of the Cranial Vault Thickness in the Krapina and Western European Neandertals

ANTOINE BALZEAU<sup>1,2</sup>

<sup>1</sup>Equipe de Paléontologie Humaine, UMR 5198 du CNRS, USM 204  
Département de Préhistoire du Muséum national d'Histoire naturelle, Paris, France

<sup>2</sup>Service Anthropologie et Préhistoire, Institut royal des Sciences naturelles de Belgique Bruxelles, Belgique.  
E-mail: abalzeau@mnhn.fr

**Key words:** Neandertals, Krapina, cranial vault thickness, CT


## Abstract

*The Krapina collection constitutes the largest sample of Neandertal individuals. However, comparisons of these fossils with other Western European Neandertals have been limited because of the fragmentary condition of the Krapina specimens and because gracility and small dimensions of the cranial remains were attributed to phylogeny and to geological age or to a sex/age bias in the composition of the sample. This study focuses on cranial vault thickness to document new evidence on its variation in the Neandertals. The results demonstrate the similarities between Krapina and the Western European Neandertals in thickness of the cranial vault along the mid-sagittal plane. Finally, Neandertals have characteristics that distinguish them from anatomically modern Homo sapiens fossils.*

## INTRODUCTION

The Krapina collection represents the largest sample of Neandertal individuals, with as many as 70 fossil individuals (1). However, there are no associated post-cranial bones or complete crania (2) and recent studies have particularly focused on the dental remains (e.g., 3, 4) or on the more complete cranial remains (e.g., 5, 6). Moreover, the age distribution and the skeletal composition of the Krapina collection are interpreted by some authors as illustrating a demographic crisis (7) or an ancestral condition for survivorship characteristics (1, 8). Finally, comparisons of the Krapina fossils with other Western European Neandertals have been limited because of the fragmentary condition of the Krapina specimens, and because gracility and small dimensions of the cranial remains were attributed to phylogeny and to geological age or to a sex/age bias in the composition of the sample (6).

Neandertals are generally considered as exhibiting thickened cranial bones (e.g., 9–13). Moreover, cranial vault thickness based on measurements obtained from few and isolated locations has often been used to discuss individual characteristics or individual taxonomic attribution (e.g., 14–22). But as yet, it is not clear to which extend Neandertals have thick bone (11, 23, 24); and what can be the characteristics of the Neandertals for this feature compared with anatomically modern *Homo sapiens* or with other fossil hominid species. It was for example previously proposed that Neandertals have the same pattern of increased vault thickness than *Homo erectus* (11).



**Figure 1.** Variation in the thickness of the cranial vault (in mm) along the mid-sagittal plane for Krapina 3, Krapina 4, Krapina 5 and Krapina 6, illustrating their mid-sagittal plane and partial 3D reconstruction in left lateral view. The cranial thickness is quantified from the sagittal glabella (Gs) to the foramen magnum, and the graphical representation of the data corresponds to the successive landmarks from left to right.

This study presents new evidence about the characteristics of the Krapina fossils compared to Western European Neandertals, as well as about the possible unique features of Neandertal cranial vault thickness.

## MATERIALS

The material includes four adult specimens from Krapina (25, 26) [Krapina 3, Krapina 4, Krapina 5 and Krapina 6 (Figure 1)]. A second sample includes «classic» Neandertal specimens from Western Europe, comprising five adult individuals: La Chapelle aux Saints 1, La Ferrassie 1, La Quina H5, Spy 1 and Spy 10. Comparative samples include anatomically modern, adult *Homo sapiens* individuals from European Upper Palaeolithic sites (Cro Magnon 1, 2, 3, Pataud 1 and Rochereil), European Mesolithic site (Téviec 8, Téviec 9 and Téviec 16) and north African Epi-Palaeolithic sites of Afalou Bou Rhummel and Taforalt (Afalou 2, 12, 13, 28, 30, 34 and Taforalt XI C1, XII C1, XV C2, XV C4, XV C5, XVII C1). These individuals are known to be robust (27–29) and are not considered representative for the complete variation in *Homo sapiens*. However, this sample is used to compare features with a possible size/robustness related component with fossil hominids.

Comparative quantitative analyses of the cranial vault thickness (CVT) were conducted using computed tomography (CT) data. The fossils were CT scanned in different institutions. Acquisition parameters varied according to individual mineralization states and size of the fossil. The Krapina sample was CT scanned with a Siemens Sensation 16 at University of Zagreb Hospital, Bolnica Sestara Milosrdnica, Department of Radiology. Settings ranged between 120 kV, 74–100 mA, 0.6–1.0 mm thick slices, with a reconstruction interval of 0.5 mm, 18–23 cm field of view, and 0.35–0.45 mm pixel size with a pixel matrix of 512\*512. Spy 1 and Spy 10 were CT scanned with a Siemens Somatom 64 at the ULB ERASME in Brussels. Settings were 0.6 mm-thick slices, with a reconstruction interval of 0.3 mm, 23.8 cm field of view, and 0.465 mm pixel size with a pixel matrix of 512\*512 (30). The Croatian and Belgian fossils were CT scanned during the Neandertal Tools project (TNT, <https://www.nespos.org/>; [31–32]). La Chapelle aux Saints 1, La Ferrassie 1 and La Quina H5, as well as the anatomically modern *Homo sapiens* fossils were CT scanned with a General Electric High Speed HAS scanner at the CHNO des Quinzevingts in Paris. Settings were 120 kV, 250 mA, 0.625 mm-thick slices, with a reconstruction interval of 0.45 mm, 23 cm field of view, and 0.45 mm pixel size with a pixel


matrix of 512\*512 (33). The CT data sets were visualized and analyzed using ArteCore 1 software. The CT data sets do not show any noticeable artifacts, even though some fossil skulls show different degrees of mineralization. Moreover, the Hounsfield values for each of the studied fossils are within the range covered by the scanner, resulting in the absence of overflow artefacts.

## METHODS


From a methodological point of view, many aspects have to be considered to perform an analysis of the cranial vault thickness (CVT). At first, attention must be paid to the reproducibility measurement, since determination of cranial landmarks varies between observers. The most easily recognisable landmarks are situated on sutural intersections. However, individual developmental stages may influence their localisation when fontanelles are opened or when sutures are obliterated. Less confidence can be placed in type II or III landmarks (*sensu* 34, 35) in terms of homology than in type I landmarks. This is important when only a few measurements are sampled and compared which are solely based on landmarks obtained by construction (e.g., frontal or parietal eminences, center of a bone in a pre-defined plane, a maximal curvature or extension). This sort of landmark may cause large inter-individual variability in the thickness values and may vary in terms of morphological significance between hominid species. In addition, anatomical landmarks on the cranial external surface do not necessarily have a counterpart on the internal surface. Cranial vault thickness is quantifiable according to various orientations: perpendicular to the external or internal cranial surface or by searching maximal or minimal extensions. Values

vary noticeably depending on this orientation, particularly at the cranial superstructures. Furthermore, thickness values may vary according to the cranial structures and position and development of the endocranial impressions (e.g., tori and keels, internal frontal and occipital crests, venous and meningeal impressions or even pathological and traumatic after-effects). Finally, the conservation state of the fossils and their diagenetical and taphonomical history must be considered since these factors can influence original thickness of the cranial vault.

We have developed a specific analytical protocol to overcome these limitations by using the mid-sagittal plane to quantify CVT (Figure 1, 2). Indeed, this plane is clearly reproducible in both Neandertal and *Homo sapiens* individuals. In addition, the definition of this plane is not influenced by the variations of size and morphology of the different bones of the cranial vault between these two species. So, we have used computed tomography (CT) data to obtain a large number of measurements concerning CVT. CT data correspond to a successive set of slices defining the whole fossil. Each image crosses the bony structures in various orientations. Thickness quantification can be done where the acquisition is perpendicular to the cranial surface and thickness. The mid-sagittal plane is the only one on the whole data set, which extends perpendicularly to the cranial thickness on its full extension. On each individual's corresponding slice, the boundary between the bone and the surrounding air was identified by manual segmentation (SMM: *Seuillage Manuel Multiple* or *Multiple Manual Thresholding*, [36]). This procedure consists of measuring the median value (or half maximum height, HMM) from the CT value of the two elements of which the interface should be defin-



**Figure 2.** Variations in the thickness of the cranial vault (in mm) along the mid-sagittal plane for the Neandertal specimens from Krapina and Western Europe (individual results). The cranial thickness is quantified from the sagittal glabella (Gs) to the foramen magnum, and the graphical representation of the data corresponds to the successive landmarks from left to right.



**Figure 3.** Combined vault thickness for the 60 landmarks along the mid-sagittal plane for Neandertal (in black), European and African anatomically modern *Homo sapiens* (in gray) and Asian *Homo erectus* (in white) individuals; and variations in the thickness of the cranial vault (in mm) along the mid-sagittal plane for the Neandertal specimens from Krapina and Western Europe (individuals results) and the variation of the mean value plus or minus one standard deviation for the fossil anatomically modern *Homo sapiens* (gray area).

ed (37–38). The manual segmentation has to be made each time the attenuation coefficient of one of the elements varies all along the interface. It allows for accurate identification of the interface between two structures, despite local fluctuation in CT numbers (36). This segmentation protocol allowed the isolation of the exact area corresponding to the bone extension in order to quantify CVT. The spatial resolution limitations of the HMH methodology are known to overestimate thickness values when very thin distances are measured (see e.g., 37). However, the elements quantified in this study do not reach this limit (e.g., thickness was always more than 1 mm thick when it was possible to quantify it). There are no overflow artefacts in the analysed datasets and we have used a specific protocol to precisely isolate the different components of the images.

The only limitation to this study is related to the spatial resolution of the datasets and to the partial volume averaging artefact, which is inherent to CT. We estimate that errors in measurements do not exceed  $\pm 0.23$  mm, and are smaller in most cases. In addition, the spatial resolution of the CT datasets was similar for all the studied individuals, resulting in a comparable precision for the measurements. Once the mid-sagittal plane is identified and the segmentation procedures are completed, four principal landmarks are defined on the external cranial surface: the sagittal glabella (noted as Gs), the sagittal bregma (Bs), the sagittal lambda (Ls) and the sagittal external occipital protuberance (Os). The term sagittal is used here to disassociate these landmarks from the classical anatomical landmarks. They do not exactly correspond to the type I landmark definition as they are defined on a 2D plane. However, they remain easily recognizable and reproducible. The following step is to use the internal projection perpendicularly to the endocranial surface of these landmarks to delimitate four endocranial chords from the sagittal glabella to the foramen magnum. Then, these chords are divided in equal segments and thickness is measured at each interval.

Gs-Bs and Bs-Ls are divided in 20, Ls-Os in 15 and Os-foramen magnum in 5. Thickness is quantified perpendicularly to the endocranial surface. Indeed, the internal cranial surface presents less topographic variations than the external cranial surface. In effect, measurements obtained perpendicular to the external surface vary widely as a result of topography changes (e.g., between glabella and the supraorbital sulcus). Thickness values may be influenced by the sutures extension at bregma and lambda or for the parietal bones. Even if some measurements are minimized in these areas, none of the studied individuals show a completely opened sagittal suture, which could have resulted in a null thickness value. Moreover, the analysed plane includes the internal frontal crest and the internal occipital crest for all the analysed Neandertals. It is also the case for the most part of the comparative sample, even if these morphological areas are less pronounced and sometimes partially outside the mid-sagittal plane or incompletely preserved. Finally, while a comparison between several individuals based on a unique metric measurement is not meaningful, our protocol permits us to present a comparative analysis of the vault thickness on 60 landmarks on the whole mid-sagittal plane.

Our whole thickness dataset is affected by the development of the frontal and occipital superstructures as well as by those of the bregmatic eminence or keels. However, the inclusion of these morphological features in our study afforded a better discussion when comparing morphologies of Neandertals with those of anatomically modern *Homo sapiens*. Figure 1 and Figure 2 give cranial vault thickness metrical data, based on the analytical protocol described above and illustrate the variation of the cranial vault thickness (in mm, Y axis) all along the mid-sagittal plane (from Gs to the foramen magnum from left to right, X axis). The left side of the Figure 3 gives the combined vault thickness for the 60 landmarks along the mid-sagittal plane. A direct measurement of the cross-sectional bone area in the mid-

sagittal CT slice was not possible, because of variations in the conservation state of the fossils and of the important influence of the suture closure state on this measurement. Because of small sample size, we do not propose any statistical approach for the variation of the CVT.

## RESULTS

### Variation within the Neandertals

Krapina 3 preserves most of the face, the right half of the frontal bone, portions of the right parietal bone, fragmentary portions of the sphenoid, a piece of the nuchal plane on the right side and the complete right temporal bone. The mid-sagittal plane goes through a very developed internal frontal crest. In this orientation, the connection between the area just above the frontal torus and the frontal squama is not perfect, causing an important and probably slightly amplified decrease in thickness between these two parts in the analysed plane. Thickness was estimated for a few measurements in the middle part of the frontal squama because of a fracture. Thickness is maximal at the sagittal glabella (24.5 mm) and decreases rapidly and regularly posteriorly (see Figure 1). Values vary between 5.5 – 6.0 mm in the middle part of the frontal squama and are greater while getting closer to the sagittal bregma (around 8.2 mm).

Krapina 4 comprises the frontal bone including the right part of the frontal torus and the partial left parietal bone. The mid-sagittal plane is preserved for the frontal bone from posteriorly to the frontal torus to around 1.5 cm anterior to the sagittal bregma. Thickness (Figure 1) is around 14 mm at the most anterior preserved point of the frontal bone and decreases as far as the middle part of the frontal squama (where the minimal value is 6.6 mm). Thickness increases in direction to the sagittal bregma, with a maximal value of 8 mm for the most posterior preserved part of the frontal bone in this orientation.

Krapina 5 includes portion of the two parietals, the right temporal bone and the occipital bone. The actual reconstitution of this skull does not allow the identification of a CT slice going through the parietal bones and the occipital bone and corresponding to the mid-sagittal plane because of the lack of connexion between the different bone fragments and their conservation state. In this context, we quantified the variation of the thickness of the parietal bones on one hand, and of the occipital bone on the other hand, on two different reconstructed CT slices with slightly different orientations. The first image goes through the junction of the parietal bones. We had to estimate some measurements because of the presence of the sagittal suture, but thickness variation can be estimated between 8.0 mm and 9.8 mm along the junction of the parietal bones (Figure 1). For the occipital bone, thickness increases from 9.0 mm to 11.0 mm just posteriorly to the sagittal lambda. Values are lower in the middle part of the occipital squama (minimal value of 9.2 mm) and gets higher while getting closer to the sagittal external occipital protuberance, with a maximal value in this area of 10.4 mm.

Krapina 6 comprises the upper face, the entire frontal torus, and some elements on the right side of the frontal squama, as well as parts of the right parietal bone and of the occipital bone. In the mid-sagittal plane, we could quantify the thickness for nearly all the frontal bone. Thickness (Figure 1) is maximal at the sagittal glabella (20.3 mm). Values vary between 6.2 mm and 6.7 mm in the middle part of the frontal squama. Thickness is more important in direction to the sagittal bregma, with a maximal value of 6.9 mm for the most posterior preserved part of the frontal bone in this orientation. We could not quantify thickness on a large number of landmarks on the parietal and occipital bones because of their preservation state. Thickness of the preserved part of the occipital bone, which corresponds to the area of the external occipital protuberance, is around 11–12 mm.

La Chapelle aux Saints 1, La Ferrassie 1, La Quina H5 and Spy 1 are preserved on nearly all the extension of their respective mid-sagittal plane. Spy 10 does not include the frontal bone in this orientation but comprises well preserved parietal and occipital bones. These individuals show some variability in terms of thickness values but maintain a similar pattern of distribution of the cranial vault thickness all along the mid-sagittal plane (36).

The preservation state of the Krapina fossils does not allow statistical comparison with the western European Neandertal for the distribution of the cranial vault thickness on the whole sagittal plane. Nevertheless, the available data permits comparisons for the thickness of the different cranial bones between these two samples (Figure 2). For the frontal bone, Krapina 3 (Figure 2) and La Chapelle aux Saints 1 have the maximal values for the thickness at the sagittal glabella (24.5 mm). Krapina 3 has great values for the thickness of the anterior part of the frontal bone because of a very developed internal frontal crest. However, thickness in this area is only slightly greater than in La Chapelle aux Saints. Krapina 6 is well into the variability of the others Neandertals for this part of the frontal bone (Figure 2), and its thickness values are close to those of Spy 1. All the analysed Neandertal specimens share a similar variation of the thickness in the area posterior to the frontal torus. Indeed, values decrease rapidly and regularly from the area of the frontal torus to the middle part of the frontal squama. The thickness for the most anterior preserved part of the frontal bone of Krapina 4 is in the middle part of the variability of the others Neandertals (Figure 2). This individual also shares with the others fossils a similar decreasing in thickness values in direction to the middle part of the frontal squama. In the middle part of the frontal squama, the thickness for all the Neandertal individuals varies between 5.5 mm – 9.0 mm, and the Krapina fossils are in the middle part of this variation. Thickness is more important in direction to the sagittal bregma with a variation for the complete sample between 5.9 mm – 8.2 mm. Krapina 3, Krapina 4 and Krapina 6 have values for the thickness of the most posterior preserved part of their frontal bone in the upper part of the variability. For the parietal bones, the available data from

Krapina concern Krapina 5. Variations of thickness at the junction of the parietal bones are influenced by the presence of the sagittal suture, resulting in varying values from one point to the neighbouring one. However, these data allow estimating general thickness of this area (Figure 2). Krapina 5 has the greatest values among the Neandertal sample with a variation between 8.0 mm and 9.8 mm. In this area, thickness varies between 6.5 mm and 8.6 mm for Spy 10 and between 4.8 mm and 7.0 mm for La Quina H5 who are the two extremes individuals, at the exclusion of Krapina 5. For the occipital bone, the available data from Krapina concern Krapina 5. This individual shares with the Neandertal individuals from Western Europe a similar pattern of distribution of the occipital bone thickness in the mid-sagittal plane. Thickness values increase from the sagittal lambda to the middle part of the occipital squama, decrease posteriorly to this area and increase again in direction to the sagittal external occipital protuberance (Figure 2). Krapina 5 has thickness values in the middle part of the occipital squama within the variability of the other Neandertals and in its upper part. The maximal thickness in this area for Krapina 5 is 11.0 mm, whereas it is around 8.0 mm for La Quina H5, 8.5 mm for La Ferrassie 1, 9.0 mm for La Chapelle aux Saints 1, 10.0 mm for Spy 10 and 11.0 mm for Spy 1. The thickness at the sagittal external occipital protuberance for Krapina 5 (10.0 mm), as well as for Krapina 6 (11.0 mm–12.0 mm), is within the variability of the other Neandertals (around 8.0 mm for La Quina H5, 10.0 mm for La Chapelle aux Saints 1, 12.0 mm for La Ferrassie 1 and Spy 10 and 14.0 mm for Spy 1). So, the thickness of Krapina 5 is greater in the middle part of the occipital squama than at the sagittal external occipital protuberance. Finally, the thickness of the nuchal plane decreases from the occipital superstructures to the foramen magnum in a similar way and with similar values for all the analysed Neandertal specimens.

### Comparison with anatomically modern *Homo sapiens*

With regard to the distribution and to the variation of the thickness all along the analysed plane, the Afalou Bou Rhummel and Taforalt skulls ( $n=12$ ) present well developed superstructures. The Cro Magnon ( $n=3$ ), Pataud ( $n=1$ ), Rochereil ( $n=1$ ) and Téviec ( $n=3$ ) individuals yielded absolute thickness values around or lesser than the mean for the Afalou Bou Rhummel and Taforalt individuals. However, all these individuals share the same pattern of cranial vault thickness distribution along the mid-sagittal plane (36). In comparison with previous studies (11, 15, 39–43), the cranial thickness values of our *Homo sapiens* sample are outside the variation of these actual populations but are within the variability observed on fossil samples (44).

Figure 3 illustrates the comparison of cranial vault thickness between Neandertals and anatomically modern *Homo sapiens*. The bar graph on the left side of this image gives values of combined vault thickness for the 60 landmarks along the mid-sagittal plane. We could not

use the Krapina fossils for this analysis because of their incomplete preservation. The analysed Neandertal individuals have values for the combined vault thickness from 396 mm for La Quina H5, 497 mm for La Ferrassie 1, 517 mm for La Chapelle aux Saints 1, to 574 mm for Spy 1. The European anatomically modern *Homo sapiens* ( $n=8$ ) have a range between 417 mm – 617 mm and the African anatomically modern *Homo sapiens* ( $n=11$ ) have a range between 528 mm – 778 mm. The mean value for the Neandertal fossils is significantly less than the mean value for the anatomically modern *Homo sapiens* ( $t = -17.705$ ,  $p < 0.01$ ). Finally, Asian *Homo erectus* fossils from the Ngandong and Sambungmacan sites ( $n = 4$ ) have values for this variable between 620 mm – 648 mm (45). So, these Neandertal specimens have individual values for the combined vault thickness in the mid-sagittal plane within the inferior variability of anatomically modern *Homo sapiens* and below the values of the Ngandong and Sambungmacan *Homo erectus*.

The right part of the Figure 3 presents the comparison of the cranial vault thickness distribution between Neandertals and anatomically modern *Homo sapiens* all along the mid-sagittal plane. The individual results are shown for the Neandertals whereas the gray area illustrates the variation of the mean value plus or minus one standard deviation for the comparative sample (the gray line in the middle part of this distribution is the mean value,  $n = 19$ ). The thickness for the Neandertal specimens exceed the variation shown for the comparative sample only for a few landmarks for Krapina 3 and La Chapelle aux Saints 1 in the anterior part of the frontal bone because of the great development of the internal frontal crest in these two individuals. One of the individual from Afalou has a value for the thickness at the sagittal glabella of 25.0 mm, which is above the values for the Neandertal fossils. On the opposite side, all the other Neandertal individuals yielded values all along the extension of the analysed plane within or below the represented variation of the comparison material.

In terms of variation of cranial vault thickness, the Neandertal specimens present some particularities compared to anatomically modern *Homo sapiens*. The analysed Neandertal and *Homo sapiens* specimens share similar thickness values in the area of the frontal superstructures. However, thickness in these Neandertals decreases rapidly and regularly from the area of the frontal torus to the middle part of the frontal squama whereas this variation is less abrupt in the comparative sample, resulting in greater value for the thickness of the middle part of the frontal squama. The variation along the junction of the parietal bones shares similar characteristics for the two analysed samples, but the Neandertals have lower values for the thickness in this area. On the occipital bone, both samples exhibit an increase of the thickness in the middle part of the occipital squama. On the opposite side, Neandertals only show a relatively reduced increase of thickness in direction to the sagittal external occipital protuberance whereas the comparative sample has an important thickening in this area.

## DISCUSSION

We have developed a specific protocol in order to avoid difficulties resulting from cranial landmarks or measurement definitions as well as limitations in interpretations when only one or few thickness data are used. CVT was quantified all along the mid-sagittal plane on a large number of landmarks. So, we compared the thickness of the frontal bone for Krapina 3, Krapina 4 and Krapina 6 with five West European Neandertals. Krapina 5 was compared with this sample for the thickness of the parietal bones and of the occipital bone. Moreover, the Neandertal sample was compared to anatomically modern *Homo sapiens*.

Krapina 3 and La Chapelle aux Saints 1 have the greater values for the thickness at the glabellar point whereas Krapina 6 is well into the variability of the whole sample of Neandertal specimens. Similarly, the thickness for the most anterior preserved part of the frontal bone of Krapina 4 is in the middle part of the variability of the other Neandertals. All the analysed Neandertals share a common pattern in the area posterior to the frontal torus with decreasing in thickness values in direction to the middle part of the frontal squama. Finally, Krapina 3, Krapina 4 and Krapina 6 have values for the thickness of the most posterior preserved part of their frontal bone in the upper part of the variability observed for the Neandertal sample. Overall, the Krapina fossils are not different from the Western European Neandertals for the values and the variation of the thickness of the frontal bone in the mid-sagittal plane. For the parietal bones, Krapina 5 has the maximal values among the Neandertal sample and a similar pattern of thickness variation. For the occipital bone, Krapina 5 shares with the Neandertal individuals from Western Europe a similar pattern of distribution of the occipital bone thickness in the mid-sagittal plane. Moreover, Krapina 5 has thickness values in the middle part of the occipital squama within the variability of the other Neandertals and in its upper part. The thickness at the sagittal external occipital protuberance for Krapina 5 (10.0 mm) is within the variability of the other Neandertals. So, the values for the cranial vault thickness in the mid-sagittal plane for Krapina 3, Krapina 4, Krapina 5 and Krapina 6 enter and complete the variation observed in the La Chapelle aux Saints 1, La Ferrassie 1, La Quina H5, Spy 1 and Spy 10 Neandertals. Moreover, all these individuals share a similar pattern of thickness variation all along the analysed plane (Figure 2).

These data also highlight the ambiguous relation between cranial vault thickness and sexual attribution of Neandertal specimens. In the analysed sample Krapina 4, Krapina 5, La Chapelle aux Saints 1, La Ferrassie 1 and Spy 10 are generally recognized as male individuals whereas Krapina 3, Krapina 6, La Quina H5 and Spy 1 are proposed to be female individuals, with some variations in sexual attribution depending on the author (e.g., 6, 9, 46–50). Our results on cranial vault thickness do not permit to put in evidence a clear distinction between these two sub-samples. For example, Krapina 3 and 6 as

well as Spy 1 have thickness values for the anterior part of the frontal bone equal or superior to those of La Chapelle aux Saints 1 and La Ferrassie 1. Similarly, for the posterior part of the frontal bone the individuals are classified in the order Krapina 3-Krapina 4-La Chapelle aux Saints 1-Krapina 6-Spy 1-Spy 10-La Ferrassie 1-La Quina H5 from high to low thickness values. Finally, Spy 10, presumably a male, has values for its cranial vault thickness below to those of Spy 1, presumably a female, on the most part of the landmarks analysed along the mid-sagittal plane. Consequently, the cranial vault thickness in the mid-sagittal plane, which however includes the frontal and occipital superstructures, appears not to be a discriminating indicator for sexual determination in Neandertals. CVT seems to be influenced by a size/robustness related component and it is this set of correlated features which is probably more meaningful to clarify sexual attribution when are considered the extreme individuals of the variation (e.g., the parietal bone of Krapina 5 (6); or La Quina H5 which has the lower value for CVT, smaller cranial dimensions and less pronounced cranial superstructures of Neandertal sample).

The anatomically modern *Homo sapiens* individuals from European superior Palaeolithic sites (Cro Magnon 1, 2, 3, Pataud 1 and Rochereil), from Mesolithic European sites (Téviec 8, 9, 16) and North African Epi-Palaeolithic sites of Afalou Bou Rummel and Taforalt are known to be robust (27–29) and are not considered representative for the whole variation in *Homo sapiens*. Nevertheless, these points do not influence the results of the hypothesis testing conducted here. Indeed, the purpose is to test if Neandertal individuals have thickened cranial bones. For the combined vault thickness in the mid-sagittal plane (Figure 3), the analysed Neandertal specimens have values within the inferior variability of European anatomically modern *Homo sapiens* and inferior to the values of the Ngandong and Sumbungmacan *Homo erectus*. When is considered the cranial vault thickness distribution all along the mid-sagittal plane, the thickness for the Neandertal specimens exceed the values obtained on the comparative sample only for a few landmarks for Krapina 3 and La Chapelle aux Saints 1 in the anterior part of the frontal bone but all the other thickness data for Neandertal individuals are within or below the represented variation of the comparison material. So, these results illustrate that Neandertal individuals cannot be considered as exhibiting thicker cranial vault than anatomically modern *Homo sapiens*, when fossil individuals are considered in this comparative sample.

In terms of variation of cranial vault thickness, the Neandertal specimens present some particularities compared to anatomically modern *Homo sapiens*. The thickness in the Neandertal individuals decreases rapidly and regularly from the area of the frontal torus to the middle part of the frontal squama, whereas this variation is less abrupt in the comparative sample, resulting in greater value for the thickness of the middle part of the frontal squama. The variation along the junction of the parietal bones shares similar characteristics for the two analysed

samples, but the Neandertals have lower value for the thickness in this area. On the occipital bone, both samples exhibit an increase of the thickness in the middle part of the occipital squama. On the opposite side, Neandertals only show a reduced increase of thickness in direction to the sagittal external occipital protuberance whereas the comparative sample has an important thickening in this area. Finally, the suprainiac fossa only induces a reduced variation of CVT and is not responsible for the important decreasing of thickness values between the middle part of the occipital squama and the occipital superstructures. So, the Neandertals are particular for the variation of the thickness in the area of the occipital superstructures whereas anatomically modern *Homo sapiens* and Ngandong and Sambungmacan *Homo erectus* share similar characteristics (45).

Additional work is needed to understand the variation of the cranial vault thickness in fossil hominids and its possible phylogenetic implications. Similar analyses and comparative studies remain to be done on all the different chronological groups of Asian *Homo erectus*, on the African, the older European and the Georgian fossils.

## CONCLUSIONS

The purpose of this study was to present new evidence about the characteristics of the Krapina fossils compared to Western European Neandertals as well as specific characters of the Neandertals about cranial vault thickness. These results illustrate that the values for the cranial vault thickness in the mid-sagittal plane for Krapina 3, Krapina 4, Krapina 5 and Krapina 6 overlap and exceed the variation observed in the La Chapelle aux Saints 1, La Ferrassie 1, La Quina H5, Spy 1 and Spy 10 Neandertals. While variable, all these individuals share a similar pattern of thickness all along the analysed plane. Based on the variation observed in this sample, the cranial vault thickness in the mid-sagittal plane appears not to be an important indicator of sex in Neandertals. Finally, our results illustrate that the Neandertals are particular for the variation of the thickness in the mid-sagittal plane whereas anatomically modern *Homo sapiens* and Ngandong and Sambungmacan *Homo erectus* share similar characteristics (45). It reflects that the cranial vault in Neandertals had overall thinner vault bones than in the comparative sample. This evidence supports previous anatomical distinctions between Neandertals and *Homo sapiens* for the skull (27, 51–53) contrary to others who argue for morphological continuity between Neandertals and *Homo sapiens* in Europe (54–56).

**Acknowledgements:** This research was supported by the TNT project (The Neandertal Tools Project), 6th Framework of the European Community. We are very thankful to the following individuals for access to fossils in their care and providing CT data: J. Radovčić, Department of Geology and Paleontology, Croatian Natural History Museum, Zagreb, Croatia; S. Bahuchet and P. Mennecier, Département Hommes, Natures, Sociétés du Muséum national d'Histoire naturelle, Paris, France; P. Semal, Laboratoire

d'Anthropologie et de Préhistoire, Institut Royal des Sciences Naturelles de Belgique, Bruxelles, Belgique; and H. de Lumley and F. Sémah, Département de Préhistoire du Muséum national d'Histoire naturelle, Paris, France. We would like to thank Professor E.A. Cabanis and J. Badawi-Fayad -CHNO des Quinze-Vingts, Paris- for the scanning procedures in Paris, and the radiology department of the University of Zagreb, hospital »Bolnica Sestara Milosrdnica«, Zagreb for the scanning procedures on the Krapina fossils. We are grateful to the technical partner (Art+com) of the TNT project for the collaborative development of the ArteCore software (<https://www.nespos.org>). Finally, we thank the editors of this volume for invitation to contribute to this special issue. J. Radovčić and D. Frayer provided invaluable comments on this paper. We are grateful to all of them.

## REFERENCES

1. WOLPOFF M H 1979 The Krapina dental remains. *Am J Phys Anthropol* 50: 67–114
2. RADOVČIĆ J, SMITH F H, TRINKAUS E, WOLPOFF M H 1988 The Krapina Hominids: An Illustrated Catalog of Skeletal Collection. Mladost Publishing House, Zagreb.
3. ROUGIER H, CREVECOEUR I, WOLPOFF M H 2006 Lower third premolar rotation in the Krapina dental sample. *Period Biol* 108: 269–278
4. GUATELLI-STEINBERG D, REID D J, BISHOP T A Did the lateral enamel of Neandertal anterior teeth grow differently from that of modern humans? *J Hum Evol* 1: 72–84
5. MINUGH-PURVIS N, RADOVČIĆ J, SMITH F H 2000 Krapina 1: a juvenile Neandertal from the early late Pleistocene of Croatia. *Am J Phys Anthropol* 111: 393–424
6. CASPARI R, RADOVČIĆ J 2006 New reconstruction of Krapina 5, a male Neandertal cranial vault from Krapina, Croatia. *Am J Phys Anthropol* 130: 394–307
7. BOCQUET-APPEL J-P, ARSUAGA J-L 1999 Age distributions of hominid samples at Atapuerca (SH) and Krapina could indicate accumulation by catastrophe. *J Archaeol Sci* 26: 327–338
8. WOLPOFF M H, CASPARI R 2006 Does Krapina reflect early Neandertal paleodemography? *Period Biol* 108: 425–432
9. THOMA A 1975 Were the Spy fossils evolutionary intermediates between classic Neandertal and modern man? *J Hum Evol* 4: 387–410
10. HUBLIN J-J 1978 Le torus occipital transverse et les structures associées: Evolution dans le genre *Homo*. Ph.D. thesis, Université Pierre et Marie Curie, Paris VI, Paris, France.
11. KENNEDY G E 1991 On the autapomorphic traits of *Homo erectus*. *J Hum Evol* 20: 375–412
12. CHURCHILL S E 1998 Cold adaptation, heterochrony, and Neandertals. *Evol Anthropol* 7: 46–61
13. THOMPSON J L 2005 Le Moustier 1 and its place among the Neandertals. In: Ullrich H (ed) The Neandertal adolescent Le Moustier 1. New aspects, new results. Staatliche Museen zu Berlin – Preußischer Kulturbesitz, Berlin, p 311–320
14. BOULE M 1911 L'Homme fossile de la Chapelle aux Saints. *Ann de Paléo* 6: 1–64; 1912 7: 65–208; 1313 8: 209–279
15. WEIDENREICH F 1943 The skull of *Sinanthropus pekinensis*: a comparative study of a primitive hominid skull. *Palaeo Sin D* 10: 1–484
16. WEIDENREICH F 1951 Morphology of Solo man. *Anthropol Papers Am Mus Nat Hist* 43: 205–290
17. LE GROS CLARK W E 1964 The endocranial cast of the Swanscombe skull. In: Ovey C D (ed) The Swanscombe skull: a survey of research on a Pleistocene site. Royal Anthropological Institute, London, p 139–144
18. SPITERY J 1982 Le frontal de l'Homme de Tautavel. 1er Congrès international de Paléontologie humaine. Nice, France, Prétirage 1, p 21–61
19. HUBLIN J J 1988 Les plus anciens représentants de la lignée préneandertalienne. In: Trinkaus E (ed) L'Homme de Neandertal, 3: l'Anatomie. E.R.A.U.L., Liège, p 75–80



20. ANTÓN S C 1997 Endocranial hyperostosis in Sangiran 2, Gibraltar 1, and Shanidar 5. *Am J Phys Anthropol* 102: 111–122
21. ARSUAGA J L, MARTÍNEZ I, GRACIA A, LORENZO C 1997 The Sima de los Huesos crania (Sierra de Atapuerca, Spain). A comparative study. *J Hum Evol* 33: 219–281
22. DEAN D, HUBLIN J J, HOLLOWAY R, ZIEGLER R 1998 On the phylogenetic position of the pre-Neandertal specimen from Reilingen, Germany. *J Hum Evol* 34: 485–508
23. LIEBERMAN D E 1995 Testing hypotheses about recent human evolution from skulls; integrating morphology, function, development, and phylogeny. *Curr Anthropol* 36: 159–197
24. LIEBERMAN D E 1996 How and why humans grow thin skulls: experimental evidence for systemic cortical robusticity. *Am J Phys Anthropol* 101: 217–236
25. GORJANOVIĆ-KRAMBERGER D 1899 Der paläolithische Mensch und sein Zeitgenossen aus dem Diluvium von Krapina in Kroatia. *Mitt Anthropol Ges Wien* 26: 65–68
26. GORJANOVIĆ-KRAMBERGER D 1906 Der diluviale Mensch von Krapina in Kroatien. Ein Beitrag zur Paläoanthropologie. In: Walkhoff O (ed) Studien über die Entwicklungsmechanik des Primatenskelletes, Volume II. Kreidel, Wiesbaden, p 59–277
27. LAHR M M 1994 The multiregional model of modern human origins: a reassessment of its morphological basis. *J Hum Evol* 26: 23–56
28. LAHR M M, WRIGHT R V S 1996 The question of robusticity and the relationship between cranial size and shape in *Homo sapiens*. *J Hum Evol* 31: 157–191
29. BALZEAU A, BADAWI-FAYAD J 2005 La morphologie externe et interne de la région supra-orbitaire est-elle corrélée à des contraintes biomécaniques? Analyses structurales des populations d'*Homo sapiens* d'Afalou Bou Rhummel (Algérie) et de Taforalt (Maroc). *Bull Mém Soc Anthropol* 17 (3–4): 185–197
30. SEMAL P, TOUSSAINT M, MAUREILLE B, ROUGIER H, CREVECOEUR I, BALZEAU A, BOUCHNEB L, LOURYAN S, DECLERCK N, RAUSIN L 2005 Numérisation des restes humains néandertaliens belges: préservation patrimoniale et exploitation scientifique. *Notae Praehist* 25: 25–38
31. MACCHIARELLI R, RADOVČIĆ J, SEMAL P, WENIGER G C 2005 A «virtual reality» for the Neandertal fossil record: «The Neandertal Tools» project. *Bull Mém Soc Anthropol (Paris)* 17: 14
32. MACCHIARELLI R, WENIGER G C 2006 NESPOS: from data accumulation to data management. Abstracts of the congress «150 years of Neandertals discoveries, continuity and discontinuity». Bonn, Germany, p 83–84
33. BADAWI-FAYAD J, YAZBECK C, BALZEAU A, NGUYEN T H, ISTOC A, GRIMAUD-HERVÉ D, CABANIS E A 2005 Multi-detector row CT scanning in Paleoanthropology at various tube current settings and scanning mode. *Surg Radiol Anat* 275: 536–543
34. BOOKSTEIN F L 1991 Morphometric tools for landmark data: geometry and biology. Cambridge University Press, Cambridge.
35. O'HIGGINS P 2000 Quantitative approaches to the study of craniofacial growth and evolution: advances in morphometric techniques. In: O'Higgins P, Cohn M (eds) Development, growth and evolution. Implications for the study of the hominid skeleton. Academic Press, London, p 165–185
36. BALZEAU A 2005 Spécificités des caractères morphologiques internes du squelette céphalique chez *Homo erectus*. Ph.D. thesis, Muséum national d'Histoire naturelle, Paris, France.
37. SPOOR F, ZONNEVELD F, MACHO G A 1993 Linear measurements of cortical bone and dental enamel by CT: applications and problems. *Am J Phys Anthropol* 91: 469–484
38. SCHWARTZ G T, THACKERAY J F, REID C, VAN REENAN J F 1998 Enamel thickness and the topography of the enamel-dentine junction in South Africa Plio-Pleistocene hominids with special reference to the Carabelli trait. *J Hum Evol* 35: 523–542
39. TODD T W 1924 Thickness of the male white cranium. *Anat Rec* 27: 45–256
40. MARTIN R 1928 Lehrbuch der Anthropologie. 2<sup>nd</sup> edition, Fisher, Jena.
41. TWIESSELMANN F 1941 Méthode pour l'évaluation de l'épaisseur des parois crâniennes. *Bull Mus Roy Hist Nat Belgique* 17: 1–33
42. ROSS A H, JANTZ R L, McCORMICK W F 1998 Cranial thickness in American females and males. *J Foren Sci* 43: 267–272
43. LYNNERUP N 2001 Cranial thickness in relation to age, sex and general body build in a Danish forensic sample. *Foren Sci Int* 117: 45–51
44. BROWN P 1994 Cranial vault thickness in Asian *Homo erectus* and *Homo sapiens*. *Cour Forsch-Inst Senckenberg* 171: 33–46
45. BALZEAU 2006 A (in press) Are thickened cranial bones and equal participations of the three structural bone layers autapomorphic traits of *Homo erectus*? *Bull Mém Soc Anthropol* 18: 145–163
46. VALLOIS H V 1937 La durée de la vie chez l'Homme fossile. *L'Anthropol* 47: 499–532
47. GENOVES S 1954 The problem of the sex of certain fossil hominids, with special reference to the Neandertal skeletons from Spy. *J R Anthropol Inst* 84: 131–144
48. SMITH F H 1976 The Neandertal remains from Krapina. *Univ Tenn Dept Anth Rprts Invests* 15: 1–359
49. HARVATI K 2003 Quantitative analysis of Neandertal temporal bone morphology using three-dimensional geometric morphometrics. *Am J Phys Anthropol* 120: 323–338
50. AHERN J C M 2006 Variation within the Krapina frontal sample and a descriptive note on the newly associated frontal specimen, Kr 27–28. *Period biol* 108: 289–297
51. STRINGER C B, HUBLIN J J, VANDERMEERSCH B 1984 The origin of anatomically modern humans in Western Europe. In: Smith H, Spencer F (eds) The origins of modern humans. Alan Liss, New York, p 51–135
52. STRINGER C B 2002 Modern human origins: progress and prospects. *Phil Trans R Soc Lond B* 357: 563–579
53. CHANG M L 2006 Taxonomic characters in middle Pleistocene systematics. *Period biol* 108: 403–415
54. SMITH F H, SIMEK J F, HARRILL M S 1989 Geographic variation in supraorbital torus reduction during the later Pleistocene (c. 80 000–15 000 BP). In: Mellars P, Stringer C (eds) The human revolution: behavioral and biological perspectives on the origins of modern humans. Edinburgh University Press, Edinburgh, p 172–193
55. FRAYER D W, WOLPOFF M H, SMITH F H, THORNE A G, POPE G G 1993 The fossil evidence for modern human origins. *Am Anthropol* 95: 399–405
56. WOLPOFF M H, HAWKS J, CASPARI R 2000 Multiregional, not multiple origins. *Am J Phys Anthropol* 112: 129–136