

ON THE STRUCTURE OF THE AUTOMORPHISM GROUP OF A MINIMAL NONABELIAN p -GROUP (METACYCLIC CASE)

IZABELA MALINOWSKA

University of Białystok, Poland

ABSTRACT. In this paper we find the complete structure for the automorphism groups of metacyclic minimal nonabelian 2-groups. This, together with [6, 7], gives the complete answer to the Question 15 from [5] (respectively Question 20 from [4]) in the case of metacyclic groups. We also correct some inaccuracies and extend the results from [13].

1. INTRODUCTION

All groups considered here are finite and the notation used is standard.

Finite p -groups are an important group class of finite groups. Since the classification of finite simple groups is finally completed, the study of finite p -groups becomes more and more active. Many leading group theorists, for example, Berkovich, Glauberman, Janko etc., turn their attention to the study of finite p -groups, see [1–4, 9, 10, 12]. Since a finite p -group has "too many" normal subgroups and, consequently, there is an extremely large number of nonisomorphic p -groups of a given fixed order, the classification of finite p -groups in the classical sense is impossible. In [1–3] Berkovich and Janko have developed some techniques for working with minimal non-abelian subgroups of finite p -groups. Roughly speaking, they show that some control over the lattice of subgroups in p -groups can be gained by considering maximal abelian subgroups together with minimal non-abelian subgroups. In [12] Janko points out that in studying the structure of non-abelian p -groups G , the minimal non-abelian subgroups of G play an important role since they generate the group G . More precisely, if A is a maximal normal abelian subgroup of G , then

2010 *Mathematics Subject Classification.* 20D45, 20D15.

Key words and phrases. Automorphisms, p -groups.

minimal non-abelian subgroups of G cover the set $G \setminus A$ (see Proposition 1.6 in [12]). A p -group G is said to be *minimal nonabelian* (for brevity, \mathcal{A}_1 -group), if G is nonabelian, but all its proper subgroups are abelian. In [5] Berkovich formulated 22 questions concerning p -groups. In Question 15 (respectively Question 20 from [4]) he proposed to describe the automorphism groups of \mathcal{A}_1 -groups. The following lemma gives the classification of \mathcal{A}_1 -groups.

LEMMA 1.1. (L. Redei) *Let G be a minimal nonabelian p -group. Then $G = \langle x, y \rangle$ and one of the following holds*

- (1) $x^{p^m} = y^{p^n} = z^p = 1$, $[x, y] = z$, $[x, z] = [y, z] = 1$, $m, n \in \mathbb{N}$, $m \geq n \geq 1$; where in case $p = 2$ we must have $m > 1$;
- (2) $x^{p^m} = y^{p^n} = 1$, $[x, y] = x^{p^{m-1}}$, $m, n \in \mathbb{N}$, $m \geq 2$, $n \geq 1$;
- (3) $a^4 = 1$, $a^2 = b^2$, $[a, b] = a^2$, $G \cong Q_8$.

In this paper we find the complete structure for the automorphism groups of metacyclic minimal nonabelian 2-groups. This, together with [6, 7], gives the complete answer to the Question 15 from [5] (respectively Question 20 from [4]) in the case of metacyclic groups. In Section 2 we generalize the results from [13] and we specify a method of finding relations in an automorphism group, that we will use in the next Sections. In the first part of Section 3 we state some results from [13], that we will use in the next part of the note, but also we specify the exact statements. Unfortunately we must point out that in Section 3 of [13] in Case A the expression " $C_K(G')$ " should be replaced by " $\Omega_{m-r}(K)$." In the end of Section 3 we state the relations in the automorphism group of a split metacyclic 2-group. In this way we remove some inaccuracies from Theorem 3.7 in [8] (see Example 1 in [13]). In Section 4 we find the complete structure of the automorphism group of a metacyclic minimal nonabelian 2-group. These relations were not considered in [8].

If L is a subgroup of a group G , then $C_{\text{Aut } G}(L)$ denotes the group of those automorphisms of G that centralize L and $N_{\text{Aut } G}(L)$ denotes the group of those automorphisms of G that normalize L . If M and N are normal subgroups of a group G , then $\text{Aut}_N(G) = C_{\text{Aut}(G)}(G/N)$ denotes the group of all automorphisms of G normalizing N and centralizing G/N . Also $\text{Aut}_N^M(G)$ denotes $\text{Aut}_N(G) \cap C_{\text{Aut } G}(M)$. If L is a subgroup of a p -group G and $l \in \mathbb{N}$ then we set $\Omega_l(L) = \langle g \in L \mid g^{p^l} = 1 \rangle$ and $\mathcal{U}_l(L) = \langle g^{p^l} \mid g \in L \rangle$.

In [15] the authors investigated the automorphism group of a semidirect product $G = H \rtimes K$. They defined the following subgroups

$$\begin{aligned} A &= \{ \theta \in \text{Aut } G \mid [K, \theta] = 1 \text{ and } H^\theta = H \}, \\ B &= \{ \theta \in \text{Aut } G \mid [H, \theta] = 1 \text{ and } [K, \theta] \subseteq H \}, \\ C &= \{ \theta \in \text{Aut } G \mid [K, \theta] = 1 \text{ and } [H, \theta] \subseteq K \}, \\ D &= \{ \theta \in \text{Aut } G \mid [H, \theta] = 1 \text{ and } K^\theta = K \}. \end{aligned}$$

By definition, we have $BD = B \rtimes D \leq C_{\text{Aut } G}(K)$ and $AC = C \rtimes A \leq C_{\text{Aut } G}(H)$.

2. CROSSED HOMOMORPHISMS AND AUTOMORPHISMS

We call an ordered triple (Q, N, θ) data if N is an abelian group, Q is a group and $\theta : Q \rightarrow \text{Aut } N$ is a homomorphism. If θ is a homomorphism of Q into $\text{Aut } N$, then Q acts on N when we define, for each $x \in Q$ and $a \in N$, a^x is the image of a under x^θ . If N is a normal subgroup of G , then the action of G/N on $Z(N)$ is given by $a^{gN} = a^{(gN)^\theta} = a^g$. Given data (Q, N, θ) a crossed homomorphism is a function $\lambda : Q \rightarrow N$ such that $(xy)^\lambda = (x^\lambda)^y y^\lambda$ for all $x, y \in Q$. We denote the set of such crossed homomorphisms by $Z^1(Q, N)$. It forms a group under the operation $q^{\lambda_1 + \lambda_2} = q^{\lambda_1} q^{\lambda_2}$; if θ is trivial, then $Z^1(Q, N) = \text{Hom}(Q, N)$.

We recall a known result ([11], Satz I,17.1) needed in the sequel:

LEMMA 2.1. *Let N be a normal subgroup of G . Then there is a natural isomorphism from $Z^1(G/N, Z(N))$ to $\text{Aut}_N^N(G)$ sending each crossed homomorphism $f : G/N \rightarrow Z(N)$ to the automorphism $\varphi_f : x \mapsto x(xN)^f$ of G .*

Lemmas 2.2–2.3 are more general versions of Lemma 2.5 and Theorem 2.6 (see also [13]).

LEMMA 2.2. *Let N be an normal subgroup of G . Let M be a normal subgroup of G such that $M \leq Z(G)$. Assume that that $L = \{\lambda \in Z^1(G/N, Z(N)) \mid (G/N)^\lambda \subseteq M\}$ and $A = N_{\text{Aut } G}(M) \cap N_{\text{Aut } G}(N)$. Then*

- (1) $A \leq \text{Aut}(G)$ and $L \leq Z^1(G/N, Z(N))$.
- (2) *If $\alpha \in A$ and $\lambda \in L$ then the function $\mu : G/N \rightarrow Z(N)$ defined by $\mu : gN \mapsto ((g^{\alpha^{-1}} N)^\lambda)^\alpha$ is a crossed homomorphism and $\mu \in L$.*

PROOF. The first part of (1) is obvious.

(2) Assume that $\alpha \in A$ and $\lambda \in L$. First let $Ng_1 = Ng_2$, then $g_2 = g_1 h$ for some $h \in N$. Then

$$(g_2 N)^\mu = ((g_2^{\alpha^{-1}} N)^\lambda)^\alpha = (((g_1 h)^{\alpha^{-1}} N)^\lambda)^\alpha = ((g_1^{\alpha^{-1}} N)^\lambda)^\alpha = (g_1 N)^\mu$$

since N is normalized by α . So μ is well defined.

Let $g_1 N, g_2 N \in G/N$. We have

$$\begin{aligned} (g_1 N \cdot g_2 N)^\mu &= (g_1 g_2 N)^\mu = (((g_1 g_2)^{\alpha^{-1}} N)^\lambda)^\alpha \\ &= ((g_1^{\alpha^{-1}} N g_2^{\alpha^{-1}} N)^\lambda)^\alpha = (((g_1^{\alpha^{-1}} N)^\lambda)^{g_2^{\alpha^{-1}}} ((g_2^{\alpha^{-1}} N)^\lambda))^\alpha \\ &= (((g_1^{\alpha^{-1}} N)^\lambda)^\alpha)^{g_2} ((g_2^{\alpha^{-1}} N)^\lambda)^\alpha = ((g_1 N)^\mu)^{g_2 N} \cdot (g_2 N)^\mu. \end{aligned}$$

It is evident that $\mu \in L$ since $(G/N)^\mu \subseteq M$. □

LEMMA 2.3. *Let G, N, M, L and A be as in Lemma 2.2. Assume that $E := \{\varphi \in \text{Aut}_N^N(G) \mid [G, \varphi] \subseteq M\}$. Then*

- (1) $E \leq \text{Aut } G$ and there is a natural isomorphism from L to E sending each crossed homomorphism $f : G/N \rightarrow M$ to the automorphism $\varphi_f : x \mapsto x(xN)^f$ of G ;
- (2) if $\alpha \in A$ and $\varphi \in E$ is determined by the crossed homomorphism $\lambda \in L$, then $\alpha^{-1}\lambda\alpha$ is determined by the crossed homomorphism $\mu \in L$ defined by $\mu : gN \mapsto ((g^{\alpha^{-1}}N)^\lambda)^\alpha$.
- (3) A normalizes E and $AE \leq \text{Aut } G$.

PROOF. (1) It is evident that $E \leq \text{Aut } G$. By definitions of M, L, E and Lemma 2.1 we get the second part of the statement.

(2)-(3) Assume that $\alpha \in A$ and $\beta \in E$. By (1) there exists $\lambda \in Z^1(G/N, Z(N))$ such that $h^\beta = h(hN)^\lambda$ ($h \in G$) and $(hN)^\lambda \in M$ for all $h \in G$. If $h \in G$ then

$$h^{\alpha^{-1}\beta\alpha} = ((h^{\alpha^{-1}})^\beta)^\alpha = (h^{\alpha^{-1}}(h^{\alpha^{-1}}N)^\lambda)^\alpha = h((h^{\alpha^{-1}}N)^\lambda)^\alpha$$

and $((h^{\alpha^{-1}}N)^\lambda)^\alpha \in M$. Hence by Lemmas 2.1 and 2.2 $\alpha^{-1}\beta\alpha \in E$, so A normalizes E . Now it is clear that $AE \leq \text{Aut } G$. \square

For the sake of completeness we recall some results from [13]. We will use them in this note.

LEMMA 2.4 ([13]). *Let N be a normal subgroup of G such that G/N is cyclic of order n . Assume that g is an element of G with $G = \langle N, g \rangle$.*

- (1) If $a \in Z(N)$ and $a^{g^{n-1}+\dots+g+1} = 1$, then the function $\lambda : G/N \rightarrow Z(N)$, defined by $(g^iN)^\lambda = a^{g^{i-1}+\dots+g+1}$ ($i \in \mathbb{N}$) and $N^\lambda = 1$, is a crossed homomorphism.
- (2) If $\lambda \in Z^1(G/N, Z(N))$ then there exists $a \in Z(N)$ such that $a^{g^{n-1}+\dots+g+1} = 1$, $(g^iN)^\lambda = a^{g^{i-1}+\dots+g+1}$ ($i \in \mathbb{N}$) and $N^\lambda = 1$.

LEMMA 2.5 ([13]). *Let G, N, g be as in Lemma 2.4. Let M be a normal subgroup of G such that $M \leq Z(N)$ and for all $a \in M$ $a^{g^{n-1}+\dots+g+1} = 1$. Assume that $L = \{\lambda \in Z^1(G/N, Z(N)) \mid (G/N)^\lambda \subseteq M\}$ and $A = N_{\text{Aut } G}(N) \cap N_{\text{Aut } G}(M)$. Then*

- (1) $A \leq \text{Aut}(G)$ and $L \leq Z^1(G/N, Z(N))$; moreover $L \cong M$.
- (2) If $\alpha \in A$ and $\lambda \in L$ then the function $\mu : G/N \rightarrow Z(N)$ defined by $\mu : hN \mapsto ((h^{\alpha^{-1}}N)^\lambda)^\alpha$ is a crossed homomorphism and $\mu \in L$.

THEOREM 2.6 ([13]). *Let G, N, L, M, g and A be as in Lemma 2.5. Assume that $E := \{\varphi \in \text{Aut}_N^N(G) \mid [G, \varphi] \subseteq M\}$. Then $E \leq \text{Aut } G$, $L \cong E \cong M$, A normalizes E , $AE \leq \text{Aut } G$ and $A \cap E \cong \{g^{-1}g^\varphi \mid \varphi \in A \cap E\}$.*

We will need the following lemma:

LEMMA 2.7. *Let G be a group, $g, h, z \in G$ and $[h, g] = z, [g, z] = 1 = [h, z]$. Assume that $i, j \in \mathbb{N}$ and $\alpha \in \text{Aut } G$. Then*

- (1) $hg^{i-1+\dots+g+1} = h^i z^{\frac{i(i-1)}{2}}$;
- (2) if $g^\alpha = g, h^\alpha = h^j, z^\alpha = z$, then $(hg^{i-1+\dots+g+1})^\alpha = h^{ij} z^{\frac{i(i-1)}{2}}$;
- (3) if $g^\alpha = g, h^\alpha = h^j, z^\alpha = z^j$, then $(hg^{i-1+\dots+g+1})^\alpha = h^{ij} z^{j\frac{i(i-1)}{2}}$;
- (4) if $g^\alpha = g^j, h^\alpha = h, z^\alpha = z^j$, then $(hg^{i-1+\dots+g+1})^\alpha = h^i z^{j\frac{i(i-1)}{2}}$;
- (5) if $g^\alpha = g^j, h^\alpha = h, z^\alpha = z$, then $(hg^{i-1+\dots+g+1})^\alpha = h^i z^{\frac{i(i-1)}{2}}$.

By Lemmas 2.3, 2.4 and 2.7 we get

LEMMA 2.8. *Let G, N, M, E, g be as in Theorem 2.6 and $i, j \in \mathbb{N}, i = j^{-1} \pmod n$. Assume that $\lambda \in Z^1(G/N, Z(N)), (gN)^\lambda = h$ for some $h \in M$ and $\beta \in E$ is an automorphism determined by λ . Assume also that $\alpha \in \text{Aut } G, [h, g] = z$ and $[g, z] = 1$. Then*

- (1) if $g^\alpha = g^j, h^\alpha = h, z^\alpha = z^j$, then $((g^{\alpha^{-1}}N)^\lambda)^\alpha = h^i z^{j\frac{i(i-1)}{2}}$;
in particular if $z = 1$, then $\beta^\alpha = \beta^i$;
- (2) if $g^\alpha = g^j, h^\alpha = h, z^\alpha = z$, then $((g^{\alpha^{-1}}N)^\lambda)^\alpha = h^i z^{\frac{i(i-1)}{2}}$;
in particular if $z = 1$, then $\beta^\alpha = \beta^i$;
- (3) if $g^\alpha = g, h^\alpha = h^j, z^\alpha = z$, then $((g^{\alpha^{-1}}N)^\lambda)^\alpha = h^j$ and $\beta^\alpha = \beta^j$.

3. A SPLIT METACYCLIC 2-GROUP

Let $G = H \rtimes K$ be a split metacyclic 2-group, where $H = \langle x \rangle$ and $K = \langle y \rangle$ and let A, B, C and D be the subgroups of $\text{Aut } G$ defined in the introduction. In this section we refer to the appropriate cases of the split metacyclic 2-groups from [8], but occasionally we repeat some known results for readers' convenience. In fact we consider only Case A.

Let $G = H \rtimes K = \langle x, y \mid x^{2^m} = y^{2^n} = 1, x^y = x^{1+2^{m-r}} \rangle$, where $m \geq 3, n \geq 1, 1 \leq r \leq \min\{m-2, n\}$.

It is convenient to consider G in the following three subcases (see [8])

- (I) $m \leq n$, (II) $n \leq m-r < m$, (III) $m-r < n < m$.

Moreover there exist two special cases. They are case (II), when $m = 2r, n = r = m-r \geq 2$ and $G = \langle x, y \mid x^{2^{2r}} = y^{2^r} = 1, x^y = x^{1+2^{2r}} \rangle$ and case (III), when $r = n > m-n \geq 2$ and $G = \langle x, y \mid x^{2^m} = y^{2^n} = 1, x^y = x^{1+2^{m-n}} \rangle$. These are referred to as exceptional cases. We will also need the following number theoretic result (see [8, 13]), which is easily established by induction.

LEMMA 3.1. *Let m, n and r be positive integers.*

- (1) For all $m \geq 2, n \geq 1, (1+2^m)^{2^n} \equiv 1+2^{m+n} \pmod{2^{2m+n-1}}$
and $(1+2^m)^{2^{n-1}} \equiv 1+2^{m+n-1} \pmod{2^{m+n}}$.

- (2) For $n \geq 2, r \geq 1$ and $m = n + r$, let $S = 1 + u + \dots + u^{2^r-1}$, where $u \equiv 1 \pmod{2^n}$. Then $S \equiv 2^r + 2^{m-1} \pmod{2^m}$ if $u \not\equiv 1 \pmod{2^{n+1}}$ and $S \equiv 2^r \pmod{2^m}$ if $u \equiv 1 \pmod{2^{n+1}}$.

Using Lemma 3.1 the following lemmas are easily established.

LEMMA 3.2.

- (1) $C_H(K) = \langle x^{2^r} \rangle$, (2) $C_K(H) = \langle y^{2^r} \rangle$,
 (3) $G' = [H, K] = \langle x^{2^{m-r}} \rangle$, (4) G is nil 2 $\Leftrightarrow 2r \leq m$.

LEMMA 3.3. $\Omega_{m-r}(K), [H, \Omega_{m-r}(K)]$ are given in the three cases as follows:

- (I) $\Omega_{m-r}(K) = \langle y^{2^{n-m+r}} \rangle \leq Z(G)$, $[H, \Omega_{m-r}(K)] = 1$;
 (II) $\Omega_{m-r}(K) = \langle y \rangle = C_K(G')$, $[H, \Omega_{m-r}(K)] = \langle x^{2^{m-r}} \rangle = G' \leq Z(G)$;
 (III) $\Omega_{m-r}(K) = \langle y^{2^{n-m+r}} \rangle \leq C_K(G')$, $[H, \Omega_{m-r}(K)] = \langle x^{2^n} \rangle \leq Z(G)$.

As in [14] when p was odd or by considering matrices of maps from [8] one could find the effect of an automorphism φ on the generators of G .

LEMMA 3.4. Let G, x, y be as above.

- (1) Assume that $n \neq r$. Then a map $\varphi : G \rightarrow G$ is an automorphism if and only if $x^{-1}x^\varphi \in \mathcal{U}_1(H)\Omega_{m-r}(K)$, $y^\varphi y^{-1} \in \Omega_n(H)C_K(H)$;
 (2) Assume that $n = r$. Then a map $\varphi : G \rightarrow G$ is an automorphism if and only if either $x^{-1}x^\varphi \in \mathcal{U}_1(H)\mathcal{U}_1(\Omega_{m-r}(K)), y^\varphi y^{-1} \in \Omega_n(H)$ or $x^{-1}x^\varphi \in \mathcal{U}_1(H)\Omega_{m-r}(K) \setminus \mathcal{U}_1(H)\mathcal{U}_1(\Omega_{m-r}(K)), y^\varphi y^{-1} \in \Omega_n(H)y^{2^{r-1}}$.

By Theorem 2.6 and the definitions of A, B and D we get the following lemma.

LEMMA 3.5. Let G, A, B, D be as above. Then

- (1) $B \cong \text{Aut}_H^H(G)$,
 (2) $AD = A \times D$ normalizes B ,
 (3) $B \cap D = 1$.

For the proofs of Theorem 3.6 and Lemma 3.7 see [13].

THEOREM 3.6. Let G be as above.

- (1) $\text{Aut } G = C_{\text{Aut } G}(H)C_{\text{Aut } G}(K)$ if and only if $r \neq n$;
 (2) $C_{\text{Aut } G}(H) = BD$;
 (3) $C_{\text{Aut } G}(K) = AC$ if and only if $m \leq n$.

We set $M := [H, \Omega_{m-r}(K)]\Omega_{m-r}(K)$, $N := G'K$ and

$$E := \{\varphi \in \text{Aut}_N^N(G) \mid [H, \varphi] \subseteq M\} \subseteq \text{Aut}_N^N(G).$$

LEMMA 3.7. Let G, M be as above and $n \neq r$.

- (1) M is abelian and normal in G .
 (2) If $a \in M$ then $a^{x^{2^{m-r-1}+\dots+x+1}} = 1$.

LEMMA 3.8. *Let G, A, D, E be as above and $n \neq r$. Then*

- (1) $E \leq \text{Aut } G$; (2) $E \cong M$;
- (3) $AD = A \times D$ normalizes E ; (4) $E \cap A \cong [H, \Omega_{m-r}(K)]$;
- (5) $C_{\text{Aut } G}(K) = AE$; (6) $D \cong \text{Aut}_{C_K(H)}(K)$.

PROOF. In the proof of Lemma 3.9 in [13] we put $\Omega_{m-r}(K)$ instead of $C_K(G')$. □

We define $c \in \text{Aut } G$ by setting $x^c = xy$, when $m - r \geq n \neq r$, and $x^c = xy^{2^{n-m+r}}$, when $m - r < n \neq r$, $y^c = y$. We also set $F := \langle c \rangle \leq E$.

THEOREM 3.9. *Let G, E, A, F be as above and $n \neq r$. Then*

- (1) $F \cong \Omega_{m-r}(K)$, $AF = AE$ and $A \cap F = 1$;
- (2) $\text{Aut } G = BD AF$ and $|\text{Aut } G| = |B||D||A||F|$.

PROOF. In the proof of Theorem 3.10 in [13] we put $\Omega_{m-r}(K)$ instead of $C_K(G')$. □

By Theorem 3.9 and Lemma 3.4 it is obvious that

THEOREM 3.10. *Let G, A, B, D, F, T be as above. Then*

- (1) $A \cong \text{Aut } H \cong C_2 \times C_{2^{m-2}}$ and $B \cong \Omega_n(H) \cong C_{2^{\min\{m,n\}}}$;
- (2) $D \cong C_K(H) \cong C_{2^{n-r}}$ except if $n > 1 = r$ when $D \cong \text{Aut } K \cong C_2 \times C_{2^{n-2}}$;
- (3) If $n \neq r$, then $F \cong \Omega_{m-r}(K) \cong C_{2^{\min\{m-r,n\}}}$;
- (4) Assume that $n = r$. Then $T \cong \Omega_{m-r}(K) \cong C_{2^{\min\{m-r,n\}}}$ except if $r = 2$ when $T \cong C_2 \times C_2$.

We define automorphisms of G on generators as follows

$$\begin{aligned} x^{a_1} &= x^{-1}, & x^{a_2} &= x^5, & y^{a_1} &= y^{a_2} = y; \\ x^b &= x, & y^b &= \begin{cases} xy, & n \geq m \\ x^{2^{m-n}}y, & n < m \end{cases}; \\ x^c &= \begin{cases} xy, & m - r \geq n, \\ xy^{2^{n-m+r}}y, & m - r < n \end{cases}, & y^c &= y. \end{aligned}$$

Now we assume that $n \neq r$ and $r \geq 2$. In this case we define

$$x^d = x, \quad y^d = y^{1+2^r}.$$

By Theorem 3.6, 3.9 and Lemma 3.8 it is clear that $\text{Aut } G = FABD$ and each automorphism φ of G can be presented uniquely as $\varphi = \alpha\beta\gamma\delta$, where $\alpha \in F, \beta \in A, \gamma \in B, \delta \in D$. It is clear that $A = \langle a_1, a_2 \rangle$, $B = \langle b \rangle$, $D = \langle d \rangle$ and AD is abelian. It is evident that $G = HK = KH$, so if $g \in G$, then $g = kh$ for some $k \in K, h \in H$. In the proof of Lemma 3.11 (2) we will use this reverse notation of elements of G .

We define i, j, k, s, t, u, w, z are such that

$$i = 0 \text{ in (I), } 5^i = 1 + 2^{m-r} \text{ mod } 2^m \text{ in (II), } 5^i = 1 + 2^n \text{ mod } 2^m \text{ in (III),}$$

$$j = 0 \text{ in (I), } 5^j = 1 - 2^{m-r+1} \text{ mod } 2^m \text{ in (II),}$$

$$5^j = 1 - 2^{n+1} \text{ mod } 2^m \text{ in (III),}$$

$$k = 1 + 2^r + 2^{m-1} \text{ in (I), } k = 1 + 2^r \text{ in (II)\&(III),}$$

$$u = 1 - 2^{n-m+r} \text{ in (I), } u = 1 - 2^{m-n} \text{ in (II), } u = 1 - 2^r \text{ in (III),}$$

$$5^t = (1 - 2^{n-1})u^{-1} \text{ mod } 2^n \text{ in (I),}$$

$$5^t = (1 - 2^{2m-r-n-1})u^{-1} \text{ mod } 2^m \text{ in (II),}$$

$$5^t = (1 - 2^{m-1})u^{-1} \text{ mod } 2^m \text{ in (III),}$$

$$s = u^{-1} \text{ mod } 2^n \text{ in (I), } s = u^{-1} \text{ mod } 2^m \text{ in (II)\&(III),}$$

$$(1 + 2^r)^w = u \text{ mod } 2^n,$$

$$z = -2^{n-m+r} + 2^{n-1} \text{ in (I), } z = -2^{m-n} + 2^{m-r+1} \text{ in (II),}$$

$$z = -2^r + 2^{n-1} \text{ in (III).}$$

LEMMA 3.11. *Let a_1, a_2, b, c, d be as above. Assume that $n \neq r$ and $r \geq 2$. Then*

- (1) $c^{a_1} = c^{-1}a_2^i, c^{a_2^{-1}} = c^5a_2^j, c^d = c^{1+2^r};$
- (2) $b^{a_1} = b^{-1}, b^{a_2} = b^5, b^{d^{-1}} = b^k;$
- (3) $c^b = c^s a_2^t b^z d^w.$

PROOF. (1) Let $N = G'K$ and $M = [H, \Omega_{m-r}(K)]\Omega_{m-r}(K)$. Then $a_1, a_2, d \in N_{\text{Aut } G}(N) \cap N_{\text{Aut } G}(M)$, $c \in \text{Aut}_N^N(G)$ and $h := x^{-1}x^c \in M$. By Lemmas 2.1, 2.4, 3.7 and 3.3 we get $g^c = g(gN)^\lambda$ ($g \in G$), $(x^i N)^\lambda = h^{x^{i-1}+\dots+x+1}$ ($i \in \mathbb{N}$). By Lemma 2.8 (3) we get the last relation. Now we use Lemma 2.8 (1) to get the first two relations: in (I) we have $[h, x] = [y^{2^{n-m+r}}, x] = 1$; in (II) since $[h, x] = [y, x] = x^{-2^{m-r}}$, we obtain

$$\begin{aligned} ((x^{a_1^{-1}}N)^\lambda)^{a_1} &= y^{-1}x^{2^{m-r}(2^m-1)(2^{m-1}-1)} = y^{-1}x^{2^{m-r}}, \\ ((x^{a_2}N)^\lambda)^{a_2^{-1}} &= y^5x^{-2^{m-r+1}}; \end{aligned}$$

in (III) since $[h, x] = [y^{2^{n-m+r}}, x] = x^{-2^n}$, by Lemma 2.8 (1) we obtain

$$((x^{a_1^{-1}}N)^\lambda)^{a_1} = x^{2^n}y^{-2^{n-m+r}}, \quad ((x^{a_2}N)^\lambda)^{a_2^{-1}} = x^{-2^{n+1}}y^{5 \cdot 2^{n-m+r}}.$$

(2) Note that $x^b = x$ and $y^b = yx^{1+2^{m-r}}$ in (I), $y^b = yx^{2^{m-n}}$ in (II), $y^b = yx^{2^{m-n}+2^{2m-n-r}}$ in (III). Let $Q = \langle x \rangle$. Then $a_1, a_2, d \in N_{\text{Aut } G}(Q)$, $b \in \text{Aut}_Q^Q(G)$ and $h := y^{-1}y^b \in Q$. By Lemmas 2.1, 2.4, 3.7 and 3.3 we get $g^b = g(gQ)^\lambda$ ($g \in G$), $(y^i Q)^\lambda = h^{y^{i-1}+\dots+y+1}$ ($i \in \mathbb{N}$). By Lemma 2.8 (3)

we obtain the first two relations. Now we use Lemma 2.8 (2) to get the last relation: in (I) since $[h, y] = [x^{1+2^{m-r}}, y] = x^{2^{m-r}(1+2^{m-r})}$ we obtain

$$\begin{aligned} ((y^d N)^\lambda)^{d^{-1}} &= (x^{1+2^{m-r}})^{1+2^r} \cdot x^{2^{m-r}(1+2^{m-r})2^{r-1}(2^r+1)} \\ &= x^{(1+2^{m-r})(1+2^r+2^{m-1})}, \end{aligned}$$

in (II) we get $[h, y] = [x^{2^{m-n}}, y] = 1$; in (III) since $[h, y] = [x^{2^{m-n}+2^{2m-n-r}}, y] = x^{2^{m-r}(2^{m-n}+2^{2m-n-r})}$ we obtain

$$\begin{aligned} ((y^d N)^\lambda)^{d^{-1}} &= (x^{2^{m-n}+2^{2m-n-r}})^{1+2^r} \cdot x^{2^{m-r}(2^{m-n}+2^{2m-n-r})(2^r+1)2^{r-1}} \\ &= (x^{2^{m-n}+2^{2m-n-r}})^{1+2^r}. \end{aligned}$$

(3) The direct computations with the help of Lemma 3.1 give the relation. \square

THEOREM 3.12. *Let G be as above and $m \geq 3$, $n \geq 1$, $1 \leq r \leq \min\{m-2, n\}$, $n \neq r$ and $r \geq 2$. Then $\text{Aut } G$ can be given by the following presentation, where the relations with commuting generators are omitted: $\text{Aut } G = \langle a_1, a_2, b, c, d \mid a_1^2 = a_2^{2^{m-2}} = b^{2^{\min\{m,n\}}} = c^{2^{\min\{m-r,n\}}} = d^{2^{n-r}} = 1, c^{a_1} = c^{-1}a_2^i, c^{a_2^{-1}} = c^5a_2^j, c^d = c^{1+2^r}, b^{a_1} = b^{-1}, b^{a_2} = b^5, b^{d^{-1}} = b^k, c^b = c^s a_2^t b^z d^w \rangle$.*

4. METACYCLIC MINIMAL NONABELIAN 2-GROUPS

In this section we will deal with groups $G = \langle x, y \mid x^{2^m} = y^{2^n} = 1, x^y = x^{1+2^{m-1}} \rangle$; where $m, n \in \mathbb{N}$, $m \geq 2, n \geq 1$. So $G = H \rtimes K$ is a split metacyclic 2-group, where $H = \langle x \rangle$ and $K = \langle y \rangle$.

First assume that $n \geq m \geq 3$. We define automorphisms of G on generators as follows

$$\begin{aligned} x^{a_1} &= x^{-1}, & x^{a_2} &= x^5, & y^{a_1} &= y^{a_2} = y; \\ x^b &= x, & y^b &= \begin{cases} xy, & n \geq m \\ x^{2^{m-n}}y, & n < m \end{cases}; \\ x^c &= \begin{cases} xy, & m > n \\ xy^{2^{n-m+1}}y, & m \leq n \end{cases}, & y^c &= y; \\ x^{d_1} &= x^{d_2} = x, & y^{d_1} &= y^{-1}, & y^{d_2} &= y^5. \end{aligned}$$

By Theorems 3.6, 3.9 and Lemma 3.8 it is clear that $\text{Aut } G = FABD$ and each automorphism φ of G can be presented uniquely as $\varphi = \alpha\beta\gamma\delta$, where $\alpha \in F, \beta \in A, \gamma \in B, \delta \in D$. It is clear that $A = \langle a_1, a_2 \rangle$, $B = \langle b \rangle$, $D = \langle d_1, d_2 \rangle$ and AD is abelian. It is evident that $G = HK = KH$, so if $g \in G$, then $g = kh$ for some $k \in K, h \in H$. In the proof of Lemma 4.1(2) we will use also this reverse notation of elements of G .

LEMMA 4.1. *Let a_1, a_2, b, c, d_1, d_2 be as above. Assume that $m \geq 3, n \geq 3$. Then*

- (1) $c^{a_1} = c^{-1}a_2^i, c^{a_2^{-1}} = c^5, c^{d_1} = c^{-1}, c^{d_2} = c^5$, where $i = 0$ when $m > n$ and $i = 2^{m-3}$ when $m \leq n$;
- (2) $b^{a_1} = b^{-1}, b^{a_2} = b^5, b^{d_1} = b^{-1}, b^{d_2^{-1}} = b^5$;
- (3) if $n - m \geq 1$, then $c^b = c^s a_2^t b^{-2^{n-m+1}} d_2^w$, where s, t, w are such that $s = 5^t = (1 - 2^{n-m+1})^{-1} \pmod{2^m}$, $5^w = 1 - 2^{n-m+1} \pmod{2^n}$;
- (4) if $m = n$, then $c^b = c^{-1} a_1 a_2^{2^{m-3}} b^{-2+2^{m-1}} d_1$;
- (5) if $m - n > 1$, then $c^b = c^s a_2^t b^{-2^{m-n}} d_2^w$, where s, t, w are such that $s = 5^t = (1 - 2^{m-n})^{-1} \pmod{2^m}$, $5^w = 1 - 2^{m-n} \pmod{2^n}$;
- (6) if $m = n + 1$, then $c^b = c^{-1} a_1 a_2^{2^{m-3}} b^{-2+2^{m-2}} d_1$.

PROOF. (1) Let $N = G'K$ and $M = [H, \Omega_{m-r}(K)]\Omega_{m-r}(K)$. Then $a_k, d_k \in N_{\text{Aut } G}(N) \cap N_{\text{Aut } G}(M)$ ($k = 1, 2$), $c \in \text{Aut}_N^N(G)$ and $h := x^{-1}x^c \in M$. By Lemmas 2.1, 2.4, 3.7 and 3.3 we get $g^c = g(gN)^\lambda$ ($g \in G$), $(x^i N)^\lambda = h^{x^{i-1} + \dots + x+1}$ ($i \in \mathbb{N}$). For the first two relations see the proof of Lemma 3.11 (1) with $r = 1$. By Lemma 2.8 (3) we obtain the last two relation.

(2) Note that $x^b = x$ and $y^b = yx^{1+2^{m-1}}$ when $n \geq m$, $y^b = yx^{2^{m-n}}$ when $m > n$. Let $Q = \langle x \rangle$. Then $a_k, d_k \in N_{\text{Aut } G}(Q)$ ($k = 1, 2$), $b \in \text{Aut}_Q^Q(G)$ and $y^{-1}y^b \in Q$. By Lemmas 2.1, 2.4, 3.7 and 3.3 we get $g^b = g(gQ)^\lambda$ ($g \in G$), $(y^i Q)^\lambda = h^{y^{i-1} + \dots + y+1}$ ($i \in \mathbb{N}$). By Lemma 2.8 (3) we obtain the first two relations. Now we will use Lemma 2.8 (2) to get the last two relations: if $m > n$ then $[y^{2^{m-n}}, y] = 1$, so we get the last two relations; if $m \leq n$, then $[x^{1+2^{m-1}}, y] = x^{2^{m-1}}$ and we get $((y^{d_1^{-1}} N)^\lambda)^{d_1} = (x^{1+2^{m-1}})^{2^n-1} x^{2^{m-1}(2^n-1)(2^{n-1}-1)} = x^{-1}$ and $((y^{d_2} N)^\lambda)^{d_2^{-1}} = (x^{1+2^{m-1}})^5 x^{2^{m-1}10} = (x^{1+2^{m-1}})^5$.

(3)-(6) The direct computations with the help of Lemma 3.1 give the relations. \square

In the next theorems the relations with commuting generators are omitted.

THEOREM 4.2. *Let G be as above and $m, n \geq 3$. Then $\text{Aut } G$ can be given by the following presentation: $\text{Aut } G = \langle a_1, a_2, b, c, d_1, d_2 \mid a_1^2 = a_2^{2^{m-2}} = b^{2^{\min\{m,n\}}} = c^{2^{\min\{m-r,n\}}} = d^{2^{n-r}} = 1, c^{a_1} = c^{-1}a_2^i, c^{a_2^{-1}} = c^5, c^{d_1} = c^{-1}, c^{d_2} = c^5, b^{a_1} = b^{-1}, b^{a_2} = b^5, b^{d_1} = b^{-1}, b^{d_2^{-1}} = b^5, c^b = \alpha \rangle$, where i is given in Lemma 4.1 and α is the appropriate relation in (3)-(4) of Lemma 4.1.*

If $m = 2$ and $n = 1$, then $G \cong \text{Aut } G$ is dihedral of order 8.

Now assume that $m > n = 2$. We define automorphisms of G on generators as follows

$$\begin{aligned} x^{a_1} &= x^{-1}, & x^{a_2} &= x^5, & y^{a_1} &= y^{a_2} = y; & x^b &= x, & y^b &= x^{2^{m-2}}y; \\ x^c &= xy, & y^c &= y; & x^d &= x, & y^d &= y^{-1}. \end{aligned}$$

THEOREM 4.3. *Let G be as above and $m > n = 2$. Then $\text{Aut } G$ can be given by the following presentation:*

- (1) *if $m > 3$, then $\text{Aut } G = \langle a_1, a_2, b, c, d \mid a_1^2 = a_2^{2^{m-2}} = b^4 = c^4 = d^2 = 1, c^{a_1} = c^{-1}a_2^{2^{m-3}}, c^d = c^{-1}, b^{a_1} = b^{-1}, b^d = b^{-1}, b^c = ba_2^t \rangle$, where $5^t = 1 - 2^{m-2} \pmod{2^m}$;*
- (2) *if $m = 3$, then $\text{Aut } G = \langle a_1, a_2, b, c, d \mid a_1^2 = a_2^2 = b^4 = c^4 = d^2 = 1, c^{a_1} = c^{-1}a_2, c^d = c^{-1}, b^{a_1} = b^{-1}, b^d = b^{-1}, c^b = c^{-1}a_1a_2d \rangle$.*

Now assume that $m \geq 3, n = 1$. We define automorphisms of G on generators as follows

$$\begin{aligned} x^{a_1} &= x^{-1}, & x^{a_2} &= x^5, & y^{a_1} &= y^{a_2} = y; \\ x^b &= x, & y^b &= x^{2^{m-1}}y; & x^c &= xy, & y^c &= y. \end{aligned}$$

THEOREM 4.4. *Let G be as above and $m \geq 3, n = 1$. Then $\text{Aut } G$ can be given by the following presentation: $\text{Aut } G = \langle a_1, a_2, b, c \mid a_1^2 = a_2^{2^{m-2}} = b^2 = c^2 = 1, c^{a_1} = ca_2^{2^{m-3}}, c^b = ca_2^{2^{m-3}} \rangle$.*

Now assume that $m = n = 2$. We define automorphisms of G on generators as follows

$$\begin{aligned} x^a &= x^{-1}, & y^a &= y; & x^b &= x, & y^b &= xy; \\ x^c &= xy^2, & y^c &= y; & x^d &= x, & y^d &= y^{-1}. \end{aligned}$$

THEOREM 4.5. *Let G be as above and $m = n = 2$. Then $\text{Aut } G$ can be given by the following presentation: $\text{Aut } G = \langle a, b, c, d \mid a^2 = b^4 = c^2 = d^2 = 1, b^a = b^{-1}, b^c = bd \rangle$.*

REFERENCES

- [1] Y. Berkovich, Groups of prime power order, vol. 1, Walter de Gruyter, 2008.
- [2] Y. Berkovich and Z. Janko, Groups of prime power order, vol. 2, Walter de Gruyter, 2008.
- [3] Y. Berkovich and Z. Janko, Groups of prime power order, vol. 3, Walter de Gruyter, 2011.
- [4] Y. Berkovich and Z. Janko, *On subgroups of finite p -groups*, Israel J. Math. **171** (2009), 29–49.
- [5] Y. Berkovich and Z. Janko, *Structure of finite p -groups with given subgroups*, Contemp. Math. **402**, 2006, 13–93.
- [6] J. N. S. Bidwell and M. J. Curran, *The automorphism group of a split metacyclic p -group*, Arch. Math. (Basel) **87** (2006), 488–497.

- [7] J. N. S. Bidwell and M. J. Curran, *Corrigendum to "The automorphism group of a split metacyclic p -group"* Arch. Math. 87 (2006), 488-497, Arch. Math. (Basel) **92** (2009), 14-18.
- [8] M. J. Curran, *The automorphism group of a split metacyclic 2-group*, Arch. Math. (Basel) **89** (2007), 10-23.
- [9] G. Glauberman, *Abelian subgroups of small index in finite p -groups*, J. Group Theory **8** (2005), 539-560.
- [10] G. Glauberman, *Centrally large subgroups of finite p -groups*, J. Algebra **300** (2006), 480-508.
- [11] B. Huppert, *Endliche Gruppen. I*, Berlin-Heidelberg-NewYork, Springer 1967.
- [12] Z. Janko, *On minimal non-abelian subgroups in finite p -groups*, J. Group Theory **12** (2009), 289-303.
- [13] I. Malinowska, *The automorphism group of a split metacyclic 2-group and some groups of crossed homomorphisms*, Arch. Math. (Basel) **93** (2009), 99-109.
- [14] F. Menegazzo, *Automorphisms of p -groups with cyclic commutator subgroup*, Rend. Sem. Mat. Univ. Padova **90** (1993), 81-101.
- [15] F. Zhou and H. Liu, *Automorphism groups of semidirect products*, Arch. Math. (Basel) **91** (2008), 193-198.

I. Malinowska
Institute of Mathematics
University of Białystok,
ul. Akademicka 2, 15-267 Białystok
Poland
E-mail: izabelam@math.uwb.edu.pl

Received: 30.11.2010.

Revised: 8.12.2010.