

THE ANCHIALINE AMPHIPODA (CRUSTACEA)
IN THE SUBTERRANEAN WATERS OF CRNA GORA
(MONTENEGRO)
(CONTRIBUTION TO THE KNOWLEDGE OF THE AMPHIPODA 261)

GORDAN S. KARAMAN*

Montenegrin Academy of Sciences and Arts, 20000 Podgorica,
Ul. Riste Stijovića 5, Montenegro

The Amphipoda (Crustacea) found in subterranean brackish waters (subterranean estuaries) in Crna Gora (Montenegro) is represented by 12+ species belonging to the families Bogidiellidae, Gammaridae, Hadziidae, Melitidae, Niphargidae, Pseudoniphargidae and Salentinellidae.

Keywords: Amphipoda, brackish waters, subterranean, cave, spring.

INTRODUCTION

Thanks to the specific geomorphologic and geological history of Crna Gora (Montenegro), as well as to the elevated precipitation (rain, snow), numerous sources of fresh water empty into the Adriatic Sea along the coast of Crna Gora [Skadar Lake through the Bojana River, numerous torrents and springs along the entire Adriatic coast of Crna Gora; some caves]. On the other hand, the karstic nature of rocks has produced numerous karstic phenomena (caves, subterranean streams and lakes, subterranean estuaries, etc.), excellent ecological conditions for the existence and diversity of numerous endemic animals, including Amphipoda.

MATERIAL & METHODS

Material was collected by various methods: the Karaman-Chappuis method, by Bou-pump, by various types of hand-nets, by use of traps, as well as by hand. Numerous data have been collected from the literature also. The collected specimens were preserved in 70 percent ethanol and studied under the microscope.

RESULTS AND DISCUSSION

Based on our investigations as well as data from literature, the present known fauna of Amphipoda collected in the subterranean brackish waters along the coast of the Adriatic Sea in Crna Gora is very rich, especially considering the relatively short coast of Crna Gora [293.5 km]. The following species were identified as being present along the coast of Crna Gora during our investigations and from the studies of other scientists (KARAMAN, S., 1950; 1953; 1955; KARAMAN, G., 1973; 1978; 2011a; 2011b, SKET, 1969; 1981, etc.):

* karaman@t-com.me

Family Bogidiellidae

Medigidiella dalmatina (Karaman, S., 1953)

Family Gammaridae

Neogammarus adriaticus Karaman, G., 1973

Rhipidogammarus karamani Stock, 1971

Family Hadziidae

Hadzia fragilis fragilis Karaman, S., 1932

Family Melitidae

Melita bulla Karaman, G., 1978

Melita valesi Karaman, S., 1955

Psammogammarus caecus Karaman, S., 1955

Family Niphargidae

Niphargus abavus Karaman, G., 2011

Niphargus kusceri Karaman, S., 1950

Niphargus spp.

Family Pseudoniphargidae

Pseudoniphargus adriaticus Karaman, S., 1955

Family Salentinellidae

Salentinella angelieri Del. Deb. & Ruffo, 1952

Among them, the species *Melita bulla*, *Melita valesi*, *Rhipidogammarus karamani* and *Neogammarus adriaticus* are semisubterranean species, because they have preserved the eyes, but they also occur in subterranean brackish waters, often together with some other, blind, anchialine amphipods.

The species *Hadzia fragilis fragilis*, described from Vjetrenica cave in Herzegovina, in pure freshwater, was collected in many localities along the Adriatic sea in brackish, anchialine waters, usually in the anchialine caves, together with some other anchialine amphipods.

The species *Salentinella angelieri* was usually collected in brackish subterranean waters (Bečići near Budva), but in Italy many populations of this species were found in pure fresh water, far from the sea [probably genetically different].

Within the members of the genus *Niphargus* collected near the coast of the Adriatic Sea in Crna Gora, only for two species have we confirmation that they were collected in freshwater as well as in subterranean brackish waters: *Niphargus kusceri* and *N. abavus*. But we believe that some other members of the genus *Niphargus*, collected in subterranean fresh water near the coast of the Adriatic sea, probably colonised brackish waters as well, like some *Niphargus* species from the North Adriatic region (SKET & KARAMAN, 1990).

Regarding endemism, it seems that only a limited number of endemic species of eastern Adriatic coast occur along the coast of Crna Gora (Montenegro): *Hadzia fragilis fragilis*, *Medigidiella dalmatina*, *Niphargus abavus*, *N. kusceri*].

Many of the new species described from the coast of Crna Gora were later discovered in other parts of the Mediterranean Sea [*Psammogammarus caecus*, *Melita valesi*, *Neogammarus adriaticus*, *Pseudoniphargus adriaticus*]. We suppose that *Melita bulla* will be discovered in some other localities along the Mediterranean Sea also.

CONCLUSION(S)

The taxonomical studies conducted by various authors showed that the subterranean brackish waters of the Adriatic coast of Crna Gora are settled by 11+ known Amphipoda species belonging to 7 families and 9 genera. We expect other freshwater species of the genus *Niphargus* to be found in the anchialine waters when ecological studies on Amphipoda of the anchialine waters of Crna Gora are undertaken.

REFERENCES

- KARAMAN, G., 1973: XLIV. Contribution to the Knowledge of the Amphipoda. On three very interesting *Neogammarus* species from Mediterranean and Adriatic Seas (Amphipoda: Gammaridae). Acta, Musei Macedonici Scientiarum Naturalium, **13**, 109–130.
- KARAMAN, G., 1978: On two Melita Species (Fam. Gammaridae) from the Mediterranean Sea, *M. bulla* n. sp. and *M. valesi* S. Kar. 1955 (Contribution to the Knowledge of the Amphipoda 85). Bollettino del Museo Civico di Storia Naturale, Verona, **5**, 221–237.
- KARAMAN, G., 2011a: Catalogue: Fauna of Gammaridean Amphipoda (Crustacea, Malacostraca) of the Adriatic Sea (Contribution to the Knowledge of the Amphipoda 252). The Montenegrin Academy of Sciences and Arts, Catalogue 2. The Section of Natural Sciences, **1**, 1–288.
- KARAMAN, G., 2011b: One new subterranean species of the genus *Niphargus* Schiödte, 1849 (family Niphargidae) from Boka Kotorska in Crna Gora (Montenegro), *Niphargus abavus*, sp. n. (Contribution to the knowledge of the Amphipoda 253). The Montenegrin Academy of Sciences and Arts, Glasnik of the Section of Natural Sciences, **19**, 197–212, 7 figs.
- KARAMAN, S., 1950: *Neogammarus rhipidiophorus* Catta iz podzemnih voda Jadranske obale. (= *Neogammarus rhipidiophorus* Catta aus unteriridischen Gewässern der Adriatischen Küste). – Srpska Akademija Nauka, Posebna Izdanja knj. 158, Odeljenje Prirodno-Matematičkih nauka, Beograd, **2**, 175–186, figs. 1–15.
- KARAMAN, S., 1953: Über subterrane Amphipoden und Isopoden des Karstes von Dubrovnik und seines Hinterlandes. Acta, Musei Macedonici Scientiarum Naturalium, Skopje, **1**, 137–167.
- KARAMAN, S., 1955: Über einige Amphipoden des Grundwasser der jugoslawischen Meeresküste. Acta, Musei Macedonici Scientiarum Naturalium, **2**, 223–242, figs. 1–51.
- SKET, B., 1969: Über die Verbreitung einiger Malacostraca (*Hadzia*, *Salentinella*, *Neogammarus*, *Jaera*) längs der jugoslawischen Adria-Küste. – Bulletin Scientifique, Conseil des Académies des sciences et des arts de la RSF de Yougoslavie, section A: Sciences naturelles, techniques et médicales, **14**, 147–148, 2 figs.
- SKET, B., 1981: Fauna of anchialine (coastal) cave waters, its origin and importance. – In: Proceeding of 8th International Congress of Speleology. Bowling Green, Kentucky, SAD: 645–647. Proc. 8th int. Congres speleol., pp. 645–647.
- SKET, B. & KARAMAN, G., 1990: *Niphargus rejici* (Amphipoda), its relatives in the Adriatic islands, and its possible relations to S.W. Asian taxa. – Stygologia, The Hague, **5**(3), 153–172.