

## Proton Affinities of Nitrogen Oxyradicals

Snježana P. Kazazić,<sup>a</sup> Leo Klasinc,<sup>a,b,\*</sup> Sean P. McGlynn,<sup>b</sup> and William A. Pryor<sup>b</sup>

<sup>a</sup>Rudjer Bošković Institute, Bijenička 54, HR-10002 Zagreb, Croatia

<sup>b</sup>Chemistry Department, Louisiana State University, Baton Rouge, LA 70803, USA

RECEIVED OCTOBER 16, 2003; REVISED DECEMBER 22, 2003; ACCEPTED DECEMBER 23, 2003

### Keywords

- atmospheric chemistry
- nitrogen oxides
- trioxonitrate radical
- *cis*- and *trans*-hydroxodinitrate radical
- dioxodinitrate(1-) radical
- proton affinity (G2) calculations

The proton affinities (PA's) of nitric oxide (NO), nitrogen dioxide (NO<sub>2</sub>), nitric acid (HNO<sub>3</sub>) as well as those of the *cis*- and *trans*-hydrogenhyponitrite (HONNO) and *cis*- and *trans*-dioxodinitrate (ONNO<sup>-</sup>) radicals were calculated by semi-empirical quantum mechanical calculations (standard Gaussian-2 procedure). The PA calculated values of NO (518.8 kJ mol<sup>-1</sup>), NO<sub>2</sub> (580.5 kJ mol<sup>-1</sup>) and HNO<sub>3</sub> (746.0 kJ mol<sup>-1</sup>) agree with the experimental values. The PA values for the NO<sub>3</sub>, ONNO<sup>-</sup> and HONNO radicals were calculated and adjusted to 580, 1330 and 770 kJ mol<sup>-1</sup>, respectively.

## INTRODUCTION

During our investigations of the protonation/proton affinity of nitrogen oxides and their conjugate acids<sup>1–4</sup> it became evident (i) that this property can be calculated reliably by advanced quantum chemical methods, and (ii) that it is also important to determine this parameter for certain critical N–O radicals. The properties of radicals usually pose problems for quantum chemical calculations, and this is compounded by the fact that experimental values are scarce and direct tests of the computations are impossible. However, the proton affinities of the typical representatives NO• and NO<sub>2</sub>• are known.<sup>5</sup> The photochemistry and reactions of NO• and NO<sub>2</sub>• with other oxy and peroxy radicals in the atmosphere are of paramount importance for tropospheric ozone production and air pollution.<sup>6–9</sup> We are particularly interested in the proton affinities of the radicals NO<sub>3</sub>•, ONNO•<sup>-</sup> and HONNO•, which are also of importance in atmospheric and combustion chemistry. Thus, NO<sub>3</sub>•, which is formed from

NO<sub>2</sub>• and ozone, represents the major nighttime oxidant in the lower atmosphere.<sup>10,11</sup> The ONNO•<sup>-</sup> is particularly interesting because calculations as well as photoelectron spectroscopy results<sup>12</sup> indicate that there are several forms of this species and that, surprisingly, the one with C<sub>2v</sub> symmetry (*i.e.*, with both oxygen atoms at the same nitrogen) is the most stable. There are numerous high quality calculations of the electronic structure of the here mentioned nitrogen oxides,<sup>13–19</sup> but not of their proton affinities. In order to gain some insight into the behavior of the proton affinity values for anions, radicals and the acid forms of nitrogen oxides, we have calculated and compared them with the existing experimental data for three related series of such species. Encouraged by the good agreement between experiment and calculation for the radicals NO• and NO<sub>2</sub>•, we believe that the predicted proton affinities for the radicals NO<sub>3</sub>•, ONNO•<sup>-</sup> and HONNO• are reliable and that they will prove useful until experimental values become available.

\* Author to whom correspondence should be addressed. (E-mail: klasinc@rudjer.irb.hr)

## CALCULATIONS

Standard Gaussian-2 (G2 procedure)<sup>20–23</sup> molecular energy calculations for 298 K were performed. The PA ( $\alpha$ )s, where  $\alpha$  represents the sites of proton attack in the numbering system of atoms in Figure 1, were calculated. Figure 1 depicts the optimized structures of some of the most stable protonated forms. This particular PA ( $\alpha$ ) procedure, which yields G2 enthalpies for 298 K, was chosen because it provides the best correspondence with experiment. We evaluated the relative stability and the optimized structures of all isomers of the protonated substrate, a tactic that provided some insight into the energetics and the fates of the adducts (*i.e.*, possible fragmentation).

## RESULTS AND DISCUSSIONS

The possibility of proton attachment to various positions, as indicated in the parentheses for individual compounds, yields different structures and PA values. The calculated PA's are given in Table I. We consider three series of compounds: anions, radicals and acids, which are mutually related by similar structures. For example, the ONNO<sup>-</sup> radical and HONNO radical are simultaneously the anion and acid radicals, respectively. Three experimental values,<sup>5</sup> the PA's for nitric acid (751.4 kJ mol<sup>-1</sup>), nitric oxide (531.8 kJ mol<sup>-1</sup>) and nitrogen dioxide (591.0 kJ mol<sup>-1</sup>) compare well with the theoretically predicted values of 746.0, 518.8 and 580.5 kJ mol<sup>-1</sup>, respectively. This agreement induces a considerable degree of confidence in the computed PA's of compounds for which experimental values are unavailable.

It is worth noting that a continuous decrease of PA is observed in the anion series whereas the radicals and acids exhibit a maximum at the middle, NO<sub>2</sub> and HNO<sub>2</sub>, species. This may be because protonation of NO<sup>•</sup> and HNO takes place preferably at the nitrogen atom; however, this is also the case of NO<sup>-</sup>. It seems that charge neutralization is the most determinative factor for the PA's in the acid series. However, if one looks only at the protonation of the oxygen atoms in single nitrogen compounds, then PA is the highest for the middle species in all the three series.

The *cis*- and *trans*- forms of the radical ONNO<sup>•-</sup> attach the proton both on nitrogen and oxygen with PA's in the range 1360–1380 kJ mol<sup>-1</sup>, which is similar to the nitrate anion. Indeed, one might expect that all protonated forms will be generated under appropriate experimental conditions. However, an isomeric NNO<sub>2</sub><sup>-</sup> radical structure also exists, as established by anion photoelectron spectroscopy<sup>24</sup> and matrix isolation experiments.<sup>25</sup> For this species, which was calculated to be of C<sub>2v</sub> symmetry, and its photoelectron spectrum studied by Arnold and Neumark,<sup>12</sup> our PA calculations predict for the N1 nitrogen atom a slightly higher value than for the two O3 and O4 oxygen atoms, *i.e.*, 1350.6 and 1344.9 kJ mol<sup>-1</sup>, respectively,




Figure 1. Structures of some of the more stable protonated forms of the compounds studied in this work. The positions of proton attack,  $\alpha$ , for which calculations were performed are designated O1, O2, N3, etc.

and no protonation of N<sub>2</sub> (calculated PA is only 968.5 kJ mol<sup>-1</sup>). Thus, these values are both lower than those found for the nitrogen and oxygen atoms protonation in the *cis*- and *trans*- forms of the isomeric ONNO<sup>•-</sup>. Because this species is just a reactive intermediate in the O<sup>-</sup> + N<sub>2</sub>O reaction,<sup>26</sup> we did not evaluate the PA of the HNNO<sub>2</sub> species. Finally, the very important HONNO radical, according to the present calculations, is preferably protonated at the nitrogen atom (N2) adjacent to the OH group and, in its most stable *trans*-form, has a very high PA value of 760.4 kJ mol<sup>-1</sup>. However, protonation of the *trans*-form at the other three positions: O4, N3 and O1 with PA's of 714.7, 707.1 and 695.0 kJ mol<sup>-1</sup>, respectively, might also occur, particularly because N2-protonation leads to a structure with a very long N–N bond of 2.2 Å, a situation that does not occur for the other three protonated forms. Very similar PA values are obtained for the N2 and N3 protonation sites of the *cis*-form (691.6 *vs.* 679.8 kJ mol<sup>-1</sup>, respectively) but, unfortunately, the calculations for O1 and O4 protonation did not converge.

In conclusion, comparison of the three available experimental values for NO, NO<sub>2</sub> and HNO<sub>3</sub> with their

TABLE I. Calculated and experimental proton affinities<sup>(a)</sup>

NO <sup>-</sup>		NO <sup>•</sup>		HNO	
↓ +H <sup>+</sup>		↓ +H <sup>+</sup>		↓ +H <sup>+</sup>	
HNO		HNO <sup>•+</sup>		H <sub>2</sub> NO <sup>+</sup>	
1516.0 (N)		518.8 (N)		693.3 (N)	
1339.3 (O)		457.5 (O)		622.0 (O)	
		Exp. 531.8			
NO <sub>2</sub> <sup>-</sup>		NO <sub>2</sub> <sup>•</sup>		HNO <sub>2</sub>	
↓ +H <sup>+</sup>		↓ +H <sup>+</sup>		↓ +H <sup>+</sup>	
HNO <sub>2</sub>		HNO <sub>2</sub> <sup>•+</sup>		H <sub>2</sub> NO <sub>2</sub> <sup>+</sup>	
1413.8 (O)		580.5 (O)		780.3 (O1)	
1379.3 (N)		435.1 (N)		621.7 (N2)	
		Exp. 591.0		648.5 (O3)	
NO <sub>3</sub> <sup>-</sup>		NO <sub>3</sub> <sup>•</sup>		HNO <sub>3</sub>	
↓ +H <sup>+</sup>		↓ +H <sup>+</sup>		↓ +H <sup>+</sup>	
HNO <sub>3</sub>		HNO <sub>3</sub> <sup>•+</sup>		H <sub>2</sub> NO <sub>3</sub> <sup>+</sup>	
1348.5		571.8 (O)		746.0 (O1)	
dissociative (N)		dissociative (N)		666.1 (O3)	
				625.5 (O4)	
				Exp. 751.4	
<sup>-</sup> ONNO <sup>-</sup>		ONNO <sup>•-</sup>		HONNO <sup>•</sup>	
↓ +H <sup>+</sup>		↓ +H <sup>+</sup>		↓ +H <sup>+</sup>	
HONNO <sup>-</sup>		HONNO <sup>•</sup>		H <sub>2</sub> ONNO <sup>•+</sup>	
<i>cis</i>	<i>trans</i>	<i>cis</i>	<i>trans</i>	<i>cis</i>	<i>trans</i>
2065.2 (O1)	2002.5 (O1)	1375.6 (O)	1355.1 (O)	691.6 (N2)	695.0 (O1)
1982.6 (N2)	1974.2 (N2)	1364.4 (N)	1384.3 (N)	679.8 (N3)	760.4 (N2)
				not converge(O)	707.1 (N3)
					714.7 (O4)
		NNO <sub>2</sub> <sup>•-</sup>			
		↓ +H <sup>+</sup>			
		HNNO <sub>2</sub> <sup>•</sup>			
		C <sub>2v</sub> -symmetry			
		1350.6 (N1)			
		968.5 (N2)			
		1344.9 (O3, O4)			

<sup>(a)</sup> The positions of protonation,  $\alpha$ , are depicted in Figure 1. All values are given in kJ mol<sup>-1</sup>.

computed counterpart values indicate a slight underestimation of ~10 kJ mol<sup>-1</sup> for the latter. Thus, the calculated proton affinities for the NO<sub>3</sub>, ONNO<sup>-</sup> and HONNO radicals should be adjusted to 580, 1380 and 770 kJ mol<sup>-1</sup>, respectively.

## REFERENCES

1. L. Klasinc, D. Srzić, Lj. Paša-Tolić, and S. Martinović, *Croat. Chem. Acta* **69** (1996) 1007–1010.
2. B. Juršić, L. Klasinc, S. Pečur, and W. A. Pryor, *Nitric Oxide: Biol. Chem.* **1** (1997) 494–501.
3. S. Kazazić, S. P. Kazazić, L. Klasinc, S. P. McGlynn, and W. A. Pryor, *Croat. Chem. Acta* **74** (2001) 271–275.
4. S. Kazazić, S. P. Kazazić, L. Klasinc, S. P. McGlynn, and W. A. Pryor, *J. Phys. Org. Chem.* **15** (2002) 728–731.
5. E. P. Hunter and S. G. Lias, *J. Phys. Chem. Ref. Data* **27** (1998) 413–656.
6. R. P. Wayne, *Chemistry of the Atmospheres*, Clarendon, Oxford, 1991.
7. J. H. Seinfeld and S. N. Pandis, *Atmospheric Chemistry and Physics*, Wiley Interscience, New York, 1998.
8. B. J. Finnlaysen-Pitts and J. N. Pitts, Jr. *Chemistry of the Upper and Lower Atmosphere*, Academic Press, San Diego, 2000.
9. P. J. Crutzen, *Pure Appl. Chem.* **70** (1998) 1319–1326.

10. S. M. Aschmann and R. Atkinson, *Atmos. Env.* **29** (1995) 2311–2316.
11. R. Atkinson, J. Arey, S. M. Aschmann, S. B. Corchnoy, and Y. H. Shu, *Int. J. Chem. Kinet.* **27** (1995) 941–955.
12. D. W. Arnold and D. M. Neumark, *J. Chem. Phys.* **102** (1995) 7035–7045.
13. L. A. Eriksson, J. Wang, R. J. Boyd, and S. Lunell, *J. Chem. Phys.* **98** (1994) 792–799.
14. R. C. Boehm and L. L. Lohr, *J. Phys. Chem.* **93** (1989) 3430–3433.
15. W. Eisfeld and K. Morokuma, *J. Chem. Phys.* **113** (2000) 5587–5597.
16. W. Eisfeld and K. Morokuma, *J. Chem. Phys.* **114** (2001) 9430–9440.
17. E. W. G. Diau, Y. He, M. C. Lin, and C. F. Melius, *Progress in Energy and Combustion Science* **21** (1995) 1–23.
18. A. L. L. East, *J. Chem. Phys.* **109** (1998) 2185–2193.
19. A. L. L. East and J. K. G. Watson, *J. Chem. Phys.* **110** (1999) 6099–6102.
20. L. A. Curtiss, K. Raghavachari, G. W. Trucks, and J. A. Pople, *J. Chem. Phys.* **94** (1991) 7221–7230.
21. L. A. Curtiss and K. Raghavachari, in: S. R. Langhoff (Ed.), *Quantum Mechanical Electronic Structure Calculations with Chemical Accuracy*, Kluwer Academic, Amsterdam, 1995.
22. K. Raghavachari and L. A. Curtiss, in: D. R. Yarkony (Ed.), *Modern Electronic Structure Theory*, World Scientific, Singapore, 1995.
23. M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, V. G. Zakrzewski, J. A. Montgomery, Jr., R. E. Stratmann, J. C. Burant, S. Dapprich, J. M. Millam, A. D. Daniels, K. N. Kudin, M. C. Strain, O. Farkas, J. Tomasi, V. Barone, M. Cossi, R. Cammi, B. Menonucci, C. Pomelli, C. Adamo, S. Clifford, J. Ochterski, G. A. Petersson, P. Y. Ayala, Q. Cui, K. Morokuma, N. Rega, P. Salvador, J. J. Dannenberg, D. K. Malick, A. D. Rabuck, K. Raghavachari, J. B. Foresman, J. Cioslowski, J. V. Ortiz, A. G. Baboul, B. B. Stefanov, G. Liu, A. Liashenko, P. Piskorz, I. Komaromi, R. Gomperts, R. L. Martin, D. J. Fox, T. Keith, M. A. Al-Laham, C. Y. Peng, A. Nanayakkara, M. Challacombe, P. M. W. Gill, B. Johnson, W. Chen, M. W. Wong, J. L. Andres, C. Gonzalez, M. Head-Gordon, E. S. Replogle, and J. A. Pople, *Gaussian 98*, Revision A.11.4, Gaussian, Inc., Pittsburgh PA, 2002.
24. L. A. Posey and M. A. Johnson, *J. Chem. Phys.* **88** (1988) 5383–5395.
25. M. E. Jacox, *J. Chem. Phys.* **93** (1990) 7622–7631.
26. J. L. Moruzzi and J. T. Dakin, *J. Chem. Phys.* **49** (1968) 5000–5006.

---

## SAŽETAK

### Protonski afiniteti dušikovih oksiradikala

Snježana Pečur Kazazić, Leo Klasinc, Sean P. McGlynn i William A. Pryor

Rabeći semi-empirijske kvantno mehaničke račune (standardna Gaussian-2 procedura) izračunani su protonski afiniteti (PA) za dušikov oksid, dušikov dioksid, nitratnu kiselinu, te za *cis*- i *trans*-hidrogenhiponitrit i *cis*- i *trans*-dioksidinitrat radikale. Izračunane PA vrijednosti za NO (518.8 kJ mol<sup>-1</sup>), NO<sub>2</sub> (580.5 kJ mol<sup>-1</sup>) i HNO<sub>3</sub> (746.0 kJ mol<sup>-1</sup>) dobro se slažu s eksperimentalnim vrijednostima. Izračunana PA vrijednost za NO<sub>3</sub> radikal je 580 kJ mol<sup>-1</sup>, za ONNO<sup>-</sup> radikal 1330 kJ mol<sup>-1</sup>, a za HONNO radikal 770 kJ mol<sup>-1</sup>.