

<b>Geologia Croatica</b>	59/1	51–63	11 Figs.	2 Tabs.		ZAGREB 2006
--------------------------	------	-------	----------	---------	--	-------------

## **Middle Miocene (Upper Badenian/Sarmatian) Palaeoecology and Evolution of the Environments in the Area of Medvednica Mt. (North Croatia)**

Davor **VRSALJKO**<sup>1</sup>, Davor **PAVELIĆ**<sup>2</sup>, Mirjana **MIKNIĆ**<sup>1</sup>, Mato **BRKIĆ**<sup>1</sup>, Marijan **KOVAČIĆ**<sup>1</sup>, Ivan **HEĆIMOVIĆ**<sup>1</sup>, Valentina **HAJEK-TADESSE**<sup>1</sup>, Radovan **AVANIĆ**<sup>1</sup> and Nenad **KURTANJEK**<sup>1</sup>

**Key words:** Middle Miocene, Palaeoecology, Environments, Stratigraphic Boundaries, Medvednica Mt., North Croatia, Central Paratethys.

### **Abstract**

In the area of Medvednica Mt., the Upper Badenian and Sarmatian deposits are divided into four facies associations: (A) deposits of a small carbonate platform represented by breccia, conglomerates, biocalcirudites, biocalcarenites and biocalclutites; (B) open-sea deposits composed of marls with intercalations of clay; (C) nearshore deposits of reduced salinity composed of conglomerates, sandstones, biocalcarenites, biocalcirudites and biocalclutites, and (D) lagoonal deposits represented by spongitic calcclutites with clay, marl and sand intercalations. Deposits of these facies associations contain numerous and very diverse fossil species with very different palaeoecological characteristics, from shallow- and deeper-water normal marine to shallow- and deeper-water environment of reduced salinity, even of fresh-water.

The transition from Late Badenian to Sarmatian deposition was characterized by three different unconformities and one conformity. Unconformities are located between different lithologies reflecting an amount of uplifting and erosion at the end of the Badenian. The occurrence of an angular unconformity suggests the influence of local tectonics.

The Middle Miocene deposition shows different local variations but generally fits with the evolution of Central Paratethys and the Pannonian Basin System.

salinity (KOCHANSKY-DEVIDE, 1944; ŠIKIĆ, 1967; ŠIKIĆ et al., 1979; BAJRAKTAREVIĆ, 1980; KOCHANSKY-DEVIDE & BAJRAKTAREVIĆ, 1981; BASCH, 1983; AVANIĆ et al., 1995; VRSALJKO et al., 1995a, b, 2005; VRSALJKO, 1999; AVANIĆ et al., 2003; PAVELIĆ et al., 2003).

The Upper Badenian and Sarmatian deposits at several localities on Medvednica Mt. illustrate specific types of sedimentation (Fig. 3). These were developed in various lithologies reflecting evolution of different environments in a relatively small area, dominated by carbonates in the Late Badenian, and siliciclastics in the Sarmatian. Additionally, the transition from the Late Badenian to the Sarmatian resulted in different types of boundary between these deposits characterized by unconformities and a conformity (Figs. 5 & 6).

Evolution of the environment may have been controlled by specifics such as characteristics of the pre-Neogene basement, erosion intensity, water energy and local tectonics. However, since the basin evolution indicates a transgressive–regressive succession in the Late Badenian and a new transgression in the Sarmatian, typical for Central Paratethys and the Pannonian Basin System, important regional external controls on sedimentation should be taken into account, including the opening and closing of an oceanic connection, climate and regional tectonic events (RÖGL & STEININGER, 1984; KOVAČ et al., 1997; HARDENBOL et al., 1998; PAVELIĆ, 2001; PAVELIĆ et al., 2003; VRSALJKO et al., 2005).

### **1. INTRODUCTION**

The Middle Miocene (Upper Badenian and Sarmatian) deposits cover a large area of north Croatia, and their lithological and stratigraphic characteristics have been studied both at the surface and in the subsurface. Their general evolutionary characteristics have been primarily based on their palaeoecological indications from the fossil fauna. In that sense, Upper Badenian sediments are characterized as marine while the Sarmatian sediments are characterized as brackish or of reduced

### **2. GEOLOGICAL SETTING**

Miocene deposits of the investigated area belong palaeogeographically to the south-western margins of Central Paratethys (Figs. 1 & 2), and geotectonically to the Pannonian Basin System (PAVELIĆ, 2001).

Miocene deposits on the Medvednica Mt. disconformably overlie Mesozoic basement, and contacts with Palaeozoic/Mesozoic basement are tectonic–erosional (ŠIKIĆ et al., 1979; BASCH, 1983; AVANIĆ et al., 2003). Triassic dolomites and limestones occur in the SW part, while the central and NE parts of Medvednica

<sup>1</sup> Croatian Geological Survey, Sachsova 2, P.O. Box 268, HR-10000 Zagreb, Croatia; e-mail: dvrsaljko@hgi-cgs.hr

<sup>2</sup> University of Zagreb, Faculty of Mining, Geology and Petroleum Engineering, Pierottijeva 6, HR-10000 Zagreb, Croatia.




Fig. 1 Palaeogeographic map of Central Paratethys: Middle Miocene (after RÖGL, 1998).

Mt. are dominated by Cretaceous deposits (magmatic-sedimentary complex).

Miocene deposits of the investigated area were divided into three parts by KOCHANSKY-DEVIDE (1944): (1) SW part called the “Dolje development”, (2) a middle part, the “Čučerje development”, and (3) the north-eastern part called the “Zelina development”. Within the “Doljanski envelopment”, the first Miocene sediments are fine-grained deposits and biogenic limestones of Badenian age transgressive over the Mesozoic basement. This succession also occurs in the northern-eastern part of Medvednica Mt., known as the “Zelina development”. In the central part of Medvednica Mt. (“Čučerje development”), the first Miocene sediments were Lower Miocene (Ottungian) freshwater sediments followed by marine (Karpatian) deposits, which are partly equivalent to the “Rzehakia layers”. These sediments were succeeded by the Badenian coarse-grained sediments, tuffitic marls and fine-grained deep marine deposits, which can be compared with isochronous Miocene deposits of the Vienna Basin (Central Paratethys), known as “Schlier”.

Upper Badenian deposits are approximately 30 to 70 m thick on the slopes of Medvednica Mt., and the maximum thickness of the Sarmatian deposits is 50 m (after BASCH, 1983).

### 3. DESCRIPTION AND INTERPRETATION OF FACIES


Four sections were studied on the slopes of Medvednica Mt. – Podsusedsko Dolje, Gornje Vrapče, Krvarić and Donje Orešje (Figs. 2, 3 & 5). The studied deposits belong to the Upper Badenian and Sarmatian, and their total thickness is 80 m. Based on their lithological characteristics and faunal content (Tables 1 & 2), these sediments were grouped into four facies associations suggesting different palaeoecological conditions (Figs. 5 & 6).

#### 3.1. Facies association A: deposits of a small carbonate platform

Deposits of this facies association (Fig. 4) were studied in the Gornje Vrapče, Krvarić and Donje Orešje columns (Fig. 5), consisting of breccia, conglomerates, biocalcirudites, biocalcarenites and biocalclutites.

Breccia and conglomerates are composed mostly of fragments of Upper Triassic dolomites and Upper Cretaceous limestones reflecting the composition of the basement. The siliciclastic content is very low.

Biocalcirudites, biocalcarenites and biocalclutites are almost structureless, but cross-bedding can be seen



**Fig. 2** Study area: (A) Croatia; (B) Northern Croatia; (C) Medvednica Mt.: location of geological columns (1 – Podsusedsko Dolje, 2 – Gornje Vrapče, 3 – Krvarić, 4 – Donje Orešje).

sporadically. These deposits consist mostly of fragments or whole skeletons of shallow marine organisms: algae *Lithotamnion* sp. which dominate, large molluscs (*Ostrea* sp., *Lucinoma borealis*, *Pecten* sp., *Isocardia cor*, *Panopea menardi*, *Conus* sp., *Chlamys elegans* and many cardiids), bryozoans, serpulids, echi-

noids, foraminifera of *Ammonia viennensis*–*Elphidium crispum* zone: *Lobatula lobatula*, *C. pseudungerianus*, *Elphidium fichtelianum*, *E. crispum* and *Ammonia viennensis*, and ostracods: *Cytheretta tenuipunctata dentata* and *Senesia* cf. *philippii*. These organisms document fully marine conditions during Late Badenian time.

SPECIES	Facies association
<b>MOLLUSCA</b>	
<i>Lucinoma borealis</i> (LINNÉ)	A, B, D
<i>Palliolium zoellikoferi</i> (BITTNER)	B, D
<i>Isocardia cor</i> (LINNÉ)	A, D
<i>Panopea menardii</i> DESHAYES	A
<i>Chlamys elegans</i> (ANDRZ.)	A, B
<i>Glycymeris cor</i> (PARTSCH)	A, B
<i>Corbula gibba</i> (OLIVI)	D
<i>Callista chione</i> (DEFRANCE)	D
<i>Ostrea</i> sp.	A
<i>Natica</i> sp.	D
<i>Venus</i> sp.	B
<b>FORAMINIFERA</b>	
<i>Ammonia viennensis</i> (d'ORBIGNY)	A
<i>Bolivina dilatata</i> REUSS	D
<i>Bulimina elongata</i> d'ORBIGNY	D
<i>Praeglobobulimina pyrula</i> (d'ORBIGNY)	B, D
<i>Globocassidulina oblonga</i> (REUSS)	B, D
<i>Lobatula lobatula</i> (WALKER & JACOB)	A
<i>Cibicidoides pseudoungerianus</i> (CUSHMAN)	A
<i>Elphidium crispum</i> (LINNÉ)	A
<i>E. fichtelianum</i> (d'ORBIGNY)	A
<i>Uvigerina semiornata</i> d'ORBIGNY	B, D
<i>U. venusta</i> FRANZENAU	B, D
<i>Pavonitina styriaca</i> SCHUBERT	B, D
<i>Heterolepa dutemplei</i> (d'ORBIGNY)	B, D
<i>Globigerina diplostoma</i> REUSS	B
<i>G. bulloides</i> d'ORBIGNY	B, D
<i>Orbulina universa</i> d'ORBIGNY	B
<i>Velapertina indigena</i> (LUCZKOWSKA)	B
<i>Pappina neudorfensis</i> (TOULA)	D
<i>Globorotalia scitula</i> (BRADY)	D
<i>Orbulina suturalis</i> BRÖNNIMANN	D
<i>Reophax</i> sp.	B, D
<i>Alveolophragmium crassum</i> (REUSS)	B, D
<i>Spirorutilus carinatus</i> (d'ORBIGNY)	D
<b>OSTRACODA</b>	
<i>Aurila angulata</i> (REUSS)	D
<i>Callistocythere canaliculata</i> (REUSS)	D
<i>Cnestocythere lamellicosta</i> TRIEBEL	D
<i>Hermanites haidingeri</i> (REUSS)	D
<i>Loxococoncha punctatella</i> (REUSS)	D
<i>L. hastata</i> (REUSS)	D
<i>Xestoleberis glabrescens</i> (REUSS)	D
<i>Senesia cf. philippii</i> (REUSS)	A
<i>Cytheretta tenuipunctata dentata</i> BRETENSKA	D, A
<b>OTHERS</b>	
Algae ( <i>Lithotamnion</i> sp.)	A, B
Hydrozoa	A, B
Bryozoa	A, B
Echinoids	A, B, D
Ostracodes	A, B, D
Fishes	A, B, D
Terrestrial flora	A, B, D

**Table 1** The important fossil association of the Late Badenian on Medvednica Mt.

The occurrence of coarse-grained sediments, i.e. breccia and conglomerates overlying the basement, and their petrographic composition, indicate erosion and reworking of the basement material during the early marine transgression, and deposition within high-energy shallow-water environments (Fig. 6).

Biocalcirudites, biocalcarenites and biocalclutites suggest lower energy conditions compared to those of the breccia and conglomerates. Their faunal association documents favourable palaeoecological conditions

SPECIES	Facies association
<b>MOLLUSCA</b>	
<i>Ervillea dissita</i> (EICHWALD)	C
<i>Musculus sarmaticus</i> GATUEV	C
<i>Modiolus</i> sp.	C
<i>Mactra</i> sp.	C
<i>Gibbula</i> sp.	C
<i>Irus</i> sp.	C
<i>Cerithium</i> sp.	C
<i>Cardium vindobonense</i> (PARTSCH)	C
<i>C. gleichenbergense</i> (PAPP)	C
<i>Planorbidae</i> sp. nov.?	D
<b>FORAMINIFERA</b>	
<i>Elphidium hauerinum</i> (d'ORBIGNY)	C
<i>E. reginum</i> (d'ORBIGNY)	C
<i>E. josephinum</i> (d'ORBIGNY)	C
<i>Rosalina obtusa</i> d'ORBIGNY	C
<i>Articulina problema</i> BOGDANOWICZ	C
<i>Sinoloculina volhynica</i> (DIDKOWSKIY)	C
<i>Sinzowella novorossica</i> (KARRER & SINZOW)	C
<i>Porosonion granosum</i> (d'ORBIGNY)	D
<b>OSTRACODA</b>	
<i>Candona cf. oblonga</i> (SARS)	D
<i>Aurila merita</i> (ZALANYI)	D
<i>A. mehesi</i> (ZALANYI)	D
<i>Loxococoncha kochi</i> (MEHES)	D
<i>Darwinula cylindrica</i> STRAUB	D
<i>Pseudocandona fertilis</i> TRIEBEL	D
<b>OTHERS</b>	
Characea; <i>Chalmasia moreleti</i> POKORNY	D
Fishes	D, C
Spongia	D, C
Serpulids	D, C
Algae	D, C
Terrestrial flora	D, C

**Table 2** The important fossil association of the Sarmatian on Medvednica Mt.

for the expansion of organisms in a shallow-marine environment formed over a carbonate basement. These organisms were reworked by strong currents as indicated by the grain-size of sediments and fragmentation of skeletons. Additionally, cross-bedding suggests deposition by traction currents generated by tides or waves.

The sedimentary environment was probably that of an isolated small carbonate platform formed over the previous land highs protected from terrestrial influence. On this platform algal banks were developed, colonized by a shallow-water fauna which was penecontemporaneously reworked and resedimented by strong currents within the same environment.

### 3.2. Facies association B: open sea deposits

These deposits were studied at the Donje Orešje locality (Fig. 7). They overlie sediments of a small carbonate platform (Figs. 5 & 6). The association is represented by marls with several intercalations of clay (Fig. 5). The bed thicknesses vary between 3 and 15 cm.

Marls are massive and strongly bioturbated, comprising a rich association of foraminifera of the *Bulimina-Bolivina* zone: *Pavonitina styriaca*, *Reophax* sp., *Cribrostomoides columbiensis*, *Alveolophragmium*




Fig. 3 Stratigraphic scheme (modified after RÖGL, 1998; HARDENBOL et al., 1998).



Fig. 4 Facies association A: deposits of a small carbonate platform; Krvarić locality.

*crassum*, *Uvigerina semiornata*, *U. venusta*, *Heterolepa dutemplei* and *Astrorhizidae*, and predominating planktonic species *Globigerina diplostoma*, *G. bulloides*, *Orbulina universa* and *Velapertina indigena* (Tables 1 & 2). Some molluscs occur sporadically, such as *Lucinoma borealis*, *Palliolium zoelikoferi*, *Glycymeris cor*, *Nassa* sp., *Venus* sp., *Chlamys* sp. and *Cardium* sp. Fragments of echinoids and bryozoans are rare. This foraminiferal association indicates deposition from the

neritic to the basinal area (about 100 m depth, according to SPEZZAFERRI et al., 2005; HOHENEGGER, 2005). In the uppermost part of this facies association agglutinated forms of foraminifera are absent, excluding *Pavonitina styriaca*. The clay intercalations may be explained as a consequence of a temporary but strong terrestrial influence. The whole aforementioned faunal association suggests an open sea environment and deposition of marls and clays during the Late Badenian.






Fig. 6 Evolution of the environments; Badenian–Sarmatian deposits (idealized profile).



Fig. 7 Facies association B: open sea deposits; Donje Orešje locality.

in the lower part of the facies association. The sorting is poor.

Biocalcarenites, biocalcrudites and biocalclutites predominate in this facies association. They are horizontal to cross-bedded. The composition is characterized by a high content of molluscs, foraminifera and

ostracods. The most frequent molluscs are *Cerithium* sp., *Gibbula* sp., *Maetra* sp., *Irus* sp., *Modiolus* sp., *Ervilia dissita*, *Cardium vindobonense*, *C. gleichenbergense* and *Musculus sarmaticus*. The foraminiferal association is represented by species of the *Elphidium reginum* zone: *Elphidium reginum*, *E. josephinum*, *Sin-*



Fig. 8 Facies association C: near-shore deposits; Krvarić locality.

*zowella novorossica* and *Rosalina obtusa*. Ostracods are *Aurila* sp. and *Cytheridea* sp. Fragments of alga occur sporadically – *Chalmasia moreletti* and characean girogonias. This fossil association indicates deposition in a shallow marine environment of reduced salinity.

Except for the Sarmatian fossils, this facies association often includes resedimented fragments of Upper Badenian algae and foraminifera. Oncoids, ooids and vadoids also occur.

Conglomerates in the lower part of the facies association indicate high-energy environments, while their structural and petrographic composition suggests strong terrestrial input and deposition in shallow water. The occurrence of the Upper Badenian fragments indicates erosion of those sediments on the land.

Biocalcarenes and biocalcrudites were probably deposited by traction currents in a shallow-water environment. Biocalclutites indicate low-energy sedimentation. Palaeoecological characteristics of the fauna additionally indicate shallow-water deposition in an environment of reduced salinity. Imperforate foraminifera *Articulina problema*, *Sinoloculina volhynica* and *Sinzowella novorossica* and encrusting foraminifera *Sinzowella caespitosa* indicate the existence of small reefs or biostromes, while ooids (oncooids, vadoids) suggest low currents and the existence of sandy bars (FRIEBE, 1994).

Strong inflow of fresh water very probably reduced the salinity of this environment. Such ecological conditions generated the development of specific organisms. Currents and oxic conditions enabled the expansion of life on the bottom on which epiphytic organisms evolved, including molluscs, foraminifera and ostracods. Most of these organisms lived as vagile benthos

which was very active on the bottom. At the same time, nubecularians built up small reefs. Species of ervilias, mactras, iruses and cardiums lived on the muddy–sandy bottom feeding from the substratum. Most of the gastropods from the family Ceritidae, and some miliolid foraminifera existed in proximity to the fresh water inflow.

#### 3.4. Facies association D: lagoonal deposits

Based on their different lithology and palaeoecology, lagoonal deposits (Fig. 9) are divided into two groups: spongitic calcclutites (Podsusedsko Dolje and Gornje Vrapče localities), and clay, marl and sand intercalations (Podsusedsko Dolje, Gornje Vrapče and Donje Orešje) (Fig. 5).

Spongitic calcclutites are horizontally bedded and intensively bioturbated with bed thicknesses between 0.1–1.5 m. Calcclutites are characterized by a high content of biogenic remnants, such as the stone cores of monaxonid and triaxonid sponges, skeletons of crabs and fish, serpulids, echinoids, and an association of molluscs and foraminifera. Some fragments of algae as well as small coal clasts also occur.

The *Spirorutilus–Praeglobobulimina–Heterolepa* foraminiferal association in spongitic calcclutites is characterized by benthic species *Semivulvulina deperdita*, *Spirorutilus carinatus*, *Pavonitina styriaca*, *Bolivina dilatata*, *Bulimina elongata*, *Uvigerina brunnensis*, *U. venusta*, *Pappina neuderfensis* and *Heterolepa dutemplei*. Planktonic species are very rare, but include *Globorotalia scitula*, *Globigerina bulloides* and *Orbulina suturalis*. The most frequent molluscs are *Lucinoma borealis*, *Isocardia cor*, *Palliolium zoelikoferi*, *Corbula*



**Fig. 9** Facies association D: lagoonal deposits; Gornje Vrapče locality.

*gibba*, *Callista chione* and *Natica* sp. The most common ostracod species include *Aurila angulata*, *Callistocythere canaliculata*, *Cnestocythere lamellicostata*, *Hermanites haidingeri*, *Loxoconcha punctatella*, *L. hastata* and *Xestoleberis glabrescens*. This fossil association documents a fully marine environment and deposition in the Late Badenian (VRSALJKO et al., 1995a; HAJEK-TADESSE, 2006).

The grain-size of the calcutites indicates deposition from suspension while the lack of coarse-grained intercalations suggests an environment protected from stronger intrabasinal influence. Benthic organisms indicate deposition on the marine bottom up to 70 m depth. The absence of strong current indicators and the rare occurrence of planktonic foraminifera can be explained as a consequence of the restricted connection with the open sea, i.e. sedimentation in a lagoon or bay.

The bottom of the lagoon was muddy-silty and most of the benthic organisms lived burrowed in the substratum. Some gastropods such as *Natica* sp. were carnivorous. There were also coprophags (serpulids, echinoids and some molluscs). A restricted marine connection enabled conditions needed for the existence of a greater quantity of planktonic foraminifera.

Clays and marls are horizontally bedded and almost laminated. Sometimes these sediments are massive. Sands occur as thin intercalations within this group of sediments. A rich fossil association is found in the marls (Table 2). An *Elphidium-Protelphidium-Porosonion* foraminiferal association is an important palaeoecological characteristic of the marls. This association is very similar to the one described in Facies association C. The most common species are *Elphidium josephinum*, *E. hauerinum*, *E. macellum*, and *Porosonion granosum*. There are also some ostracods, remnants of fish skeletons, molluscs and a terrestrial flora. Molluscs

*Ervilia* sp., *Maetra* sp. and *Cardium* sp. also occur in both Facies associations C and D, but are of smaller dimensions in the Facies association D. Ostracod species *Aurila merita*, *A. mehesi*, *Loxoconcha kochi* infer reduced salinity, while *Candona* cf. *oblonga*, *Darwinula cylindrica* and *Pseudocandona fertilis* are fresh-water species. The whole fossil association documents generally reduced salinity within Sarmatian environments. Some fresh-water ostracods may reflect stratification of the water, similar to the recent Black Sea. Characean girogonias and gastropods *Planorbidae* occur sporadically.

Clays and marls settled out from suspension while thin sandy intercalations may represent turbidites deposited from weak gravity flows. The lack of planktonic foraminifera suggests a restricted connection with the open sea (ŠIKIĆ, 1967; KOCHANSKY-DEVIDE & BAJRAKTAREVIĆ, 1981; VRSALJKO et al., 1995a). This facies association and its palaeoecological characteristics indicate deposition in a lagoon or bay, which had an input of fine-grained terrestrial material.

The foraminiferal association is strongly reduced in the number of constituent species and diversity compared to that of the Late Badenian. This can be explained by the formation of an environment of reduced salinity (Tables 1 & 2), and is supported by the observed increase in the number of ostracods and the occurrence of fresh-water species.

#### 4. CHARACTERISTICS OF THE BADENIAN-SARMATIAN BOUNDARY

The transition from the Late Badenian to the Sarmatian is very variable in the relatively small studied area of Mt. Medvednica. It is characterized by three types of unconformities, and a conformity.




Fig. 10 The Badenian–Sarmatian boundary; Krvarić locality.

The Gornje Vrapče locality was characterized by very strong erosion of the Upper Badenian deposits and their resedimentation in the beginning of the Sarmatian (Figs. 5 & 6). The unconformity is very clear, marked by desiccation cracks in the uppermost Badenian deposits and reworked fragments of marls with Upper Badenian fauna within transgressive Sarmatian deposits.

The Krvarić locality is very specific (Fig. 5). The lithology of the Upper Badenian and Sarmatian sediments is similar and represented by biocalcarenes that partly hides the unconformity. The hiatus is recognizable by the occurrence of a thin irregular clay–bauxite interbed which documents short-lived emersion before the onset of the Sarmatian transgression (Fig. 10).

In the Podsusedsko Dolje locality (Figs. 5 & 11) a typical angular disconformity between the Upper Badenian and Sarmatian is found. The dip of the Sarmatian deposits is ca. 10° less than the dip of the Upper Badenian sediments. The Sarmatian succession begins with conglomerates and sandstones which represent a transgressive lag. It is overlain by fine-grained sediments deposited in a lagoon of reduced salinity.

The boundary between the Upper Badenian and Sarmatian sediments in the Donje Orešje locality is characterized by its conformity (Fig. 5). The sedimentation style is similar in the Late Badenian to the Sarmatian. The transition is characterized by a gradual change of palaeoecological characteristics from a normal marine to reduced salinity fauna. This conformable transition indicates sedimentation in the deeper part of the basin which was not significantly affected by the sea-level fall at the end of the Badenian.

## 5. EVOLUTION OF THE ENVIRONMENTS

The onset of Late Badenian deposition is characterized by transgression and formation of a normal marine environment. This transgression was very probably a consequence of the opening of a connection between Central Paratethys and the Indo–Pacific Ocean due to eustatic sea-level rise (STEININGER et al., 1978; RÖGL & STEININGER, 1984; KOVAČ et al., 1997; HARDENBOL et al., 1998; RÖGL, 1998). Pre-existing land could have been formed due to fault-block rotation and uplifting in the Early Badenian, representing the end of the syn-rift phase (PAVELIĆ, 2001; PAVELIĆ et al., 2003). This uplift enabled strong erosion of all syn-rift deposits and the occurrence of pre-Neogene basement on the surface, including Upper Triassic dolomites and Upper Cretaceous limestones. These rocks provided favourable conditions for the evolution of small carbonate platforms on which different shallow-water organisms started to develop. Algal banks with *Lithotamnion* sp. predominated, but ecological conditions also supported the expansion of many different shallow-water organisms, such as bivalves, gastropods, bryozoans, echinoids, foraminifera etc. These carbonate platforms were protected from terrestrial influence. However, strong currents generated by tides or waves caused reworking of material on the bottom and resedimentation. Similar interpretations have been made for deposits from central Medvednica Mt., eastern Croatia and the Styrian Basin, supporting the ideas that in SW Central Paratethys, several small carbonate platforms existed in the beginning of the Late Badenian (SCHMID et al., 2001; AVANIĆ et al., 2003; PAVELIĆ et al., 2003; VRSALJKO, 2003; VRSALJKO et al., 2005).



**Fig. 11** The Badenian–Sarmatian boundary; Podsusedsko Dolje locality.

A continuation of sea-level rise generated deepening and the evolution of two different environments with fine-grained deposition. In NE Medvednica Mt. (Donje Orešje locality; Fig. 2) open marine environments were formed, characterized by the deposition of marls on the steep slope reaching depths of 100 m or more. Open marine conditions enabled the evolution of numerous and diverse planktonic foraminifera. In contrast, in SW Medvednica Mt. (Podsusedsko Dolje and Gornje Vrapče localities; Fig. 2) ca. 70 m deep lagoon or bay formed in the shallow sea. This environment was partly protected from the open sea, and enabled the existence primarily of benthic organisms, while the number and diversity of planktonic organisms was restricted (Tables 1 & 2). This difference in two contemporaneous environments could be explained by palinspastic reconstruction. The position of the Podsusedsko Dolje and Gornje Vrapče localities was close to the Dinarides, i.e. to the land which may have surrounded the lagoon or bay (Figs. 1 & 2). The Donje Orešje locality was more distal from the land, i.e. closer to the central part of Central Paratethys, and there the influence of the open sea was more important (Figs. 1 & 2). Upper Badenian deposition in the central part of Medvednica Mt. also occurred in an open marine environment, which supports this palinspastic reconstruction (AVANIĆ et al., 2003).

The end of the Badenian was characterized by sea-level fall causing shallowing. It has been explained by the beginning of the isolation of Central Paratethys from the Ocean (PAPP & SENEŠ, 1974; SENEŠ, 1974; RÖGL & STEININGER, 1984; KOVAČ et al., 1997; RÖGL, 1998; VRSALJKO, 1999, 2003; PAVELIĆ, 2001; PAVELIĆ et al., 2003; VRSALJKO et al., 2005). In north-eastern Medvednica Mt., shallowing of the open sea did not have a significant influence on sedi-

mentation, suggesting that it was deep enough to maintain relatively similar deposition. In its SW part, shallowing caused emergence and the formation of small islands, i.e. an archipelago, and was followed by a new transgression in the Sarmatian. This emergence indicates external controls on deposition. In the Gornje Vrapče locality the unconformity between the Upper Badenian and Sarmatian deposits was marked by strong erosion and resedimentation of lagoonal marl fragments up to 30 cm long, together with the first transgressive deposits. Sarmatian sediments indicate significant sea-level fall that exposed relatively deep-water Upper Badenian deposits on the surface. At the Krvarić locality, the unconformity is marked by a thin clay–bauxite interbed between shallow-water deposits, suggesting weaker erosion, and insignificant sea-level fall at the end of the Late Badenian (Figs. 5 & 10). The unconformity which occurs at the Podsusedsko Dolje locality may be used for interpretation of this differential shallowing. Since this unconformity is of an angular type, it was very probably caused by uplift due to tectonics (Figs. 5 & 11). Strong uplift which primarily caused the exposure of relatively deep, lagoonal deposits on the surface, additionally resulted in the inclination of the Upper Badenian beds. Upper Badenian sedimentation belongs to the post-rift phase, which is generally characterized by subsidence due to cooling of the lithosphere, but local tectonics may also be expected in its earliest part (KOVAČ et al., 1997; PAVELIĆ, 2001; PAVELIĆ et al., 2003). So, it can be summarized that local tectonics in the earliest post-rift phase probably controlled sedimentation in that area of Medvednica Mt. (AVANIĆ et al., 2003).

The onset of Sarmatian deposition was a result of the beginning of a transgression (ŠIKIĆ et al., 1979;

BASCH, 1983; VRSALJKO et al., 1995a, b; VRSALJKO, 1999). The connection of Central Paratethys with the Ocean was restricted, but a change of climate to more humid one, could generate sea-level rise (PIKIJA et al., 1989; FRIEBE, 1994). Erosion of terrestrial material and its resedimentation was an important characteristic during that transgression. After shallow-water deposition, continued sea-level rise formed an environment approximately 100 m deep characterized by dramatically different ecological conditions. Due to the restricted connection with the sea, salinity was reduced, resulting in the occurrence of new species, particularly benthic organisms: foraminifera, molluscs, ostracods and characeans (RÖGL & STEININGER, 1984).

## 6. CONCLUSIONS

On the slopes of Medvednica Mt., the Late Badenian was characterized by formation of fully marine conditions, and the development of small carbonate platforms over pre-existing land due to the marine transgression. The sea-level rise caused deepening and evolution of the open sea environment in its NE part, and a lagoon or bay in its SW part. The formation of the lagoon or bay may be a consequence of the proximity of the Dinarides which represented land at that time, while an open sea environment developed closer to the central part of Central Paratethys. Deposition was strongly controlled by eustacy, and was mostly protected from terrestrial influence.

At the end of the Late Badenian, local tectonics generated differential uplifting, shallowing and exposure of the Upper Badenian deposits on the surface in the SW part of Mt. Medvednica, while in its NE part sedimentation continued into the Sarmatian. The Late Badenian represents the onset of the post-rift phase, generally characterized by thermal subsidence, but tectonics still controlled sedimentation locally (RÖGL & STEININGER, 1984; PAVELIĆ, 2001; PAVELIĆ et al., 2003).

The beginning of the Sarmatian was characterized by a new transgression, erosion and resedimentation of the Upper Badenian deposits, and formation of a relatively deep-water environment of reduced salinity. Such a specific environment, influenced by the input of fine-grained terrestrial material enabled development of species of different palaeoecological characteristics.

## Acknowledgements

This work represents a part of the multidisciplinary geological investigation which was performed from 1991 to 1997 within the project entitled "Clastic-Carbonate Neogene Deposits of Medvednica Mt.", which formed part of the project "Geological Map of the Republic of Croatia, Scale 1:50000", financed by the Ministry of Science and Technology of the Republic of Croatia

(Project No. 0181001). The authors are grateful to reviewers, Stjepan ČORIĆ (Vienna) and Alfred DULAI (Budapest), who helped to improve the manuscript.

## 7. REFERENCES

- AVANIĆ, R., BRKIĆ, M., MIKNIĆ, M., ŠIMUNIĆ, AI. & PAVELIĆ, D. (1995): Gornjobadske i sarmatske naslage Gornje Kustošije [*The Upper Badenian and Sarmatian deposits of the Gornja Kustošija* – in Croatian].– In: ŠIKIĆ (ed.): Geološki vodič Medvednice, 71–73, Institute of Geology, Zagreb.
- AVANIĆ, R., KOVAČIĆ, M., PAVELIĆ, D., MIKNIĆ, M., VRSALJKO, D., BAKRAČ, K. & GALOVIĆ, I. (2003): The Middle and Upper Miocene Facies of Mt. Medvednica (Northern Croatia).– In: VLAHOVIĆ, I. & TIŠLJAR, J. (eds.): Evolution of Depositional Environments from the Palaeozoic to the Quaternary in the Karst Dinarides and the Pannonian Basin. 22nd IAS Meeting of Sedimentology, Opatija – September 17–19, 2003, Field Trip Guidebook, 167–172.
- BAJRAKTAREVIĆ, Z. (1980): Mikropaleontološka i biostratigrafska istraživanja tortona (badena) nekih lokaliteta Samoborskog gorja, Plešivice i Marija-goričkih brda (Micropaleontologic and Biostratigraphic investigation of the Tortonian (Badenian) of some localities of Samoborsko gorje Mts., Plešivica Mt. and Marija-gorička brda Mts.).– Geol. vjesnik, 31, 37–47.
- BASCH, O. (1983): Osnovna geološka karta SFRJ 1:100.000. Tumač za list Ivanić-Grad, L 33–81 (Basic geological map 1:100.000, Geology of the Ivanić-Grad sheet).– Geološki zavod Zagreb (1980), Savezni geol. zavod, Beograd, 81 p.
- FRIEBE, J.G. (1994): Serpulid–bryozoan–foraminiferal biostromes controlled by temperate climate and reduced salinity: Middle Miocene of the Styrian Basin, Austria.– Facies, 30, 51–62.
- HAJEK-TADESSE, V. (2006): Miocenski ostrakodi sjeverne Hrvatske (Miocene ostracods of Northern Croatia – in Croatian with an English Abstract).– Unpublished PhD Thesis, Faculty of Science, University of Zagreb, 179 p.
- HARDENBOL, J., THIERRY, J., FARLEY, B., JACQUIN, T. & VAIL, P. (1998): Mesozoic and Cenozoic Sequence Chronostratigraphic Framework of European Basins.– In: GRACIANSKY, P., HARDENBOL, J., JACQUIN, T. & VAIL, P. (eds.): Mesozoic and Cenozoic Sequence Stratigraphy of European Basins. SEPM Special Publication 60.
- HOHENEGGER, J. (2005): Estimation of environmental paleogradient values based on presence/absence data: a case study using benthic foraminifera for paleodepth estimation.– Palaeogeography, Palaeoclimatology, Palaeoecology, 217, 115–130.
- KOCHANSKY-DEVIDÉ, V. (1944): Fauna marinskog miocena južnog pobočja Medvednice–Zagrebačke gore (Miocene marine Fauna des südlichen Abhanges der Medvednica, Zagreber Gebirge) [*Fauna of marine Miocene of southern flanks of the Medvednica Mt. and the Zagrebačka gora Mt.* – in Croatian and German].– Vjestnik Hrv. drž. geol. zavoda i Hrv. drž. geol. muzeja, 2/3, 171–272.

- KOCHANSKY-DEVIDÉ, V. & BAJRAKTAREVIĆ, Z. (1981): Miocen (baden i sarmat) najzapadnijeg ruba Medvednice (Miozän (Baden und Sarmat) des westlichen Randes von Medvednica Gebirge, Kroatien, Jugoslawien) [*Miocene (Badenian and Sarmatian) of the westernmost rim of the Medvednica Mt.* – in Croatian and German].– Geol. vjesnik, 33, 43–48.
- KOVAČ, M., BARATH, I. & NAGYMAROSY, A. (1997): The Miocene collapse of the Alpine–Carpathian–Pannonian junction.– *Acta Geol. Hungarica*, 49, 445–458.
- PAPP, A. & SENEŠ, J. (1974): Grunzuge der Entwicklung der Fauna und die Biozonen im Sarmatien s. str. Der Zentralen Paratethys [*Characteristics of the Fauna Development and Biozones in the Sarmatian s.str. of the Central Paratethys* – in German].– In: PAPP, A., MARINESCU, F. & SENEŠ, J. (eds.): Chronostratigraphie und Neostatotypen – Miozän der Zentralen Paratethys – Sarmatien. VEDA, Slowakische Akademie der Wissenschaften, Bratislava, 1974, Bd. IV, 41–44.
- PAVELIĆ, D. (2001): Tectonostratigraphic model for the North Croatian and North Bosnian sector of the Miocene Pannonian Basin System.– *Basin Research*, 13, 359–376.
- PAVELIĆ, D., AVANIĆ, R., KOVAČIĆ, M., VRSALJKO, D. & MIKNIĆ, M. (2003): An Outline of the Evolution of the Croatian Part of the Pannonian Basin System.– In: VLAHOVIĆ, I. & TIŠLJAR, J. (eds.): Evolution of Depositional Environments from the Paleozoic to the Quaternary in the Karst Dinarides and the Pannonian Basin. 22nd IAS Meeting of Sedimentology, Opatija–September 17–19, 2003, Field Trip Guidebook, 155–161.
- PIKIJA, M., ŠIKIĆ, K., TIŠLJAR, J. & ŠIKIĆ, L. (1989): Biolititni i prateći karbonatni facijesi sarmata u području Krašić–Ozalj, središnja Hrvatska [*Sarmatian biolithites and associated carbonate deposits in the Krašić–Ozalj area, central Croatia* – in Croatian].– Geol. vjesnik, 42, 15–28.
- RÖGL, F. (1998): Paleogeographic considerations for Mediterranean and Paratethys seaways (Oligocene to Miocene).– *Ann. Naturhist. Mus. Wien*, 99A, 279–310.
- RÖGL, F. & STEININGER, F. (1984): Neogene Paratethys, Mediterranean and Indo–pacific seaways.– In: BRENCHY, P. (ed.): Fossils and Climate. Wiley and Sons, 171–200, London.
- SCHMID, H.P., HARZHAUSER, M. & KROH, A. (2001): Hypoxic events on a Middle Miocene carbonate platform of the Central Paratethys (Austria, Badenian, 14 Ma).– *Ann. Naturhist. Mus. Wien*, 102A, 1–50.
- SENEŠ, J. (1974): Palaeogeographie der Zentralen Paratethys im Sarmatien s.str. [*Palaeogeography of the Central Paratethys in the Sarmatian s.str.* – in German].– In: PAPP, A., MARINESCU, F. & SENEŠ, J. (eds.): Chronostratigraphie und Neostatotypen – Miozän der Zentralen Paratethys – Sarmatien. VEDA, Slowakische Akademie der Wissenschaften, Bratislava, 1974, Bd. IV, 140–154.
- SPEZZAFERRI, S., RÖGL, F., ČORIĆ, S. & HOHENEGGER, J. (2005): Paleoenvironmental changes and agglutinated foraminifera across the Karpatian/Badenian (early/middle Miocene) boundary in the Styrian Basin (Austria, Central Paratethys).– In: BUBIK, M. & KAMINSKI, M.A. (eds.): Proceedings of the Sixth International Workshop on Agglutinated Foraminifera. Spec. Publ., 8, 423–459.
- STEININGER, F., RÖGL, F. & MÜLLER, C. (1978): Geodynamic und palaeogeographische Entwicklung des Badenien [*Geodynamic and Palaeogeographic Development of the Badenian* – in German].– In: PAPP, A., CICHÁ, I., SENEŠ, J. & STEININGER, F. (eds.): Chronostratigraphie und Neostatotypen – Miozän der Zentralen Paratethys – Badenien. VEDA, Slowakische Akademie der Wissenschaften, Bratislava, 1978, Bd. VI, 110–127.
- ŠIKIĆ, K., BASCH, O. & ŠIMUNIĆ, A. (1979): Osnovna geološka karta SFRJ 1:100.000. Tumač za list Zagreb, L 33–80 (Basic geological map 1:100.000, Geology of the Zagreb sheet).– Institut za geološka istraživanja, Zagreb (1972), Savezni geološki zavod, Beograd.
- ŠIKIĆ, L. (1967): Torton i sarmat jugozapadnog dijela Medvednice na osnovu faune foraminifera (Torton und Sarmat des sudlichen Teiles der Medvednica auf Grund der Foraminiferenfaunen) [*Tortonian and Sarmatian of south-western part of the Medvednica Mt. on the basis of foraminifera fauna* – in Croatian and German].– Geol. vjesnik, 20, 127–135.
- VRSALJKO, D. (1999): The Pannonian palaeoecology and biostratigraphy of molluscs from Kostanjek–Medvednica Mt., Croatia.– *Geol. Croatica*, 52/1, 9–27.
- VRSALJKO, D. (2003): Biostratigrafija miocenskih naslaga Žumberačkog i Samoborskog gorja na temelju mekušaca (Biostratigraphy of Miocene deposits from Žumberačko and Samoborsko gorje Mts. on the basis of Molluscs – in Croatian, with an English Abstract).– Unpubl. PhD Thesis, Faculty of Science, University of Zagreb, 147 p.
- VRSALJKO, D., ŠIKIĆ, K., PIKIJA, M., GLOVACKI-JERNEJ, Ž. & MIKNIĆ, M. (1995a): Miocenske naslage Gornjeg Vrapča [*The Miocene deposits of the Gornje Vrapče* – in Croatian].– In: ŠIKIĆ (ed.): Geološki vodič Medvednice. Institute of Geology, Zagreb, 61–66.
- VRSALJKO, D., ŠIMUNIĆ, A. & MIKNIĆ, M. (1995b): Naslage sarmata i panona kraj Krvarića [*Sarmatian and Pannonian deposits near Krvarić* – in Croatian].– In: ŠIKIĆ (ed.): Geološki vodič Medvednice. Institute of Geology, Zagreb, 69–71.
- VRSALJKO, D., PAVELIĆ, D. & BAJRAKTAREVIĆ, Z. (2005): Stratigraphy and palaeogeography of Miocene Deposits from the Marginal Area of Žumberak Mt. and Samoborsko Gorje Mts. (Northwestern Croatia).– *Geologia Croatica*, 58/2, 133–150.

Manuscript received March 14, 2006.

Revised manuscript accepted June 9, 2006.

