

## Compactly generated rectifiable spaces or paratopological groups\*

FUCAI LIN<sup>1,†</sup>

<sup>1</sup> School of Mathematics and Statistics, Minnan Normal University, Zhangzhou 363 000, P. R. China

Received December 17, 2012; accepted June 21, 2013

---

**Abstract.** A rectifiable space (or a paratopological group)  $G$  is compactly generated if  $G = \langle K \rangle$  for some compact subset  $K$  of  $G$ . In this paper, we mainly discuss compactly generated rectifiable spaces or paratopological groups. The main results are that: (1) each  $\sigma$ -compact metrizable rectifiable space containing a dense compactly generated rectifiable subspace is compactly generated; (2) a metriable rectifiable space is compactly generated if and only if it is  $\sigma$ -compact and finitely generated modulo open sets; (3) any  $\sigma$ -compact paratopological group can be embedded as a closed paratopological subgroup in some compactly generated paratopological group. Finally, we consider generalized metric properties of compactly generated rectifiable spaces.

**AMS subject classifications:** 22A05, 22A15, 22A30, 54C05, 54E35, 54H11

**Key words:** compactly generated, rectifiable spaces, paratopological groups,  $\sigma$ -compact, perfect mappings, finitely generated modulo open sets, compactifications, metrizable spaces, quasi- $G_\delta$ -diagonal

---

### 1. Introduction

Recall that a *topological group*  $G$  is a group  $G$  with a (Hausdorff) topology such that the product map from  $G \times G$  onto  $G$  is jointly continuous and the inverse map of  $G$  onto itself associating  $x^{-1}$  with arbitrary  $x \in G$  is continuous. A *paratopological group*  $G$  is a group  $G$  with a topology such that the product maps of  $G \times G$  into  $G$  is jointly continuous. A topological space  $G$  is said to be a *rectifiable space* [4] provided that there are a surjective homeomorphism  $\varphi : G \times G \rightarrow G \times G$  and an element  $e \in G$  such that  $\pi_1 \circ \varphi = \pi_1$  and for every  $x \in G$  we have  $\varphi(x, x) = (x, e)$ , where  $\pi_1 : G \times G \rightarrow G$  is the projection to the first coordinate. If  $G$  is a rectifiable space, then  $\varphi$  is called a *rectification* on  $G$ . It is well known that rectifiable spaces and paratopological groups are all good generalizations of topological groups. It is easy to see that a topological group  $G$  with the neutral element  $e$  has a rectification  $\varphi(x, y) = (x, x^{-1}y)$ . However, there exists a paratopological group which is not a rectifiable space; Sorgenfrey line ([8, Example 1.2.2]) is such an example. Also, the 7-dimensional sphere  $S_7$  is rectifiable but not a topological group [21, § 3]. In fact, it

---

\*This work was supported by the NSFC (No. 11201414, 10971185) and the Natural Science Foundation of Fujian Province (No. 2012J05013) of China.

†Corresponding author. *Email address:* linfucai2008@aliyun.com (F. Lin)

is even not a semitopological group, because each (locally) compact semitopological group is a topological group [7]. Further, it is easy to see that both paratopological groups and rectifiable spaces are homogeneous.

Recently, the study of rectifiable spaces has become an interesting topic in topological algebra, see [1, 11, 13, 14, 15, 16, 20, 21].

## 2. Preliminaries

The following theorem was announced for the first time in [4], and the readers can see the proof in [5, 11, 20].

**Theorem 1** (see [4]). *A topological space  $G$  is rectifiable if and only if there exist an element  $e \in G$  and two continuous maps  $p : G^2 \rightarrow G$ ,  $q : G^2 \rightarrow G$  such that for any  $x \in G, y \in G$  the next identities hold:*

$$p(x, q(x, y)) = q(x, p(x, y)) = y, q(x, x) = e.$$

In fact, we can assume that  $p = \pi_2 \circ \varphi^{-1}$  and  $q = \pi_2 \circ \varphi$  in Theorem 1. If we fix a point  $x \in G$ , then  $f_x, g_x : G \rightarrow G$  defined with  $f_x(y) = p(x, y)$  and  $g_x(y) = q(x, y)$ , for each  $y \in G$ , are homeomorphisms. We denote  $f_x, g_x$  by  $p(x, G), q(x, G)$ , respectively.

If  $G$  is a rectifiable space, then we shall call the map  $p$  the multiplication on  $G$ . Moreover, sometimes we shall write  $x \cdot y$  instead of  $p(x, y)$  and  $A \cdot B$  instead of  $p(A, B)$  for any  $A, B \subset G$ . Therefore,  $q(x, y)$  is an element such that  $x \cdot q(x, y) = y$ ; since  $x \cdot e = x \cdot q(x, x) = x$  and  $x \cdot q(x, e) = e$ , it follows that  $e$  is a right neutral element for  $G$  and  $q(x, e)$  is a right inverse for  $x$ . Hence a rectifiable space  $G$  is a topological algebraic system with binary operations  $p, q$ , 0-ary operation  $e$  and identities as above. It is easy to see that this algebraic system need not satisfy the associative law about the multiplication operation  $p$ . Clearly, every topological loop is rectifiable.

If  $G$  is a rectifiable space (or a paratopological group) and  $X \subset G$ , then we use  $\langle X \rangle$  to denote the smallest rectifiable subspace of  $G$  which contains  $X$ . A set  $X$  algebraically generates  $G$  if  $G = \langle X \rangle$ .

Recall that a rectifiable space  $G$  (a paratopological group) is:

- (1)  $\sigma$ -compact if  $G = \bigcup \{K_n : n \in \mathbb{N}\}$ , where each  $K_n$  is compact, and
- (2) compactly generated if  $G = \langle K \rangle$  for some compact subset  $K$  of  $G$ .

**Note 1.** (a): *Obviously, each compactly generated rectifiable space is  $\sigma$ -compact. However, there exists a compactly generated paratopological group which is not  $\sigma$ -compact. Indeed, let  $X$  be an uncountable compact space, and let  $AP(X)$  be a free Abelian paratopological group. Then  $-X$  is closed discrete in  $AP(X)$  [17], which implies that  $AP(X)$  is not  $\sigma$ -compact. Moreover,  $AP(X)$  is not a topological group.*

(b): *There exists a countable, metrizable, and compactly generated paratopological group which is not a topological group. Indeed, let the rational number  $\mathbb{Q}$  with the subspace topology of Sorgenfrey line. Then  $\mathbb{Q}$  is a countable, metrizable paratopological group which is not a topological group. Put  $\mathcal{S} = \{0\} \cup \{\frac{1}{n} : n \in \mathbb{N}\}$ ; then  $Q = \langle \mathcal{S} \rangle$ . Therefore,  $\mathbb{Q}$  is compactly generated.*

(c): Sorgenfrey line is not a compactly generated paratopological group since each compact subset of Sorgenfrey line is countable [2, 3.3.b].

All spaces considered in this paper are supposed to be  $T_1$  and regular unless stated otherwise. The notation  $\mathbb{N}$  denotes the set of all positive integer numbers. The letter  $e$  denotes the neutral element of a group or the right neutral element of a rectifiable space. Readers may refer to [2, 8, 10] for notations and terminology not explicitly given here.

### 3. Compactly generated rectifiable spaces

In this section, we mainly discuss compactly generated rectifiable spaces. Firstly, we give some technical lemmas.

**Lemma 1** (see [9]). *Let  $\{U_n : n \in \mathbb{N}\}$  be a local base at point  $e$  of a topological space  $G$  such that  $\overline{U_{n+1}} \subset U_n$  for all  $n \in \mathbb{N}$ . Assume that  $\{F_n : n \in \mathbb{N}\}$  is a sequence of subsets of  $G$  such that*

1. *each  $F_n$  is compact, and*
2.  *$F_n \subset \overline{U_n}$ .*

*Then  $K = \bigcup\{F_n : n \in \mathbb{N}\} \cup \{e\}$  is compact. Moreover, if each  $F_n$  is finite, then for each enumeration  $i : \mathbb{N} \rightarrow K$  a sequence  $\{i(n) : n \in \mathbb{N}\}$  converges to  $e$ .*

Let  $A$  be a subspace of a rectifiable space  $G$ . Then  $A$  is called a *rectifiable subspace* [14] of  $G$  if we have  $p(A, A) \subset A$  and  $q(A, A) \subset A$ .

**Lemma 2** (see [14]). *Let  $G$  be a rectifiable space. If  $V$  is an open rectifiable subspace of  $G$ , then  $V$  is closed in  $G$ .*

**Lemma 3.** *Let  $H$  be a dense rectifiable subspace of a rectifiable space  $G$ . Then for each open rectifiable subspace  $E$  of  $H$  there exists an open rectifiable subspace  $E'$  of  $G$  such that  $E' \cap H = E$ .*

**Proof.** Let

$$E' = \bigcup\{V : V \text{ is open in } G \text{ and } \text{cl}_G(V) \cap H \subset E\}.$$

Obviously,  $E'$  is an open subset of  $G$  and  $E' \cap H = E$ . Now, we shall prove that  $E'$  is a rectifiable subspace of  $G$ .

Indeed, suppose that  $a, b \in E'$ . It follows from the definition of  $E'$  that there exist open sets  $U$  and  $V$  in  $G$  such that  $a \in U, b \in V, \text{cl}_G(U) \cap H \subset E$  and  $\text{cl}_G(V) \cap H \subset E$ . By the density of  $H$  in  $G$ , we have  $\text{cl}_G(U \cap H) = \text{cl}_G(U)$  and  $\text{cl}_G(V \cap H) = \text{cl}_G(V)$ . Therefore, it follows from the continuity of  $p$  in  $G$  that

$$p(U, V) \subset p(\text{cl}_G(U), \text{cl}_G(V)) = p(\text{cl}_G(U \cap H), \text{cl}_G(V \cap H)) \subset \text{cl}_G(p(U \cap H, V \cap H)).$$

Then we have  $\text{cl}_G(p(U, V)) = \text{cl}_G(p(U \cap H, V \cap H))$ , and

$$\text{cl}_G(p(U, V)) \cap H = \text{cl}_G(p(U \cap H, V \cap H)) \cap H \subset \text{cl}_G(p(E, E)) \cap H = \text{cl}_G(E) \cap H = E$$

since  $E$  is closed in  $H$  by Lemma 2. Therefore,  $p(a, b) \in p(U, V) \subset E'$ .

Suppose that  $c, d \in E'$ . Then there exist open sets  $O, W$  in  $G$  such that  $c \in O, d \in W, \text{cl}_G(O) \cap H \subset E$  and  $\text{cl}_G(W) \cap H \subset E$ . Obviously,  $q(O, W)$  is open in  $G$ . Moreover, it is also easy to see that  $\text{cl}_G(q(O, W)) = \text{cl}_G(q(O \cap H, W \cap H))$ . Since

$$\text{cl}_G(q(O, W)) \cap H = \text{cl}_G(q(O \cap H, W \cap H)) \cap H \subset \text{cl}_G(q(E, E)) \cap H \subset \text{cl}_G(E) \cap H = E,$$

it follows that  $q(c, d) \in q(O, W) \subset E'$ .  $\square$

**Corollary 1.** *A dense rectifiable subspace of a connected rectifiable space has no proper open rectifiable subspaces.*

**Proof.** By Lemma 2, each open rectifiable subspace of a rectifiable space is closed, and therefore, a connected rectifiable space cannot have proper open rectifiable subspaces. Now the result follows from Lemma 3.  $\square$

**Lemma 4** (see [14]). *Let  $G$  be a rectifiable space. If  $Y$  is a dense subset of  $G$  and  $U$  is an open neighborhood of the right neutral element  $e$  of  $G$ , then  $G = Y \cdot U$ .*

**Theorem 2.** *If a  $\sigma$ -compact metrizable rectifiable space  $G$  contains a dense compactly generated rectifiable subspace, then  $G$  is also compactly generated.*

**Proof.** Let  $H$  be a dense rectifiable subspace of  $G$  such that  $H$  is generated by some compact set  $E$ , and let  $G = \bigcup \{K_n : n \in \mathbb{N}\}$ , where each  $K_n$  is compact. Since  $G$  is metrizable, the point  $e$  has a countable local base  $\{U_n : n \in \mathbb{N}\}$ , where  $\overline{U_{n+1}} \subset U_n$  for each  $n \in \mathbb{N}$ . By the density of  $H$  in  $G$ , it follows from Lemma 4 that  $p(H, U_n) = G$  for each  $n \in \mathbb{N}$ . For each  $n \in \mathbb{N}$ , there exists a finite subset  $F_n$  of  $H$  such that  $K_n \subset p(F_n, U_n)$ , and put  $L_n = \overline{U_n} \cap q(F_n, K_n)$ , then each  $K_n \subset p(F_n, L_n)$  since  $K_n \subset p(F_n, q(F_n, K_n))$ . Obviously, each  $L_n$  is compact and, by Lemma 1,  $L = \bigcup \{L_n : n \in \mathbb{N}\}$  is also compact. Therefore,

$$G = \bigcup \{K_n : n \in \mathbb{N}\} \subset \bigcup \{p(F_n, L_n) : n \in \mathbb{N}\} \subset \bigcup \{p(H, L_n) : n \in \mathbb{N}\} \subset p(H, L).$$

Since  $H$  is generated by  $E$ ,  $G$  is generated by the compact set  $E \cup L$ . Therefore,  $G$  is compactly generated.  $\square$

**Corollary 2.** *If a  $\sigma$ -compact metrizable rectifiable space  $G$  contains a dense finitely generated rectifiable subspace, then  $G$  is also compactly generated.*

Next, we define the notion of finitely generated modulo open sets in rectifiable spaces which contains all compactly generated rectifiable spaces.

**Definition 1.** *We will say that a rectifiable space (or a paratopological group)  $G$  is finitely generated modulo open sets if for each non-empty open rectifiable subspace  $H$  of  $G$  there exists a finite subset  $F$  of  $G$  such that  $G = \langle F \cup H \rangle$ .*

**Proposition 1.** *Let  $G$  be a rectifiable space. Then the following conditions are equivalent:*

1.  $G$  is finitely generated modulo open sets;

2. for each non-empty open subset  $V$  of  $G$  there exists a finite subset  $F$  of  $G$  such that  $G = \langle F \cup V \rangle$ .

**Proof.** Obviously, (2) $\Rightarrow$ (1).

(1) $\Rightarrow$ (2). Let  $V$  be a non-empty open subset  $V$  of  $G$ . Let  $H$  be the rectifiable subspace generated by  $V$ , that is,  $H = \langle V \rangle$ . Obviously,  $H$  is open in  $G$ , and so by (2) there is a finite set  $F \subset G$  such that

$$G = \langle F \cup H \rangle = \langle F \cup \langle V \rangle \rangle = \langle F \cup V \rangle.$$

□

**Theorem 3.** *If a rectifiable space  $G$  is compactly generated, then it is finitely generated modulo open sets.*

**Proof.** Assume that  $G = \langle K \rangle$ , where  $K$  is a compact set. Let  $H$  be an open rectifiable subspace of  $G$ . Then  $\mathcal{H} = \{g \cdot H : g \in G\}$  is an open covering of  $G$ . Since  $K$  is compact, there exist finitely many elements of  $\mathcal{H}$ , say  $g_1 \cdot H, \dots, g_n \cdot H$ , which cover  $K$ . Put  $F = \{g_1, \dots, g_n\}$ . Then  $G = \langle F \cup H \rangle$ . □

**Theorem 4.** *Let  $G$  be a metrizable rectifiable space  $G$  and  $A$  a countable subset of  $G$ . Suppose that  $G$  is finitely generated modulo open sets. Then  $G$  contains a sequence  $\mathcal{S}$  converging to  $e$  of  $G$  such that  $A \subset \langle \mathcal{S} \rangle$ .*

**Proof.** Let  $A = \{a_n : n \in \omega\}$ . Since  $G$  is metrizable, let  $\{U_n : n \in \omega\}$  be a local base at  $e$  such that

$$G = U_0 \supseteq U_1 \supseteq \dots \supseteq U_n \supseteq \dots$$

Since  $G$  is finitely generated modulo open sets, for each  $n \in \omega$  we can fix a finite set  $F_n$  such that  $G = \langle F_n \cup U_{n+1} \rangle$ .

By induction on  $n$ , we will define a sequence  $\{B_n : n \in \omega\}$  of finite subsets of  $G$  with the following properties:

- (a)  $B_n \subset U_n$ ;
- (b)  $G = \langle B_0 \cup B_1 \cup \dots \cup B_n \cup U_{n+1} \rangle$ , and
- (c)  $a_n \in \langle B_0 \cup B_1 \cup \dots \cup B_n \rangle$ .

To begin with, let  $B_0 = F_0 \cup \{a_0\}$ ; then  $B_0$  satisfies all three conditions (a)-(c). Assume that we have already defined finite sets  $B_0, B_1, \dots, B_{n-1}$  satisfying all three conditions (a)-(c). By (b),  $F_n \cup \{a_n\} \subset \langle B_0 \cup B_1 \cup \dots \cup B_{n-1} \cup U_n \rangle$ . Since  $F_n$  is finite, we can find a finite set  $B_n \subset U_n$  such that

$$F_n \cup \{a_n\} \subset \langle B_0 \cup B_1 \cup \dots \cup B_{n-1} \cup B_n \rangle.$$

Clearly, (a)-(c) are satisfied.

Put  $\mathcal{S} = \cup\{B_n : n \in \omega\}$ . By (c),  $A \subset \langle \mathcal{S} \rangle$ . By Lemma 1 and (a),  $\mathcal{S}$  can be enumerated as a sequence converging to  $e$ . □

**Theorem 5.** *Let  $G$  be a  $\sigma$ -compact metrizable rectifiable space  $G$ . Then  $G$  is compactly generated if and only if  $G$  is finitely generated modulo open sets.*

**Proof.** By Theorem 3, we only need to prove the sufficiency. Suppose that for each open rectifiable subspace  $H$  of  $G$  there exists a finite set  $F$  such that  $G = \langle F \cup H \rangle$ . Obviously,  $G$  is separable, and let  $D$  be a countable dense subset of  $G$ . By Theorem 4,  $G$  has a dense compactly generated rectifiable subspace, and by Theorem 2,  $G$  is compactly generated.  $\square$

**Corollary 3.** *A metrizable rectifiable space  $G$  is compactly generated if and only if  $G$  is  $\sigma$ -compact and finitely generated modulo open sets.*

A rectifiable space without proper open rectifiable subspaces trivially satisfies condition (2) of Proposition 1. Therefore, we have the following corollary.

**Corollary 4.** *A  $\sigma$ -compact metrizable rectifiable space  $G$  without proper open rectifiable subspaces is compactly generated.*

By Corollaries 1 and 4, we also have the following corollary.

**Corollary 5.** *A  $\sigma$ -compact dense rectifiable subspace of a connected metrizable rectifiable space  $G$  is compactly generated.*

**Corollary 6.** *A  $\sigma$ -compact connected metrizable rectifiable space  $G$  is compactly generated.*

By Theorems 3 and 4, it is easy to prove the following theorem.

**Theorem 6.** *A countable metrizable rectifiable space is compactly generated if and only if it is compactly generated by a sequence converging to the right neutral element  $e$ .*

#### 4. Compactly generated paratopological groups

In this section, we mainly discuss compactly generated paratopological groups.

**Lemma 5.** *Let  $G$  be a paratopological group. If  $Y$  is a dense subset of  $G$  and  $U$  is an open neighborhood of the neutral element  $e$  of  $G$ , then  $G = Y^{-1} \cdot U$ .*

**Proof.** For arbitrary  $g \in G$ , since  $Y$  is a dense subset of  $G$ , we have  $Ug^{-1} \cap Y \neq \emptyset$ . Take  $x \in Ug^{-1} \cap Y$ . Then  $g \in x^{-1}U \subset Y^{-1} \cdot U$ .  $\square$

The proof of the following theorem is similar to that of Theorem 2.

**Theorem 7.** *If a  $\sigma$ -compact first-countable paratopological group  $G$  contains a dense compactly generated subgroup, then  $G$  is also compactly generated.*

**Proof.** Let  $H$  be a dense subgroup of  $G$  such that  $H$  is generated by some compact set  $E$ , and let  $G = \bigcup \{K_n : n \in \mathbb{N}\}$ , where each  $K_n$  is compact. Since  $G$  is first-countable, the point  $e$  has a countable local base  $\{U_n : n \in \mathbb{N}\}$ , where  $\overline{U_{n+1}} \subset U_n$  for each  $n \in \mathbb{N}$ . By the density of  $H$  in  $G$ , it follows from Lemma 5 that  $HU_n = G$  for each  $n \in \mathbb{N}$ . For each  $n \in \mathbb{N}$ , there exists a finite subset  $F_n$  of  $H$  such that  $K_n \subset F_nU_n$ , and put  $L_n = \overline{U_n} \cap (F_n)^{-1}K_n$ , then each  $K_n \subset F_nL_n$  since  $K_n \subset$

$F_n(F_n)^{-1}K_n$ . Obviously, each  $L_n$  is compact and, by Lemma 1,  $L = \bigcup\{L_n : n \in \mathbb{N}\}$  is also compact. Therefore,

$$G = \bigcup\{K_n : n \in \mathbb{N}\} \subset \bigcup\{F_nL_n : n \in \mathbb{N}\} \subset \bigcup\{HL_n : n \in \mathbb{N}\} \subset HL.$$

Since  $H$  is generated by  $E$ ,  $G$  is generated by the compact set  $E \cup L$ . Therefore,  $G$  is compactly generated.  $\square$

**Note 2.** Under the class of paratopological groups, we can obtain all results from Proposition 1 to Theorem 5 and Corollary 4 to Theorem 6 in Section 3 by similar proofs. In fact, the respective counterparts also hold for first-countable paratopological groups and this condition is weaker than the metrizability.

Since a compactly generated rectifiable space  $G$  is  $\sigma$ -compact,  $G$  has Souslin property, see [18] or [19]. Moreover, E.A. Reznichenko showed that every  $\sigma$ -compact Hausdorff paratopological group has Souslin property, see [2, Theorem 5.7.12]. However, the following question is still open.

**Question 1.** Let  $G$  be a compactly generated paratopological group. Does  $G$  have Souslin property?

**Theorem 8.** Any  $\sigma$ -compact paratopological group  $G$  can be embedded as a closed paratopological subgroup in some compactly generated paratopological group.

**Proof.** Let  $\sigma\Pi = \sigma\Pi\{G_n : n \in \mathbb{Z}\}$  be the  $\sigma$ -product of copies of  $G$  with the topology induced from Tikhonov power  $G^{\mathbb{Z}}$ , where  $\sigma\Pi$  is a  $\sigma$ -product with the neutral element  $e$  as a distinguished point. Then  $\sigma\Pi$  is also a paratopological group. For each  $n \in \mathbb{Z}$ , let  $i_n : G \rightarrow G_n$  be a topological isomorphism, and we can identify  $G_n$  with its image in  $\sigma\Pi$  under the natural embedding. Suppose that  $G = \bigcup\{K_n : n \in \mathbb{Z}\}$ , where each  $K_n$  is compact. Let  $K$  denote the subspace  $\bigcup_{n \in \mathbb{Z}} i_n(K_n)$  of the paratopological group  $\sigma\Pi$ . Since  $K$  is closed in the compact subspace  $\Pi\{K_n : n \in \mathbb{Z}\}$  of the paratopological group  $G^{\mathbb{Z}}$  under the natural embedding  $\sigma\Pi \rightarrow G^{\mathbb{Z}}$ ,  $K$  is compact in  $\sigma\Pi$ .

The group  $\mathbb{Z}$  of integers with the discrete topology acts on the paratopological group  $\sigma\Pi$  by shifting coordinates: for  $x = (x_n)_{n \in \mathbb{Z}} \in \sigma\Pi$  and  $k \in \mathbb{Z}$ ,  $k \cdot x$  is the element of  $\sigma\Pi$  whose  $n$ -th coordinate is  $x_{n+k}$ . Let  $G'$  denote the semidirect product  $\sigma\Pi \rtimes \mathbb{Z}$ . Assume  $1_z$  is the smallest positive element of  $\mathbb{Z}$ . Then the space  $K \cup \{1_z\}$  is a compact subspace of  $G'$  and  $G' = \langle K \cup \{1_z\} \rangle$ . Indeed, let  $H = \langle K \cup \{1_z\} \rangle$  in  $G'$ . Clearly,  $\mathbb{Z} \subset H$ . Next, we shall prove that, for each  $m \in \mathbb{Z}$ ,  $G_m \subset H$ . Take arbitrary  $x \in G_m$ . Then  $i_m^{-1}(x) \in K_n$  for some  $n \in \mathbb{Z}$ . Let  $a$  be the element  $(i_n i_m^{-1}(x), 0)$  and  $b$  the element  $(e, m - n)$  of the semidirect product  $G'$ . Clearly,  $a \in K \subset H$  and  $b \in H$ , and hence  $ba$  belongs to  $H$ . However, it is easy to see that  $ba = x$ .  $\square$

If  $G$  be countable, then each of the sets  $K_n$  can be assumed finite. A simple analysis of the topological structure of the space  $K \cup \{1_z\}$  enables us to obtain

**Theorem 9.** Any countable paratopological group  $G$  can be embedded as a closed paratopological subgroup in some paratopological group algebraically generated by a subspace homeomorphic to the one-point compactification  $\partial\mathbb{N}$  of a countable discrete space.

**Question 2.** Can any  $\sigma$ -compact rectifiable space  $G$  be embedded as a closed rectifiable subspace in some compactly generated rectifiable space?

## 5. Generalized metrizable properties of compactly generated rectifiable spaces

A closed mapping  $f$  is called *perfect* if each fiber is compact.

**Proposition 2.** *Suppose that  $F$  is a compact subspace of a rectifiable space  $G$ . Then the restriction  $p$  and  $q$  to the subspace  $F \times G$  is a perfect and open mapping of  $F \times G$  onto  $G$ .*

**Proof.** We firstly prove that the restriction  $p$  to the subspace  $F \times G$  is a perfect and open mapping of  $F \times G$  onto  $G$ .

Let  $f : F \times G \rightarrow F \times G$  be defined by  $f(x, y) = (x, p(x, y))$  for each  $(x, y) \in F \times G$ . Obviously,  $f$  is continuous, one-to-one, and  $f(F \times G) = F \times G$ . Moreover,  $f^{-1}(x, y) = (x, q(x, y))$ . Therefore,  $f^{-1}$  is also continuous. Thus  $f$  is a homeomorphism. For  $i = 1, 2$ , denote by  $\pi_i$  the projection of  $F \times G$  onto the  $i$ -th factor. Since  $p(x, y) = \pi_2(x, p(x, y)) = \pi_2 f(x, y)$  for all  $x \in F$  and  $y \in G$ ,  $p$  is the composition of  $f$  and  $\pi_2$ , that is,  $p = \pi_2 \circ f$ . Since  $F$  is compact, it follows from [8, Theorem 3.1.16] that  $\pi_2$  is closed. Then  $p$  is closed since  $f$  is a homeomorphism and  $\pi_2$  is closed. For each  $y \in G$ ,  $p^{-1}(y) = f^{-1}(F \times \{y\}) = \bigcup \{(x, q(x, y)) : x \in F\}$  is closed in the compact subspace  $F \times q(F, y)$ . Indeed, let  $(x, q(z, y)) \in (F \times q(F, y)) \setminus p^{-1}(y)$ , where  $x, z \in F$ . Then  $q(x, y) \neq q(z, y)$ , and thus there exist two open sets  $U$  and  $V$  in  $G$  such that  $q(x, y) \in U$ ,  $q(z, y) \in V$  and  $U \cap V = \emptyset$ . Since  $q$  is continuous, there exists an open neighborhood  $W$  of  $e$  such that  $q(x \cdot W, y \cdot W) \subset U$  and  $q(z \cdot W, y \cdot W) \subset V$ . Then  $(x \cdot W, V)$  is an open neighborhood of  $(x, q(z, y))$ . However, since  $q(x \cdot w, y) \subset U$  for each  $w \in W$ , it follows that  $(x \cdot W, V) \cap p^{-1}(y) = \emptyset$ . Therefore,  $p^{-1}(y)$  is closed in  $F \times q(F, y)$ , and thus it is compact. Then  $p$  is perfect.

Let  $O$  be an open subset of  $F \times G$ . Put  $O' = \pi_1(O)$ . For each  $x \in O'$ , let  $U_x = \{y \in G : (x, y) \in O\}$ ; then  $O_x$  is open in  $G$  as the projection of the open subset  $O \cap \pi_1^{-1}(x)$  of  $\{x\} \times G$  onto the second factor. Therefore,  $p(O) = \bigcup_{x \in O'} p(x, O_x)$  is open in  $G$ , which implies that  $p$  is an open mapping.

As for the mapping  $q$ , we only redefine the mapping  $f$  by  $(x, y) = (x, q(x, y))$  for each  $(x, y) \in F \times G$ , and the rest of the proof is immediate.  $\square$

**Corollary 7.** *Suppose that  $F$  is a compact subspace of a rectifiable space  $G$ , and that  $M$  is a closed subspace of  $G$ . Then  $p(F, M)$  and  $q(F, M)$  are all closed in  $G$ .*

**Note 3.** *Corollary 7 gives an affirmative answer to the following question. Recently, L.X. Peng and S.J. Guo [16] have also obtained Corollary 7. However, we prove Corollary 7 by a different method.*

**Question 3** (see [15]). *Let  $G$  be a rectifiable. If  $F, P$  are compact and closed subsets of  $G$ , respectively, is  $P \cdot F$  or  $F \cdot P$  closed in  $G$ ?*

Since the restriction of a perfect mapping to a closed subspace is again a perfect mapping, it follows from Corollary 7 and Proposition 2 that we have the following corollary.

**Corollary 8.** *Suppose that  $F$  is a compact subspace of a rectifiable space  $G$ , and that  $M$  is a closed subspace of  $G$ . Then the restriction  $p$  and  $q$  to the subspace  $F \times M$  is a perfect mapping of  $F \times M$  onto a closed subspace of  $G$ .*



A space  $G$  is of *countable tightness* if for each subset  $A$  of  $G$  and each point  $x \in \text{cl}(A)$  there exists a countable subset  $D$  of  $A$  such that  $x \in \text{cl}(D)$ .

**Theorem 10.** *Suppose that  $F$  is a compact subspace of a rectifiable space  $G$  and that  $M$  is a closed subspace of  $G$ . Suppose also that both  $F$  and  $M$  have countable tightness. Then both spaces  $p(F, M)$  and  $q(F, M)$  have countable tightness, too.*

**Proof.** Since perfect mappings do not increase the tightness and the tightness of the product  $F \times M$  is countable by [8, 3.12.8(a)], it follows from Corollary 8 that both spaces  $p(F, M)$  and  $q(F, M)$  have countable tightness, too.  $\square$

**Theorem 11.** *Suppose that  $F$  is a compact metrizable subspace of a rectifiable space  $G$ , and that  $M$  is a closed metrizable subspace of  $G$ . Then both spaces  $p(F, M)$  and  $q(F, M)$  are metrizable, too.*

**Proof.** By Corollary 7,  $p(F, M)$  and  $q(F, M)$  are closed in  $G$ . Since perfect mappings preserve the metrizability [8, Theorem 4.4.15], it follows from Corollary 8 that  $p(F, M)$  and  $q(F, M)$  are metrizable.  $\square$

A *network* for a space  $X$  is a collection  $\mathcal{F}$  of subsets of  $X$  such that whenever  $x \in U$  with  $U$  open, there exists  $F \in \mathcal{F}$  with  $x \in F \subset U$ .

**Theorem 12.** *Let  $G$  be a rectifiable space, and let  $H$  be a rectifiable subspace of  $G$  compactly generated by a compact metrizable space  $F$ . Suppose further that  $G = p(H, M)$ , where  $M$  is a closed metrizable subspace of  $G$ . Then  $G$  is the union of a countable family of closed metrizable subspaces.*

**Proof.** By induction on  $n$ , we can define a sequence  $\{A_n : n \in \omega\}$  of subsets of  $G$  such that:

- (1)  $A_0 = F \cup p(F, F) \cup q(F, F)$ ;
- (2)  $A_1 = p(A_0, A_0) \cup q(A_0, A_0)$ ;
- (3)  $A_n = p(A_{n-1}, A_{n-1}) \cup q(A_{n-1}, A_{n-1})$ .

Obviously, each  $p(A_n, A_n), q(A_n, A_n), A_n$  are compact. Since compact space with a countable network is metrizable [10], it follows from Theorem 11 that each  $A_n$  is also metrizable. Since  $H = \langle F \rangle$ ,  $H = \bigcup_{n \in \omega} A_n$ . Since  $G = p(H, M)$ , it follows from Theorem 11 again that  $G$  is the union of a countable family of closed metrizable subspaces.  $\square$

A *neighborhood assignment* for a space  $X$  is a function  $\varphi$  from  $X$  to the topology of  $X$  such that  $x \in \varphi(x)$  for each point  $x \in X$ . A space  $X$  is a *D-space*[6], if for any neighborhood assignment  $\varphi$  for  $X$  there is a closed discrete subset  $D$  of  $X$  such that  $X = \bigcup_{d \in D} \varphi(d)$ .

**Corollary 9.** *Let  $G$  be a rectifiable space, and let  $H$  be a rectifiable subspace of  $G$  compactly generated by a compact metrizable space  $F$ . Suppose further that  $G = p(H, M)$ , where  $M$  is a closed metrizable subspace of  $G$ . Then  $G$  is a D-space.*

**Proof.** It is well known that each metrizable space is a  $D$ -space. Hence  $M$  is a  $D$ -space, and then each  $p(h, M)$  is a  $D$ -space, too. Since a countable infinite union of closed  $D$ -subspaces is  $D$  [3], it follows that  $G = p(H, M) = \bigcup_{h \in H} p(h, M)$  is a  $D$ -space.  $\square$

Recall that a space  $X$  has a *quasi- $G_\delta$ -diagonal* provided there is a sequence  $\{\mathcal{G}(n) : n \in \mathbb{N}\}$  of collections of open subsets of  $X$  such that for any distinct points  $x, y \in X$  there is a number  $n$  with  $x \in st(x, \mathcal{G}(n)) \subset X \setminus \{y\}$ .

**Theorem 13.** *Let  $G$  be a compactly generated Tychonoff rectifiable space, and  $Y = bG \setminus G$  have locally quasi- $G_\delta$ -diagonal, where  $bG$  is a Hausdorff compactification of  $G$ . Then  $G$  satisfies one of the following conditions:*

- (1)  $G$  is locally compact;
- (2)  $G$  is separable and metrizable.

**Proof.** Suppose that  $G$  is nowhere locally compact. Since  $G$  is  $\sigma$ -compact, it follows from [14, Theorem 7.3] that  $G$  is separable and metrizable.  $\square$

A space  $X$  is said to have a *regular  $G_\delta$ -diagonal* if the diagonal  $\Delta = \{(x, x) : x \in X\}$  can be represented as the intersection of the closures of a countable family of open neighborhoods of  $\Delta$  in  $X \times X$ .

Since a rectifiable space with a countable pseudocharacter has a regular  $G_\delta$ -diagonal [14] and a paracompact space with a  $G_\delta$ -diagonal is submetrizable [10], we have the following proposition.

**Proposition 3.** *If  $G$  is a compactly generated rectifiable space with a countable pseudocharacter, then  $G$  is submetrizable.*

**Proposition 4.** *Let  $G$  be a compactly generated Tychonoff rectifiable space with a countable pseudocharacter, and let  $Y = bG \setminus G$  be Lindelöf, where  $bG$  is a Hausdorff compactification of  $G$ . Then  $G$  is separable and metrizable.*

**Proof.** Since  $Y = bG \setminus G$  is Lindelöf,  $G$  is countable type [12], and thus  $G$  is a  $p$ -space [1]. Then  $G$  is a  $\sigma$ -compact  $p$ -space with a  $G_\delta$ -diagonal, hence it is separable and metrizable [10, Corollaries 3.8 and 3.20].  $\square$

## Acknowledgement

The author wishes to thank the referee for the detailed corrections and suggestions to the paper and all his/her efforts made in order to improve the paper.

## References

- [1] A. V. ARHANGEL'SKIĬ, M. M. CHOBAN, *On remainders of rectifiable spaces*, Topology Appl. **157**(2010), 789-799.
- [2] A. V. ARHANGEL'SKIĬ, M. TKACHENKO, *Topological Groups and Related Structures*, Atlantis Press and World Sci., Paris, 2008.
- [3] C. R. BORGES, A. WEHRLY, *A study of  $D$ -spaces*, Topology Proc. **16**(1991), 7-15.

- [4] M. M. ČOBAN, *On topological homogeneous algebras*, in: *Interim Reports of Prague Topol. Symp. II*, Prague, 1987, 25–26.
- [5] M. M. ČOBAN, *The structure of locally compact algebras*, *Serdica* **18**(1992), 129–137.
- [6] E. K. DOUWEN, W. F. PFEFFER, *Some properties of the Sorgenfrey line and related spaces*, *Pacific J. Math.* **81**(1979), 371–377.
- [7] R. ELLIS, *Locally compact transformation groups*, *Duke Math. J.* **24**(1957), 119–125.
- [8] R. ENGELKING, *General Topology* (revised and completed edition), Heldermann Verlag, Berlin, 1989.
- [9] H. FUJITA, D. SHAKHMATOV, *A characterization of compactly generated metric spaces*, *Proc. Amer. Math. Soc.* **131**(2002), 953–961.
- [10] G. GRUENHAGE, *Generalized metric spaces*, *Handbook of Set-Theoretic Topology*, North Holland, Amsterdam, 1984.
- [11] A. S. GUL'KO, *Rectifiable spaces*, *Topology Appl.* **68**(1996), 107–112.
- [12] M. HENRIKSEN, J. ISBELL, *Some properties of compactifications*, *Duke Math. J.* **25**(1958), 83–106.
- [13] F. LIN, *Topologically subordered rectifiable spaces and compactifications*, *Topology Appl.* **159**(2012), 360–370.
- [14] F. LIN, C. LIU, S. LIN, *A note on rectifiable spaces*, *Topology Appl.* **159** (2012), 2090–2101.
- [15] F. LIN, R. SHEN, *On rectifiable spaces and paratopological groups*, *Topology Appl.* **158**(2011), 597–610.
- [16] L. X. PENG, S. J. GUO, *Two questions on rectifiable spaces and related conclusions*, *Topology Appl.* **159**(2012), 3335–3339.
- [17] N. M. PYRCH, A. V. RAVSKY, *On free paratopological groups*, *Matematychni Studii* **25**(2006), 115–125.
- [18] E. A. REZNIČENKO, V. V. USPENSKIJ, *Pseudocompact Mal'tsev spaces*, *Topology Appl.* **86**(1998), 83–104.
- [19] V. V. USPENSKIJ, *On continuous images of Lindelof topological groups*, *Dokl. Akad. Nauk SSSR* **285**(1985), 824–827, in Russian; English transl.: *Soviet Math. Dokl.* **32**(1985), 802–806.
- [20] V. V. USPENSKIJ, *The Mal'tsev operation on countably compact spaces*, *Comments. Math. Univ. Carolin.* **30**(1989), 395–402.
- [21] V. V. USPENSKIJ, *Topological groups and Dugundji compacta*, *Mat. Sb.* **180**(1989), 1092–1118, in Russian; English transl. in: *Math. USSR-Sb.* **67**(1990), 555–580.