

THREATENED FISHES OF THE WORLD: *Heteropneustes fossilis* (Bloch, 1794) (Siluriformes: Heteropneustidae)

Md. Yeamin Hossain¹*, Rafiqul Islam¹, Zoarder Faruque Ahmed², Md. Mosaddequr Rahman³, Md. Alomgir Hossen¹, S. M. Abu Naser¹, Raisul Islam Rasel⁴

¹Department of Fisheries, Faculty of Agriculture, University of Rajshahi, Rajshahi 6205, Bangladesh

²Department of Fisheries Management, Faculty of Fisheries, Bangladesh Agricultural University, Mymensingh 2202, Bangladesh

³Faculty of Fisheries, Kagoshima University, 4-50-20 Shimoarata, Kagoshima 890-0056, Japan

⁴Department of Zoology, Anando Mohan College, National University, Gazipur 1704, Bangladesh

*Corresponding Author, E-mail: yeamin.fish@ru.ac.bd

ARTICLE INFO

Received: 30 October 2014

Received in revised form: 7 February 2015

Accepted: 10 February 2015

Available online: 11 February 2015

Keywords:

Heteropneustes fossilis

Stinging catfish

Vulnerable

Asia

ABSTRACT

The threatened stinging catfish *Heteropneustes fossilis* is a nutritionally valuable food in Asian countries but its natural populations have declined due to over-exploitation, habitat loss and pollution, thus deserving high conservation importance for the remaining isolated, wild populations of *H. fossilis* in Asian countries.

The results would be useful for the protection of the presence of endemic species and sustainable fisheries in the lake.

How to Cite

Hossain, M. Y., Islam, R., Ahmed, Z. F., Rahman, M. M., Hossen, M. A., Naser S. M. A., Rasel, R. I. (2015): Threatened fishes of the world: *Heteropneustes fossilis* (Bloch, 1794) (Siluriformes: Heteropneustidae). Croatian Journal of Fisheries, 73, 77-79. DOI: 10.14798/73.2.796

COMMON NAME

Stinging catfish and shingi in Bangladesh (Rahman, 1989); Singee and Bitchu in India (Nath and Dey, 1989; Daniels, 2002); Hunga in Sri Lanka (Pethiyagoda, 1991); Lahoord, Nullie and Lohar in Pakistan (Misra, 1976).

extensively fished because of its invigorating quality that includes taste and its nutritional and medicinal values (Jha and Rayamajhi, 2010). Its muscle contains high amounts of protein, iron (226 mg/100g) and calcium (Saha and Guha, 1939; Alok et al., 1993). Being a lean fish (fat content only 2.57 ± 0.24%), it is conducive to people on low-fat diets (Rahman et al., 1982).

CONSERVATION STATUS

Least concern in Bangladesh according to IUCN (2014), Vulnerable in India and Sri Lanka according to Dahanukar et al. (2004) and IUCN Sri Lanka (2000), respectively.

IDENTIFICATION

H. fossilis is elongated and highly compressed (Fig. 1) with maxillary barbels extending to the end of pectoral fins and mandibular pairs extended up to the base of pelvis. Caudal fin is rounded and distinctly separated. Fin formula: D. 6-7; P1. 1/6-7; P2. 6; A. 62-70 (Rahman, 1989). Male has thinner ventral line which is absent in female.

IMPORTANCE

Stinging catfish *H. fossilis* is commercially and aquaculturally a significant species in many Asian countries. It is



Fig 1. *Heteropneustes fossilis* sample and photo were taken by the senior author (Md. Yeamin Hossain) from the Ganges River (known as Padma in Bangladesh) on 3 February 2015.

DISTRIBUTION

H. fossilis is distributed throughout south and southeast Asian countries including Bangladesh, India, Laos, Myanmar, Nepal, Pakistan, Sri Lanka and Thailand (Talwar and Jhingran, 1991). It is also found in Iran and Iraq (Coad, 1996; FAO, 1997).

ABUNDANCE

This fish was previously abundant in open (rivers and streams) and closed water ecosystems like ponds, floodplains, canals, *beel* and *haors* (Hussain and Mazid, 2001) but has now been reduced dramatically (Rahman et al., 2012).

HABITAT AND ECOLOGY

Stinging catfish inhabits freshwater and also (though rarely) brackish waters. This is mainly a fish of ponds, ditches, *beels*, swamps and marshes but is sometimes also found in muddy rivers (Jha and Rayamajhi, 2010). It is omnivorous during its life cycle (Breder and Rosen, 1966). The fry are planktivorous; juveniles feed on crustacea, plant, miscellaneous matter and insects; adults feed on insects, detritus and plant matter (Pal et al., 1996).

REPRODUCTION

Size at sexual maturity of male and female *H. fossilis* was estimated as 5.50 cm total length (TL) (age ~ 1 year) and 12.00 cm TL (age ~ 1 year), respectively (Talwar and Jhingran, 1991). Spawning season extends from July to August (Joy and Tharakan, 1999). Fecundity ranged from 1375 to 46737 (Bhatt et al., 1977).

THREATS

Population have declined due to reckless fishing, habitat destruction and conversion, injudicious application of pesticides on agricultural fields, release of industrial effluents, disease and effects of climate change (Dahanukar et al., 2004; Jha and Rayamajhi, 2010).

CONSERVATION ACTION

Several studies on the ecology and biology of *H. fossilis* have been conducted in Bangladesh (e.g., Pal et al., 1996; Kohinoor et al., 2012). In addition, induced breeding of this species has been done in Bangladesh (Rahman et al., 2013) and India (Alok et al., 1993; Joy and Tharakan, 1999).

CONSERVATION RECOMMENDATIONS

Studies on the reproduction and population biology are urgently needed (Hossain, 2014; Hossain and Alam, 2015). Legal and institutional reforms are necessary to engage local communities in *in situ* conservation and for engaging the ornamental fish industry in global conservation (Pethiyagoda, 2006). Establishment of suitable sanctuaries in selected rivers, streams, floodplains and swamplands has been suggested (Hossain et al., 2008; 2009; 2015a). Fishing for immature fish during the spawning season and indiscriminate use of pesticides on the agricultural fields should be strictly banned (Hossain et al., 2015b). *H. fossilis* would also benefit from effective habitat preservation and increased public motivations.

ACKNOWLEDGEMENTS

The authors would like to acknowledge TWAS (GRA No.: 14-028 RG/BIO/AS_I-UNESCO FR: 324028574) for the support of sampling in the Ganges River (Padma), Bangladesh.

Sažetak

UGROŽENE VRSTE RIBA U SVIJETU: *Heteropneustes fossilis* (Bloch, 1794) (Siluriformes: Heteropneustidae)

Ugroženi bodljasti som, *Heteropneustes fossilis*, je nutritivno vrijedna namirnica u azijskim zemljama, ali njegove prirodne populacije opadaju zbog pretjeranog iskorištavanja, gubitka staništa i zagađenja zbog kojih zaslužuje visoku važnost za zaštitu preostalih izoliranih, divljih populacija *H. fossilis* u azijskim zemljama.

Ključne riječi: *Heteropneustes fossilis*, bodljasti som, osjetljiva vrsta, Azija

REFERENCES

- Alok, D., Krishnan, T., Talwar, G. P., Garg, L. C. (1993): Induced spawning of catfish, *Heteropneustes fossilis* (Bloch), using D-Lys6 salmon gonadotropin-releasing hormone analog. *Aquaculture*, 115, 159-167.
- Bhatt, V. S., Dalal, S. G., Abidi, S. A. H. (1977): Fecundity of

- the freshwater catfishes *Mystus seenghala* (Sykes), *Mystus cavasius* (Hamilton), *Wallago attu* (Bloch) and *Heteropneustes fossilis* (Bloch) from the plains of Northern India. *Hydrobiologia*, 54, 219-224.
- Breder, C. M., Rosen, D. E. (1966): Modes of Reproduction in Fishes. T. F. H. Publications, Neptune City, New Jersey. 941 pp.
- Coad, B. W. (1996): Exotic fish species in the Tigris-Euphrates basin. *Zoology in the Middle East*, 13, 71-83.
- Dahanukar, N., Raut, R., Bhat, A. (2004): Distribution, endemism and threatened status of freshwater fishes in the western Ghats of India, *Journal of Biogeography*, 31, 123-136.
- Daniels, R. J. R. (2002): Freshwater fishes of Peninsular India. Madhav Gadgil (ed.) India-A lifescape 2. Universities Press, Hyderabad.
- FAO. (1997): FAO database on introduced aquatic species. FAO Database on Introduced Aquatic Species, FAO, Rome.
- Hossain, M. Y. (2014): Threatened Fishes of the World: *Mystus vittatus* (Bloch, 1794) (Siluriformes: Bagridae). *Croatian Journal of Fisheries*, 72, 183-185.
- Hossain, M. Y., Ahmed, Z. F., Al-Kady, M. A. H., Ibrahim, A. H. M., Ohtomi, J., Fulanda, B. (2008): Threatened Fishes of the World: *Wallago attu* (Bloch, M. E. and Schneider, J. G., 1801) (Siluriformes: Bagridae). *Environmental Biology of Fishes*, 82, 277-278.
- Hossain, M. Y., Alam, M. J. (2015): Threatened Fishes of the World: *Plotosus canius* (Hamilton, 1822) (Siluriformes: Plotosidae). *Croatian Journal of Fisheries*, 73, 35-36.
- Hossain, M. Y., Hossain, M. A., Islam, R., Hossen, M. A., Rahman, O., Rahman, M. M. (2015b): Threatened Fishes of the World: *Pethia ticto* (Hamilton, 1822) (Cypriniformes: Cyprinidae). *Croatian Journal of Fisheries*, 73, 37-39.
- Hossain, M. Y., Islam, R., Hossain, M. A., Hossen, M. A., Rahman, M. M., Ohtomi, J. (2015a): Threatened Fishes of the World: *Cirrhinus reba* (Hamilton 1822) (Cypriniformes: Cyprinidae). *Croatian Journal of Fisheries*, 73, 40-42.
- Hossain, M. Y., Rahman, M. M., Mollah, M. F. A. (2009): Threatened Fishes of the World: *Pangasius pangasius* Hamilton-Buchanan, 1822 (Pangasiidae). *Environmental Biology of Fishes*, 84, 315-316.
- Hussain, M. G., Mazid, M. A. (2001): Genetics Improvement and Conservation of Carp Species in Bangladesh. Bangladesh Fisheries Research Institute, Mymensingh, Bangladesh, 56 pp.
- IUCN Sri Lanka. (2000): The 1999 List of Threatened Fauna and Flora of Sri Lanka. Colombo: IUCN Sri Lanka. 114 pp.
- IUCN. (2014): IUCN Red List of Threatened Species. Version 2014.2. IUCN 2014. IUCN Red List of Threatened Species. www.iucnredlist.org. (Accessed on 10 October, 2014).
- Jha, B. R., Rayamajhi, A. (2010): *Heteropneustes fossilis*. The IUCN Red List of Threatened Species. Version 2014.2. www.iucnredlist.org. (Accessed on 10 October, 2014).
- Joy, K. P., Tharakan, B. (1999): Induced spawning of the Indian catfish, *Heteropneustes fossilis*, by GnRH analogue alone or in combination with dopamine-affecting drugs. *Journal of Applied Aquaculture*, 9, 23-32.
- Kohinoor, A. H. M., Khan, M. M., Yeasmine, S., Mandol, P., Islam, M. S. (2012): Effects of stocking density on growth and production performance of indigenous stinging catfish, *Heteropneustes fossilis* (Bloch). *International Journal of Agricultural Research, Innovation and Technology*, 2, 9-14.
- Misra, K. S. (1976): Pisces. Vol. 3. Delhi, Manager of Publications, 3, 351 p.
- Nath, P., Dey, S. C. (1989): Fish and fisheries of North East India. Arunachal Pradesh, 1, 1-143p.
- Pal, M., Kohli, S., Goswami, U. C. (1996): Food and feeding habits of *Heteropneustes fossilis* (Bloch) from the Brahmaputra River system, Assam. *Indian Journal of Fisheries*, 43, 97-101.
- Pethiyagoda, R. (1991): Freshwater Fishes of Sri Lanka. The Wildlife Heritage Trust of Sri Lanka, Colombo, 362 pp.
- Pethiyagoda, R. (2006): Conservation of Sri Lankan Freshwater Fishes. The Fauna of Sri Lanka, 103-112.
- Rahman, M. A., Gheyasuddin, S., Rashid, M. H., Chowdhury M. F. Z. (1982): Proximate composition and nutritive quality of freshwater *zeol* fishes of Bangladesh. *Bangladesh Journal of Fisheries* 2-5, 37-43.
- Rahman, A. K. A. (1989): Freshwater Fishes of Bangladesh. Zoological Society of Bangladesh. Department of Zoology, University of Dhaka, Dhaka 1000, 364 p.
- Rahman, M. M., Hossain, M. Y., Ahamed, F., Fatematuzzhura, Subba, B. R. Abdallah, E. M., Ohtomi, J. (2012): Biodiversity in the Padma (Distributary of the Ganges) River, northwestern Bangladesh: Recommendations for conservation. *World Journal of Zoology*, 7, 328-337.
- Rahman, M. M., Hossain, M. Y., Hossain, M. I., Provhat, S. J., Islam, M. S., Hossain, M. B. (2013): Induced breeding of the stinging catfish, *Heteropneustes fossilis*: comparison among different inducing agents. *Turkish Journal of Fisheries and Aquatic Sciences*, 13, 532-537.
- Saha, K. C., Guha B. C. (1939): Nutritional investigation of Bengal fish. *The Indian Journal of Medical Research*, 26, 921-927.
- Talwar P. K., Jhingran A. G. (1991): Inland Fishes of India and Adjacent Countries. Vol. 2. Published by Oxford & IBH Publishing Co. Pvt. Ltd. New Delhi, Bombay, Calcutta, 1158 pp.