

PREVALENCE AND RISK FACTORS FOR DEVELOPING ORAL ALLERGY SYNDROME IN ADULT PATIENTS WITH SEASONAL ALLERGIC RHINITIS

Nada Ivičević¹, Željka Roje², Zlatko Kljajić², Lovro Bojić³, Goran Kardum⁴, Lada Omero² and Goran Račić²

¹School of Medicine, University of Split; ²ENT Department, ³Department of Ophthalmology, Split University Hospital Center; ⁴Faculty of Humanities and Social Sciences, University of Split, Split, Croatia

SUMMARY – The aim of this study was to assess the prevalence of oral allergy syndrome (OAS) in patients with seasonal allergic rhinitis (SAR) and the possible risk factors for its development. This cross-sectional study was conducted in primary care offices in the Split-Dalmatia County during the period from March 1 to September 30, 2012. Data sources were medical history with confirmation of SAR (positive skin-prick test to seasonal inhalant allergens: grass, tree and weed pollens), anthropometric patient data (age, sex, weight and height), and a questionnaire in which patients evaluated their nasal and ocular symptoms, comorbidities and lifestyle. The χ^2 -test, Pearson χ^2 -test, Spearman's rho correlation coefficient and Kolmogorov-Smirnov test were used on statistical analysis. The prevalence of OAS was 45.7%. The risk factors for OAS development were diabetes ($p < 0.001$), severity of nasal symptoms ($p < 0.05$) and severity of ocular symptoms ($p < 0.001$). In conclusion, the prevalence of OAS in the Split-Dalmatia County is very high. The risks factors for OAS in patients with SAR are diabetes and severity of nasal and ocular symptoms.

Key words: *Oral allergy syndrome, risk factors; Rhinitis, allergic, seasonal; Croatia*

Introduction

Oral allergy syndrome (OAS) represents a set of localized oral symptoms such as itching, tingling, and swelling of lips, tongue, palate and pharynx in patients suffering from seasonal allergic rhinitis (SAR), asthma, or both. These symptoms are usually self-limiting and last for only a few seconds to a few minutes¹.

Symptoms occur as a consequence of specific food allergies caused by cross-reactivity between pollen and protein from fresh fruits and vegetables¹⁻⁴. Allergic reactions to proteins in fruit and vegetables, which show cross-reactivity with pollen antigens in

sensitive individuals, behave according to hypersensitivity type I and are mediated by immunoglobulin E (IgE)¹⁻³. In contrast to other types of food allergy in which drivers are specific IgE to antigens from food, in OAS drivers are specific IgE to antigens from grass pollens, trees and weeds^{3,5}. In addition to food, OAS can also occur in patients who are hypersensitive to latex the basic origin of which is also vegetable (rubber)^{6,7}. Due to the fact that incriminated antigens for the pathogenesis of OAS are mostly thermo labile and subjected to decomposition by digestive enzymes, dominant symptoms occur in the upper gastrointestinal tract (mouth and throat) and immediately after eating the offending fruit and vegetables. Even though systemic reactions are very rare, OAS has a considerable potential for anaphylaxis¹. Diagnosis is based on the history of SAR and typical symptoms that occur immediately after eating certain food of plant origin.

Correspondence to: Prof. Željka Roje, MD, PhD, ENT Department, Split University Hospital Center, Spinčićeva 1, HR-21000 Split, Croatia
E-mail: zroje@kbsplit.hr

Received October 28, 2013, accepted February 3, 2015

Differential diagnoses include many diseases (such as burning mouth syndrome, angioedema, hay fever, various other oral diseases, etc.), and for this reason a multidisciplinary approach is necessary, as different specialists need to be involved in the diagnostic procedure². Prevention and treatment are carried out through avoiding certain fruits and vegetables and/or their thermal processing, and antihistamines. The frequency of OAS is difficult to estimate. Many patients have mild symptoms that are eliminated by simply avoiding the incriminated food and have no need for further consultation with the physician³. Therefore, data in the literature significantly vary (by 8%-70%)⁸⁻¹⁴. It is also unclear in which patients with SAR we can expect OAS. For these two reasons, this study was conducted to estimate the prevalence of OAS in patients with SAR and to identify the possible risk factors for its development.

Patients and Methods

This cross-sectional study was conducted in primary care offices in the Split-Dalmatia County in the period from March 1 to September 30, 2012. The study was approved by the Ethics Committee of the Split University School of Medicine. A total of 600 questionnaires were distributed, of which 284 were not filled out or were filled out partially and were excluded from the study. There were 316 completed questionnaires, of which 16 were excluded because the respondents were under age 18 (n=15), and one was filled out by a pregnant woman. The final sample for analysis included 300 questionnaires.

The survey included 300 adult patients with SAR, which was confirmed by skin-prick test to seasonal inhalant allergens (grass, tree and weed pollens) and by symptoms of allergic rhinitis according to ARIA guidelines (runny nose, nasal obstruction, itchy nose and eyes, sneezing, watery and red eyes)^{15,16}. Exclusion criteria were positive skin-prick test for perennial allergens (dust mites, feathers, and dander) and food allergy. Data were collected through a questionnaire that quantified the severity of nasal and ocular symptoms of SAR (0 – no symptoms, 1 – mild, 2 – moderate, and 3 – severe) and oral symptoms of OAS in the same manner^{14,15}. A score of individual symptoms of OAS more than 3 is noted as positive in terms of the diagnosis of OAS.

Anthropometric parameters, other chronic diseases and lifestyle as the possible risk factors for OAS were also noted in the questionnaire (age, gender, weight, height, presence of asthma, diabetes, high blood pressure or some other chronic disease, smoking habits, positive family history of allergy, and duration of SAR/asthma symptoms).

Statistical analysis

We analyzed differences between the frequency of different categories and frequencies between categorical variables (different symptoms) using the χ^2 -test. Of statistical tests, we used Pearson χ^2 -test to determine correlation between the Total Nasal Symptom Score (TNSS) and Total Ocular Symptom Score (TOSS), TNSS and Total Oral Symptom Score (TORSS), and correlation between TOSS and TORSS. The same test was used to determine the possible association of anthropometric parameters, presence of asthma, diabetes, high blood pressure, smoking, family history, and duration of suffering from SAR/asthma as the possible risk factors for the development of OAS. Spearman's rho correlation coefficient was used to determine the association among TNSS, TOSS and TORSS. Kolmogorov-Smirnov test was used to determine normality of distribution. Odds ratios and 95% confidence intervals were estimated. Differences were considered significant at $p < 0.05$.

Results

In our study, 300 adult patients with SAR, 168 (56%) male and 132 (44%) female, completed the survey. All investigated clinical characteristics and demographic data on patients with and without OAS were compared (Table 1).

The average age was 42 (18-85) years, average weight 78 (± 13.98) kilograms and average height 175 (± 11.24) cm. Of all patients, 127 (42.3%) were active smokers and 54 (18%) former smokers. Elevated blood pressure was present in 87 (29%) patients. Diabetes mellitus was present in 25 (8.3%) patients with SAR but 17 (68%) of them had OAS, which was statistically significant ($p < 0.001$). Other chronic illnesses were present in 38 (12.7%) patients, mostly gastrointestinal (n=11; 3.7%) and cardiovascular (n=7; 2.3%) diseases. Asthma was diagnosed in 65 (21.7%) patients, while

Table 1. Comparison of patients with and without oral allergy syndrome

	SAR n=300	OAS positive n=137	OAS negative n=163	Odds ratio	95% Confidence interval	
Gender:				0.93		
male	168	78 (46.4%)	90 (53.6%)		0.59-1.47	**
female	132	59 (44.7%)	73 (55.3%)			**
Smoker:						
yes	127	61 (48.0%)	66 (52.0%)			**
no	119	51 (42.9%)	68 (57.1%)			**
ex	54	25 (46.3%)	29 (53.7%)			**
Positive skin-prick test:						
grass pollen	191	103 (53.9%)	88 (46.1%)	0.38	0.23-0.63	**
<i>Parietaria</i>	145	83 (57.2%)	62 (42.8%)	0.39	0.25-0.63	**
weed pollen	163	84 (51.5%)	79 (48.5%)	0.59	0.37-0.94	**
Mediterranean area tree pollen	216	107 (49.5%)	109 (50.5%)	0.56	0.33-0.95	**
Hypertension	87	39 (44.8%)	48 (55.2%)	1.04	0.63-1.73	**
Diabetes mellitus	25	17 (68%)	8 (32%)	0.36	0.15-0.87	p<0.001
Asthma	65	36 (55.4%)	29 (44.6%)	0.60	0.34-1.05	**
Atopic diathesis	166	81 (48.8%)	85 (51.2%)	0.75	0.47-1.19	**
Suffered from asthma or SAR:						
up to one year	9	3 (33.3%)	6 (66.7%)			**
more than one year	47	18 (38.3%)	29 (61.7%)			**
up to 5 years	74	28 (37.8%)	46 (62.2%)			**
more than 5 years	170	88 (51.8%)	82 (48.2%)			**

**Pearson χ^2 -test (nonsignificant); OAS = oral allergy syndrome; SAR = seasonal allergic rhinitis

166 (55.3%) patients had atopic diathesis (positive family history). At the time of survey, 170 (56.7%) respondents had suffered from SAR or asthma for more than 5 years. The majority of patients were allergic to Mediterranean area tree pollen (n=216; 72%), grass pollen (n=191; 63.7%), weed pollen (n=163; 54.3%) and *Parietaria* (n=145; 48.3%). There were 78 (26%) monosensitized subjects and the others were polysensitized. The leading nasal symptom was sneezing, which occurred in 276 (92%) patients. The most common ocular symptom was watery eyes that occurred in 254 (84.7%) patients. The frequency of OAS in our study was 45.7% (n=137). The leading oral symptom of tingling throat, palate and mouth after eating offending raw plants was present in 176 (58.7%) patients.


The severity of nasal and ocular symptoms was assessed by TNSS and TOSS (mild (0-4), moderate (5-8) and severe (9-12). The incidence of OAS symptoms in-

creased with the severity of TNSS (rs=0.115; p<0.05) (Fig. 1a) and severity of TOSS (rs=0.260; p<0.001) (Fig. 1b). Patients with more severe nasal symptoms had more severe ocular symptoms (rs=0.248; p<0.001).

Of the potential risk factors for developing OAS in patients with SAR investigated in the present study, only three parameters were statistically significant: 1) diabetes (Pearson χ^2 -test); 2) severity of nasal symptoms (Spearman's rho correlation coefficient); and 3) severity of ocular symptoms (Spearman's rho correlation coefficient).

Discussion

Oral allergy syndrome is caused by cross-reactivity between proteins from fresh fruits or vegetables and pollens. That is why it can develop only in patients



TNSS = Total Nasal Symptom Score mild (0-4), moderate (5-8) and severe (9-12); Spearman's rho correlation coefficient ($r_s=0.115$; $p=0.047$)
 TOSS = Total Ocular Symptom Score mild (0-4), moderate (5-8) and severe (9-12); Spearman's rho correlation coefficient ($r_s=0.260$; $p<0.001$)

Fig. 1. (a) Incidence of oral allergy syndrome due to severity of nasal symptoms; (b) incidence of OAS due to severity of ocular symptoms.

with SAR. It is common, but not all pollen allergic patients develop OAS. The prevalence of OAS has not been precisely established (i.e. varying across very wide ranges). It is affected by many factors including the form of making it more sensitive in certain geographic areas and the increasing prevalence of allergic rhinitis¹. Therefore, literature data on the frequency of OAS significantly differ (by 8%-70%)^{8-12,17-19}. According to our results, the prevalence was 45.7%.

Since the symptoms of OAS are usually mild and transient, many patients do not associate them with pollen allergy and rhinitis, and do not report it to physician on visiting the office. For this reason, OAS symptoms should be interviewed in all patients with rhinitis due to pollen allergy. If the diagnosis of OAS is established, patients should be informed about the possibility of hypersensitivity to certain fruits and vegetables¹⁴. In our study, gender was not a statistically significant risk factor for developing OAS symptoms. Men were slightly predominant, which is in contrast to literature data, where female gender seems to be the risk factor for OAS^{14,20,21}.

In this study, the severity of ocular and nasal symptoms was associated with OAS occurrence, i.e. the more severe ocular/nasal symptoms implied greater likelihood for the patient to develop OAS. It was the most striking result with great influence on everyday allergology practice and patient education; however, it should be noted that correlations were low but

significant. We can speculate that the same pattern of neuro-regulatory mechanisms included in ocular symptoms, like naso-ocular reflex, is involved in oral symptoms throughout the trigeminal nerve mediated reflexes on oral mucosa. Eyes and oral mucosa share the same innervation pattern and it can be proper explanation of our results. Further investigations are needed to exclude or confirm our speculations.

In spite of all the possible risk factors for OAS that we expected to be statistically significant, such as asthma, smoking or atopic diathesis, only diabetes was a statistically significant risk factor for developing OAS in patients with SAR. We found no literature data on this issue and have no idea on the possible mechanisms of this type of association.

References

1. Ausucua M, Dublin I, Echebarria MA, Aguirre JM. Oral allergy syndrome (OAS). General and stomatological aspects. *Med Oral Patol Oral Cir Bucal*. 2009;14(11):568-72.
2. Kelava N, Lugović-Mihić L, Duvančić T, Romić R, Šitum M. Oral allergy syndrome – the need of a multidisciplinary approach. *Acta Clin Croat*. 2014;53(2):210-9.
3. Katelaris CH. Food allergy and oral allergy or pollen-food syndrome. *Curr Opin Clin Immunol*. 2010;10:246-51.
4. Bokszezanin KL, Przybyła AA. Molecular aspects of allergy to plant products. Part I. Class I and II allergens and cross reactivity of IgE antibodies. *Pol Merkur Lekarski*. 2012;32(188):129-34.

5. Kondo Y, Urisu A. Oral allergy syndrome. *Allergol Int.* 2009;58:485-91.
6. Blanco C, Carrillo T, Castillo R, Quirarte J, Cuevas M. Latex allergy: clinical features and cross-reactivity with fruits. *Ann Allergy.* 1994;73:309-13.
7. Garcia Ortiz JC, Moyano JC, Alvarez M, Bellido J. Latex allergy in fruit-allergic patients. *Allergy* 1998;53:532-5.
8. Ma S, Sicherer SH, Nowak-Węgrzyn A. A survey on the management of pollen-food allergy syndrome in allergy practices. *J Allergy Clin Immunol.* 2003;112:784-7.
9. Cuesta-Herranz J, Lazaro M, Figueredo E, Igea JM, Umpierrez A, de-las-Heras M. Allergy to plant-derived fresh foods in a birch- and ragweed-free area. *Clin Exp Allergy.* 2000;30:1411-6.
10. Cuesta-Herranz J, Lazaro M, Martinez A, Figueredo E, Palacios R, de-Las-Heras M, *et al.* Pollen allergy in peach-allergic patients: sensitization and cross reactivity to taxonomically unrelated pollens. *J Allergy Clin Immunol.* 1999;104(3 Pt 1):688-94.
11. Fernandez-Rivas M, Bolhaar S, Gonzalez-Mancebo E, Asero R, van Leeuwen A, Bohle B, *et al.* Apple allergy across Europe: how allergen sensitization profiles determine the clinical expression of allergies to plant foods. *J Allergy Clin Immunol.* 2006;118:481-8.
12. Rodriguez J, Crespo JF, Lopez-Rubio A, De La Cruz-Bertolo J, Ferrando-Vivas P, Vives R, *et al.* Clinical cross-reactivity among foods of the Rosaceae family. *J Allergy Clin Immunol.* 2000;106:183-9.
13. Kondo Y, Tokuda R, Urisu A, Matsuda T. Assessment of cross-reactivity between Japanese cedar (*Cryptomeria japonica*) pollen and tomato fruit extracts by RAST inhibition and immunoblot inhibition. *Clin Exp Allergy.* 2002;32:590-4.
14. Caliskaner Z, Naiboglu B, Kutlu A, Kartal O, Ozturk S, Onem Y, *et al.* Risk factors for oral allergy syndrome in patients with seasonal allergic rhinitis. *Med Oral Patol Oral Cir Bucal.* 2011;16(3):312-6.
15. Fokkens W, Jogi R, Reinartz S, Sidorenko I, Sitkauskienė B, van Oene C, *et al.* Once daily fluticasone furoate nasal spray is effective in seasonal allergic rhinitis caused by grass pollen. *Allergy.* 2007;62(9):1078-84.
16. Bousquet J, Khaltaev N, Cruz AA, Denburg J, Fokkens WJ, Togias A, *et al.* Allergic Rhinitis and its Impact on Asthma (ARIA) 2008. Update (in collaboration with the World Health Organization, GA (2) LEN and AllerGen). *Allergy.* 2008;63 Suppl 86:8-160.
17. Ortolani C, Ispano M, Pastorello EA, Ansaloni R, Magri GC. Comparison of results of skin prick tests (with fresh foods and commercial food extracts) and RAST in 100 patients with oral allergy syndrome. *J Allergy Clin Immunol.* 1989;83(3):683-90.
18. Sloane D, Sheffer A. Oral allergy syndrome. *Allergy Asthma Proc.* 2001;22:231-5.
19. Roje Ž, Račić G. Oral allergy syndrome. *Medicus.* 2011;20(2):251-6.
20. Ono E, Maeda Y, Tanimoto H, Fukutomi Y, Oshikata C, Sekiya K, *et al.* Clinical features of oral allergy syndrome to plant foods allergens in Kanto regions. *Arerugi.* 2007;56:587-92.
21. Inomata N, Morita A, Kirino M, Yamazaki H, Yamaguchi J, Yamane Y, *et al.* Oral allergy syndrome due to plant-derived foods: a clinical review of 63 patients over a period of 6 years. *Arerugi.* 2007;56:1276-84.

Sažetak

UČESTALOST I RIZIČNI ČIMBENICI ZA RAZVOJ ORALNOG ALERGIJSKOG SINDROMA U ODRASLIH BOLESNIKA SA SEZONSKIM ALERGIJSKIM RINITISOM

N. Ivičević, Ž. Roje, Z. Kljajić, L. Bojić, G. Kardum, L. Omero i G. Račić

Cilj našega istraživanja bio je procijeniti učestalost oralnog alergijskog sindroma (OAS) u bolesnika sa sezonskim alergijskim rinitisom (SAR) te moguće rizične čimbenike za njegov nastanak. Ovo presječno istraživanje provedeno je u ordinacijama primarne zdravstvene zaštite Splitsko-dalmatinske županije u razdoblju od 1. ožujka 2012. do 30. rujna 2012. godine. Izvori podataka su bili: medicinska dokumentacija bolesnika ("karton") u kojemu je potvrda dijagnoze SAR-a (pozitivan kožni-ubodni test na sezonske inhalacijske alergene: poleni trava, stabala i korova), antropometrijski podatci o bolesniku (spol, dob, težina i visina) i upitnik u kojemu bolesnici ocjenjuju nosne i očne simptome, komorbiditet i stil života. Od statističkih testova koristili smo χ^2 -test, Pearsonov χ^2 -test, Spearmanov test korelacije i Kolmogorov-Smirnovljevi test. Učestalost OAS-a bila je 45,7%. Rizični čimbenici za njegov nastanak bili su: dijabetes ($p < 0,001$) te ozbiljniji nosni ($p < 0,05$) i očni simptomi ($p < 0,001$). Može se zaključiti da je učestalost OAS-a u Splitsko-dalmatinskoj županiji visoka. Rizični čimbenici za nastanak OAS-a u bolesnika sa SAR-om su dijabetes i veća izraženost očnih i nosnih simptoma.

Ključne riječi: Oralni alergijski sindrom; Rizični čimbenici; Rinitis, alergijski, sezonski; Hrvatska