

Pannonian ostracods from the southwestern Transylvanian basin

Erika Kovács¹, Imre Magyar^{2,3}, Orsolya Sztanó⁴ and Radovan Pipík¹

¹ Slovak Academy of Sciences, Earth Science Institute, Ďumbierska 1, 974 01 Banská Bystrica, Slovak Republic; (kovacs@savbb.sk; pipik@savbb.sk)

² MOL Hungarian Oil and Gas Plc., Október 23. utca 18, 1117 Budapest, Hungary; (corresponding author: immagyar@mol.hu)

³ MTA-MTM-ELTE Research Group for Paleontology, H-1431 Budapest, POB 137.

⁴ Eötvös Loránd University, Department of Physical and Historical Geology, Pázmány Péter sétány 1/C, H-1117 Budapest, Hungary; (sztano@ludens.elte.hu)

doi: 10.4154/gc.2016.16



Abstract

Pannonian (Late Miocene) ostracods were investigated from 7 outcrops, exposing deep-water lacustrine sediments along the western margin of the Transylvanian Basin. Sedimentological patterns in the outcrops indicate deposition in sublittoral to profundal environments where fine-grained, suspension fall-out sediments are intercalated with various types of mass transport deposits, most commonly turbidites. The ostracod fauna, consisting of 30 taxa, is dominated by endemic species of the brackish Lake Pannon. The assemblages indicate a mixture of deep-water (sublittoral to profundal) species, such as Paratethyan *Candoninae*, and shallow-water (littoral) species, belonging to *Cyprideis*, *Loxococoncha*, *Amnicythere* and *Hemicytheria*. The mixed character of the assemblages is most probably a consequence of the reworking of littoral specimens into deeper environments, a phenomenon also observed in the molluscan fauna from some of the investigated outcrops. Alternatively, some endemic and extinct species of the littoral genera might have adapted to the deep-water environment. A significant decrease in the abundance and diversity of ostracods from west to east is interpreted reflecting increasingly distal environments. The ostracod fauna indicates the *Hemicytheria hungarica* (2 outcrops), *Hemicytheria tenuistriata* (3 outcrops), and *Propontoniella candeo* Zones (2 outcrops) of the lower Pannonian Slavonian Substage.

Article history:

Manuscript received July 18, 2015

Revised manuscript accepted April 17, 2016

Available online June 01, 2016

Keywords: Ostracoda, biostratigraphy, Late Miocene, Pannonian, Transylvanian Basin

1. INTRODUCTION

Study of ostracods of the Neogene Paratethys has substantially contributed to the stratigraphic, environmental, and palaeogeographic understanding of this huge and complex marine to lacustrine depositional system (DANIELOPOL et al., 2008a). Of the large brackish lakes that repeatedly formed during the history of the Paratethys, the Late Miocene-Early Pliocene Lake Pannon was probably the longest-lived (MÜLLER et al., 1999; HARZHAUSER & MANDIC, 2008). As a consequence, this lake was inhabited by an astonishingly diverse lacustrine biota (e.g. PAPP et al., 1985; STEVANOVIĆ et al., 1990).

The dominantly endemic ostracods of Lake Pannon have been studied and described by generations of palaeontologists, starting with REUSS (1850), and leading to the introduction of ca. 500 new species. The taxonomy, environmental requirements, and palaeogeographic distribution of these ostracods, however, is far from being well understood, and needs further study (DANIELOPOL et al., 2008a). In particular, the Transylvanian Basin (TB), representing the easternmost part of the early Lake Pannon, lacks modern and well-illustrated documentation of the endemic ostracod taxa. The objective of this paper is to present the Late Miocene (Pannonian) ostracod faunas from several outcrops of the southwestern TB, and to explore their palaeoecological, palaeobiogeographic and biostratigraphic context.

2. NEOGENE EVOLUTION OF THE TRANSYLVANIAN BASIN

The TB is bordered by the Eastern and Southern Carpathians, and the Apuseni Mountains (Fig. 1). The basin fill can be divided into several tectono-stratigraphic megasequences (KRÉZSEK, 2005; KRÉZSEK & BALLY, 2006), the uppermost one of which

comprises the Middle and Upper Miocene (Badenian, Sarmatian and Pannonian Central-Paratethys stages). The Middle Miocene to Quaternary evolution of the TB is related to the Carpathian orogene (ROYDEN, 1988; SCHMID et al., 2008). During this period it evolved as a compressional back-arc basin (DE BROUCKER et al., 1998; CIULAVU, 1999; SANDERS et al., 1999; KRÉZSEK et al., 2010), and its sedimentary history was connected to the subduction along the Carpathian arc. Although a very high rate of subsidence is encountered during this period, no major extensional faults are reported; instead, most structures are related to salt-tectonics (KRÉZSEK & BALLY, 2006). Middle Badenian evaporites are widespread in the subsurface of the entire TB (GHERGARI et al., 1991; KOVÁČ et al., 2007), and salt tectonics was most effective from the Sarmatian to the Pliocene (KRÉZSEK & BALLY, 2006).

Following the fully marine conditions of the Badenian and restricted marine environment in the Sarmatian, the uplift of the Carpathians closed the gateways between the Central and Eastern Paratethys at the end of the Middle Miocene and thus isolated the intra-Carpathian region from the rest of the Paratethys (TERBORGH et al., 2013). With the onset of the Late Miocene Pannonian age, the TB became part of the large brackish Lake Pannon (MAGYAR et al., 1999). The Sarmatian–Pannonian boundary in the TB was traced in a large outcrop of marls and turbiditic sandstones in Oarba de Mureş (SZTANÓ et al., 2005), on the basis of dinoflagellate biostratigraphy (SÜTŐ-SZENTAI & SZEGŐ, 2008), magnetostratigraphy and radiometric dating (VASILIEV et al., 2010), and foraminifer and ostracod biostratigraphy (FILIPESCU et al., 2011). The boundary was unanimously marked on the basis of microfossils. Although the magnetostratigraphic interpretation of the measured polarity pattern is contradictory, high-precision radiometric dating of a volcanic ash deposit from the very top of

the one-km-thick Sarmatian sequence (VASILIEV et al., 2010) indicates that the Sarmatian-Pannonian boundary is about 11.6 Ma old, in accordance with recent magnetostratigraphic interpretations from the Vienna basin (PAULISSEN et al., 2011) and the Pannonian basin proper (TER BORGH et al., 2013).

After the isolation event, the water level of Lake Pannon gradually rose, controlled both by increased precipitation and subsidence all over the Pannonian Basin system (MAGYAR et al., 1999; GARCIA-CASTELLANOS, 2006; HARZHAUSER et al., 2008). Pulses of tectonic uplift with short wave lengths are recorded by coarse-grained fan delta systems along the basin margins (MAŢENCO et al., 2010). These fan deltas prograding from the Carpathians and from the Apuseni Mts. toward the basin centre fed the deep-lacustrine turbidite systems interfingering with hemipelagics/profundal marls. Sedimentary facies of the Pannonian deposits thus reflect deposition in a deep lacustrine setting, including lacustrine deep-water fans (KRÉZSEK & FILIPESCU, 2005; SZTANÓ et al., 2005; KRÉZSEK et al., 2010). Shallow-water deposits are subordinate and were preserved mostly in the eastern part of the basin. The final infill of the basin did not take place until the end of the Miocene, but these deposits are seldom preserved, except where covered by back-arc volcanics (KRÉZSEK and FILIPESCU, 2005; MAŢENCO et al., 2010; KRÉZSEK et al., 2010).

Apart from Pliocene lacustrine sediments in the southeastern corner of the TB (for a modern review see HARZHAUSER & MANDIĆ, 2008), Late Miocene Lake Pannon sediments are the youngest deposits in the basin fill succession. Their present-day thickness is usually less than 500 m (KRÉZSEK et al., 2010). Due to Pliocene to Quaternary uplift, more than 500 m of sediments have been eroded particularly in the northernmost part of the TB (DE BROUCKER et al., 1998; KRÉZSEK & FILIPESCU, 2005). Erosion is still active today in the basin, where the altitude varies between 300 and 600 m above sea level (MAŢENCO et al., 2010).

3. PANNONIAN OSTRACODS FROM THE TRANSYLVANIAN BASIN

The first contribution to the knowledge of Pannonian ostracods of the TB was provided by HÉJJAS (1894), who considered this fauna special and isolated. His observation that the ostracod fauna indicates deep water environment was ahead of his time (WANEK, 1992a). CHINTĂUAN (1971) investigated Pannonian ostracods from the western part of Munţii Bârgăului. CLICHICI (1980) described an ostracod association from near the Aiud River. This fauna, representing the C-D-E zones of the Pannonian (FILIPESCU, 1999) included: *Erpetocypris abscissa* (POKORNÝ), *Candona trapezoidea* (MÉHES), *Candona* sp., *Loxococoncha granifera* (REUSS) and *Leptocythere auriculata* (MÉHES).

WANEK (1992b) investigated the Sarmatian and Pannonian ostracod fauna of the TB and compared his results with the data of HÉJJAS (1894). He concluded that the complete Sarmatian and the Slavonian Substage of the Pannonian could be identified in the TB.

OLTEANU (in POPESCU, 1995) mentioned some „upper Pannonian–Pontian” ostracod taxa from Lopadea Veche: *Amplocypris abscissa* (REUSS), *A. recta* (REUSS), *Pannonocypris auriculata* [= *Herpetocyprilla auriculata*] (REUSS), *Lineocypris trapezoidea* (ZALÁNYI), *Hemicytheria josephinae* (KOLLMANN), *Caspiocypris pontica* (SOKAČ), *Cyprideis pannonica* (MÉHES).

WANEK & FILIPESCU (1994) reported an upper Slavonian ostracod assemblage from Moldoveneşti. A diverse lower Pan-

nonian ostracod fauna was described from Ormeniş by WANEK (in FILIPESCU, 1996); the fauna was composed of *Hungarocypris auriculata* [= *Herpetocyprilla auriculata*] (REUSS), *Amplocypris reticulata* (HEJJAS), *Candona* (*Typhlocypris*) cf. *redunca* (ZALÁNYI), *C. (Caspiocypris) transilvanica* HÉJJAS, *C. (Caspiocypris) improba* KRSTIĆ, *Leptocythere (Amnicythere) paralella* (MÉHES), *Leptocythere (Amnicythere) rodrigera* (POKORNÝ), *Leptocythere (Amnicythere) servica* KRSTIĆ, *Loxococoncha rhombovalis* POKORNÝ, *L. porosa* (MÉHES), *Loxocorniculina hodonica* POKORNÝ, *Graptocythere loerentheyi* MÉHES, *Hemicytheria folliculosa* (REUSS), and *Limnocythere* aff. *aquaensis* CARBONNEL. At Oarba de Mureş (section A), FILIPESCU et al. (2011) marked the Sarmatian–Pannonian boundary at the first occurrence of ostracod assemblages containing deep-water species of the *Amplocypris-Candona* group.

DE LEEUW et al. (2013) tried to obtain better age constraints on the Middle to Late Miocene Central Paratethyan regional stages (Badenian, Sarmatian and Pannonian) through integrated magnetostratigraphic, radio-isotopic and biostratigraphic research in Transylvania. At Mediaş, Sighişoara, and Târnăveni, the ostracod assemblages indicated a Pannonian age. In the case of Viforoasa and Şoimuş, 3 ostracod biozones were identified following the zonation of KRSTIĆ (1985): the *Propontoniella candeo* Zone, indicating the uppermost part of the Slavonian Substage, and the *Amplocypris abscissa* and *Hemicytheria croatica* Zones, indicating the lower part of the Serbian Substage.

4. MATERIAL AND METHODS

We have selected 7 outcrops from the southwestern part of the TB, where description of the sedimentary facies and interpretation of the depositional environment was followed by sampling for ostracods, preferably from fine-grained sediments where in situ burial of the shells seemed more probable.

Altogether 10 micropaleontological samples (250 g air-dried material) were collected. These samples were washed in hydrogen peroxide (10%), sieved using a 150 µm sieve and oven dried. After a preliminary observation under stereomicroscope, the ostracods were examined and photographed using a scanning electron microscope. A JEOL JEE 4B vacuum evaporator was used for deposition of gold on the surface of specimens. Pictures were taken with SEM (AMRAY 1830) at an accelerating voltage of 20kV in the SEM laboratory of the Department of Petrology and Geochemistry, Eötvös Loránd University, Budapest.

All the ostracod valves, both adults and juveniles, whole and broken specimens were counted from each sample.


5. THE INVESTIGATED OUTCROPS AND THEIR OSTRACOD FAUNA

5.1. The Lopadea Veche/Oláhlapád locality

This sampled outcrop is located 1 km east of the Lopadea stream valley (46°22'45.36"N, 23°39'55.74"E) (Fig. 1). Lopadea Veche belongs to the best studied Pannonian exposures in the TB (LÖRENTHEY, 1893; KOCH, 1900; PÁVAY-VAJNA, 1910; GILLET, 1943; ILIE, 1952; LUBENESCU & LUBENESCU, 1976; OLTEANU in POPESCU, 1995).

Today several small outcrops can be found in the hilly outskirts of the village. They expose an alternation of argillaceous marl and yellow, fine-medium-grained sands, building a 7–8 m thick succession (Fig. 2). Given the lack of specific sedimentary structures it is difficult to decide if the sand was transported by gravity-driven processes or just currents/waves/storms. Both the marls and the sandstones are friable and the carbonate cementa-

Figure 1. Geological map of the Transylvanian Basin with the distribution of Pannonian sediments and location of the sampled outcrops (adapted from DE LEEUW et al., 2013).



tion is poor. The thickness of sand layers increases upwards from 2–3 cm to 10–15 cm. Along the bedding planes plant fragments, branches, and poorly preserved leaf imprints occur together with a rich mollusc assemblage equally consisting of deep-water (*Congeria banatica* R. HÖRNES, *Paradacna* cf. *lenzi* (R. HÖRNES),

Undulotecha pancici (BRUSINA)) and shallow-water forms (*Melanopsis vindobonensis* FUCHS). Our sample was collected from a marl bed in the lower part of the outcrop (Fig. 2).

The ostracod association is well preserved and evenly distributed in the argillaceous marl, reflecting autochthonous or al-






Figure 2. Lopadea Veche; A – alternation of argillaceous marl and yellowish, fine- and medium-grained sand layers, several cm in thickness. White dot indicates sampling location. B – close-up view of the sampled argillaceous marl.




Gârbova de Jos



B

Gârbova de Jos



C

Figure 3. Gârbova de Jos; A – alternation of clast-supported, graded conglomerates, pebbly mudstones, and mud-clast conglomerates; the outcrop is cut by a large normal fault. White dot indicates sampling location. B and C – thin bedded turbidites (fine-grained, graded sandstone beds and mudstone with mud clasts).

most autochthonous burial. This material consists of 173 valves and carapaces, the latter representing 10% of the specimens; juvenile individuals are subordinate in the sample. The assemblage is rich in species, including *Amplocypris* cf. *reticulata* (Pl. 1, Figs. 20–22), *A. sp.*, *Herpetocyprilla hieroglyphica* (MÉHES), *Candona (Caspionella) praebalcanica* KRSTIĆ, *Hemicytheria folliculosa*, *Cyprideis pannonica* (Pl. 3, Figs. 5–8), *Loxocorniculina hodonica*, *Candona (Typhlocypris) trigonella* (HÉJJAS) (Pl. 1, Figs. 29–32), *Candona (Propontoniella) sp.*, and *Candona (Caspio-*

pris) transilvanica (Fig. 8). The most common genera are *Amplocypris* and *Herpetocyprilla*.

Species of *Cyprideis*, *Loxoconcha* and related genera typify marginal ostracod assemblages world-wide (BOOMER & EISENHAUER, 2002; BOOMER et al., 2005). According to KRSTIĆ and STANCHEVA (1990), *Cyprideis* is common in the marginal parts of the Pannonian basin, but missing from the deep basins. *Cyprideis* is therefore an indicator of shallow waters. It inhabits mostly brackish (mesohaline) environments, showing the great-




Figure 4. Gârbovița; A – alternation of marl beds and medium- to very fine-grained, graded sandstone layers. B – close-up view of the marl layers, stained by organic matter. White dot indicates sampling location.



Figure 5. Mihaiț. Decimetre to metre thick, massive or laminated marl layers intercalated with thin sandstone beds. White dot indicates sampling location.

est abundance at salinities of 2–16.5‰ (WAGNER, 1964), but certain species of the genus may adapt to hypersaline environments (MEISCH, 2000). Pannonian assemblages with *Cyprideis*, *Amplocypris*, *Amnicythere*, *Hemicytheria* and *Loxococoncha* were probably highly tolerant of fluctuations in salinity: they occupied estuaries, i.e. semi-enclosed coastal water bodies with a salinity gradient from brackish to freshwater, in the southern part of the Pannonian basin (e.g. Kolubara basin in northwestern Serbia; RUNDIĆ, 2006). *Loxocorniculina* is known from brackish water ostracod assemblages of the Paratethys and the Eastern Mediterranean (PIPIK & BODERGAT, 2004). According to KORECZ (1991), *Candona (Caspiolla) praebalcanica* inhabited plio-mesohaline environments, with 15–80 m water depth, and *Candona (Caspiocypris) transilvanica* inhabited shallow but calm environments.

The entire ostracod assemblage shows a mixture of shallow- and deep-water species, probably as a result of redeposition of littoral forms into deeper parts of the basin. *Candona (Caspiolla) praebalcanica* appears in our material with complete carapaces filled with sediment, a phenomenon indicating rapid burial (HOLMES, 2001). Otoliths and fish teeth also occur in the microfauna (Pl. 3, Figs. 9–16).

5.2. The Gârbova de Jos/Alsóorbó locality

The 20 m high outcrop is located near the village of Gârbova de Jos (46°27'25.58"N, 24°17'10.57"E) (Fig. 1). The outcropping succession is cut by a large normal fault of several metres offset, therefore a relatively thick sedimentary succession can be studied (Fig. 3). It comprises clast-supported, graded conglomerates, thick pebbly mudstones, mud-clast conglomerate, pebbly sandstones, and coarse- to medium-grained and medium- to fine-grained graded sandstone beds intercalated with mudstones. Individual beds are 1–50 cm thick. Poor normal grading and plane- to cross-lamination were recognized. Clast size occasionally attains 50 cm in diameter. These boulders are covered by horizontally laminated, deformed mudstone layers. Higher up, mudstone with a few centimetre thick graded sandstone intercalations becomes dominant. The top of the fine-grained sandstones rarely contains small vertical burrows. These beds are the products of sandy-silty turbidity currents, gravelly-sandy turbidity currents, and debris flows. The latter indicates the proximity of the basin-margin slope. This interpretation is supported by the composition of the gravel, indicating the Apuseni Mts. as a source area. According to KRÉZSEK et al. (2010), a shallow marine ramp supplied coarse clastics to the open shelf; the frequent, large-volume mass wasting events, however, may suggest a steeper slope, such as that of a fan-delta, and to deeper water, probably below the „shelf-edge”.

The studied sample was collected from the lower part of the outcrop, from the dark grey suspension fall out mudstone, which also contained thin shells of deep-water bivalves (*Paradacna* cf. *lenzi*, *Congeria* sp.).

The ostracod association is very poorly preserved with thin and translucent valves. In addition to the fragments, only 48 complete valves were observed in our material. The fauna includes *Amplocypris* sp., *Candona (Typhlocypris)* sp., *Candona (Propontoniella)* sp. and *Hemicytheria folliculosa* (Fig. 8).

Except for the small and ornamented valves of *H. folliculosa*, which are well-preserved, the valves are damaged and broken. In general, ostracods with smooth, thin and elongated carapaces suggest a calm, relatively deep environment (DANIELOPOL, 1980; RUNDIĆ, 2006). Specimens of the littoral *Hemicytheria folliculosa* were probably transported into this environment.

5.3. The Gârbovița / Középorbó locality

This 20 m high outcrop is situated on the right bank of the Gârbova stream valley, across from the village (46°16'51.66"N, 23°38'07.26"E) (Fig. 1). The fossils of this outcrop were briefly treated by WANEK & FILIPESCU (1994); they mention fossil leaves of the tropical plant *Daphnogene*, molluscs, and ostracods, the latter indicating the Pannonian „C” zone of PAPP (1951).

The outcrop exposes marl beds and medium- to very fine-grained, graded, planar to cross-laminated sandstone layers of 5–10 cm thickness; the latter are interpreted as turbidites. The marl layers are dark in colour due to plant remains (organic matter) (Fig. 4). The depositional setting can be best understood by comparing this outcrop with the previous one, where the same marls and sandstones crop out together with various types of conglomerate (KRÉZSEK, 2005).

Congeria partschi CZJZEK and *Melanopsis vindobonensis* – molluscs that lived in shallow-water lacustrine environments – are common in the redeposited turbidite layers. The silty marl layers contain a rich sublittoral mollusc assemblage: *Paradacna* cf. *lenzi*, *Lymnocardium* cf. *majeri* (M. HÖRNES), *Lymnocardium* cf. *otiophorum* (BRUSINA), *Congeria banatica*, „*Gyraulus*” *ponticus* (LÖRENTHEY), *Micromelania* sp. occur together with fish teeth and otoliths. Our sample was taken from this deep-water marl facies.

The ostracod assemblage from Gârbovița is abundant and diverse; 225 ostracod valves were picked from the dried material. The fauna includes the following species (Fig. 8): *Amplocypris firmus* KRSTIĆ (Pl. 1, Figs. 15–19), *A. matejici* KRSTIĆ (Pl. 1, Figs. 11–14), *Herpetocyprilla auriculata* (REUSS), *H. hieroglyphica*, *Candona (Caspiolla) praebalcanica* (Pl. 1, Figs. 23, 24), *C. (Typhlocypris) trigonella*, *C. (Caspiocypris) transilvanica*, *C. (Propontoniella)* sp., *Hemicytheria folliculosa* (Pl. 2, Figs. 27–32), *Loxocorniculina hodonica* (Pl. 3, Figs. 2, 3), and *Cyprideis* sp.

Three fossil species of *Herpetocyprilla* have been recognised in the deposits of Lake Pannon, and they were previously assigned to *Hungarocypris* (DANIELOPOL et al., 2008b). A single extant representative of the genus, *Herpetocyprilla mongolica* DADAY, is endemic to Lake Issyk-Kul, Kyrgyzstan, Central Asia. It lives together with recent thalassic and athalassic *Cyprideis torosa* and *Tyrrhenocythere amnicola* at salinity of 6‰ (BRONSTEIN, 1947; RICKETTS et al., 2001). Other shallow-water forms, such as *Hemicytheria*, *Loxococoncha*, and *Cyprideis*, are subordinate in the sample (8, 6 and 3% of the total specimens, respectively).

5.4. The Mihaiț/Mihálcfalva locality

The sampled outcrop, approximately 60–80 m high and 200–300 m long, is situated 1 km N of the village, along the road DJ142L, on the left bank of the Mureș river (46°10'35.05"N, 23°42'20.70"E) (Figs. 1, 5). Decimetre to metre thick, massive or laminated marl and calcareous marl comprise the sediment succession with some rare, thin, medium- to fine-grained sandstone intercalations. The marls are mostly suspension fall out deposits accumulated on the basin plain. Thin-shelled deep-water molluscs, such as *Paradacna* cf. *lenzi*, *Congeria banatica*, *Congeria* sp., *Undulotheca pancici*, *Micromelania* sp. are relatively common in the massive marl from which the sample for ostracods was collected.

The material consists of 56 ostracod valves, including different indeterminate species of Paratethyan Candoninae, and *Loxococoncha rhombovalis*, *Hemicytheria folliculosa*, and *Amnicythere miscere* (KRSTIĆ) (Fig. 8; Pl. 2, Figs. 25, 26, Pl. 3, Fig. 1).

Smooth, thin and translucent valves of *Candona* have the most significant role in the association, but they are unidentifiable



Figure 6. Tău. An erosional surface divides the outcrop into a lower unit, dominated by mudstones formed as levee to overbank deposits intercalated with rare overspilling thin-bedded turbidites, and an upper unit with the complex, migrational deep-water channel fill. White dot indicate sampling locations within the lower unit.



Figure 7. Oarba de Mureș, outcrop E. Massive and bioturbated clay marls. White dot indicates sampling location.




Figure 8. Pie charts of species distribution in each sample (percentage of specimens in all specimens from 250 g sediment).

to species level due to their very poor preservation. The small and ornamented valves of the other taxa are well preserved. *Loxocochna* occurs in meso- to euhaline, littoral environments (MORKHOVEN, 1963). Numerous *Amniccythere* species live in the littoral and sublittoral zones of the Black and Caspian Seas at brackish salinity, but some species have been observed in freshwater bays and estuaries. *Amniccythere multituberculata* (LIVENTAL) lives today in the central and southern Caspian Basins at salinities of 11.5–13.5‰. It is missing in the northern Caspian Basin, where freshwater input from Volga River effectively lowers salinity (SCHORNIKOV, 1964). The freshwater Lake Ohrid, however, also hosts some endemic species of the genus, such as *Amniccythere karamani* (KLIE) (NAMIOTKO et al., 2012).

5.5. The Tău/Székástóhát locality

The outcrop is almost 1 km wide, where a 12–15 m high continuous succession can be followed over a distance of 300 m. It occurs along road DC203 (46°5'20.14"N, 23°50'33.36"E) (Fig. 1).

The central part of the outcrop can be divided into two units (Fig. 6). The lower unit consists of thin beds of mudstone intercalating with very thin, fine- or medium-grained sandstone beds. This unit contains a deep-water mollusc association with *Paradacna* cf. *lenzi*, *P.* cf. *abichi* (R. HÖRNES), *Lymnocardium* sp., *Congeria czjzeki* M. HÖRNES, *C.* sp., *Velutinopsis* sp., *Undulotherca pancici*, *Micromelania* sp. The upper unit starts with a wide erosional scour and is composed up of thick, often amalgamated, massive beds of medium- to very coarse-grained, granular sandstone. Rip-up mud clasts and water-escape structures are common. Shallow and laterally stepping erosional surfaces are also characteristic.

Sand beds in both units are the products of turbidity currents. The lower unit dominated by mudstones was formed as levee to overbank deposits with rare overspilling thin-bedded turbidites. The upper sandstone represents a complex, laterally migrating deep-water channel fill (KRÉZSEK et al., 2010; TÓKÉS, 2013) (Fig. 5). As a whole, a major isolated channel-levee system in a deep, profundal environment can be reconstructed in the outcrop.

Two samples were taken for ostracods; one from the lower part and another from the upper part of the fine-grained overbank sediments (lower unit).

The assemblage from the lower part of the marl unit is poorly preserved and consists of 88 ostracod valves (Fig. 8). The fauna includes *Amplocypris* sp., *Candona* sp., *Hemicytheria folliculosa*, *H. reticulata* (SOKAČ) (Pl. 2, Fig. 24), and *Amniccythere* sp. The assemblage from the upper part of the marl unit is poor in species and individuals (the sample consists of 57 ostracod valves; Fig. 8). The fauna includes *Hemicytheria folliculosa*, *H. reticulata*, *Amniccythere* sp., and *Amplocypris* sp.

Candona is rare in the lower sample and completely missing from the upper one. *Hemicytheria* and *Amplocypris* are dominant in the upper marl bed. *Amniccythere* occurs in both samples, but it is more abundant in the upper one. *Hemicytheria*, a representative of the littoral environment, indicates resedimentation processes.

5.6. The Cunța/Konca locality

The sampling location is about 1 km west of the village of Cunța at the western border of its administrative area (45°92'63.63"N, 23°70'21.23"E) (Fig. 1). LÖRENTHEY (1893) published a list of molluscs from Cunța (at that time spelled as Koncza), including *Melanopsis fossilis* (GMELIN), *M. pygmaea* HÖRNES, *M. bouei* FÉRUSSAC, *Plagiodacna carinata* (DESHAYES), *Congeria subglobosa* PARTSCH, *C. partschi*, *Congeria* sp., *Unio* sp., and

Theodoxus crenulatus (KLEIN). LÖRENTHEY (1893) thought that this fauna represented the „*Lyrcaea*-horizon”, the uppermost stratigraphic unit out of the three ones that he recognized in the Transylvanian basin. The exact location where fossils of these shallow-water species were collected, however, remains unknown.

The sampled outcrop displays decimetre to metre thick coarse-grained sandstone and pebbly sandstone beds, up to cobble-sized clasts. These are mostly well sorted with horizontal lamination. The sandstones alternate with bluish grey, bioturbated marls. Some marl layers are laminated. Fish remnants (otoliths, teeth) are common in the studied beds. Gravity mass flows and suspension settling alternated in the depositional environment. Three samples were collected from different levels of the bluish grey marl layers.

The sample from the lower part of the outcrop consists of 154 well-preserved ostracod valves. The most frequent genera are *Amplocypris* and *Herpetocyprilla*. The fauna includes the following species (Fig. 8): *Amplocypris matejici*, *A.* sp., *Herpetocyprilla auriculata* (Pl. 1, Figs. 6–10), *Candona (Thaminocypris)* sp., *C. (Caspicypris)* sp., *C. (Propontoniella)* sp., *C. (Typhlocypris) trigonella*, *C. (Lineocypris)* sp., *Cypria* sp. and *Loxocochna rhombovalis* (Pl. 3, Fig. 4).

The ostracod fauna from the middle part of the outcrop comprises a rich and well-preserved association. We found 344 juvenile and adult ostracod valves, representing the following species: *Candona (Lineocypris) dorsobrevis* KRSTIĆ (Pl. 1, Figs. 25–28), *C. (Caspicypris)* sp., *C. (Lineocypris)* sp., *C. (Propontoniella) macra*, *C. (Propontoniella)* sp., *C. (Caspicypris) transilvanica*, *C. (Caspioilla)* sp., *C. (Thaminocypris)* sp., *Amplocypris* sp., *A. firmus*, *Herpetocyprilla auriculata*, *C. (Caspicypris) aspera* (HÉJJAS), and *Hemicytheria folliculosa* (Fig. 8).

The ostracod fauna from the upper part of the Cunța outcrop is the richest and most diverse association in this study; we found 665 well-preserved ostracod valves in the material. The association includes *Amplocypris reticulata*, *A. matejici*, *Herpetocyprilla hieroglyphica* (Pl. 1, Figs. 1–5), *Candona (Caspicypris) transilvanica* (Pl. 2, Figs. 5–8, 13–15), *C. (Propontoniella) macra* (Pl. 2, Figs. 9–12), *C. (Caspioilla)* sp., *C. (Caspicypris)* cf. *aspera* (Pl. 2, Figs. 1–4), *C. (Lineocypris)* sp., *C. (Lineocypris) dorsobrevis*, *Hemicytheria folliculosa*, *Cypria pannonica* (KRSTIĆ) (Pl. 2, Figs. 20–23), *Cypria* sp., and *Amniccythere* sp. (Fig. 8).

The Cunța assemblages, especially from the middle and upper parts of the section, differ from our other samples in the diversity and dominance of various Paratethyan Candoninae species. In general, the majority of *Candona* species are freshwater, tolerating a slight increase of salinity, but the endemic Lake Pannon Candoninae adapted to brackish water (MEISCH, 2000; CZICZER et al., 2008). Typically trapezoid-shaped species of the subgenus *Lineocypris* occur predominantly in the deep parts of large lakes (MORKHOVEN, 1963).

The appearance of *Cypria* in both the lower and upper samples is also a unique feature of the Cunța outcrop. Most species of *Cypria* live in freshwater (lakes, pools) preferring environments with abundant vegetation, although they tolerate mesohaline conditions as well (MEISCH, 2000). The appearance of this genus therefore may indicate freshwater influence, probably proximity of a deltaic environment, in accord with the presence of *Unio* in the Cunța mollusc fauna (LÖRENTHEY, 1893).

5.7. The Oarba de Mureș/Marosorbó locality

The investigated outcrop is on the right bank of the river Mureș near Oarba de Mureș village (46°27'25.58"N, 24°17'10.57"E)

Table 1. Occurrences of Pannonian ostracod species in the investigated outcrops and the stratigraphic distribution of each species according to the literature (POKORNÝ, 1944; KRSTIĆ, 1971, 1972a, b, 1973a, b, 1985; SOKAČ, 1972, 1990; JIŘIČEK, 1985; WANĚK, 1992a; FILIPESCU, 1996; RUNDIĆ, 1998, 2006; RUNDIĆ et al., 2011; FILIPESCU et al., 2011).

Regional Chronostratigraphy		PANNONIAN																					
		Slavonian Substage						Serbian Substage															
Age		11.6 Ma						ca. 9 Ma															
OSTRACODA BIOZONES	POKORNÝ, 1944	α			β			γ		δ													
	PAPP, 1951	A	B	C	D		E																
	KOLLMANN, 1960	A	B	C	D		E		F	G	H												
	JIŘIČEK, 1985	A		B		C		D	E1	E2	E3												
	KRSTIĆ, 1985	Hemicytheria lörentheyi	Hemicytheria hungarica	Hemicytheria tenuistriata	Propontoniella candeo		Amplocypris abscissa	Hemicytheria croatica	Serbiella sagittosa	Typhlocyprina lineocypriformis													
Ostracod species	<i>Amplocypris reticulata</i> (HÉJJAS, 1894)			o									•										
	<i>Amplocypris firmus</i> KRSTIĆ, 1973				•								•										
	<i>Amplocypris matejci</i> KRSTIĆ, 1973				•																		
	<i>Amplocypris</i> sp.			o		o	•	o	o	•													
	<i>Herpetocyprina auriculata</i> (REUSS, 1850)				•									•	•								
	<i>Herpetocyprina hieroglyphica</i> (MÉHES, 1907)			•	o										•								
	<i>Candona (Thaminocypris) sp.</i>												•	•									
	<i>Candona (Typhlocypris) trigonella</i> (HÉJJAS, 1894)			•	•		o				o												
	<i>Candona (Typhlocypris) sp.</i>					o																	
	<i>Candona (Caspiocypris) aspera</i> (HÉJJAS, 1894)													•	o								
	<i>Candona (Caspiocypris) transilvanica</i> (HÉJJAS, 1894)				o	o								o	•								
	<i>Candona (Caspiocypris) sp.</i>							o					o	•									
	<i>Candona (Caspiolla) praebalcanica</i> (KRSTIĆ, 1972)				o	o		o															
	<i>Candona (Caspiolla) sp.</i>							o						o	•								
	<i>Candona (Propontoniella) macra</i> (KRSTIĆ, 1972)													o	•								
	<i>Candona (Propontoniella) sp.</i>				o	o	o		o				o	o	•								
	<i>Candona (Lineocypris) dorsobrevis</i> (KRSTIĆ, 1972)													o	•								
	<i>Candona (Lineocypris) sp.</i>													o	o	o							
	<i>Candona</i> sp.														•								
	<i>Hemicytheria folliculosa</i> (REUSS, 1850)			o	o	•	o		•	o	o	o	o										
	<i>Hemicytheria reticulata</i> (SOKAČ, 1972)										o	o											
	<i>Cyprideis pannonica</i> (MÉHES, 1908)			o																			
	<i>Cyprideis</i> sp.				o																		
	<i>Cypria pannonica</i> (KRSTIĆ, 1973)																						
	<i>Cypria</i> sp.																						
<i>Ilyocypris cf. expansa</i> (REUSS, 1850)										•													
<i>Amnicothera miscere</i> (KRSTIĆ, 1973)							o																
<i>Amnicothera</i> sp.											o	o											
<i>Loxocorniculina hodonica</i> (POKORNÝ, 1952)				o	o																		
<i>Loxocochla rhombovalis</i> (POKORNÝ, 1952)							o																

Outcrops

(Figs. 1, 7). The Upper Sarmatian to Lower Pannonian outcrops at Oarba de Mureş received much attention recently (SZTANÓ et al., 2005; SÜTÖ-SZENTAI & SZEGŐ, 2008; VASILIEV et al., 2010; FILIPESCU et al., 2011). The outcrop sampled (and designated as outcrop „E”) is located in the horizontal continuation of outcrop D of SZTANÓ et al. (2005), exposing massive and bioturbated clay marl with up to 0.5–1 m thick beds, and with intercalations of a few cm thick calcareous marls (Fig. 7). In this part of the succession sediments indicate deep basin-floor environment without mass-flow events. In one of the calcareous marl

layers, molluscs („*Lymnocardium*” *undatum* (REUSS), *Congerina* cf. *banatica*) were preserved as moulds. The sample was collected from a massive clay marl bed from the lower part of the outcrop.

The investigated ostracod fauna, consisting of 100 valves, has low diversity (Fig. 8). The assemblage includes the following brackish water ostracods: *Amplocypris* sp., *Candona (Caspiocypris)* sp., *C. (Caspiolla)* sp., *C. (Propontoniella)* sp., *C. (Typhlocypris) trigonella*, and *Ilyocypris cf. expansa* (REUSS) (Pl. 2, Figs. 16–19).

The most common species in the assemblage belong to *Amplocypris* and *Candona*. Their valves are thin, translucent and often broken. The oblong and elongated carapace shape may indicate benthic dwellers (DE DECKKER, 2002).

Ilyocypris cf. *expansa* seems to be an exotic element in the assemblage, because *Ilyocypris* is known to live in freshwater, tolerating slowly running water (MEISCH, 2000). Valves of the modern species *Ilyocypris lacustris* KAUFMANN, 1900 were recorded from freshwater lakes at depths of 35–139 m, but the presence of living specimens in deep water environment is uncertain (MEISCH, 2000). Its presence in this deep-water assemblage is probably due to reworking from a shallow, fresh-water-influenced environment.

6. DISCUSSION AND CONCLUSIONS

6.1. Palaeoenvironment and palaeogeography

The overall palaeoecological interpretation of the investigated ostracod faunas is difficult; probably much more samples would be required to understand the ecological significance of these endemic species. In general, the assemblages indicate a mixture of deep-water and shallow-water species. Thin-shelled, smooth Paratethyan Candoninae occupied the sublittoral to profundal zones of Lake Pannon, from the wave base down to the foot of the slope. The littoral zone was inhabited by forms with thicker and ornamented shells, such as, for instance, *Cyprideis*, *Loxococoncha*, *Amnicythere*, and *Hemicytheria*. In our samples, the deep-water and littoral taxa occur in various ratio.

Interestingly, the mollusc fauna shows similar characteristics across some of the investigated outcrops (Lopadea Veche, Gârbovița, and, according to the literature, Cunța). Thin-shelled bivalves, known to be confined to deep water environments elsewhere (e.g. *Paradacna abichi*, *Congerina banatica*, *Undulotheca pancici*), occur together (or in very close proximity to) typical littoral, herbivorous snails, such as *Melanopsis*. Sedimentological patterns in all the outcrops indicate various kinds of mass transport deposits, most commonly interpreted as turbidites. Although the ostracod samples were taken from fine-grained, suspension fall-out sediments, reworking and re-deposition of littoral ostracod valves into the deep-water environment is the most plausible explanation for the mixed character of the ostracod (and mollusc) faunas. In this case, the actual ostracod assemblages from the investigated part of the TB highly depend on local taphonomic factors. Alternatively, some species of the otherwise littoral ostracod genera may have adapted to living at greater depths; such examples are recorded from the Caspian Sea (BOOMER et al. 2005).

In terms of the abundance and diversity of ostracods, these outcrops can be divided into two distinct groups: Lopadea, Gârbovița, and Cunța (all the three samples) yielded more than 100 specimens (154 to 665) and at least 10 species (10 to 14), whereas Gârbova de Jos, Mihalț, Tau (both samples), and Oarba de Mureș yielded consistently fewer specimens (48–100) and species (4–6) from the same amount of sediment (Fig. 8). In addition, reworked littoral molluscs occur in the first group, but they were not observed in the second. The first group of outcrops lie closer to the basin margin, directly on the foothills of the Apuseni Mts, whereas the second group is located in a more basinward position (although the distance between Gârbovița and Gârbova de Jos is only a few kilometres). We hypothesize that these differences in abundance and diversity of the ostracods reflect environmental differences, such as deeper water and/or decreasing

levels of oxygen and nutrients, i.e. more distal environments towards the basin interior.

In the Caspian Sea (a Paratethyan remnant) which is the best modern analogue of Lake Pannon, BOOMER et al. (2005) reported a steep gradient in ostracod diversity with increasing water depth within the deep water realm. They found that the number of ostracod species decreased from 17 to 3 between 315 and 479 m. In addition, their samples showed that total specimen abundance declined exponentially with increasing water depth. We speculate that these patterns may be analogous with the diversity and abundance drop that we observed in our samples, although there are no means of estimating the absolute water depth in the early Pannonian TB.

6.2. Biostratigraphic interpretation and age

Various ostracod biostratigraphic systems have been erected in different parts of the Pannonian Basin System, but they usually reflect local characteristics and significantly differ from each other (e.g. POKORNÝ, 1944; KOLLMANN, 1960; SOKAČ, 1972; KRSTIĆ, 1973a, b, 1985; JIŘIČEK, 1985; RUNDIĆ, 1998, 2002, 2006; RUNDIĆ et al., 2011). GANIĆ et al. (2010) and TER BORGH et al. (2013) came to the conclusion that the „classical” Lake Pannon ostracod biozones are difficult to identify in the otherwise diverse and abundant ostracod record of the Beočin claypit (Fruška Gora, Serbia), which is roughly coeval with these Transylvanian samples, possibly because earlier biostratigraphic research underestimated the role of palaeoecological conditions in the occurrence and distribution of ostracods. A similar view was expressed by OLTEANU (2011).

As no special zonation for the TB has been currently produced, the biostratigraphic zonation of KRSTIĆ (1971, 1972a, b, 1973a, b, 1985) and SOKAČ (1972, 1990) originally elaborated for the southern parts of the Pannonian Basin is applied here. Recent stratigraphic observations from the TB (WANEK, 1992a; FILIPESCU, 1996; FILIPESCU et al., 2011) were also taken into account. For instance, WANEK (1992b) found that *Amplocypris reticulata* is probably an endemic species for the TB, and is present only in the „ α phase” of POKORNÝ (1944), corresponding to the lowermost Pannonian, i.e. lower part of the Slavonian Substage; *Loxococoncha rhombovalis* determines the final part of the Slavonian Substage; *Cyprideis pannonica* appears in the uppermost Sarmatian and is present in the „ α phase” of the Pannonian.

KRSTIĆ (1985) erected 8 stratigraphic ostracod zones for the Pannonian Stage: four zones in the older Slavonian and four in the younger Serbian Substages. The four Slavonian zones (from bottom to top) are the *Hemicytheria loerentheyi*, *Hemicytheria hungarica*, *Hemicytheria tenuistriata*, and *Propontoniella candeo* Zones. Characteristic species of the three latter zones from the present material include:

Hemicytheria hungarica Zone: *Cyprideis pannonica*, *Amplocypris reticulata*, *Amnicythere miscere*, *Herpetocyprilla auriculata*, *Candona (Casiocypris) transilvanica*.

Hemicytheria tenuistriata Zone: *Loxocorniculina hodonica*, *Herpetocyprilla auriculata*, *Hemicytheria folliculosa*, *Candona (Typhlocypris) trigonella*, *C. (Propontoniella) macra*, *C. (Casiocypris) transilvanica*, *Ilyocypris* cf. *expansa*.

Propontoniella candeo Zone: *Candona (Propontoniella) macra*, *C. (Thaminocypris) sp.*, *Herpetocyprilla hieroglyphica*, *Hemicytheria reticulata*, *Loxococoncha rhombovalis*.

The appearance of *Hemicytheria reticulata* in the latter biozone requires some discussion because SOKAČ (1972) and OL-




Plate 1. Pannonian ostracod assemblages from the southwestern part of the TB.

- 1–5 *Herpetocyprilla hieroglyphica* (MÉHES), Cunța; right valve (RV), internal lateral view (1), RV, external lateral view (2), left valve (LV), juvenile (juv.) (3), central muscle scars (4), pore canals on the ventral part of the RV (5).
- 6–10 *Herpetocyprilla auriculata* (REUSS), Cunța; LV, fragment (6), RV juv. (7), RV juv. (8), LV juv. (9), RV juv. (10).
- 11–14 *Amplocypris matejici* KRSTIĆ, Gârbovița; LV (11), RV (12), RV (13), LV (14).
- 15–19 *Amplocypris firmus* KRSTIĆ, Gârbovița; LV (15), RV (16), RV (17), RV (18), RV (19).
- 20–22 *Amplocypris cf. reticulata* (HÉJJAS), Lopadea Veche; LV (20), LV (21), RV (22).
- 23, 24 *Candona (Casiolla) praebalcanica* KRSTIĆ, Gârbovița; carapace, lateral view of LV (23), carapace, lateral view of RV (24).
- 25–28 *Candona (Lineocypris) dorsobrevis* KRSTIĆ, Cunța; LV (25), carapace, lateral view of RV (26), LV (27), RV (28).
- 29–32 *Candona (Typhlocypris) trigonella* (HÉJJAS), Lopadea Veche; carapace, lateral view of LV (29), carapace, lateral view of LV (30), carapace, lateral view of RV (31), carapace, lateral view of RV (32).




Plate 2. Pannonian ostracod assemblages from the southwestern part of the TB.

- 1–4 *Candona (Caspiocypris) cf. aspera* (HÉJJAS), Cunța; LV (1), RV (2), RV (3), LV (4).
 5–8 *Candona (Caspiocypris) transilvanica* (HÉJJAS), Cunța; RV (5), RV (6), LV (7), RV (8).
 9–12 *Candona (Propontoniella) macra* KRSTIĆ, Cunța; LV (9), RV (10), LV (11), RV (12).
 13–15 *Candona (Caspiocypris) transilvanica* (HÉJJAS), Cunța; LV (13), RV (14), carapace, lateral view of RV (15).
 16–19 *Ilyocypris cf. expansa* (REUSS), Oarba de Mureș; internal lateral view of RV (16), LV (17), RV (18), LV (19).
 20–23 *Cyprina pannonica* KRSTIĆ, Cunța; LV (20), LV (21), RV (22), LV (23).
 24 *Hemicytheria reticulata* SOKAČ, Tău; LV.
 25, 26 *Amnicythere miscere* (KRSTIĆ), Mihălț; LV (25), LV (26).
 27–32 *Hemicytheria folliculosa* (REUSS), Gârbovița; LV juv. (27), RV juv. (28), LV juv. (29), RV juv. (30), LV juv. (31), LV juv. (32).




Plate 3. Pannonian micropalaeontological assemblages from the southwestern part of the TB.

- 1 *Amnicythere miscere* (KRSTIĆ), Mihalt; LV.
 2, 3 *Loxocorniculina hodonica* POKORNÝ, Gârbovița; LV (2), LV (3).
 4 *Loxococoncha rhombovalis* POKORNÝ, Cunța; RV.
 5–8 *Cyprideis pannonica* (MÉHES), Lopadea Veche; RV, internal lateral view (5), RV, external lateral view (6), LV (7), RV (8).
 9–16 Fish skeletal fragments; fish teeth, Gârbovița (9–10), fish tooth, Mihalt (11), fish teeth, Cunța (12–13), fish teeth, Lopadea Veche (14, 16), fish squama, Lopadea Veche (15).

TEANU (2011) claim that this species occurs only in the Serbian substage „Late Pannonian”). Indeed, the specimens presented here differ from the original description of SOKAČ (1972) in lacking the sharp, high ventral rib and the pronounced posterior-ventral part. Such differences in the expression of ornamentation have been extensively discussed (for more details see KEEN, 1982), and they can be related to ecological or genetic factors, especially in the brackish environment of Lake Pannon in which ecological conditions were highly variable (PIPIK, 1998). Lake Pannon is well known for its extensive intralacustrine evolution (e.g. GEARY, 1990; PIPÍK 2007). This discovery of *H. reticulata* may represent an ancestor of *H. reticulata* in the sense of SOKAČ (1972), but this relationship should be demonstrated by detailed morphometric and evolutionary study of the subfamily Hemicytherinae in Lake Pannon.

In conclusion, the three ostracod biozones above are represented in the samples from the southwestern part of the TB (Table 1). Lopadea Veche and Gârbovița belongs to the *Hemicytheria hungarica* Zone. Samples from Gârbova de Jos, Oarba de Mureș, and Mihălț can be assigned into the *Hemicytheria tenuistriata* Zone. Finally, the ostracod fauna from Tău and Cunța indicates the *Propontoniella candeo* Zone. All these zones belong to the lower Pannonian Slavonian Substage, but the oldest Pannonian ostracod biozone, the *Hemicytheria loerentheyi* Zone was not identified in any of the samples. This interpretation is roughly supported by the mollusc fauna. The age of the outcrops is thus approximately between 10 and 11.3 Ma (cf. VASILIEV et al., 2010).

ACKNOWLEDGMENT

The authors thank Ferenc WANEK for discussions on the Transylvanian ostracods, and Dániel BOTKA and Csaba KRÉZSEK for the photo courtesy of the Mihălț and Tau outcrops, respectively. Ilaria MAZZINI (Roma Tre) kindly reviewed an earlier version of the manuscript. The reviewers of the present paper were Valentina HAJEK-TADESSE (Croatian Geological Survey, Zagreb) and Ljupko RUNDIĆ (University of Belgrade); their careful reviews are acknowledged herein. This is MTA-MTM-ELTE Paleo contribution No. 220, partly funded by the Hungarian National Research, Development and Innovation Office – NKFI 116618.

REFERENCES

- BOOMER, I. & EISENHAEUER, G. (2002): Ostracod faunas as palaeoenvironmental indicators in marginal marine environments. – In: HOLMES, J.A. & CHIVAS, A.R. (eds.): *The Ostracoda: Applications in Quaternary research*. Geophysical Monograph Series, 131, American Geophysical Union, Washington DC, 135–149. doi: 10.1029/131GM07.
- BOOMER, I., GRAFENSTEIN VON, U., GUICHARD, F. & BIEDA, S. (2005): Modern and Holocene sublittoral ostracod assemblages (Crustacea) from the Caspian Sea: A unique brackish, deep-water environment. – *Palaeogeography, Palaeoclimatology, Palaeoecology*, 225, 173–186. doi:10.1016/j.palaeo.2004.10.023.
- BRONSTEIN, Z.S. (1947): Fauna SSSR. Rakoobraznye, Tom II, Vyp. 1. Ostracoda Presnykh Vod. Academy of Sciences of the USSR Publishers, Moscow, 5–339. (English translation (1988): *Freshwater Ostracoda—Fauna of the USSR: Crustaceans, vol. II, No.1*. AA Balkema, Rotterdam, 1–455).
- CHINTĂUAN, I. (1971): Study of the ostracods from the Neogene sediments from western part of Munții Bărgăului. – *Studia Universitatis Babeș-Bolyai, Series Geologia-Mineralogia*, 16, 71 p.
- CIULAVU, D. (1999): Tertiary tectonics of the Transylvanian Basin. – PhD Thesis, Vrije Universiteit, Amsterdam, 152 p.
- CLICHICI, O. (1980): Cercetări geologice în perimetrul Măgina-Aiud-Ciumburd (Jud. Alba) [*Geological research in area Măgina-Aiud-Ciumburd (Alba County)* – in Romanian]. – *Studia Universitatis Babeș-Bolyai, Series Geologia-Geographia*, 25, 21–32.
- CZICZER, I., MAGYAR, I., PIPÍK, R., BÖHME, M., ČORIĆ, S., BAKRAČ, K., SÜTŐSZENTAI, M., LANTOS, M., BABINSZKI, E. & MÜLLER, P. (2008): Life in the sublittoral zone of long-lived Lake Pannon: paleontological analysis of the Upper Miocene Szák Formation, Hungary. – *International Journal of Earth Sciences*, 98, 1741–1766. doi: 10.1007/s00531-008-0322-3.
- DANIELOPOL, D. (1980): On the carapace shape of some European freshwater interstitial Candoninae (Ostracoda). – *Proceedings of the Biological Society of Washington*, 93, 743–756.
- DANIELOPOL, D.L., GROSS, M., PILLER, W.E. & BALTANAS, A. (2008a): Ostracods of the Paratethys Sea and Lake Pannon – Perspectives for renewal of cooperative projects. – *Senckenbergiana lethaea*, 88, 141–145. doi: 10.1007/BF03043985.
- DANIELOPOL, D.L., BUTTINGER, R., PIPÍK, R., GROSS, M., OLTEANU, R. & KNOBLECHNER, J. (2008b): Miocene “*Hungarocypris*” species of Lake Pannon (Central and South-Eastern Europe) transferred to *Herpetocyrella* Daday, 1909 (Ostracoda, Cyprididae). – *Senckenbergiana lethaea*, 88, 147–160. doi: 10.1007/BF03043986.
- DE BROUCKER, G., MELLIN, A. & DUINDAM, P. (1998): Tectonostratigraphic Evolution of the Transylvanian Basin, Pre-Salt Sequence, Romania. – In: DINU, C. & MOCANU, V. (eds.): *Geological and Hydrocarbon Potential of the Romanian areas*, Bucharest Geosciences Forum, 1, 36–70.
- DE DECKKER, P. (2002): Ostracod Palaeoecology. – In: HOLMES, J.A. & CHIVAS, A.R. (eds.): *The Ostracoda: Applications in Quaternary research*. Geophysical Monograph Series, 131, American Geophysical Union, Washington DC, 121–134. doi: 10.1029/131GM06.
- DE LEEUW, A., FILIPESCU, S., MATENCO, L., KRIJGSMAN, W., KUIPER, K. & STOICA, M. (2013): Paleomagnetic and chronostratigraphic constraints on the middle to late Miocene evolution of the Transylvanian basin (Romania): Implications for Central Paratethys stratigraphy and emplacement of the Tisza-Dacia plate. – *Global and Planetary Change*, 103, 82–98. doi:10.1016/j.gloplacha.2012.04.008.
- FILIPESCU, S. (1996): Stratigraphy of the Neogene from the Western border of the Transylvanian Basin. – *Studia Universitatis Babeș-Bolyai, Series Geologia*, 41, 3–69.
- FILIPESCU, S. (1999): The significance of foraminifera fauna from the Feleac Formation (Transylvanian Basin, Romania). – *Studia Universitatis Babeș-Bolyai, Series Geologia*, 44, 125–131.
- FILIPESCU, S., WANEK, F., MICLEA, A., DE LEEUW, A. & VASILIEV, I. (2011): Micropaleontological response to the changing paleoenvironment across the Sarmatian-Pannonian boundary in the Transylvanian Basin (Miocene, Oarba de Mureș section, Romania). – *Geologica Carpathica*, 62, 91–102. doi: 10.2478/v10096-011-0008-9.
- GANIĆ, M., RUNDIĆ, L., KNEŽEVIĆ, S., CVETKOV, V. (2010): The Upper Miocene Lake Pannon marl from the Filijala Open Pit (Beocin, northern Serbia): new geological and paleomagnetic data. – *Geološki Anali Balkanskoga Poluostrva*, 71, 95–108. doi: 10.2298/GABP1071095G.
- GARCIA-CASTELLANOS, D. (2006): Long-term evolution of tectonic lakes: Climatic controls on the development of internally drained basins. – In: WILLETT, S.D., HOVIUS, N., BRANDON, M.T. & FISHER, D.M. (eds.): *Tectonics, Climate, and Landscape Evolution*. Geological Society of America Special Paper, 398, 283–294. doi: 10.1130/2006.2398(17).
- GEARY, D.H. (1990): Patterns of evolutionary tempo and mode in the radiation of *Melanopsis* (Gastropoda: Melanopsidae). *Paleobiology*, 16, 492–511. doi: 10.1017/S0094837300010216.
- GHERGARI, L., MÉSZÁROS, N., HOSU, A., FILIPESCU, S. & CHIRA, C. (1991): The gypsiferous formation at Cheia (Cluj County). – *Studia Universitatis Babeș-Bolyai, Series Geologia*, 36, 13–28.
- GILLET, S. (1943): Les limnocardiides des couches à congéries de Roumanie. – *Memoire Institutului Geologic al României*, 4, 1–114.
- HARZHAUSER, M. & MANDIĆ, O. (2008): Neogene lake systems of Central and South-East Europe: Faunal diversity, gradients and interrelations. – *Palaeogeography, Palaeoclimatology, Palaeoecology*, 260, 417–434. doi:10.1016/j.palaeo.2007.12.013.
- HARZHAUSER, M., KERN, A., SOLIMAN, A., MINATI, K., PILLER, W.E., DANIELOPOL, D.L. & ZUSCHIN, M. (2008): Centennial- to decadal-scale environmental shifts in and around Lake Pannon related to a major Late Miocene lake-level rise. – *Palaeogeography, Palaeoclimatology, Palaeoecology*, 270, 102–115. doi:10.1016/j.palaeo.2008.09.003.
- HÉJJAS, I. (1894): Új adatok Erdélyi fossil ostracoda-faunájához [*New data on the fossil ostracod fauna of Transylvania* – in Hungarian]. – *Orvos- Természettudományi Értesítő, II. Természettudományi Szak, Kolozsvár*, 16, 35–68.
- HOLMES, J.A. (2001): Ostracoda. – In: SMOL, J.P., BIRKS, H.J. & LAST, H.J.B. (eds.): *Tracking environmental change Using Lake Sediments Zoological Indicators*, 4, Kluwer Academic Publisher, Dordrecht, 125–151. doi: 10.1007/0-306-47671-1.
- ILIE, M. (1952): Cercetări geologice în regiunea Cluj-Cojocna-Turda-Ocna Mureșului-Aiud [*Geological research in Cluj-Cojocna-Turda-Ocna Mureșului-Aiud region* – in Romanian]. – *Anuarul Comitetului Geologic*, 24, 303–369.
- JIRIČEK, R. (1985): Die ostracoden des Pannonien. – In: PAPP, A., JÁMBOR, Á. & STEININGER, F.F. (eds.): *Chronostratigraphie und Neostatotypen, Miozän der Zentralen Paratethys 7, Pannonien*. Akadémiai Kiadó, Budapest, 378–425.

- KEEN, M.C. (1982): Intraspecific variation in Tertiary ostracods. – In: BATE, R.H., ROBINSON, E. & SHEPPARD, L.M. (eds.): Fossil and recent ostracods. Ellis Horwood Limited, Chichester, 381–405.
- KOCH, A. (1900): Die Tertiärbildungen des Beckens des Siebenbürgischen Landestheile, II. Neogene Abtheilung. – Budapest, 369 p.
- KOLLMANN, K. (1960): Cytherideinae und Schulerideinae n. subfam. (Ostracoda) aus dem Neogen des östl. Oesterreich. – Mitteilungen der Österreichischen Geologischen Gesellschaft in Wien, 51, 89–195.
- KORECZ, A. (1991): A DK-dunántúli pannóniai s.l. ostracoda fauna vizsgálatának eredményei [The Pannonian s.l. ostracod fauna from SE Transdanubia – in Hungarian]. – PhD thesis, Eötvös Loránd University, Department of Paleontology, Budapest, 231 p.
- KOVÁČ, M., ANDREYEVA-GRIGOROVICH, A., BAJRAKTAREVIĆ, Z., BRZOBOHATÝ, R., FILIPESCU, S., FODOR, L., HARZHAUSER, M., NAGYMAROSY, A., OSZCZYPKO, N., PAVELIĆ, D., RÖGL, F., SAFTIĆ, B., SLIVA, I. & STUDENCKA, B. (2007): Badenian evolution of the Central Paratethys Sea: paleogeography, climate and eustatic sea-level changes. – Geologica Carpathica, 58, 579–606.
- KRÉZSEK, C. (2005): Sedimentologia și arhitectura depozitelor panoniene din estul Bazinului Transilvaniei [Sedimentology and architecture of the Pannonian sediments in the Eastern Transylvanian Basin – in Romanian]. – PhD thesis, Babeș-Bolyai University, Faculty of Biology and Geology, Cluj-Napoca, 165 p.
- KRÉZSEK, C. & BALLY, A.W. (2006): The Transylvanian Basin (Romania) and its relation to the Carpathian fold and thrust belt: insights in gravitational salt tectonics. – Marine and Petroleum Geology, 23, 405–442. doi:10.1016/j.marpetgeo.2006.03.003
- KRÉZSEK, C. & FILIPESCU, S. (2005): Middle to late Miocene sequence stratigraphy of the Transylvanian Basin (Romania). – Tectonophysics, 410, 437–463. doi:10.1016/j.tecto.2005.02.018.
- KRÉZSEK, C., FILIPESCU, S., SILYE, L., MAȚENCO, I. & DOUST, H. (2010): Miocene facies associations and sedimentary evolution of the Southern Transylvanian Basin (Romania): Implications for hydrocarbon exploration. – Marine and Petroleum Geology, 27, 191–214. doi:10.1016/j.marpetgeo.2009.07.009.
- KRSTIĆ, N. (1971): Ostrakodi kongerjskih slojeva: 9. *Cytherura*, *Cytherissa*, *Limnocythere* i *Loxocauda* [Ostracodes des couches Congériennes: 9. *Cytherura*, *Cytherissa*, *Limnocythere* et *Loxocauda* – in Serbian with French summary]. – Glasnik Prirodnjačkog Muzeja, Serija A, Beograd, 26, 143–185.
- KRSTIĆ, N. (1972a): Rod *Candona* (Ostracoda) iz kongerjskih slojeva južnog dela Panskog Basena [Genus *Candona* (Ostracoda) from Congeria beds of southern Pannonian Basin – in Serbian with English summary]. – The Serbian Academy of Sciences and Arts, Monographs, 450, The Section of Natural and Mathematical Sciences, Beograd, 39, 145 p.
- KRSTIĆ, N. (1972b): Ostrakodi kongerjskih slojeva: 10. *Loxoconcha* [Ostracodes from the Congerian Beds: 10. *Loxoconcha* – in Serbian with English summary]. – Glasnik Prirodnjačkog Muzeja, Serija A, Beograd, 27, 243–275.
- KRSTIĆ, N. (1973a): Biostratigrafija kongerjskih slojeva okoline Beograda na osnovu ostrakoda sa opisom vrsta roda *Amplocypris* [Biostratigraphy of the Congerian beds in the Belgrade region on the basis of ostracoda – in Serbian with English summary]. – Institute for Geological and Mining Exploration and Investigation of Nuclear and Other Mineral Raw Materials, Monographs, 4, 158 p.
- KRSTIĆ, N. (1973b): Ostrakodi kongerjskih slojeva 11: *Amnocythere* [Ostracoda from the Congerian beds: 11. *Amnocythere* – in Serbian with English summary]. – Radovi IGRI, 8, Beograd, 53–99.
- KRSTIĆ, N. (1985): Ostracoden im Pannonien der Umgebung von Belgrad. – In: PAPP, A., JÁMBOR, Á. & STEININGER, F.F. (eds.): Chronostratigraphie und Neostatotypen, Miozän der Zentralen Paratethys 7, Pannonien. Akadémiai Kiadó, Budapest, 103–144.
- KRSTIĆ, N. & STANCHEVA, M. (1990): Ostracod of Eastern Serbia and Northern Bulgaria with notices on a Northern Turkey assemblage and some Mediterranean assemblages. – In: STEVANOVIĆ, P., NEVESSKAJA, L.A., MARINESCU, FL., SOKAČ, A. & JÁMBOR, Á. (eds.): Chronostratigraphie und Neostatotypen: Neogen der Westlichen ("Zentrale") Paratethys 8. Pontien (1989). JAZU and SANU, Zagreb-Beograd, 753–819.
- LÖRENTHEY, I. (1893): Adatok Szilágy megye és az erdélyi részek alsó pontusi lérakódásainak ismeretéhez. II. Az erdélyi-medence alsó pontusi képződményeinek faunája [Data for the lower Pontian sediments in Salaj County and Partium. II. Fauna of the Lower Pontian sediments from Transylvania – in Hungarian]. – Orvos- Természettudományi Értesítő, II. Természettudományi Szak, Kolozsvár, 15, 213–230.
- LUBENESCU, V. & LUBENESCU, D. (1976): Observații biostratigrafice asupra Pannonianului de la Lopadea Veche (Depresiunea Transilvaniei) [Biostratigraphical observations on the Pannonian from Lopadea Veche (Transylvanian Depression) – in Romanian]. – Dări de seamă ale ședințelor, 63, 57–64.
- MAGYAR, I., GEARY, H.D. & MÜLLER, P. (1999): Paleogeographic evolution of the Late Miocene Lake Pannon in Central Europe. – Palaeogeography, Palaeoclimatology, Palaeoecology, 147, 151–167. doi:10.1016/S0031-0182(98)00155-2.
- MAȚENCO, L., KRÉZSEK, CS., MERTEN, S., SCHMID, S.M., CLOETINGH, S. & ADRIESSEN, P. (2010): Characteristics of collisional orogens with low topographic build-up: an example from the Carpathians. – Terra Nova, 22, 155–165. doi: 10.1111/j.1365-3121.2010.00931.x.
- MEISCH, C. (2000): Freshwater Ostracoda of Western and Central Europe. – Spektrum Akademischer Verlag, Heidelberg–Berlin, 522 p.
- MORKHOVEN, VAN F.P.C.M. (1963): Post-Paleozoic Ostracoda, their morphology, taxonomy and economic use, Part 2. – Elsevier, New York, 478 p. doi: 10.2108/zsj.26.846.
- MÜLLER, P., GEARY, D.H. & MAGYAR, I. (1999): The endemic molluscs of the Late Miocene Lake Pannon: their origin, evolution, and family-level taxonomy. – Lethaia, 32, 47–60. doi: 10.1111/j.1502-3931.1999.tb00580.x.
- NAMIOTKO, T., DANIELOPOL, D.L., BELMECHERI, S., GROSS, M. & VON GRAFENSTEIN, U. (2012): On the Leptocytheridae ostracods of the long-lived Lake Ohrid: a reappraisal of their taxonomic assignment and biogeographic origin. – International Review of Hydrobiology, 97, 356–374. doi: 10.1002/iroh.201211496.
- OLTEANU, R. (2011): Atlas of the Pannonian and Pontian ostracods from the eastern area of the Pannonian basin. Geo-Eco-Marina 17, 135–177.
- PAPP, A. (1951): Das Pannon des Wiener Beckens. – Mitteilungen der geologischen Gesellschaft in Wien, 39–41, 99–193.
- PAPP, A., JÁMBOR, Á. & STEININGER, F. (eds.) (1985): Chronostratigraphie und Neostatotypen, Miozän der Zentralen Paratethys 7, Pannonien. Akadémiai Kiadó, Budapest, 636 p.
- PAULISSEN, W., LUTHI, S.M., GRUNERT, P., ČORIĆ, S. & HARZHAUSER, M. (2011): Integrated high-resolution stratigraphy of a Middle to Late Miocene sedimentary sequence in the central part of the Vienna Basin. – Geologica Carpathica, 62, 155–169. doi: 10.2478/v10096-011-0013-z.
- PÁVAY-VAJNA, F. (1910): Die geologischen Verhältnisse der Umgebung von Olah-Lapad. – Földtani Közlöny, 40, 420–434.
- PIPIĆ, R. (1998): Salinity changes recorded by ostracoda assemblages found in Pannonian sediments in the western margin of the Danube Basin. – Bulletin des centres de recherches exploration-production Elf-Aquitaine, Pau, Mém. 20, 167–177.
- PIPIĆ, R. (2007): Phylogeny, palaeoecology, and invasion of non-marine waters by the late Miocene hemicytherid ostracod *Tyrrhenocythere* from Lake Pannon. – Acta Palaeontologica Polonica, 52, 351–368.
- PIPIĆ, R. & BODERGAT, A.M. (2004): Euxinocythere (Ostracoda, Cytheridae, Leptocytherinae) du Miocène supérieur du Basin de Turiec (Slovaquie): taxonomie et paléoécologie. – Revue de Micropaléontologie, 47, 36–52. doi:10.1016/j.revmic.2004.01.001.
- POKORNÝ, V. (1944): La microstratigraphie du Pannonien entre Hodonin et Mikulčice (Moravie meridionale, Tchécoslovaquie). – Bull. Inter. Acad. tcheque. Sci., 23, 1–25.
- POPESCU, B.M. (1995): Romania's petroleum systems and their remaining potential. – Petroleum Geosciences, 1, 337–350. doi: 10.1144/petgeo.1.4.337.
- REUSS, A.E. (1850): Die fossile Entomostraceen des österreichischen Tertiärbeckens. – Naturwissenschaftliche Abhandlungen, Wien, 2/1, 41–92.
- ROYDEN, L.H. (1988): The Pannonian Basin: a Study in Basin Evolution. – American Association of Petroleum Geologists, Memoir, 45, 17–25.
- RICKETTS, R.D., JOHNSON, TH.C., BROWN, E.T., RASMUSSEN, K.A. & ROMANOVSKY, V.V. (2001): The Holocene paleolimnology of Lake Issyk-Kul, Kyrgyzstan: trace element and stable isotope composition of ostracodes. – Palaeogeography, Palaeoclimatology, Palaeoecology, 176, 207–227. doi: 10.1016/S0031-0182(01)00339-X.
- RUNDIĆ, LJ. (1998): The carapace of fossil ostracodes: paleoenvironmental indicators. – Geološki Anali Balkanskoga Poluostrva, 64, 137–151.
- RUNDIĆ, LJ. (2002): Some species of the genus *Hemicytheria*, Pokorny (Ostracoda, Crustacea) from the Upper Miocene of Serbia. – Annales Géologiques de la Péninsule Balkanique, 64, 137–151. doi:10.2298/GABP0264137R.
- RUNDIĆ, LJ. (2006): Late Miocene ostracodes of Serbia: morphologic and palaeoenvironmental considerations. – Geološki Anali Balkanskoga Poluostrva, 67, 89–100. doi: 0350-0608/2006/0350-06080667089R.
- RUNDIĆ, LJ., GANIĆ, M., KNEŽEVIĆ, SL. & SOLIMAN, A. (2011): Upper Miocene Pannonian sediments from Belgrade (Serbia): new evidence and paleoenvironmental considerations. – Geologica Carpathica, 62, 267–278. doi: 10.2478/v10096-011-0021-z.
- SANDERS, C.A.E., ANDRIESSEN, P. & CLOETINGH, S. (1999): Life cycle of the East Carpathian orogen: erosion history of a doubly vergent critical wedge assessed by fission track thermochronology. – Journal of Geophysical Research, 104 (B12), 29, 1–112. doi: 10.1029/1998JB900046.
- SCHMID, S.M., BERNOULLI, D., FÜGENSCHUH, B., MAȚENCO, L., SCHEFER, S., SCHUSTER, R., TISCHLER, M. & USTASZEWSKI, K. (2008): The Alpine-Carpathian-Dinaric orogenic system: correlation and evolution of tectonic units. – Swiss Journal of Geosciences, 101, 139–183. doi: 10.1007/s00015-008-1247-3.

- SCHORNIKOV, E.I. (1964): An experiment on the distinction of the Caspian elements of the ostracod fauna in the Azov-Black Sea Basin.– *Zoologicheskii Zhurnal*, 43, 1276–1293.
- SOKAČ, A. (1972): Pannonian and Pontian Ostracode Fauna of the Mt. Medvednica.– *Paleontologia Jugoslavica*, 11, 1–140.
- SOKAČ, A. (1990): Pontian ostracod fauna in the Pannonian Basin – In: STEVANOVIĆ, P.M., NEVESSKAYA, L.A., MARINESCU, F., SOKAČ, A. & JÁMBOR, Á. (eds.): *Chronostratigraphie und Neostatotypen, Neogen der Westlichen* (“Zentrale”) Paratethys 8, Pontien. JAZU and SANU, Zagreb–Beograd, 672–721.
- STEVANOVIĆ, P.M., NEVESSKAYA, L.A., MARINESCU, F., SOKAČ, A. & JÁMBOR, Á. (eds.) (1990): *Chronostratigraphie und Neostatotypen, Neogen der Westlichen* (“Zentrale”) Paratethys 8, Pontien. JAZU and SANU, Zagreb–Beograd, 952 p.
- SÜTŐ-SZENTAI, M. & SZEGŐ, É. (2008): Organic-walled microplankton studies from beds of the Sarmatian-Pannonian stratotype section at Marosorbó (Oarba de Mureș), Transylvanian Basin.– *Földtani Közlöny*, 138, 279–296.
- SZTANÓ, O., KRÉZSEK, CS., MAGYAR, I., WANEK, F. & JUHÁSZ, GY. (2005): Sedimentary cycles and rhythms in a Sarmatian to Pannonian (Late Miocene) transitional section at Oarba de Mureș / Marosorbó, Transylvanian Basin.– *Acta Geologica Hungarica*, 48, 235–257. doi: 10.1556/AGeol.48.2005.3.1
- TER BORGH, M., VASILIEV, I., STOICA, M., KNEŽEVIĆ, S., MAŢENCO, L., KRIGSMAN, W., RUNDIĆ, L. & CLOETINGH, S. (2013): The isolation of the Pannonian basin (Central Paratethys): new constraints from magnetostratigraphy and biostratigraphy.– *Global and Planetary Change*, 103, 99–118. doi:10.1016/j.gloplacha.2012.10.001.
- TÖKÉS, L.B. (2013): Pannon-tavi turbiditek dél-erdélyi feltárásokban: szedimentológiai és gamma szelvényezés eredményei [*The results of sedimentological and gamma-ray logging of Upper Miocene lacustrine turbidites of the Transylvanian Basin – in Hungarian*].– MSc Thesis, Eötvös Loránd University, Department of Geology, Budapest, 90 p.
- VASILIEV, I., DE LEEUW, A., FILIPESCU, S., KRIGSMAN, W., KUIPER, K., STOICA, M. & BRICEAG, A. (2010): The age of the Sarmatian–Pannonian transition in the Transylvanian Basin (Central Paratethys).– *Palaeogeography, Palaeoclimatology, Palaeoecology*, 297, 54–69. doi:10.1016/j.palaeo.2010.07.015.
- WAGNER, C.W. (1964): Ostracods as environmental indicators to Recent and Subrecent estuarine deposits of the Netherlands.– In: PURI, S. (ed.): *Ostracoda as ecological and paleoecological indicators*. Pubblicazioni della Stazione Zoologica di Napoli, 33, 480–495.
- WANEK, F. (1992a): *Kagylósrák-faunák az Erdélyi-medence felső neogén üledékeiben* [*Ostracod faunas in the Upper Neogene sediments of the Transylvanian basin – in Hungarian*].– Erdélyi Múzeum-Egyesület Természettudományi és Matematikai Szakosztályának Közleményei, 1, 93–98.
- WANEK, F. (1992b): *Biostratigrafia ostracodelor Sarmatiene și Pannoniene din Bazinul Transilvaniei* [*Biostratigraphy of Sarmatian and Pannonian Ostracods in the Transylvanian basin – in Romanian*].– Raport BBU, Arhiva Romgaz, manuscript, 49 p.
- WANEK, F. & FILIPESCU, S. (1994): *O fauna cu ostracode din Pannonianul inferior de la Moldovenesti* [*A fauna with ostracods from the Lower Pannonian of Moldovenesti – in Romanian*].– Symposium “75 ani de la înființarea Universității Daciei Superioare, Cluj-Napoca” Programme & Abstracts.