

## IMPACT OF MONOCROTOPHOS AND NEEM OIL MIXTURE ON DEFOLIATOR MANGMENT IN GROUNDNUT

K. SAHAYARAJ\* & A. AMALRAJ

Plant protection Research Unit, Department of Botany, St. Joseph's College, Trichy- 620002, Tamil Nadu 620 002, India.  
\*Crop Protection Research Centre, Department of Zoology, St. Xavier's college, Palayamkottai 627 002, Tamil Nadu, India,  
Ph. No. 0461 2361957, e-mail ttn\_ksraj@sancharnet.in

Manuscript received: MJESec DAN, 2005; Reviewed: MJESec DAN, 2005; Accepted for publication: MJESec DAN, 2005

### ABSTRACT

The integrated effect of intercropping, a synthetic pesticide (monocrotopas) (M) and neem based biopesticide (neem oil - 2%) (NO) on three-groundnut defoliators damage and also the groundnut production was studied. The monocrotopas and neem oil combination was found to be very effective in reducing the defoliator infestation. Defoliator's incidence was significantly higher in untreated plots, resulting in significantly lower yield (1539.03 Kg h<sup>-1</sup>). The groundnut yield was increased (2011.18 Kg h<sup>-1</sup>) when monocrotophos and neem oil mixture was applied than monocrotophos (1877.77 Kg h<sup>-1</sup>) and control categories. The estimated avoidable groundnut and black gram yield loss were lower in monocrotopas.

**KEY WORDS:** monocrotophos, neem oil, intercropping, groundnut production

## INTRODUCTION

Groundnut (*Arachis hypogaea* Linn.) is the premier oilseed crop of India. It holds a 35 per cent share of the total oilseed area (24 million ha) and contributes nearly 40 per cent of the total oil seeds production (20 million tones). The low level of productivity in India is largely because of the crop is rainfed, exposed to various abiotic and biotic stresses. An estimated annual loss of Rs. 150 crores in groundnut due to pests has been reported [1,2]. Among the major pests reported in groundnut, *Aproaerma modicella* Dev., *Helicoverpa armigera* Hubner and *Spodoptera litura* Fabricius are the major defoliators of groundnut [3 – 7]. Unrestrained application of chemical pesticides for pest control has created pesticide resistance on both *H. armigera* [8] and *S.litura* [7, 9]. This grown consciousness toward the environment has made to imperative to the crop production specialists to search for viable and economical alternatives to the chemicals. Therefore, the strategy should aim at economically sound Integrated Pest Management (IPM) where plant product and intercrops place an important role. Growing mixed crops or intercropping is an important feature of Indian agriculture [7, 10]. Moreover, intercropping affects the microclimate in the agroecosystem and is highly relevant in the pest management strategy and also plays a very important role in population dynamics of insect pests. Further more, it was reported that integration of groundnut with grams increased the yield and gross economic return [11]. However, information on the incidence of defoliator in relation to monocrotophos and neem oil mixture application is lacking. The present study was conducted during in the farmers filed to evaluate the impact of monocrotophos and neem oil mixture on the groundnut defoliators infestation, groundnut and black gram yield, avoidable loss, cost of cultivation and net income.

## MATERIAL AND METHODS

Groundnut (PLO - Red) and Blackgram (T-9) in 6:1 ratio were raised under controlled irrigation conditions at Regunathapuram, Pudukottai district, Tamil Nadu, India. The experiments were laid in randomized block design in a plot size of 20 x 24 m<sup>2</sup> for each treatment. Three replications were maintained for each treatment. Three applications with monocrotophos alone (M) (2%) and monocrotophos with neem oil (M + NO) (2%) were made at 33, 48 and 63 days after seeding sowing (ASS). Observations on the defoliator infestation level were recorded after 4 days of spray from thirty randomly selected plants from each replication. Infestations were recorded in morning hours from top, middle and bottom leaflets in each selected plant. The mean values were taken

into account. Symptoms define by Wightman and Amin [3] has been followed to identify the infestation of each defoliator. The percentage of defoliation was calculated by using Kapadia et. al. [12] procedure. At harvest, the yield of main crop and intercrop in different treatments were recorded. The data were analysed statistically and expressed as Kg h<sup>-1</sup>. Percentage of available loss was also calculated with standard procedure [13]. The data on pest infestation level, mean yield, available loss, cost of cultivation and net income separately was pooled and subjected to statistically analysed with ANOVA and DMRT by using system statistics version 6.

## RESULT AND DISCUSSION

### Defoliators infestation level

The results revealed that all the three-defoliator population was effectively controlled by the monocrotophos and neem oil mixture followed by monocrotophos alone. Among the three defoliators studied, the intensity of *H. armigera* was highest in all the categories followed by *S. litura* and *A. modicella* (Table 1). Previously it has been reported that among the defoliating insects of groundnut, leaf miner has been causing serious damage to the foliage, resulting in 23 to 89 per cent loss in pod yield at national level [14]. Nandagopal et. al [15] reported that monocrotophos was effective against leaf miners and also too to other pests [16]. Neem leaf extract has more effect on *A. modicella* than other two plants extract test (*Pongamia glabra* and *Calotropis gigantea*) under laboratory condition [17]. The percent defoliation of *S. litura* was higher in control (1.23 ±0.16, 5.90±0.36 and 7.43 ± 0.42 for 37, 52 and 67 DASS, respectively) and it was reduced to 0.3,1.9 and 3.5 times in monocrotophos spray. Further reduction (0.35, 2.14 and 4 times, for 37, 52 and 67 DASS, respectively) was observed in the monocrotophos and neem oil mixture. This might be due to the action of neem oil. The neem based insecticide suppressed the *S. litura* damage as there was least foliage damage in groundnut and also gave higher pod yield says Anon [2]. Recently it was reported that the intercrop groundnut and black gram showed higher incidence and percent leaf damage by *S.litura* [7].

In monocrotophos alone-sprayed field, *H. armigera* infestation level was gradually decreased from 37 days after sowing (6.13±0.29 %) to 52 (7.06±0.27 %) and 67 (7.80±0.21 %) DASS. This clearly shows the pesticide resistant capacity of *H. armigera*. The resistant level was probably driven by high selection pressure created by increased usage of insecticide [18]. The total reduction of leaf infestation due to monocrotophos spray was significantly lower ( $p<0.05$ ) (10.63±0.36 %)

IMPACT OF MONOCROTOPHOS AND NEEM OIL MIXTURE ON DEFOLIATOR MANGMENT IN GROUNDNUT

Table 1. Percent of infestation caused by defoliator in untreated and two IBM components treated in groundnut field

Treatment	Pest Counting after sowing (days)	Percent of defoliation/Plant			Total defoliation (%)
		<i>A.modicella</i>	<i>H.armigera</i>	<i>S.litura</i>	
Control	37	0.50 ± 0.11 <sup>aA</sup>	6.70±0.33 <sup>aB</sup>	1.23±0.16 <sup>aAC</sup>	8.00±0.53 <sup>aD</sup>
	52	3.03 ± 0.29 <sup>bA</sup>	12.50±0.55 <sup>bB</sup>	5.90±0.36 <sup>bC</sup>	21.60±0.72 <sup>bD</sup>
	67	3.96 ± 0.33 <sup>bcCA</sup>	17.76±0.63 <sup>cbB</sup>	7.43±0.42 <sup>cC</sup>	29.13±0.83 <sup>cd</sup>
Monocrotophos 36EC	37	0.48 ± 0.12 <sup>aA</sup>	6.13±0.29 <sup>aB</sup>	1.16±0.14 <sup>aC</sup>	7.73±0.39 <sup>aBD</sup>
	52	1.86 ± 0.22 <sup>bA</sup>	7.06±0.27 <sup>aB</sup>	3.16±0.29 <sup>bC</sup>	11.90±0.43 <sup>bD</sup>
	67	1.10 ± 0.14 <sup>cA</sup>	7.80±0.21 <sup>aB</sup>	2.70±0.22 <sup>cC</sup>	10.63±0.36 <sup>bcBD</sup>
Monocrotophos 36EC + Neem Oil	37	0.46 ± 0.13 <sup>aA</sup>	6.63±0.37 <sup>aB</sup>	1.13±0.14 <sup>aC</sup>	7.36±0.48 <sup>aD</sup>
	52	1.13 ± 0.27 <sup>bA</sup>	6.63±0.33 <sup>aB</sup>	2.80±0.35 <sup>aAC</sup>	10.56±0.44 <sup>bBD</sup>
	67	1.03 ± 0.15 <sup>bcCA</sup>	5.76±0.25 <sup>aB</sup>	2.50±0.31 <sup>bcC</sup>	9.38±0.48 <sup>bcD</sup>

Values carrying same small alphabet in a column of each treatment separately and capital Alphabets in a row are not statistically significant at 5 % using DMRT.

Table 2. Yield, percent available loss Kg h<sup>-1</sup>, Cost of cultivation and new income (Rs./ha) in different IPM components

Treatment	Mean Yield		% Available loss		Cost of cultivation	Net income
	G	BG	G	BG		
Monocrotophos (M)	1877.77 <sup>a</sup>	29.015 <sup>a</sup>	0.18 <sup>a</sup>	0.23 <sup>a</sup>	9955 <sup>a</sup>	3549 <sup>a</sup>
Monocrotophos + Neem oil (M + NO)	2011.18 <sup>b</sup>	33.55 <sup>b</sup>	0.23 <sup>a</sup>	0.33 <sup>b</sup>	10134 <sup>b</sup>	4361 <sup>b</sup>
Control	1506.03 <sup>c</sup>	22.30 <sup>c</sup>	-	-	9044 <sup>c</sup>	2006 <sup>c</sup>

G - groundnut; BG - Black gram, - indicates available loss was not observed.

Values in the column with letters in common are not significantly different at p = 0.05 using the DMRT

when compared to control (29.13±0.83 %) at 67 DASS (Table 1). This results confirms the earlier findings of Das [19] and Nandagopal et. al. [15]. They reported that monocrotophos reduced the groundnut leaf miner damage. As observed in the *A. modicella* and *H. armigera*, the per cent infestation caused by *S. litura* was also gradually decreased from control to monocrotophos and to M+NO mixture. Efficacy of neem products in insect control was reviewed by many scientists [20 - 21]. According to them not much work has been done on the use of neem products in the control of groundnut insect pests. The role of neem products either alone or blend with a synthetic insecticide, on the biopesticidal and biological activities of leaf miner was studied [22]. The results indicated that crude neem oil suppressed

oviposition, whereas larval and pupal mortality of leaf miner were not affected in desired extent. The effect of partially purified neem seed extract did not cause any juvenometric activity in the treated larvae [21].

**Cost effectiveness of neem based IPM**

Economic analysis of different treatments brought out the need for IPM and non-IPM methods in groundnut. The pod yield was significantly high (2011.18 Kg h<sup>-1</sup>) in the M+NO plot followed with monocrotophos (1877.77 Kg h<sup>-1</sup>) and theses two were superior to untreated control (1539.03 Kg h<sup>-1</sup>) (Table 2). It is in confirmation with the observation of Nandagopal et. al. [15] and Reddy [23]. As observed in the groundnut, the blackgram production was also high in the M+NO treatment (33.55 Kg h<sup>-1</sup>)

followed by monocrotophos plot (29.19 Kg h<sup>-1</sup>) and least in control field (22.30 Kg h<sup>-1</sup>). Senthilvel [24] reported similar result in groundnut (TMV 12) and blackgram (TMV5) intercropping system. He pointed out that raising groundnut with a blackgram in 6:1 ratio was a more remunerative and economic system for intercropping system. The net return per ha. was highest in the M + NO mixture plots followed by monocrotophos (Table 2). This is in accordance with the suggestion of Mullen [25]. They reported that IPM components increased the annual production. Table 2 clearly indicates that the percent available loss was higher in blackgram of experimental plots. Thus, on the basis of this study it could be concluded that the effectiveness of M + NO mixture was found to be optimum in controlling groundnut defoliators when applied three times at 15 days interval starting from 37 days after seedling sowing.

#### ACKNOWLEDGMENT

The senior author thanks Department of Science and Technology, Govt. of India for the financial assistance. He also grateful to Rev. Fr. A. Antonyswamy, S.J., and Prof. M. Thomas Punithan for encouragements and support.

#### REFERENCES

- [1]. SINGH, V. 1980. All India co-ordinated research project on oilseeds with special reference to groundnut. In: Proc. International Workshop on groundnut held at the ICRISAT center. Patancheru, A.P. during 13 – 17 October, 1980.
- [2]. AMIN, P.W., 1983. Major field insect pests of groundnut in India and associated crop losses. Proceedings of the All India Workshop on Crop losses due to insects. ICRISAT, Patancheru. India. 52p.
- [3]. WIGHTMAN, J.A. and AMIN, P.W., 1988. Groundnut pests and their control in the semi-arid tropics. Tropical pest Management, 34, 218-266.
- [4]. REDDY, P.S., and GHTWANDE, M.P., 1986. Major insect pests of groundnut and their management. Pesticides. 20(5): 52-56.
- [5]. MUTHIAH, C and KAREEM, A.A., 2000. survey of groundnut leaf miner and its natural enemies in Tamil Nadu. International Arachis Newsletter. 20: 62 – 66.
- [6]. DHARNE, P.K AND PATEL, S.K., 2000. Screening of promising groundnut genotypes for their reaction to Spodoptera litura. International Arachis Newsletter. 20: 67 – 69.
- [7]. Singh, A. K., Nath, P. 2003. Effect of intercropping

on the population and groundnut leaf damage by Spodoptera litura (Fab.). Shashpa., 10 (1): 33 – 38.

[8]. VENUGOPAL RAO, N., RAJA SEKHAR, P., VENKATAIAH, M., and RAJASRI, M., 1994. Estimation of insecticide resistance in Helicoverpa armigera in Andhra Pradesh. Indian Journal of plant protection. 22: 33-37.

[9]. MEHROTR., K.N., 1989. Pesticide resistance in insect pests. Indian Scenario pesticide Research Journal. 1:93-103.

[10]. AIYER, A.K.Y.N., 1949. Mixed cropping in India. Indian Journal of Agricultural Science, 19,: 439-543.

[11]. SINGH, T.V.K., SINGH, K.M. and SINGH, R.N., 1991. Impact of intercropping: IV Yields of groundnut. Indian Journal of Entomology. 53(3): 369 – 372.

[12]. KAPADIA, M.N., BHARODIA, R.D. and VERA, V.J., 1982. Biology and estimation of incidence of groundnut leafminer S. subscivella. Gujarat University Journal. 8. 37-39.

[13]. GOVINDU, H.C., VEERESH, G.K., WALKER, P.T. and JENKYN, J.F., 1980. Assessment of crop losses due to pests and diseases. UAS. Technical series no.32. University of Agricultural Sciences, Bangalore, India. pp.1-300.

[14]. MOHAMMED, A., 1981. The groundnut leafminer S. subscivella (Lepidoptera: Gelachidae). A review of world literature. Occasional paper 3. Groundnut Improvement program. International Crops Research Institute for Semi-arid Tropics (ICRISAT), Patancheru. A.P., India. 21p.

[15]. NANDAGOPAL, V., SONI, V.C., and HALL, D.R. and GEDIA, M.V., 1995. Effect of some components of IPM on the insect pest incidence and yields in groundnut. In National Seminar on IPM in Agriculture, Agriculture College Nagpur. 20p.

[16]. SHERASIYA, R.A. and BUTANI, P.G., 1998. Assessment of avoidable yield loss due to insect pests in groundnut. Journal of oilseed Research. 15(2): 390 – 392.

[17]. SAHAYARAJ, K., and PAULRAJ, M.G., 1998. Relative toxicity of some plant extracts to groundnut leaf miner, Aproaerema modicella Dev. International Arachis Newsletter, 18, 27-29.

[18]. SEKAR, P.V., RAO, N.V., RAO B.R., and VENKATAIAH, M., 1995. Current status of synthetic pyrethroid resistance in Helicoverpa armigera population on groundnut. International Arachis Newsletters. 15, 62-63.

[19]. DAS, B.B., 1988. Efficacy of some synthetic

pyrethroids and conventional insecticides spray against groundnut pest complex. *Pesticides*, 22, 10-12.

[20]. SAXENA, R.C., 1989. Insecticides from neem. In *Insecticides from neem* (Eds. J.T. Arasan., J.R. Philogene and P. Morand) ACS Symposium series 387, Washington, DC, pp. 110-135.

[21]. SCHEMUTTERER, H., 1990. Properties and potential of natural pesticides from the neem tree *Azadirachta indica*. *Annual Review of Entomology*, 35, 271-197.

[22]. GHEWANDE, M.P., NANDAGOPAL, V. and DESAI, S., 1996. Management of major insect pests and diseases of groundnut (*Arachis hypogaea* L.) using neem and its products in India. In *Neem*

sustainable agriculture. Basic and applied aspect-II (Ed. S.S. Nanval). CCS Haryana Agricultural University Dept. of Agronomy, Hisar. (in press).

[23]. REDDY, Y.U.R., 1986. Economics of increasing cropping intensity in dry land agriculture in India. *Andhra Agricultural Journal*, 33(4), 303-13.

[24]. SENTHILVEL, T., 1989. An ideal intercropping system for rainfed groundnut (*Arachis Hypogaea*) in Tamil Nadu. *Indian Journal of Agricultural Science*, 59(7), 435-437.

[25]. MULLEN, J.R., NORTON, G.W. and REAVES, D.W. 1997. Economic analysis of environmental benefits of IPM. *Journal of Agriculture and Applied Economics*, 29(2): 243 – 253.

