

THE SPANS OF FIVE STAR-LIKE SIMPLE CLOSED CURVES

THELMA WEST

University of Louisiana at Lafayette, USA

ABSTRACT. Let X be a continuum, that is a compact, connected, nonempty metric space. The span of X is the least upper bound of the set of real numbers r which satisfy the following conditions: there exists a continuum, C , contained in $X \times X$ such that $d(x, y)$ is larger than or equal to r for all (x, y) in C and $p_1(C) = p_2(C)$, where p_1, p_2 are the usual projection maps. The following question has been asked. If X and Y are two simple closed curves in the plane and Y is contained in the bounded component of the plane minus X , then is the span of X larger than the span of Y ? We define a set of simple closed curves, which we refer to as being five star-like. We answer this question in the affirmative when X is one of these simple closed curves. We calculate the spans of the simple closed curves in this collection and consider the spans of various geometric objects related to these simple closed curves.

1. INTRODUCTION

The span of a metric continuum was defined in 1964 by A. Lelek (see [3, p.209]). Variations of the span have been defined since then (cf [4, 5, 2]). In general it is difficult to evaluate the spans of a geometric object. It is of interest how these various spans, for a particular object, are related to each other. Also of interest is how the spans of related objects compare to each other. The following question by H. Cook has been particularly interesting.

If X_1 and X_2 are two simple closed curves in the plane and X_2 is contained in the bounded component of X_1 , then is the span of X_1 larger than the span of X_2 ? ([1])

This question has not yet been answered in general. Given some specific conditions on either X_1 or X_2 , the answer has been shown to be yes. We

2000 *Mathematics Subject Classification.* 54F20, 54F15.

Key words and phrases. Span, simple closed curve.

give a short summary of the results for these special cases that have been determined.

In the following results we assume that:

1. X_2 is a simple closed curve,
2. X_3 is a continuum that is contained in the bounded component of $R^2 - X_2$,
3. X_1 is a plane separating continuum that contains X_2 in one of its bounded components, and
4. the span of X is denoted by $\sigma(X)$.

If X_2 is the boundary of a convex region in the plane and X_1 is a simple closed curve, then $\sigma(X_2) < \sigma(X_1)$ ([7], see also [6]).

If X_2 is the boundary of a convex region in the plane, then $\sigma(X_3) < \sigma(X_2) < \sigma(X_1)$ [11].

Also, $\sigma(X_3) < \sigma(X_2) < \sigma(X_1)$ when X_2 is either an “indented circle” ([8, 9, 13]), or a “concave upward symmetric” simple closed curve [10], or a simple closed curve as defined in [12]. In this paper we define a set of simple closed curves that we refer to as five star-like and show that if X_2 is a five star-like simple closed curve and X_1 and X_3 are as defined previously, then $\alpha(X_3) < \alpha(X_2) < \alpha(X_1)$, where α represents the span σ , semispan σ_0 , surjective span σ^* , surjective semispan σ_0^* , symmetric span s , and surjective symmetric span s^* .

2. PRELIMINARIES

Let X be a continuum, that is a compact, connected metric space. The span of X , $\sigma(X)$, is the least upper bound of the set of real numbers r which satisfy the following conditions: there exist a continuum, C , and continuous functions $f, g : C \rightarrow X$, such that

$$d\min(f, g) = \min\{d(f(c), g(c)) \mid c \in C\} \geq r$$

and

$$f(C) = g(C) \qquad \sigma$$

span

To obtain the various other spans, we replace the preceding equation with the following:

$$f(C) \subseteq g(C) \qquad \sigma_0$$

semispan

$$f(C) = g(C) = X \qquad \sigma^*$$

surjective span

$$f(C) \subseteq g(C) = X \qquad \sigma_0^*$$

surjective semispan

$$f(C) = g(C) \qquad s$$

symmetric span

and $\forall c \in C, \exists c'$ such that $f(c) = g(c')$ and $f(c') = g(c)$

$$f(C) = g(C) = X \quad s^*$$

surjective symmetric span

and $\forall c \in C, \exists c'$ such that $f(c) = g(c')$ and $f(c') = g(c)$.

The inequalities below follow immediately from the definitions,

$$\begin{aligned} 0 \leq \sigma^*(X) &\leq \sigma(X) \leq \sigma_0(X) \leq \text{diam } X, \\ 0 \leq \sigma^*(X) &\leq \sigma_0^*(X) \leq \sigma_0(X) \leq \text{diam } X, \\ 0 &\leq s(X) \leq \sigma(X), \\ 0 &\leq s^*(X) \leq \sigma^*(X). \end{aligned}$$

The following results are easy consequences of the various definitions.

1. If J is an arc, then $\alpha(J) = 0$, where $\alpha = \sigma, \sigma_0, \sigma^*, \sigma_0^*, s, s^*$.
2. If X is a simple closed curve, then $\sigma(X) = \sigma^*(X)$, $\sigma_0(X) = \sigma_0^*(X)$, and $s(X) = s^*(X)$.

We utilize the following theorem from [3] in the proof of Corollary 2.

Theorem L: If Y is a closed subset of the Hilbert cube I^ω and $\rho : Y \rightarrow S$ is an essential mapping of Y onto the circumference S , then

$$\inf_{s \in S} (\rho^{-1}(s), \rho^{-1}(-s)) \leq \sigma(Y).$$

3. MAIN RESULTS

- (a) Let Q be a five-sided, convex polygon with sequentially labeled vertices Q'_i for $i = 0, 1, 2, 3, 4$, with all interior angles $\angle Q'_i, i = 0, 1, 2, 3, 4$, larger than 90° . Extend each side $\overline{Q'_j Q'_{j+1}}$ of the polygon.
- (b) Let Q_{j+2} be the point of intersection of $\overleftrightarrow{Q'_j Q'_{j+1}}$ and $\overleftrightarrow{Q'_{j+2} Q'_{j+3}}$ for $j = 0, 1, 2, 3, 4$, where all indices are taken modulo 5.
- (c) Let X be the star shaped simple closed curve defined by $X = \cup_{j=0}^4 (\overline{Q'_j Q'_{j+1}} \cup \overline{Q'_{j+1} Q_{j+1}})$. We refer to X as a five star-like simple closed curve.
- (d) Let t_j be the point on $\overline{Q_{j-1} Q_{j-3}}$ that is the closest to Q_j . Note that $t_j \in (\overline{Q'_{j-2} Q_{j-1}} - \{Q'_{j-2}\})$, since $Q_j \in \overleftrightarrow{Q'_{j-2} Q'_{j-1}}$ and $\angle Q'_{j-2} > 90^\circ$.
- (e) Let r_j be the point on $\overline{Q_{j+1} Q_{j+3}}$ that is closest to Q_j . Note that $r_j \in (\overline{Q_{j+1} Q'_{j+1}} - \{Q'_{j+1}\})$, since $Q_j \in \overleftrightarrow{Q'_{j+1} Q'_j}$ and $\angle Q'_{j+1} > 90^\circ$.
- (f) Note that $t'_j = d(Q_j, t_j) < d(Q_j, Q'_{j+2}) = q'_j$ and $r'_j = d(Q_j, r_j) < d(Q_j, Q'_{j+2}) = q'_j$.

(g) Suppose that $q'_3 = \min\{q'_i\}_{i=0}^4$. We can make this assumption since we can relabel the vertices so that this is true. We refer to the number

$$\text{fss}(X) = \max\{q'_3, \min\{r'_0, r'_1, t'_1, t'_2\}, \min\{r'_1, t'_2, q'_4\}, \min\{r'_4, r'_0, t'_0, t'_1\}, \min\{r'_4, t'_0, q'_2\}\}$$

as the five star-like spread of X , where X is a five star-like simple closed curve.

THEOREM 3.1. *Let X be a five star-like simple closed curve. Then $\sigma(X) = \sigma_0(X) = \sigma^*(X) = \sigma_0^*(X) = s(X) = s^*(X) = \text{fss}(X)$.*

PROOF. Let $f, g : I \rightarrow X$ be continuous functions from I , the unit interval, onto X , such that their movements are always clockwise on X and one function is constantly Q_j for some j , while the other function moves from Q_k to Q_{k+1} , passing through the point Q'_k , where $k \neq j, j-1$ and $f[I] = g[I] = X$. Let \mathcal{P} be the set consisting of all pairs of functions (f, g) that satisfy these conditions. We claim that $\alpha(X) = \max\{\text{dmin}(f, g) \mid (f, g) \in \mathcal{P}\} = \text{fss}(X)$, where $\alpha = \sigma, \sigma_0, \sigma^*, \sigma_0^*, s, s^*$. We consider two cases.

CASE A: $\max\{\text{dmin}(f, g) \mid (f, g) \in \mathcal{P}\} = q'_3$

Consider the pair of functions given in Table 1.

TABLE 1. $f(t)$ and $g(t)$ versus t .

t	$f(t)$	$g(t)$
0	Q_0	Q_3
0.1	Q_1	Q_3
0.2	Q_1	Q_4
0.3	Q_2	Q_4
0.4	Q_2	Q_0
0.5	Q_3	Q_0
0.6	Q_3	Q_1
0.7	Q_4	Q_1
0.8	Q_4	Q_2
0.9	Q_0	Q_2
1.0	Q_0	Q_3

Note that

$$d(Q_j, \overline{Q_{j+2}Q'_{j+2}} \cup \overline{Q'_{j+2}Q_{j+3}}) = d(Q_j, Q'_{j+2}) = q'_j.$$

Given this observation and assumption (g) in the construction of X , we see that $\text{dmin}(f, g) = q'_3$.

We observe that $\alpha(X) \geq d(Q_3, Q'_0) = q'_3$ for $\alpha = \sigma, \sigma_0, \sigma^*, \sigma_0^*, s, s^*$. This is true since, $\sigma(X) = \sigma^*(X)$, $\sigma_0(X) = \sigma_0^*(X)$, $f(I) = g(I) = X$, and for all $t \in [0, 1]$ there is a $t' \in [0, 1]$ such that $f(t) = g(t')$ and $g(t) = f(t')$.

In particular, (f, g) , from Table 1, is a “better” pair than the pairs (f_1, g_1) and (f_2, g_2) given in Table 2 and Table 3.

TABLE 2. $f_1(t)$ and $g_1(t)$ versus t .

t	$f_1(t)$	$g_1(t)$
0	Q_0	Q_3
0.1	Q_0	Q_4
0.2	Q_1	Q_4
0.3	Q_2	Q_4
0.4	Q_2	Q_0
0.5	Q_3	Q_0
0.6	Q_4	Q_0
0.7	Q_4	Q_1
0.8	Q_4	Q_2
0.9	Q_0	Q_2
1.0	Q_0	Q_3

TABLE 3. $f_2(t)$ and $g_2(t)$ versus t .

t	$f_2(t)$	$g_2(t)$
0	Q_0	Q_2
0.1	Q_1	Q_2
0.2	Q_1	Q_3
0.3	Q_1	Q_4
0.4	Q_2	Q_4
0.5	Q_2	Q_0
0.6	Q_2	Q_1
0.7	Q_3	Q_1
0.8	Q_4	Q_1
0.9	Q_4	Q_2
1.0	Q_0	Q_2

Consequently, since

$$\text{dmin}(f, g) \geq \max\{\text{dmin}(f_1, g_1), \text{dmin}(f_2, g_2)\}$$

it must be the case that

$$\text{dmin}(f, g) = q'_3 \geq \min\{r'_4, t'_0\}$$

and

$$\text{dmin}(f, g) = q'_3 \geq \min\{t'_2, r'_1\}$$

(*)

This implies that $\text{fss}(X) = q'_3$.

We define a continuous function $p_j : Y \rightarrow \overrightarrow{Q'_j Q_j} \cup \overrightarrow{Q'_j Q_{j+1}}$ where $j = 0, 1, 2, 3, 4$, and Y is the closure of the bounded component of $R^2 - X$. First we define p_j on $Y \cap Y_l$ where Y_l is the closure of the bounded component of $R^2 - (\overrightarrow{Q'_j Q_{j+1}} \cup \overrightarrow{Q_{j+1} Q'_{j+1}} \cup \overrightarrow{Q'_{j+1} Q_{j+2}} \cup \overrightarrow{Q_{j+2} Q'_{j+2}} \cup \overrightarrow{Q'_{j+2} Q_{j+3}} \cup \overrightarrow{Q_{j+3} Q'_j})$. We consider two cases in defining p_j on Y_l for $j = 0, 1, 2, 3, 4$.

CASE $p_j, Y_l, 1: t'_{j+2} \leq r'_{j+1}$

Let $t \in \overrightarrow{Q_{j+2} Q_{j+1}}$ and let L_t be the line that is parallel to $\overrightarrow{Q_{j+2} t_{j+2}}$ and passes through the point t . For each $y \in Y \cap L_t$, let $p_j(y) = t$. It may be the case $Q_{j+1} = t_{j+2}$. If so, then $L_t \cap Y = \{Q_{j+1}, Q_{j+2}\}$ and $\text{diam}(p_j^{-1}\{t\}) = d(Q'_{j+2}, t_{j+2}) = t'_{j+2}$. Otherwise, $\overrightarrow{Q_{j+1} t_{j+2}} \cup \overrightarrow{t_{j+2} Q_{j+2}} \cup \overrightarrow{Q_{j+2} Q_{j+1}}$ forms a right triangle and for each t , $\text{diam}(p_j^{-1}\{t\}) \leq t'_{j+2}$.

Let $h_1 : [0, 1] \rightarrow \overrightarrow{Q'_j t_{j+2}}$ be the homeomorphism such that $h_1(t) = (1 - t)t_{j+2} + tQ'_j$.

Let $h_2 : [0, 1] \rightarrow \overrightarrow{Q_{j+2} Q_{j+3}}$ be the homeomorphism such that $h_2(t) = (1 - t)Q_{j+2} + tQ_{j+3}$.

Let L_t be the line connecting $h_1(t)$ and $h_2(t)$ for each $t \in [0, 1]$. For each $y \in Y \cap L_t$ let $p_j(y) = h_1(t)$. Note that $\text{diam}(p_j^{-1}(\{h_1(t)\})) \leq \max\{t'_{j+2}, q'_{j+3}\}$.

CASE $p_j, Y_l, 2: r'_{j+1} < t'_{j+2}$

Let $t \in \overrightarrow{r_{j+1} Q_{j+2}}$ and let L_t be the line that is parallel to $\overrightarrow{r_{j+1} Q_{j+1}}$ and passes through the point t . Let t' be the point such that $\{t'\} = L_t \cap ((\overrightarrow{Q'_j Q_{j+1}} - \overrightarrow{Q'_j Q_{j+1}}) \cup Q_{j+1})$. For each $y \in Y \cap L_t$ let $p_j(y) = t'$. Note that for each t' , $\text{diam}(p_j^{-1}\{t'\}) \leq r'_{j+1}$.

Let $h_1 : [0, 1] \rightarrow \overrightarrow{Q'_j Q_{j+1}}$ be the homeomorphism such that $h_1(t) = (1 - t)Q_{j+1} + tQ'_j$.

Let $h_2 : [0, 1] \rightarrow \overrightarrow{r_{j+1} Q_{j+3}}$ be the homeomorphism such that $h_2(t) = (1 - t)r_j + tQ_{j+3}$.

Let L_t be the line connecting $h_1(t)$ and $h_2(t)$. For each $y \in L_t \cap Y$ let $p_j(y) = h_1(t)$. Note that for each t , $\text{diam}(p_j^{-1}(\{h_1(t)\})) \leq \max\{r'_{j+1}, q'_{j+3}\}$.

The definition of p_j on $Y \cap Y_r$, where Y_r is the closure of the bounded component of $R^2 - (\overrightarrow{Q'_j Q_j} \cup \overrightarrow{Q_j Q'_{j+4}} \cup \overrightarrow{Q'_{j+4} Q_{j+4}} \cup \overrightarrow{Q_{j+4} Q'_{j+3}} \cup \overrightarrow{Q'_{j+3} Q_{j+3}} \cup \overrightarrow{Q_{j+3} Q'_j})$ is defined in a similar manner in the two corresponding cases.

CASE $p_j, Y_r, 1: r'_{j+4} \leq t'_j$.

In this case for $y \in Y \cap Y_r$, $\text{diam}(p_j^{-1}(\{p_j(y)\})) \leq \max\{r'_{j+4}, q'_{j+3}\}$

CASE $p_j, Y_r, 2: t'_j < r'_{j+4}$

In this case for $y \in Y \cap Y_r$, $\text{diam}(p_j^{-1}(\{p_j(y)\})) \leq \max\{t'_j, q'_{j+3}\}$.

We see that for $y \in Y_l$, $\text{diam}(p_j^{-1}(\{p_j(y)\})) \leq \max\{\min\{r'_{j+1}, t'_{j+2}\}, q'_{j+3}\}$. For $y \in Y_r$, $\text{diam}(p_j^{-1}(\{p_j(y)\})) \leq \max\{\min\{r'_{j+4}, t'_j\}, q'_{j+3}\}$. Consequently, for $y \in Y$,

$$\text{diam}(p_j^{-1}(\{p_j(y)\})) \leq \max\{\min\{r'_{j+1}, t'_{j+2}\}, \min\{r'_{j+4}, t'_j\}, q'_{j+3}\}.$$

Let $f^*, g^* : C \rightarrow Z$ be any two continuous functions from a continuum C into a continuum $Z \subseteq Y$, such that $f^*(C) \subseteq g^*(C) \subseteq Z$. Consider $p_j \circ f^*, p_j \circ g^* : C \rightarrow (Q'_j Q_j \cup Q'_j Q_{j+1})$. The image of $p_j \circ g^*(C)$ is an arc and $p_j \circ f^*(C) \subseteq p_j \circ g^*(C)$. Since all the spans of an arc are zero, there is a $c \in C$ such that $p_j \circ f^*(c) = p_j \circ g^*(c)$. Consequently, $d(f^*(c), g^*(c)) \leq \text{diam}(p_j^{-1}(\{p_j(g^*(c))\})) \leq \max\{\min\{r'_{j+1}, t'_{j+2}\}, \min\{r'_{j+4}, t'_j\}, q'_{j+3}\}$ and

$$\sigma_0(Z) \leq \max\{\min\{r'_{j+1}, t'_{j+2}\}, \min\{r'_{j+4}, t'_j\}, q'_{j+3}\}.$$

In this case (i.e. case A) when $\text{fss}(X) = \text{dmin}(f, g)$ where f and g are defined in Table 1, we conclude that $\sigma_0(X) \leq q'_{j+3}$ by taking $Z = X \subseteq Y$, $j = 0$, and using (*). Given the inequalities relating the various spans and the fact that for each $t \in [0, 1]$ there is a $t' \in [0, 1]$ such that $g(t) = f(t')$ and $f(t) = g(t')$, we conclude that $\sigma(X) = \sigma_0(X) = \sigma^*(X) = \sigma_0^*(X) = s(X) = s^*(X) = \text{fss}(X) = q'_3$. This completes case A.

CASE B: $\max\{\text{dmin}(f, g) \mid (f, g) \in \mathcal{P}\} > q'_3$

Let $(f, g) \in \mathcal{P}$ such that $\max\{\text{dmin}(f, g) \mid (f, g) \in \mathcal{P}\} = \text{dmin}(f, g) > q'_3$. Since $r'_3 < q'_3$ and $t'_3 < q'_3$, the pair (f, g) can not include the following (forbidden) steps as given in Table 4 and Table 5.

TABLE 4. Forbidden steps

t	$f(t)$	$g(t)$	t	$f(t)$	$g(t)$	t	$f(t)$	$g(t)$
t_1	Q_4	Q_3	t_1	Q_0	Q_3	t_1	Q_1	Q_3
t_2	Q_0	Q_3	t_2	Q_1	Q_3	t_2	Q_2	Q_3

TABLE 5. Forbidden steps

t	$f(t)$	$g(t)$	t	$f(t)$	$g(t)$	t	$f(t)$	$g(t)$
t_1	Q_3	Q_4	t_1	Q_3	Q_0	t_1	Q_3	Q_1
t_2	Q_3	Q_0	t_2	Q_3	Q_1	t_2	Q_3	Q_2

For any pair of functions, f and g , we consider all the possible starting values of f and g , that is $f(0)$ and $g(0)$, and determine what steps are possible for $t > 0$ given the restriction $\text{dmin}(f, g) > d(Q_3, Q'_1) = q'_3$. Because of the symmetry of this process, we consider either $f(0) = Q_j$ and $g(0) = Q_k$ where $k \neq j$, or $f(0) = Q_k$ and $g(0) = Q_j$ for $k \neq j$, but not both. All

these possibilities are summarized in the tables below. For $n \in N$, we let $0 = t_0 < t_1 < t_2 < \dots < t_n < 1$.

TABLE 6. Left to right: Patterns I, II, III, IV .

t	$f(t)$	$g(t)$									
0	Q_0	Q_2									
t_1	Q_1	Q_2	t_1	Q_1	Q_2	t_1	Q_0	Q_3	t_1	Q_0	Q_3
t_2	Q_1	Q_3	t_2	Q_1	Q_3	t_2	Q_0	Q_4	t_2	Q_0	Q_4
t_3	Q_1	Q_4									
t_4	Q_1	Q_0	t_4	Q_2	Q_4	t_4	Q_1	Q_0	t_4	Q_2	Q_4
t_5	Q_2	Q_0									

TABLE 7.

t	$f(t)$	$g(t)$
0	Q_0	Q_1
t_1	Q_0	Q_2

TABLE 8. Left to right: Patterns V and VI.

t	$f(t)$	$g(t)$	t	$f(t)$	$g(t)$
0	Q_4	Q_0	0	Q_4	Q_0
t_1	Q_4	Q_1	t_1	Q_4	Q_1
t_2	Q_4	Q_2	t_2	Q_0	Q_1
t_3	Q_0	Q_2	t_3	Q_0	Q_2

TABLE 9.

t	$f(t)$	$g(t)$
0	Q_3	Q_0
t_1	Q_4	Q_0

We start with $f(0) = Q_0$ and $g(0) = Q_2$ in Table 6. We see that there are four possibilities: pattern I, II, III, or IV. In Table 7, where $f(0) = Q_0$ and $g(0) = Q_1$, there is only one possible second step. After this step, the functions would follow one of the patterns I, II, III, or IV in Table 6. In Table 8, we start with $f(0) = Q_4$ and $g(0) = Q_0$. We see that there are two possibilities, patterns V and VI. Each of these ends with $f(t_3) = Q_0$ and $g(t_3) = Q_2$. These functions would then follow either pattern I, II, III, or IV

TABLE 10. Left to right: Patterns VII and VIII.

t	$f(t)$	$g(t)$	t	$f(t)$	$g(t)$
0	Q_4	Q_1	0	Q_4	Q_1
t_1	Q_4	Q_2	t_1	Q_0	Q_1
t_2	Q_0	Q_2	t_2	Q_0	Q_2

TABLE 11.

t	$f(t)$	$g(t)$
0	Q_2	Q_1
t_1	Q_3	Q_1
t_2	Q_4	Q_1

TABLE 12.

t	$f(t)$	$g(t)$
0	Q_3	Q_1
t_1	Q_4	Q_1

TABLE 13.

t	$f(t)$	$g(t)$
0	Q_3	Q_2
t_1	Q_4	Q_2
t_2	Q_0	Q_2

in Table 6. In Table 9 we have $f(0) = Q_3$ and $g(0) = Q_0$. There is only one possible second step, that is $f(t_1) = Q_4$ and $g(t_1) = Q_0$. These functions then would follow either pattern V or VI in Table 8. After this, the functions would follow one of the patterns, I, II, III, or IV in Table 6. In Table 10, we have $f(0) = Q_4$ and $g(0) = Q_1$. There are two possibilities, patterns VII or VIII. Both of these patterns end with $f(t_2) = Q_0$ and $g(t_2) = Q_2$. These functions would then follow one of the patterns I, II, III, or IV in Table 6. Similarly, the functions in Table 11 would subsequently follow one of the patterns VII or VIII in Table 10, then follow one of the patterns I, II, III, or IV in Table 6. The functions in Table 12 would follow either pattern VII or VIII in Table 10, then either pattern I, II, III, or IV in Table 6. The pairs of functions in tables 13 and 14 would each follow one of the patterns I, II, III, or IV in Table 6.

We have considered all possible combinations for $f(0)$ and $g(0)$. Note that $f(0) = Q_3$ and $g(0) = Q_4$ is not possible, since for t_1 , $f(t_1) = Q_3$, $g(t_1) = Q_0$, and $d_{\min}(f, g) \leq r_3 < d(Q_3, Q'_0)$. We see that all pairs of functions f and g , satisfying our conditions, contain one of the patterns I, II, III, or IV. So, if

TABLE 14.

t	$f(t)$	$g(t)$
0	Q_4	Q_2
t_1	Q_0	Q_2

there is a pair of functions $(f, g) \in \mathcal{P}$ such that $d_{\min}(f, g) > d(Q_3, Q'_0) = q'_3$, then f and g contain one of the patterns I, II, III, or IV. So, for any pair, $(f, g) \in \mathcal{P}$, we are now considering, we see that:

$$d_{\min}(f, g) \leq \max\{\min \text{ I, min II, min III, min IV}\}$$

where

$$\begin{aligned} d(Q_i, t_i) &= t'_i, & d(Q_i, r_i) &= r'_i, & d(Q_i, Q'_{i+2}) &= q'_i, & \text{and} \\ \min \text{ I} &= \min\{r'_0, r'_1, t'_1, t'_2, q'_1\} = \min\{r'_0, r'_1, t'_1, t'_2\} \\ \min \text{ II} &= \min\{r'_1, t'_2, q'_1, q'_2, q'_4\} = \min\{r'_1, t'_2, q'_4\} \\ \min \text{ III} &= \min\{r'_0, r'_4, t'_0, t'_1, q'_0\} = \min\{r'_0, r'_4, t'_0, t'_1\} \\ \min \text{ IV} &= \min\{r'_4, t'_0, q'_0, q'_2, q'_4\} = \min\{r'_4, t'_0, q'_2\}. \end{aligned}$$

Note that patterns I and III are mirror images of each other, as are II and IV.

If the “best” f and g is not as given in Table 1, then since all other pairs, f and g , contain one of the patterns I, II, III, or IV, then the best f and g must be one of the pairs of functions given in Table 15. Hence, the “best” pair f and g will be from either Table I, II, III, or IV in Table 15, and

$$d_{\min}(f, g) = \max\{\min \text{ I, min II, min III, min IV}\},$$

where $d_{\min}(f, g) = \min \text{ I}$ for Table I, $d_{\min}(f, g) = \min \text{ II}$ for Table II, $d_{\min}(f, g) = \min \text{ III}$ for Table III, and $d_{\min}(f, g) = \min \text{ IV}$ for Table IV. Consequently, $\alpha(X) \geq \max\{\min \text{ I, min II, min III, min IV}\}$, where $\alpha = \sigma, \sigma_0, \sigma^* \sigma_0^*, s, s^*$ since the functions f and g satisfy the requirements for all of the spans. We will show that $\sigma_0(X) \leq \max\{\min \text{ I, min II, min III, min IV}\}$ and consequently that $\alpha(X) = \max\{\min \text{ I, min II, min III, min IV}\}$ for $\alpha = \sigma, \sigma_0, \sigma^*, \sigma_0^*, s, s^*$.

Since I and III are mirror images of each other as are II and IV, we need to examine only two cases,

$$\min \text{ I} = \max\{\min \text{ I, min II, min III, min IV}\}$$

and

$$\min \text{ II} = \max\{\min \text{ I, min II, min III, min IV}\}$$

CASE 1: $\min \text{ II} = \max\{\min \text{ I, min II, min III, min IV}\} = \min\{r'_1, t'_2, q'_4\}$.

Suppose that $\max\{\min \text{ I, min II, min III, min IV}\} = \min \text{ II}$, that is, the best pair of functions f and g is from Table II. So, $d_{\min}(f, g) = \min$

TABLE 15. Left to right Table I, Table II, Table III, Table IV.

t	$f(t)$	$g(t)$									
0	Q_0	Q_2									
0.1	Q_1	Q_2	0.1	Q_1	Q_2	0.1	Q_0	Q_3	0.1	Q_0	Q_3
0.2	Q_1	Q_3	0.2	Q_1	Q_3	0.2	Q_0	Q_4	0.2	Q_0	Q_4
0.3	Q_1	Q_4									
0.4	Q_1	Q_0	0.4	Q_2	Q_4	0.4	Q_1	Q_0	0.4	Q_2	Q_4
0.5	Q_2	Q_0									
0.6	Q_2	Q_1	0.6	Q_2	Q_1	0.6	Q_3	Q_0	0.6	Q_3	Q_0
0.7	Q_3	Q_1	0.7	Q_3	Q_1	0.7	Q_4	Q_0	0.7	Q_4	Q_0
0.8	Q_4	Q_1									
0.9	Q_0	Q_1	0.9	Q_4	Q_2	0.9	Q_0	Q_1	0.9	Q_4	Q_2
1.0	Q_0	Q_2									

$\text{II} = \min\{r'_1, t'_2, q'_4\}$ and $\alpha(X) \geq \min\{r'_1, t'_2, q'_4\}$, where $\alpha = \sigma, \sigma_0, \sigma^*, \sigma_0^*, s$, and s^* . We will now show that $\alpha(X) = \min\{r'_1, t'_2, q'_4\} = \text{dmin}(f, g)$, where $\alpha = \sigma, \sigma_0, \sigma^*, \sigma_0^*, s$, and s^* .

Let $f^*, g^* : C \rightarrow Z$ be continuous functions from a continuum C into a continuum $Z \subseteq Y$ such that $f^*(C) \subseteq g^*(C)$. We consider three subcases.

CASE 1A: $\min \text{II} = r'_1, r'_1 \leq t'_2$, and $r'_1 \leq q'_4$.

Observe that $\min \text{IV} = \min\{r'_4, t'_0, q'_2\} \leq \min \text{II} = r'_1 \leq t'_2 < q'_2$, so $\min \text{IV} \neq q'_2$ and $\min \text{IV} = \min\{r'_4, t'_0\}$. So, either $r'_4 \leq r'_1 = \text{dmin}(f, g)$ or $t'_0 \leq r'_1 = \text{dmin}(f, g)$. We define $p_0 : Y \rightarrow Q'_0, Q_0 \cup Q'_0 Q_1$ based on $r'_1 \leq t'_2$ and either $r'_4 \leq r'_1$ or $t'_0 \leq r'_1$. Since $r'_1 \leq t'_2$, we define p_0 on Y_l by using case $p_0, Y_l, 1$ when $r'_1 = t'_2$ and case $p_0, Y_l, 2$ when $r'_1 < t'_2$. Also, either $r'_4 \leq t'_0$ or $t'_0 < r'_4$. We define p_0 on Y_r by case $p_0, Y_r, 1$ when $r'_4 \leq t'_0$ and by case $p_0, Y_r, 2$ when $t'_0 < r'_4$. Consider $p_0 \circ f^*, p_0 \circ g^* : C \rightarrow Q'_0 Q_0 \cup Q'_0 Q_1$. Since the range of $p_0 \circ g^*$ is an interval and $p_0 \circ f^*(C) \subset p_0 \circ g^*(C)$, we see that there is a $c \in C$ such that $p_0 \circ f^*(c) = p_0 \circ g^*(c)$ and $d(f^*(c), g^*(c)) \leq \max\{\min\{t'_0, r'_4\}, q'_3, r'_1\} = r'_1$.

CASE 1B: $\min \text{II} = t'_2, t'_2 < r'_1$, and $t'_2 \leq q'_4$.

Observe that $\min \text{IV} = \min\{r'_4, t'_0, q'_2\} \leq \min \text{II} = t'_2 < q'_2$, so $\min \text{IV} \neq q'_2$, and $\min \text{IV} = \min\{r'_4, t'_0\}$. So, either $r'_4 \leq t'_2 = \text{dmin}(f, g)$ or $t'_0 \leq t'_2 = \text{dmin}(f, g)$. We define $p_0 : Y \rightarrow Q'_0 Q_0 \cup Q'_0 Q_1$ based on $t'_2 < r'_1$ and either $r'_4 \leq t'_2$ or $t'_0 \leq t'_2$. Since $t'_2 < r'_1$, we define p_0 on Y_l by using case $p_0, Y_l, 1$. Also, either $r'_4 \leq t'_0$ or $t'_0 < r'_4$. We define p_0 on Y_r by case $p_0, Y_r, 1$ when $r'_4 \leq t'_0$ and by case $p_0, Y_r, 2$ when $t'_0 < r'_4$. Again we conclude that there is a $c \in C$ such that $d(f^*(c), g^*(c)) \leq \max\{\min\{t'_0, r'_4\}, q'_3, t'_2\} = t'_2$.

CASE 1C: $\min \text{II} = q'_4, q'_4 < r'_1$, and $q'_4 < t'_2$.

Observe that $\min I = \min\{r'_0, r'_1, t'_1, t'_2\} \leq \min II = q'_4 < r'_1$ and $q'_4 < t'_2$, $\min I = \min\{r'_0, t'_1\}$, so either $r'_0 \leq q'_4 = \overrightarrow{\text{dmin}(f, g)}$ or $t'_1 \leq q'_4 = \overrightarrow{\text{dmin}(f, g)}$. We define $p_1 : Y \rightarrow \overrightarrow{Q'_1 Q_1} \cup \overrightarrow{Q'_1 Q_2}$ based on either $r'_0 \leq t'_1$ or $t'_1 < r'_0$ for $y \in Y_r$ and either $t'_3 \leq r'_2$ or $r'_2 < t'_3$ for $y \in Y_l$. Note that $t'_3 < q'_3 < q'_4$. So we see that there is a $c \in C$ such that $d(f^*(c), g^*(c)) \leq \max\{\min\{r'_0, t'_1\}, \{\min\{r'_2, t'_3\}, q'_4\}\} = q'_4$.

Therefore, in case 1 we conclude that $\sigma_0(X) \leq \text{dmin}(f, g) = \min II$. Since X is a simple closed curve we know that $\sigma(X) = \sigma^*(X)$ and $\sigma_0(X) = \sigma_0^*(X)$. Also, we see that for all $t \in [0, 1]$ there is a $t' \in [0, 1]$ such that $f(t) = g(t')$ and $f(t') = g(t)$. Consequently, we see that $\sigma(X) = \sigma_0(X) = \sigma^*(X) = \sigma_0^*(X) = s(X) = s^*(X) = \min II$.

CASE 2: $\min I = \max\{\min I, \min II, \min III, \min IV\} = \min\{r'_0, r'_1, t'_1, t'_2\}$.

Suppose that $\min I = \max\{\min I, \min II, \min III, \min IV\}$, that the best pair of functions f and g is from Table I. So, $\text{dmin}(f, g) = \min I = \min\{r'_0, r'_1, t'_1, t'_2\}$ and $\alpha(X) \geq \min\{r'_0, r'_1, t'_1, t'_2\}$ where $\alpha = \sigma, \sigma_0, \sigma^*, \sigma_0^*, s$ and s^* . We will now show that $\alpha(X) = \min\{r'_0, r'_1, t'_1, t'_2\} = \text{dmin}(f, g)$ where $\alpha = \sigma, \sigma_0, \sigma^*, \sigma_0^*, s$ and s^* .

Let $f^*, g^* : C \rightarrow Z$ be continuous functions from a continuum C into a continuum $Z \subseteq Y$ such that $f^*(C) \subseteq g^*(C)$. We consider four subcases.

CASE 2A: $\min I = r'_0, r'_0 \leq t'_1, r'_0 \leq r'_1$, and $r'_0 \leq t'_2$.

Observe that $\min II = \min\{r'_1, t'_2, q'_4\} \leq \min I = r'_0$. We can assume that $r'_0 \neq r'_1$, since this case has been covered in case 1A, so $r'_0 < r'_1$. We can assume that $r'_0 \neq t'_2$, since this case has been covered in case 1B, so $r'_0 < t'_2$. So, we assume that $\min II = q'_4 \leq \min I = r'_0$. So, we have that $q'_4 \leq r'_0 = \text{dmin}(f, g)$. We can assume that $q'_4 \neq r'_0$, since this is covered in case 1C. So, $q'_4 < r'_0 = \text{dmin}(f, g)$. We define $p_1 : Y \rightarrow \overrightarrow{Q'_1 Q_1} \cup \overrightarrow{Q'_1 Q_2}$ based on case $p_1, Y_r, 1$ for $y \in Y_r$ since $r'_0 \leq t'_1$ and based on either case $p_1, Y_l, 1$ if $t'_3 \leq r'_2$ or case $p_1, Y_l, 2$ if $r'_2 < t'_3$ for $y \in Y_l$. Consider $p_1 \circ f^*, p_1 \circ g^* : C \rightarrow \overrightarrow{Q'_1 Q_1} \cup \overrightarrow{Q'_1 Q_2}$. Since the range is an interval there is a $c \in C$ such that $p_1 \circ f^*(c) = p_1 \circ g^*(c)$ and $d(f^*(c), g^*(c)) \leq \max\{\min\{t'_3, r'_2\}, q'_4, r'_0\} = r'_0$.

CASE 2B: $\min I = t'_1, t'_1 < r'_0, t'_1 \leq r'_1$, and $t'_1 \leq t'_2$.

Observe that $\min II = \min\{r'_1, t'_2, q'_4\} \leq \min I = t'_1$. We can assume that $t'_1 \neq r'_1$, since this is already covered in case 1A, so $t'_1 < r'_1$. We can assume that $t'_1 \neq t'_2$, since this is already covered in case 1B, so $t'_1 < t'_2$. So we assume that $\min II = q'_4 \leq \min I = t'_1$. So, we have that $q'_4 \leq t'_1 = \text{dmin}(f, g)$. We can assume that $q'_4 \neq t'_1$, since this is covered in case 1C. So, $q'_4 < t'_1 = \text{dmin}(f, g)$.

We define $p_1 : Y \rightarrow \overrightarrow{Q'_1 Q_1} \cup \overrightarrow{Q'_1 Q_2}$ based on case $p_1, Y_r, 2$ for $y \in Y_r$ since $t'_1 < r'_0$ and based on either case $p_1, Y_l, 1$ if $t'_3 \leq r'_2$ or case $p_1, Y_l, 2$ if $r'_2 < t'_3$ for $y \in Y_l$. Consider $p_1 \circ f^*, p_1 \circ g^* : C \rightarrow \overrightarrow{Q'_1 Q_1} \cup \overrightarrow{Q'_1 Q_2}$. Since the

range is an interval there is a $c \in C$ such that $p_1 \circ f^*(c) = p_1 \circ g^*(c)$ and $d(f^*(c), g^*(c)) \leq \max\{\min\{t'_3, r'_2\}, q'_4, t'_1\} = t'_1$.

CASE 2C: $\min I = r'_1, r'_1 < r'_0, r'_1 < t'_1$, and $r'_1 \leq t'_2$.

This case has been covered in case 1A.

CASE 2D: $\min I = t'_2, t'_2 < r'_0, t'_2 < r'_1$, and $t'_2 < t'_1$.

This case has already been covered in case 1B.

Therefore in case 2 we conclude that $\sigma_0(X) \leq \text{dmin}(f, g) = \min I$. Also, $\sigma(X) = \sigma^*(X)$ and $\sigma_0(X) = \sigma_0^*(X)$ since X is a simple closed curve and for all $t \in [0, 1]$ there is a $t' \in [0, 1]$ such that $f(t) = g(t')$ and $f(t') = g(t)$. Consequently, $\sigma(X) = \sigma_0(X) = \sigma^*(X) = \sigma_0^*(X) = s(X) = s^*(X) = \min I$.

The other three cases can be proved in a similar manner.

Hence, $\sigma(X) = \sigma_0(X) = \sigma^*(X) = \sigma_0^*(X) = s(X) = s^*(X) = \max\{q'_{j+3}, \min I, \min II, \min III, \min IV\} = \text{fss}(X)$. □

THEOREM 3.2. *Let Z be a continuum such that $Z \subseteq Y$, where Y is the closure of the bounded component of $R^2 - X$ and X is a five star-like simple closed curve, then $\alpha(Z) \leq \text{fss}(X)$, where $\alpha = \sigma, \sigma_0, \sigma^*, \sigma_0^*, s, s^*$.*

PROOF. Since each of the functions p_i for $i = 0, 1, 2, 3, 4$ were defined for any continuum $Z \subseteq Y$ where Y is the closure of the bounded component of $R^2 - X$, we see that for any such Z , $\alpha(Z) \leq \text{fss}(X)$, where $\alpha = \sigma, \sigma_0, \sigma^*, \sigma_0^*, s$, and s^* . □

COROLLARY 3.3. *Let Z be a simple closed curve such that $Z \subseteq Y$ where Y is the closure of the bounded component of $R^2 - X$ and X is a five star-like simple closed curve, then*

$$\sigma(Z) \leq \sigma(X) .$$

So, in this situation, the question by Howard Cook is answered in the affirmative.

COROLLARY 3.4. *Let Y be a plane separating continuum and X be a five star-like simple closed curve contained in the closure of a bounded component of $R^2 - Y$, then*

$$\sigma(X) \leq \sigma(Y) .$$

PROOF. Let Y be a plane separating continuum such that X is contained in one of the bounded components of $R^2 - Y$. We can assume that the origin, \mathcal{O} , is in the bounded component of $R^2 - Y$. There is an $r > 0$ such that the circle with center the origin and radius r , $C(\mathcal{O}, r)$, is such that Y is contained in the bounded component of $R^2 - C(\mathcal{O}, r)$. We will show that $\sigma(Y) \geq \sigma(X)$.

We will consider the case where $\sigma(X) = \min II$. The lines $\overleftrightarrow{Q_2Q_1}$ and $\overleftrightarrow{Q_2Q_4}$ separate the bounded component of $R^2 - C(\mathcal{O}, r)$ into four wedges. We let V_0 be the closure of the wedge that contains Q'_4 and Q'_0 and W_0 be the closure

of the wedge that is opposite V_0 . The set $V_0 - (\overline{Q_4Q'_4} \cup \overline{Q'_4Q_0} \cup \overline{Q_0Q'_0} \cup \overline{Q'_0Q_1})$ has two components. Let W'_0 be the closure of the component that intersects $C(\mathcal{O}, r)$. For each $z \in W_0 \cap Y$, let $r(z) \in \overrightarrow{Q_2z} \cap W_0 \cap C(\mathcal{O}, r)$ and for each $z \in W'_0 \cap Y$, let $r(z) \in \overrightarrow{Q_2z} \cap W'_0 \cap C(\mathcal{O}, r)$. The lines $\overrightarrow{Q_4Q_2}$ and $\overrightarrow{Q_4Q_1}$ separate the bounded component of $R^2 - C(\mathcal{O}, r)$ into four wedges. Let V_1 be the closure of the wedge that contains Q'_1 . Let W_1 be the closure of the wedge that is opposite V_1 . The set $\overline{Q_1Q'_1} \cup \overline{Q'_1Q_2}$ separates V_1 into two components. Let W'_1 be the closure of the component that intersects $C(\mathcal{O}, r)$. For $z \in W_1 \cap Y$, let $r(z) \in \overrightarrow{Q_4z} \cap W_1 \cap C(\mathcal{O}, r)$, and for $z \in W'_1 \cap Y$, let $r(z) \in \overrightarrow{Q_4z} \cap W'_1 \cap C(\mathcal{O}, r)$. Similarly, using the lines $\overrightarrow{Q_1Q_4}$ and $\overrightarrow{Q_1Q_2}$, we define the sets W_2 and W'_2 . Note that $Q_1 \in W_2$ and $Q'_2, Q'_3 \in W'_2$. We define r for $z \in W_2 \cap Y$ by $r(z) \in \overrightarrow{Q_1z} \cap W_2 \cap C(\mathcal{O}, r)$ and for $z \in W'_2 \cap Y$ by $r(z) \in \overrightarrow{Q_1z} \cap W'_2 \cap C(\mathcal{O}, r)$. Observe that for either $x \in W_0$ and $y \in W'_0 \cap \overrightarrow{Q_2x}$, or $x \in W_1$ and $y \in W'_1 \cap \overrightarrow{Q_4x}$, or $x \in W_2$ and $y \in W'_2 \cap \overrightarrow{Q_1x}$, $d(x, y) \geq \min \text{ II}$.

We can rotate $X \cup Y$ in the plane about the origin so that the ray $\overrightarrow{\mathcal{O}x}$ corresponds to the positive x-axis, where $\{x\} = W_0 \cap W'_2 \cap \overrightarrow{Q_1Q_2} \cap C(\mathcal{O}, r)$. Let θ_j for $j = 0, 1, 2, 3, 4, 5, 6$, be angles in the rotated position such that $0 = \theta_0 < \theta_1 < \theta_2 < \theta_3 < \theta_4 < \theta_5 < \theta_6 = 2\pi$, where $re^{i\theta_0}, re^{i\theta_3} \in \overrightarrow{Q_1Q_2} \cap C(\mathcal{O}, r)$, $re^{i\theta_1}, re^{i\theta_4} \in \overrightarrow{Q_2Q_4} \cap C(\mathcal{O}, r)$, and $re^{i\theta_2}, re^{i\theta_5} \in \overrightarrow{Q_4Q_1} \cap C(\mathcal{O}, r)$. Let $f : [0, 2\pi] \rightarrow [0, 2\pi]$ be a piecewise linear, surjective function, such that $f(\theta_j) = j(\frac{\pi}{3})$, and for $\theta \in [\theta_j, \theta_{j+1}]$, $f(\theta) = j(\frac{\pi}{3}) + (\frac{\theta - \theta_j}{\theta_{j+1} - \theta_j})(\frac{\pi}{3})$ for each $j = 0, 1, 2, 3, 4, 5$.

Let $h : C(\mathcal{O}, r) \rightarrow C(\mathcal{O}, r)$ be the homeomorphism given by $h(re^{i\theta}) = h(re^{if(\theta)})$.

We define $p : Y \rightarrow C(\mathcal{O}, r)$ by $p(y) = h \circ r(y) = h(re^{i\theta y}) = re^{if(\theta y)}$ where $r(y) = re^{i\theta y}$. We see that p is an essential map from Y onto $C(\mathcal{O}, r)$. Also, for each $y, z \in Y$ such that $p(y)$ and $p(z)$ are diametrically opposed to each other on $C(\mathcal{O}, r)$, $d(p^{-1}\{p(y)\}, p^{-1}\{p(z)\}) \geq \min \text{ II}$.

So in this case, by Theorem L, $\sigma(Y) \geq \sigma(X) = \min \text{ II}$. The cases where $\sigma(X) = \min\{q_i\}_{i=0}^4$, min I, min III, or min IV can be shown in a similar manner. We conclude that in all cases $\sigma(Y) \geq \sigma(X)$. □

When Y is a simple closed curve, the question by Howard Cook is answered in the affirmative in this situation.

REFERENCES

- [1] H. Cook, W. T. Ingram and A. Lelek, A list of problems known as Houston problem book. Continua (Cincinnati, OH, 1994), 365-398, Lecture Notes in Pure and Appl. Math. 170, Dekker, New York, 1995.
- [2] James F. Davis, *The equivalence of zero span and zero semispan*, Proc. Amer. Math. Soc. **90** (1984), 133-138.

- [3] A. Lelek, *Disjoint mappings and the span of spaces*, Fund. Math. **55** (1964), 199-214.
- [4] A. Lelek, *An example of a simple triod with surjective span smaller than span*, Pacific J. Math. **64** (1976), 207-215.
- [5] A. Lelek, *On the surjective span and semispan of connected metric spaces*, Colloq. Math **37** (1977), 35-45.
- [6] K. Tkaczynska, *The span and semispan of some simple closed curves*, Proc. Amer. Math. Soc. **111** 1991.
- [7] K. Tkaczynska, *On the span of simple closed curves*, Houston J. Math. **20** (1994), 507-528.
- [8] T. West, *Spans of simple closed curves*, Glas. Mat. Ser. III **24**(1989), 405-415.
- [9] T. West, *A bound for the span of certain plane separating continua*, Glas. Mat. Ser. III **32**(1997), 291-300.
- [10] T. West, *Spans of certain simple closed curves and related spaces*, Topology Proc. **23**, Summer 1998, 363-378.
- [11] T. West, *Concerning the spans of certain plane separating continua*, Houston J. Math. **25** (1999), 697-708.
- [12] T. West, *Spans of various two cells, surfaces, and simple closed curves*, Glas. Mat. Ser. III **37** (2002), 383-392.
- [13] T. West, *Spans of continua related to indented circles*, Glas. Mat. Ser. III **39** (2004), 171-183.

T. West
Department of Mathematics
University of Louisiana at Lafayette
LA 70504-1010
USA
E-mail: ThelmaWest@Yahoo.com

Received: 23.5.2006.

Revised: 4.12.2006.