

A fixed point theorem on asymptotic contractions

K. P. R. SASTRY*, G. V. R. BABU†, S. ISMAIL‡ AND M. BALAIAH§

Abstract. *The aim of this paper is to prove a fixed point theorem on asymptotic contractions with hypotheses slightly different from that of Chen [1], Theorem 2.2.*

Key words: *asymptotic contractions, complete metric space, fixed point*

AMS subject classifications: 54H25, 47H10

Received March 8, 2007

Accepted July 17, 2007

1. Introduction

Throughout this paper, (M, d) denotes a complete metric space, R^+ the set of all nonnegative reals, and Φ the class of all mappings $\varphi : R^+ \rightarrow R^+$ satisfying (i) φ is continuous, and (ii) $\varphi(t) < t$ for $t > 0$. For $x \in M$, the orbit of x is $\{x, Tx, T^2x, \dots\}$ and is denoted by $O(x)$.

In 2003, Kirk [2] introduced a new class of mappings namely *asymptotic contractions* and established the existence of fixed points for such mappings by using ultra filter methods.

Definition 1.1 (Kirk [2]). *A mapping $T : M \rightarrow M$ is said to be an asymptotic contraction, if there exists a sequence $\{\varphi_n\}_{n=1}^\infty$, $\varphi_n : R^+ \rightarrow R^+$ and $\varphi \in \Phi$ such that for all $n \in N$*

$$d(T^n x, T^n y) \leq \varphi_n(d(x, y)) \quad (1)$$

for all $x, y \in M$, where $\varphi_n \rightarrow \varphi$ uniformly on the range of d .

Theorem 1.2 (Kirk [2]). *Suppose $T : M \rightarrow M$ is an asymptotic contraction for which the mappings φ_n in (1) are also continuous. Assume that some orbit of T is bounded. Then T has a unique fixed point $x_* \in M$, and moreover the Picard sequence $\{T^n x\}_{n=1}^\infty$ converges to x_* for each $x \in M$.*

*8-28-8/1, Tamil street, Chinna Waltair, Visakhapatnam-530017, India, e-mail: kprsastry@hotmail.com

†Department of Mathematics, Andhra University, Visakhapatnam-530003, India, e-mail: gvr_babu@hotmail.com

‡Department of Mathematics, Andhra University, Visakhapatnam-530003, India, e-mail: shaik_1949@hotmail.com

§Department of Mathematics, Andhra University, Visakhapatnam-530003, India, e-mail: balaiah_m19@hotmail.com

In 2005, Chen [1] proved *Theorem 1.2*, under weaker assumptions without using ultra filter methods.

Theorem 1.3 (Chen [1], Theorem 2.2). *Let $T : M \rightarrow M$ be a mapping satisfying*

$$(1.3.1) \quad d(T^n x, T^n y) \leq \varphi_n(d(x, y))$$

for all $x, y \in M$, where $\varphi_n : R^+ \rightarrow R^+$ and $\lim_{n \rightarrow \infty} \varphi_n = \varphi$ uniformly on any bounded interval $[0, b]$. Suppose that φ is upper semicontinuous and $\varphi(t) < t$ for $t > 0$,

(1.3.2) *there exists a positive integer n_* such that φ_{n_*} is upper semicontinuous and $\varphi_{n_*}(0) = 0$, and*

(1.3.3) *there exists $x_0 \in M$ such that $O(x_0)$ is bounded.*

Then T has a unique fixed point $x_ \in M$ such that $\lim_{n \rightarrow \infty} T^n x = x_*$ for all $x \in M$.*

The following example shows that T may not have a fixed point if condition (1.3.2) is dropped in *Theorem 1.3*.

Example 1.4. *Let $M = \{0, 1, 2^{-1}, 2^{-2}, \dots\}$ with the usual metric. Define $T : M \rightarrow M$ by $T0 = 1$; $T(2^{-n}) = 2^{-(n+1)}$ for $n = 0, 1, 2, \dots$. Define φ_n on R^+ by $\varphi_n(t) = 2^{-1}t + n^{-1}$ and φ on R^+ by $\varphi(t) = 2^{-1}t$, $t \in R^+$. Then $\{\varphi_n\}$ and φ satisfy conditions (1.3.1) and (1.3.3). Here we observe that $\varphi_n(0) \neq 0$ for every n , so that condition (1.3.2) does not hold and T has no fixed points.*

In this paper, we prove that *Theorem 1.3* holds well if (1.3.2) is replaced by (1.3.2)': There exists $x_0 \in M$ such that $O(x_0)$ is bounded and there exists $y \in \overline{O(x_0)}$ such that $O(Ty)$ is closed.

2. Main result

Theorem 2.1. *Let $T : M \rightarrow M$ be a mapping satisfying (1.3.1) and (1.3.2)'. Then T has a unique fixed point $x_* \in M$ and $\lim_{n \rightarrow \infty} T^n x_0 = x_*$. In fact $\lim_{n \rightarrow \infty} T^n x = x_*$ for all $x \in M$.*

Proof. For $x_0 \in M$, the sequence $\{T^n x_0\}$ is Cauchy, which follows from the proof of *Theorem 1.3* (see [1], Theorem 2.2) and hence converges, say, to x_* . Thus

$$(2.1.1) \quad \overline{O(x_0)} = \{x_*\} \cup O(x_0).$$

Since $O(x_0)$ is bounded, there exists $b > 0$ such that $d(T^n x_0, T^m x_0) \leq b$ for all n, m . Now, by using (1.3.1), we have

$$d(T^n x_0, T^n x_*) \leq \varphi_n(d(x_0, x_*)).$$

Since $\varphi_n \rightarrow \varphi$ as $n \rightarrow \infty$, there exists a natural number N such that

$$\varphi_n(d(x_0, x_*)) < \varphi(d(x_0, x_*)) + 1 \quad \text{for all } n \geq N.$$

Take

$$a = \max_{1 \leq k \leq N-1} \{d(T^k x_0, T^k x_*), \varphi(d(x_0, x_*)) + 1\},$$

so that $d(T^n x_0, T^n x_*) \leq a$ for all n . Now

$$\begin{aligned} d(T^n x_*, T^m x_*) &\leq d(T^n x_*, T^n x_0) + d(T^n x_0, T^m x_0) + d(T^m x_0, T^m x_*) \\ &\leq a + b + a = 2a + b \quad \text{for all } n, m. \end{aligned}$$

Hence $O(x_*)$ is bounded. Thus $\{T^n x_*\}$ converges, say, to y_* .

We now show that $y_* = x_*$. If possible, suppose that $y_* \neq x_*$.

For a fixed positive integer N , we have

$$d(T^{n+N} x_0, T^{n+N} x_*) \leq \varphi_n(d(T^N x_0, T^N x_*)).$$

By letting $n \rightarrow \infty$, we get

$$d(x_*, y_*) \leq \varphi(d(T^N x_0, T^N x_*)).$$

Now, by letting $N \rightarrow \infty$, we have

$$d(x_*, y_*) \leq \varphi(d(x_*, y_*)) < d(x_*, y_*),$$

a contradiction. Thus $T^n x_* \rightarrow x_*$ as $n \rightarrow \infty$.

By condition (1.3.2)' and (2.1.1) the following two cases arise.

Case (i): $y = x_*$

Since $T^n x_* \rightarrow x_*$, $x_* \in O(Tx_*)$. Therefore there exists a positive integer N such that $x_* = T^N x_*$. Hence $T^{nN+1} x_* = Tx_*$ for $n = 1, 2, \dots$. Consequently, the subsequence $\{T^{nN+1} x_*\}$ of $\{T^n x_*\}$ converges to Tx_* . Thus $Tx_* = x_*$.

Case (ii): $y \neq x_*$

In this case $y \in O(x_0)$ by (2.1.1) and $O(Ty)$ is closed by (1.3.2)'. Also observe that $O(Ty)$ is bounded. Hence the conclusion follows by case (i).

Now, for $x \in M$ we have

$$d(T^n x, x_*) = d(T^n x, T^n x_*) \leq \varphi_n(d(x, x_*)) < \varphi(d(x, x_*)) + 1 \quad \text{for a large } n.$$

Hence $\{T^n x\}$ is bounded. Thus $T^n x \rightarrow x_*$ as $n \rightarrow \infty$. This completes the proof of the theorem. \square

The following is an example which satisfies (1.3.2)' but not (1.3.2).

Example 2.2. Let $M = \{0, 1, 2^{-1}, 3^{-1}, \dots\}$ with the usual metric. Define $T : M \rightarrow M$ by $T0 = 0$; $T(n^{-1}) = (n+1)^{-1}$ for $n = 1, 2, \dots$. For $t \in R^+$, define $\varphi_n(t) = n^{-1}$ for all n , and $\varphi(t) = 0$ for all t . Clearly $\varphi(t) < t$ for $t > 0$ and φ_n converges to φ uniformly on M . It is easy to verify the condition (1.3.1). Put $x_0 = 1$. Then, $O(x_0) = \{1, 2^{-1}, 3^{-1}, \dots\}$ is bounded. We have $0 \in \overline{O(x_0)}$ and $T^k x_0 \rightarrow 0$ as $k \rightarrow \infty$. Hence $0 \in O(Tx_0)$ so that $O(Tx_0)$ is closed. Consequently (1.3.2)' holds. Thus T satisfies all the conditions of Theorem 2.1 and 0 is the only fixed point of T .

However, we observe that $\varphi_n(0) \neq 0$ for every n , so that condition (1.3.2) fails to hold.

References

- [1] Y. Z. CHEN, *Asymptotic fixed points for nonlinear contractions*, Fixed Point Theory and Applications **2005**(2005), 213-217.
- [2] W. A. KIRK, *Fixed points of asymptotic contractions*, J. Math. Anal. Appl. **277**(2003), 645-650.