

## ERRATA

**Novel Syntheses of Thiazolo[3,2-a]pyrimidin-7-ones** [*Croat. Chem. Acta* 57 (1984) 183—185] by Vinko Škarić and Đurđica Škarić.

Page 183, line 19. The sentence should be completed as

The synthesis of II appeared to proceed *via* the intermediary 1-(2,3-di-iodo-propyl)-2-thiouracil (III), followed by a thiazolo-ring formation,...

**Solid/Liquid Equilibria in Aqueous Systems of Dodecyl Benzene Sulphonate and Alkaline Earth Ions** [*Croat. Chem. Acta* 57 (1984) 93—107] by Đ. Težak, F. Strajnar, D. Šarčević, O. Milat, and M. Stubičar.

Page 94, line 10 from the top: ref. 1, 2, 3 should be changed to 11, 12, 13; last line: name Lundlum should be changed to Ludlum and also in the reference 15., page 106; page 102, Fig. 7c should be changed in 7d, and 7d should be changed in 7c. Figure captions: Fig. 1, line two: dimenzation should be changed to dimerization; Fig. 7, line four: dorplets should be changed to droplets; Fig. 9, line five: srystalline should be changed to crystalline.

## ARTICLE

**Topological Localized Molecular Orbitals of Hückel and Möbius Annulenes**, [*Croat. Chem. Acta* 56 (1983) 491—497] by Albert Moyano and Juan Carlos Paniagua.

In page 494 the text from line 6 to line 12 should read:

LMO's of this kind appear therefore in annulenes with  $2p + pm$  carbon atoms and net charge  $pm+$ , where  $m = 0, 1, 2, \dots$  and  $p$  is the number of  $\pi$  bonds, which is odd or even for Hückel or Möbius closed shell systems, respectively.

A subset of the LMO's is said to be continuously degenerate if their localization sum is invariant under a non-discrete set of orthogonal transformations. There is therefore an infinite number of sets of LMO's corresponding to a plateau in the localization sum hypersurface. Although according to...

In page 495, in the second figure, a minus sign should be added to each drawing:

