

CCA-909

YU ISSN 0011-1643

541.126

Note

On the Stability Constants of Copper(II) Formato, Acetato, Propionato, Butyrato and Glycolato Complexes Determined by the Spectrophotometric Method

B. Grabarić and I. Filipović

Laboratory of Inorganic Chemistry, Faculty of Technology, University of Zagreb,
POB 179, 41001 Zagreb, Croatia, Yugoslavia

Received May 13, 1975

In an earlier paper¹ the stability constants of Co(II), Ni(II), Zn(II), Cd(II) and Pb(II) formato, acetato, propionato, butyrato, glycolato and chloroacetato complexes, determined by potentiometric¹, polarographic^{2,3} and spectrophotometric⁴ methods, were summarized. Good agreement between stability constants determined by different methods were obtained, except for the complexes specified in the title, which were examined spectrophotometrically and polarographically.

In the meantime we published an improved technique and numerical treatment in determining stability constants by the spectrophotometric method of corresponding solutions⁵, so we investigate these systems more rigorously.

The measurements were carried out at 25.0 ± 0.2 °C in buffer solutions of constant acid to salt ratio, being 1 : 2 for formate and glycolate, 5 : 1 for acetate and 6 : 1 for propionate and butyrate. Ionic strength of all solutions was kept at a constant value of 2 mol dm⁻³ by adding sodium perchlorate. Measurements of absorption were performed at 760 nm. The change of pH between 3 to 5 had no influence on absorption. The stability constants obtained under these experimental conditions and with the apparatus, which were same as previously⁴, but with the new technique and numerical treatment, are given in Table, together with their 95 per cent confidence intervals.

It must be emphasized, however, that the polarographic method is preferred over the spectrophotometric one, because the experimental conditions required for the former allow very low metal ion concentrations to be used. This is advantageous when precipitate can be formed. *E. g.* in the spectrophotometric investigation of the specified complexes (except for Cu(II) formato complexes), the required experimental conditions do not allow investigation at ligand concentrations higher than 0.1 mol dm⁻³ due to the precipitation of a slightly soluble Cu(II) complex.

The present results confirm all conclusions in the earlier papers¹⁻⁴, but are in much better agreement with those obtained by polarographic^{2,3,6} and potentiometric⁷ method, than the earlier ones⁴. This was expected because of the improvements in technique and numerical treatment.

TABLE
Stability constants of Cu(II) complexes

Ligand	$\log \beta_1$	$\log \beta_2$	$\log \beta_3$	$\log \beta_4$
Formate	1.59 ± 0.08	2.48 ± 0.11	2.92 ± 0.22	3.58 ± 0.12
Acetate	1.91 ± 0.01	2.46 ± 0.01		
Propionate	1.94 ± 0.01	2.76 ± 0.08		
Butyrate	1.89 ± 0.02	4.13 ± 0.02		
Glycolate	2.46 ± 0.04			

REFERENCES

1. I. Filipović, T. Matusinović, B. Mayer, I. Piljac, B. Bach-Dragutinović, and A. Bujak, *Croat. Chem. Acta*, **42** (1970) 541.
2. I. Filipović, I. Piljac, A. Medved, J. Savić, A. Bujak, B. Bach-Dragutinović, and B. Mayer, *Croat. Chem. Acta* **40** (1968) 131.
3. I. Filipović, A. Bujak, and V. Vukićević, *Croat. Chem. Acta*, **42** (1970) 493.
4. B. Grabarić and I. Filipović, *Croat. Chem. Acta*, **42** (1970) 479.
5. B. Grabarić, I. Piljac, and I. Filipović, *Anal. Chem.*, **45** (1973) 1932.
6. B. Grabarić, M. Tkalčec, I. Piljac, I. Filipović, and Vl. Simeon, *Anal. Chim. Acta* **74** (1975) 147.
7. B. Grabarić, B. Mayer, I. Piljac, and I. Filipović, *Electrochim. Acta* **20** (1975) 799.

SAŽETAK

Određivanje konstanti stabilnosti formijato-, acetato-, propionato-, butirato- i glikolato-kompleksa bakra(II) spektrofotometrijskom metodom

B. Grabarić i I. Filipović

Konstante stabilnosti kompleksa navedenih u naslovu određene su usavršenom spektrofotometrijskom metodom korespondirajućih otopina obzirom na tehniku mjerenja i numeričko-statističku obradu. Dobivene konstante stabilnosti mnogo se bolje slažu sa konstantama stabilnosti dobivenim polarografskom i potenciometrijskom metodom od ranije navedenih.

ZAVOD ZA ANORGANSKU KEMIJU
TEHNOLOŠKI FAKULTET
SVEUČILIŠTA U ZAGREBU

Prispjelo 13. svibnja 1975.

CCA-910

YU ISSN 0011-1643

539.19:516

Original Scientific Paper

Graph Theory and Molecular Orbitals. XVI. on π -Electron Charge Distribution¹

I. Gutman and N. Trinajstić

The Rugjer Bošković Institute, P.O.B. 1016, 41001 Zagreb, Croatia, Yugoslavia

Received May 16, 1975

An integral formula is derived exhibiting the dependence of the π -electron charge density (in the framework of Hückel theory) on the coefficients of the characteristic polynomial of the molecular graph and one of its subgraphs. Several simple consequences of this formula are demonstrated.

The π -electron charge density is one of the most important pieces of information about a conjugated molecule which can be obtained from simple Hückel MO calculations². This quantity was first introduced by Coulson³ and was later successfully correlated with a number of experimental findings (*e. g.*, reactivity towards electrophilic and nucleophilic reagents in aromatic substitution reactions⁴, proton NMR shifts⁵, *etc.*). Besides, there is a close relationship between the π -electron charge distribution and the dipole moment of a conjugated molecule⁶.

A well-known result is a uniform charge distribution in alternant hydrocarbons regardless any detail in their structure.⁷ No analogous regularity exists for non-alternant hydrocarbons. The above results were also quite recently obtained using graph-theoretical considerations and proof-techniques^{8,9}.

In this work we would like to present some additional results leading to the further understanding of the dependence of π -electron charge density on structural features of the molecule. We use here the formalism of graph theory as it was developed in earlier papers of this series¹⁰. Thus, in case the molecular graph G possesses N vertices, $\mathbf{A} \equiv \mathbf{A}(G)$, its adjacency matrix will be defined as

$$A_{pq} = \begin{cases} 1 & \text{if vertices } p \text{ and } q \text{ are adjacent} \\ 0 & \text{otherwise} \end{cases} \quad (1)$$

The characteristic polynomial of G is

$$P(x) \equiv \det |x\mathbf{1} - \mathbf{A}| \quad (2)$$

Its roots (*i. e.* graph spectrum) are denoted as x_1, x_2, \dots, x_N . For graphs without self-loops, as are the molecular graphs corresponding to conjugated hydrocarbons,

$$\sum_{j=1}^N x_j = 0 \quad (3)$$

Let G_r be a subgraph of G obtained after removal of the vertex r and all incident edges to it from the graph G . Let further G_h be a graph obtained by introducing a self-loop with a weight h on the vertex r of the graph G^{11-13} . The weight h is to be understood as a variable parameter. In matrix notation this means that

$$A_{pq}(G_h) = A_{pq}(G) \quad (4)$$

for all p and q , except for $p = q = r$, when

$$A_{rr}(G_h) = h \text{ and } A_{rr}(G) = 0 \quad (5)$$

Therefore, for a graph with a self-loop with weight h ,

$$\sum_{j=1}^N x_j = h \quad (6)$$

Of course,

$$G_h = G \text{ for } h = 0 \quad (7)$$

We denote the characteristic polynomial of G_r and G_h by P_r and P_h , respectively. The following relation

$$P_h = P - h P_r \quad (8)$$

was proved elsewhere^{12,13}.

The characteristic polynomial P may be written in the form

$$P \equiv P(x) = \sum_{n=0}^N a_n x^{N-n} \quad (9)$$

This enables us to define another polynomial, H ,

$$H \equiv H(x) = (ix)^N P(1/ix) \quad (10)$$

where $i = \sqrt{-1}$. Then

$$H = \sum_n i^n a_n x^n \quad (11)$$

which can be also presented as

$$H = U + i V \quad (12)$$

where

$$U \equiv U(x) = \sum_n (-)^n a_{2n} x^{2n} \quad (13)$$

$$V \equiv V(x) = \sum_n (-)^n a_{2n+1} x^{2n+1} \quad (14)$$

In an analogous way we define the polynomials H_r , U_r , V_r , and H_h , U_h , V_h ,

$$H_r = (ix)^{N-1} P_r(1/ix) = U_r + i V_r \quad (15)$$

$$H_h = (ix)^N P_h(1/ix) = U_h + i V_h \quad (16)$$

Because of the relation (8),

$$H_h = H - i h x H_r \quad (17)$$

and therefore,

$$U_h = U + h x V_r \quad (18)$$

$$V_h = V - h x U_r \quad (19)$$

which finally gives

$$|H_h| = [(U + h x V_r)^2 + (V - h x U_r)^2]^{1/2} \quad (20)$$

We note that the dependence of the coefficients a_n on the molecular graph structure is nowadays completely elucidated¹⁴⁻¹⁸. Therefore, the topological properties of the polynomials H , U , V , etc., are also well understood¹. For later discussion it will be important that the polynomials U and U_r are positive for all values of x . This is a consequence of the relation

$$(-)^n a_{2n} \geq 0 \quad (21)$$

which holds for an arbitrary graph¹.

A Topological Formula for π -Electron Charge Density

In Hückel theory the π -electron charge density (Q_r) on the atom r is related to the π -electron energy¹⁹ as follows

$$Q_r = \frac{\partial E_\pi}{\partial a_r} \quad (22)$$

where a_r is the Coulomb integral^{2,4,6} corresponding to the atom (vertex) r , and E_π is the total π -electron energy. When the corresponding graph-theoretical terminology is used¹⁰, the Eq. (22) is given by,

$$Q_r = \frac{\partial E_\pi(G)}{\partial A_{rr}} \quad (23)$$

The above equation may be further transformed by taking into account Eqs. (4), (5), and (7):

$$Q_r = \left[\frac{\partial E_\pi(G_h)}{\partial h} \right]_{h=0} \quad (24)$$

In¹ it is shown that the following equation is valid for graphs having no self-loops (that is, for graphs whose spectra fulfill Eq. (3)):

$$E_\pi(G) = (1/\pi) \int_{-\infty}^{+\infty} \frac{\ln |H|}{x^2} dx \quad (25)$$

Using Eq. (6) the above formula may be easily extended to include graphs possessing self-loops. Namely,

$$E_\pi(G_h) = h + (1/\pi) \int_{-\infty}^{+\infty} \frac{\ln |H_h|}{x^2} dx \quad (26)$$

A detailed analysis shows that

$$\frac{\partial E_\pi(G_h)}{\partial h} = 1 + (1/\pi) \int_{-\infty}^{+\infty} \frac{\partial}{\partial h} \frac{\ln |H_h|}{x^2} dx \quad (27)$$

which combined with Eq. (20) and substituted back into Eq. (23) yields a topological formula for π -electron charge density:

$$Q_r = 1 + (1/\pi) \int_{-\infty}^{+\infty} \frac{UV_r - VU_r}{U^2 + V^2} \frac{dx}{x} \quad (28)$$

Expression (28) exhibits the exact analytical form of the dependence of Q_r on the coefficients of the characteristic polynomials P and P_r . Another form of Eq. (28) is

$$Q_r = 1 + (1/\pi) \int_{-\infty}^{+\infty} \frac{P_r(ix)}{P(ix)} dx \quad (29)$$

This elegant formula was first obtained by Coulson and Longuet-Higgins¹⁹, but using a very different way of reasoning. It can be shown that Eqs. (28) and (29) are completely equivalent. However, the form of Eq. (28) is more convenient for the following discussion.

It results from Eqs. (28) and (29) that Q_r is uniquely determined with (and that it can be calculated from) the spectra of graph G and G_r . It is interesting to compare this conclusion with the original definition of charge density³, which is based solely on the coefficients of MO's (*i. e.* eigenvectors of the graph G).

DISCUSSION

The form of Eq. (28) indicates that the dependence of Q_r on the molecular topology is rather complicated. This is indeed the case^{8,9,20}. However, in the present discussion we wish to show some properties of the charge distribution which follow straightforwardly from our formula and the topological behaviour of the polynomials H and H_r .

First, for all alternant systems it is always^{17,18},

$$a_{2n+1} = 0 \quad (30)$$

and therefore it follows from Eq. (13),

$$V(x) = V_r(x) \equiv 0 \quad (31)$$

This leads, then, to

$$Q_r = 1 \quad (32)$$

Thus, the charge distribution in all alternant hydrocarbons (including free radicals) is uniform.

In non-alternants the charge density of a π -centre obviously depends both on the hydrocarbon itself and the π -centre position in the molecule. This fact is well reflected in formula (28). Namely, Eq. (28) may be also written as

$$Q_r = 1 + A_r + B_r \quad (33)$$

where

$$A_r = -(1/\pi) \int_{-\infty}^{+\infty} \frac{VU_r}{U^2 + V^2} \frac{dx}{x} \quad (34)$$

$$B_r = (1/\pi) \int_{-\infty}^{+\infty} \frac{U V_r}{U^2 + V^2} \frac{dx}{x} \quad (35)$$

Since the polynomials U , U_r (and of course, $U^2 + V^2$) are positive for all values of x , the sign of the integral A_r is determined with $V(x)$ only; that is to say solely from the molecular graph G . By analogy, the sign of the term B_r depends on the graph G_r only. Hence, A_r may be understood as being the »molecular« and B_r as being the »positional« contribution to the charge density. The actual numerical value of the π -electron charge density results from the delicate balance of these two contributions. As a simple illustration let us consider a non-alternant hydrocarbon molecule having only one odd-membered cycle Γ , while all other structural details are arbitrary. Let the size of Γ be γ . The coefficients a_{2n+1} of the characteristic polynomial of such graphs have the property

$$\text{sign} [(-)^n a_{2n+1}] = \begin{cases} 0 & \text{for } 2n+1 < \gamma \\ (-)^{(\gamma+1)/2} & \text{for } 2n+1 \geq \gamma \end{cases} \quad (36)$$

Therefore, from Eq. (13),

$$\text{sign} \left[\frac{V(x)}{x} \right] = (-)^{(\gamma+1)/2} \quad (37)$$

and finally from Eq. (34),

$$\text{sign} [A_r] = (-)^{(\gamma-1)/2} = \begin{cases} +1 & \text{if } \gamma = 4m+1 \\ -1 & \text{if } \gamma = 4m+3 \end{cases} \quad (38)$$

it is seen that the sign of A_r is not depending on r .

Let us now consider an atom r belonging to the cycle Γ . Graph G_r then contains no odd cycles and therefore^{17,18} $V_r(x) \equiv 0$. Consequently, $B_r = 0$, and the only contribution to Q_r stems from A_r . The following rule follows, therefore, immediately from Eq. (38).

Rule

In a conjugated hydrocarbon containing only one odd-membered cycle all atoms belonging to this cycle are negatively charged ($Q_r > 1$) if the size of the cycle is $4m+1$ and positively charged ($Q_r < 1$) if the size of the cycle is $4m+3$.

This result is independent of any other detail in the molecular structure.

The case when the considered π -centre does not belong to Γ may be treated by analogy. However, G_r contains now the cycle Γ and therefore,

$$\text{sign} \left[\frac{V_r(x)}{x} \right] = (-)^{(\gamma+1)/2} \quad (39)$$

which combined with Eq. (35) yields,

$$\text{sign} [B_r] = \begin{cases} -1 & \text{if } \gamma = 4m+1 \\ +1 & \text{if } \gamma = 4m+3 \end{cases} \quad (40)$$

a result which is expected since the total charge of the neutral hydrocarbon is zero.

REFERENCES

1. Part XV: I. Gutman and N. Trinajstić, *J. Chem. Phys.*, in press.
2. K. Higasi, H. Baba, A. Rembaum, *Quantum Organic Chemistry*, Wiley, New York 1965, Chapter V.
3. C. A. Coulson, *Proc. Roy. Soc. (London)* **A 169** (1939) 413.
4. L. Salem. *The Molecular Orbital Theory of Conjugated Systems*, Benjamin, New York 1966, Chapter 6.
5. R. B. Herman, *Int. J. Quantum Chem.* **2** (1968) 165.
6. A. Streitwieser, Jr., *Molecular Orbital Theory for Organic Chemists*, Wiley, New York 1961, Chapter 6.
7. C. A. Coulson and G. S. Rushbrooke, *Proc. Cambridge Phil. Soc.* **36** (1940) 193; G. G. Hall, *Proc. Roy. Soc. (London)* **A 191** (1955) 251.
8. Part V: I. Gutman and N. Trinajstić, *Chem. Phys. Lett.* **20** (1973) 25.
9. Part VI: I. Gutman, N. Trinajstić, and T. Živković, *Tetrahedron* **29** (1973) 3449.
10. For review see: I. Gutman and N. Trinajstić, *Topics Curr. Chem.* **42** (1973) 49; *Croat. Chem. Acta* **45** (1975).
11. R. B. Mallion, A. J. Schwenk, and N. Trinajstić, *Croat. Chem. Acta* **46** (1974) 171.
12. R. B. Mallion, N. Trinajstić, and A. J. Schwenk, *Z. Naturforsch.* **29a** (1974) 1481.
13. R. B. Mallion, A. J. Schwenk, and N. Trinajstić, in M. Fiedler (Ed.), *Recent Advances in Graph Theory*, Academia, Prague, 1975, p. 345.
14. C. A. Coulson, *Proc. Cambridge Phil. Soc.* **46** (1950) 202.
15. H. Sachs, *Publ. Math. Debrecen* **11** (1963) 119.
16. H. Hosoya, *Theoret. Chim. Acta* **25** (1972) 215.
17. Part I: A. Graovac, I. Gutman, N. Trinajstić, and T. Živković, *Theoret. Chim. Acta* **26** (1972) 67.
18. Part II: D. Cvetković, I. Gutman, and N. Trinajstić, *Croat. Chem. Acta* **44** (1972) 365.
19. C. A. Coulson and H. C. Longuet-Higgins, *Proc. Roy. Soc. (London)* **A 192** (1947) 39; *ibid.* **A 192** (1947) 16.
20. I. Gutman, *Ph. D. Thesis*, University of Zagreb 1973.

SAŽETAK

Teorija grafova i molekularne orbitale. XVI. O raspodjeli π -elektronskog naboja*I. Gutman i N. Trinajstić*

Izvedena je integralna formula koja ukazuje na vezu između raspodjele π -elektronskog naboja u konjugiranom ugljikovodiku i koeficijenata karakterističnog polinoma pripadnog molekularnog grafa i jednog njegovog podgraфа. Prikazano je također i nekoliko jednostavnih posljedica ove topološke formule.

INSTITUT »RUĐER BOŠKOVIĆ«,
41001 ZAGREB, HRVATSKA

Prispjelo 16. svibnja 1975.