
Uncommon Metastasis of Melanoma of the Face after Sixteen Years - Case Presentation

Irina Filipović-Zore¹, Aleksandar Milenović², Zoran Ivasović³, Jurica Šiljeg⁴

¹Department of Oral Surgery, School of Dental Medicine University of Zagreb

²Department of Maxillofacial Surgery, University Hospital "Dubrava", Zagreb

³Department of Oral Surgery, University Hospital Centre Zagreb

⁴Private Oral Surgery "Dr Tihomir Švajhler", Zagreb

Summary

We present the case of unusual metastasis of a facial melanoma in a lymph node of the neck, after sixteen years.

A 40-year-old male patient was referred to the oral surgeon because of an enlarged lymph node on the right side, region one. Due to the fact that he had a number of untreated teeth and radiating pain on the right side of the face he was admitted for examination with suspected odontogenic lymphadenitis. From the case history it was learnt that the patient had had an operation for melanoma in the right nasolabial cheek sixteen years previously. An examination revealed a postoperative scar on the right cheek, which was hard and painless, and a lymph node, two centimetres in size in region one, and also several untreated teeth. The patient was first sent for cytopuncture of the node, which showed metastasis of the melanoma. For preoperative preparation radical dissection was performed of the right side of the neck and all teeth extracted which could not be treated. Only one positive lymph node was found in the PHD specimen. The patient was included in the protocol for melanoma and all tests were carried out according to clinical guidelines.

As all findings are presently normal no further therapy is necessary.

The case shows that even in such situations we cannot disregard enlarged lymph nodes of the neck in differential diagnostics.

Neobična klinička slika i lokalizacija karcinoma gornje usnice - prikaz slučaja

Irina Filipović-Zore¹, Aleksandar Milenović², Zoran Ivasović³, Jurica Šiljeg⁴

¹Zavod za oralnu kirurgiju Stomatološkog fakulteta Sveučilišta u Zagrebu

²Klinika za kirurgiju čeljusti i lica KB "Dubrava", Zagreb

³Klinički zavod za oralnu kirurgiju KBC Zagreb

⁴Specijalistička stomatološka ordinacija za oralnu kirurgiju prim.dr.sc. Tihomira Švajhlera, Zagreb

Sažetak

U radu se prikazuje neobična klinička slika i lokalizacija karcinoma gornje usnice.

Pacijentica K. A., sedamdesetčetiri godine, dolazi u ambulantu za oralnu medicinu radi gnojnog infiltrata u gornjoj usnici sa sumnjom na ubod kukca. Uputi se u ambulantu za oralnu kirurgiju gdje se učini incizija i drenaža te se postavi dijagnoza pioge-noga granuloma.

Kako promjena u međuvremenu od mjesec-dva ne prolazi već postaje sve tvrđa i vidljivija, i vestibularno i labijalno, naruči se na ekciziju. U međuvremenu pacijentica odlazi na Kliniku za dermatovenerologiju zbog bazocelularnih promjena na licu. U tijeku ekscizija promjena lica učini se i površna biopsija s kožne strane usnice te se dobije PHD nalaz planocelularnoga karcinoma. Pacijentici se na klinici za dermatovenerologiju savjetuje radioterapija promjene, no pacijentica se ponovno javlja u ambulantu oralne kirurgije. Zbog sumnje u površno uzet nalaz promjene i eventualne zamjene s mioblastomom zrnatih stanica, učini se ekscizija promjene u cijelosti te se dobije identičan PHD nalaz planocelularnoga karcinoma. Tada se pacijentica dogovorno premjesti u KKLČU KB "Dubrava" gdje se učini proširena ekscizija, te se PHD-om utvrdi da je promjena u predhodnom zahvatu ekscidirana u cijelosti.