

Stela Jokić<sup>1\*</sup>, Krunoslav Aladić<sup>2</sup>, Đurđica Ačkar<sup>1</sup>, Antun Jozinović<sup>1</sup>, Jurislav Babić<sup>1\*\*</sup>, Drago Šubarić<sup>1\*\*</sup>

## SUPERCRITICAL CO<sub>2</sub> EXTRACTION – A NEW PERSPECTIVE IN THE UTILISATION OF FOOD INDUSTRY BY-PRODUCTS

<sup>1</sup>Faculty of Food Technology, Josip Juraj Strossmayer University of Osijek, F. Kuhača 20, 31000 Osijek, Croatia (\*stela.jokic@ptfos.hr)

<sup>2</sup>Croatian Veterinary Institute, Branch – Veterinary Institute Vinkovci, J. Kozarca 24, 32100 Vinkovci, Croatia

\*\*Croatian Academy of Engineering, Zagreb, Croatia

### Abstract

*The food industry, same as any other industry, aims to utilise raw materials as efficiently as possible in the production process, but it also tries to create minimum amounts of waste, where waste is not necessarily “waste”, but a by-product or a raw material for some subsequent process. Today there is great interest in utilising food industry by-products for various purposes because they contain many potentially useful substances, and they could become significant raw materials in the production/development of new products due to that fact. This paper places special emphasis on the influence of modern green extraction techniques, like supercritical CO<sub>2</sub> extraction, on the utilisation and extractability of certain biologically active components from selected by-products which have an application in food, cosmetics, and pharmaceutical industries.*

### 1. INTRODUCTION

The human awareness regarding the protection of nature and the environment is growing daily. Sustainable development, one that meets today's requirements and does not jeopardise the needs of future generations, has become an important issue, and all of the research on reducing the use of toxic chemicals, conserving energy, and managing waste and by-products is welcome at the international level. All over the world, there is research dedicated to this area, demanding that each researcher follows the tradition of sustainable development directed at a certain area, which also includes long-term decision making in the agriculture and food production sectors. The high quality of food production and the commitment to environmental protection has contributed to the successful marketing of agriculture and food products (Perretti, 2006). Large quantities of solid (shells, peels, seeds, leaves) and liquid waste (wastewater) are created during the processing of raw materials from plants in the food industry every year, and their storage, processing, or management present a serious ecological and economic problem. The large quantities of by-products created every day are mostly managed through disposal sites or by making cattle fodder. Therefore every industry, including the food industry, has the goal of fully using their raw materials in the production process, while creating

as little waste as possible, where waste is not necessarily “waste”, but a by-product or a raw material in some future process. Today there is great interest in using by-products from the food industry for various purposes because they contain many potentially useful substances, and they could represent significant raw materials in the production/development of new products (Schieber et al., 2001; O'Shea et al., 2012).

The use of food industry by-products has become extremely interesting because they represent cheap and nutritionally valuable raw materials, and that is enough to affect the reduction in waste. From the economic standpoint, large amounts of by-products, especially grape pomace and olive pomace, are still not being used in the best possible way in the Republic of Croatia. Furthermore, there is no competition on the domestic market in the area of highly profitable final products that could be created from those initial raw materials, and they are used for some of the most highly sought after food, cosmetics, and pharmaceutical products in the world.

Modern extraction techniques, like supercritical CO<sub>2</sub> (SC-CO<sub>2</sub>) extraction, that represent new, mild, and selective food processing technologies, with the purpose of environmental protection, are becoming more prominent in the world. In recent years, this technique has been applied in the extraction of potentially useful substances from food by-products (Table 1).

### 2. SELECTED FOOD INDUSTRY BY-PRODUCTS

#### 2.1. Grape pomace

Grape pomace created in the process of wine production, as well as olive pomace created in olive oil production, are just some of the potentially valuable by-products that could be successfully used in the production of some functional food products, but also in creating various extracts rich in active substances (Palma and Taylor, 1999; Ibáñez et al., 2000; Marti et al., 2001; Louli et al., 2004). Grape pomace represents a major problem due to its quantity and properties, primarily the leftover acids that make processing, usually composting, more difficult. If the pomace is not treated properly, it can represent a serious risk for the environment, starting with surface and deep pollution, to unpleasant odours created if it is left unprocessed. Na-

**Table 1.** Selected examples of SC-CO<sub>2</sub> extraction of high-value food industry by-products of plant origin

| Sample                 | Extraction parameters |                |           |                      |  |                         | Analyte                              | Reference                            |
|------------------------|-----------------------|----------------|-----------|----------------------|--|-------------------------|--------------------------------------|--------------------------------------|
|                        | Temperature (K)       | Pressure (MPa) | Time (h)  | Flow rate (g/min)    | Co-solvent (%)   | Particle size (mm)      |                                      |                                      |
| Grape seed             | 313                   | 16<br>18<br>20 | ---       | 10.2                 | ---  | 0.75                    | Grape seed oil                       | Passos et al. (2008)                 |
| Grape seed             | 310<br>353            | 25             | 7         | ---                  | ---  | 1.125<br>0.638<br>0.363 | $\alpha$ -tocopherol enriched oil    | Bravi et al. (2007)                  |
| Grape seed             | 313<br>323<br>333     | 20<br>30<br>40 | 1.5       | 32.33                | ---  | 0.380                   | Grape seed oil                       | Jokić et al. (2016)                  |
| Grape pomace           | 308<br>328            | 10<br>40       | 3         | 0.8                  | Ethanol (5)  | 0.165<br>0.261<br>0.319 | Resveratrol                          | Casas et al. (2010)                  |
| Grape pomace           | 318                   | 10<br>15<br>25 | ---       | 18.3                 | Methanol (5)   | ---                     | Phenolic compounds                   | Louli, Ragoussis and Magoulas (2004) |
| Grape skin             | 313                   | 15             | 0.25      | 2                    | Ethanol (7.5)  | ---                     | Resveratrol                          | Marti et al. (2001)                  |
| Elderberry pomace      | 313                   | 21             | 1.6       | ---                  | CO <sub>2</sub> (0-90)<br>Ethanol (0.5-100)<br>H <sub>2</sub> O (0-95) | ---                     | Anthocyanins                         | Seabra et al. (2010)                 |
| Apricot by-products    | 316-350               | 13.3-47.3      | 1.5       | 1                    | Ethanol (2-28)   | 0.07-0.6                | $\beta$ -carotene                    | Sanal et al. (2005)                  |
| Apricot seed           | 313-343               | 30-60          | ---       | 1-5                  | Ethanol (0-3)  | 0.425-1.5               | Apricot seed oil                     | Özkal, Yener and Bayındırlı (2005)   |
| Peach seed             | 303<br>313<br>323     | 10<br>20<br>30 | 2.5       | 8.3                  | Ethanol (2.5)  | 0.25-0.35               | Peach seed oil                       | Mezzomo et al. (2010)                |
| Tomato skin            | 313-373               | 20-40          | 1.5       | 1-2                  | ---  | ---                     | Lycopene                             | Chun et al. (2009)                   |
| Tomato skin            | 313<br>343            | 25<br>45       | 0.17-0.33 | 6.38                 | Ethanol (5.15)   | 0.5-1                   | Lycopene                             | Kassama, Shi and Mittal (2008)       |
| Tomato skin and seed   | 313<br>333<br>353     | 30<br>38<br>46 | ---       | ---                  | ---  | 0.3<br>0.4<br>0.6       | Carotenoids, tocopherols sitosterols | Vagi et al. (2007)                   |
| Apple and peach pomace | 313-333               | 20-60          | 0.17-0.67 | 2                    | Ethanol (14-20)  | 0.638                   | Phenolic compounds                   | Adil et al. (2007)                   |
| Cherry seed            | 313-333               | 18-22          | ---       | ---                  | ---  | 1.25-2.25               | Cherry seed oil                      | Bernardo-Gil et al. (2001)           |
| Orange peel            | 293 - 323             | 8-28           | ---       | 8.3-58.3             | ---  | 0.1-10                  | Essential oil                        | Mira et al. (1999)                   |
| Orange peel            | 313                   | 10             | 1         | 29.3                 | ---  | 0.324                   | Essential oil                        | Jerković et al. (2015)               |
| Citrus peel            | 333                   | 9.5            | 0.75      | ---                  | Ethanol (15)   | ---                     | Naringin                             | Giannuzzo et al. (2003)              |
| Olive pomace           | 323                   | 35             | 3         | 33                   | Ethanol (10)   | ---                     | Tocopherol                           | Ibáñez et al. (2000)                 |
| Olive pomace           | 313-323               | 10-30          | 2.5       | 1.8-2.7              | Ethanol (10)   | 0.30-0.55               | Olive oil                            | De Lucas, Rincon and Gracia (2003)   |
| Cocoa shell            | 323<br>358            | 15-45          | ---       | ---                  | ---  | 2-4                     | Theobromine                          | Rossi (1996)                         |
| Hemp pressed cake      | 313<br>323<br>333     | 20<br>30<br>40 | 1.5-7.5   | 11.7<br>29.1<br>46.7 | ---  | ---                     | Hemp oil<br>Defatted cake            | Aladić (2015)                        |
| Rice by-products       | 353                   | 68             | ---       | 1.082                | ---  | ---                     | Tocochromanols<br>Oryzanols          | Perretti et al. (2003)               |

mely, large quantities of pomace attract vermin and flies, and can lead to occurrences and spreading of various diseases. Tannin solutions and solutions of other wine pomace components, which are separated during resting, can cause the reduction of oxygen levels in the soil, but they can also penetrate into the soil and groundwater (Voća, 2010). If the pomace is used as fertilizer in vineyards, the soil in the vineyards becomes acidic over time, which is also a problem. Because of that, more and more attention is dedicated to processing waste or by-products from wineries that mostly consist of solid bio products (skins, stems, and seeds), which comprise on average 20-30% of the mass of the processed grapes, and 15% of the solid waste are grape seeds. Its use in the food industry may contribute to the reduction of production costs, as well as to the development of new products. The extraction process using SC-CO<sub>2</sub> can be used, for example, to completely extract oil from grape seeds, which is very significant in industrial processes, because it creates minimal losses in the production process and the full utilization of the initial raw material (Jokić et al., 2016). Oil obtained by using SC-CO<sub>2</sub> extraction is greenish-yellow in colour with a characteristic aroma; it is rich in vitamin E and can be used for cooking, medicine, and in pharmaceutical and cosmetic industry. The proportion of linoleic acid is higher when compared to any other oil, like sunflower or corn oil, which makes grape seed oil suitable for storage because it is highly stable. Defatted grape seed flour, which is created as a by-product of SC-CO<sub>2</sub> extraction, can also be used for other purposes (in making enriched extruded products, bakery industry, etc.). Grape seed pomace is also characterised by high phenolic compounds content due to low extraction during the wine production process. In the total amount of phenolic substances that can be extracted from grapes, 10% is in the pulp, 60-70% in the grape seeds, and 28-35% in the grape skin (Palma and Taylor, 1999; Murga et al., 2000; Louli et al., 2004) so the tablets and capsules based on extracts rich in polyphenolic compounds with antioxidant properties could be developed.

## 2.2. Olive pomace

Another by-product created in the process of olive oil extraction is olive pomace that includes skins and seeds. By-products of olive processing are rich in antioxidants, and various extraction techniques are used to isolate them from the plant material (Ibáñez et al., 2000; De Lucas et al., 2003). The use of the appropriate green technology, such as SC-CO<sub>2</sub> extraction, can result in the production of oil from olive pomace, where the quality and utilization of oil would be significantly higher than with using the standard pressing procedure. The goal is to get completely defatted olive pomace created as a by-product of SC-CO<sub>2</sub> extraction, which could then be used for the development of new functional products, i.e. in the process for the production of corn snack products enriched with defatted pomace.

## 2.3. Apricot kernels

Apricot kernels as by-products of processing are a valuable source of amygdalin (in literature often referred to as vitamin B17). Aside from that, apricot kernel seeds contain a high proportion of oil rich in mono- and polyunsaturated fatty acids, and a series of ingredients with lower proportions, like tocopherols and phenolic compounds (Turan et al., 2007). The oil extracted from apricot kernels using SC-CO<sub>2</sub> may have significant applications in cosmetics.

## 2.4. Cherry seeds

Cherry processing generates significant amounts of kernels for which there is no further use. However, since cherry seeds are a valuable source of oil (which can be obtained by SC-CO<sub>2</sub> extraction) rich in essential fatty acids, as well as in compounds like carotenoids, tocopherols, squalene, phytosterols, etc., they can be utilised as a potential natural source of nutraceutical ingredients in the pharmaceutical industry (Bak et al., 2009; Górnas et al., 2015).

## 2.5. Citrus peel

Three main by-products of citrus are dry pulp, molasses, and citrus oil (from the peel). Essential oils from citrus have many applications as aromas and for aromatising various products (Kesterson and Hendrickson, 1958). Essential oils can be produced from selected varieties of citrus peels by using various extraction techniques, where SC-CO<sub>2</sub> extraction is the preferred technique (Jerković et al., 2015).

## 2.6. Coffee processing by-products

By-products of coffee processing, such as mucilage and parchment, have been less studied, despite the fact that they contain important bioactive components (Esquivel and Jimenez, 2012). They can be an alternative source of natural antioxidants. Cocoa shells, which are removed during the processing of cocoa beans (*Theobroma cacao* L.), represent at least 10% of the weight of cocoa fruit (Owusu-Domfeh, 1972). In countries that produce cocoa beans, processing of such waste can provide an economic advantage and reduce the extent of environmental problems. The cocoa bean shell is also a potential source of theobromine (Hartati, 2010), which can be extracted using SC-CO<sub>2</sub>, since it has a great pharmacological function, it is a powerful diuretic, it promotes diuresis and stimulates circulation to the kidneys, and consequently helps to remove harmful substances from the urinary tract.

## 2.7. Tomato waste

A significant amount of waste is accumulated during the tomato canning process; it consists of seeds, skins (peel), and a small quantity of pulp (Avelino et al., 1997). Some of the created by-products are mostly used as additives to cattle fodder, and the unused remains are not managed, and as such represent a problem for the environment

(collection, disposal, and processing). The oil that can be created from tomato seeds by applying SC-CO<sub>2</sub> is attracting interest because of the abundance of unsaturated fatty acids, especially linolenic acid (Roy et al., 1996), as well as lycopene, a carotenoid responsible for the red colour of the tomato, the highest concentrations of which are in the skin (Favati et al., 1997; Baysal et al., 2000; Ollanketo et al., 2001; Rozzi et al., 2002).

### 2.8. Oilseed cakes

During the production of oil from oilseeds, it is of great importance to obtain higher extraction yields and oil quality. After oilseed processing, the remaining cake is commonly used in animal nutrition (Sorin-Stefan et al., 2013). The by-product (cake) from the process of oil pressing has great potential applications in food production due to the high content of residual oil, protein, fibre, minerals, and other substances. By using SC-CO<sub>2</sub>, it is possible to get a totally defatted cake (Aladić, 2015) with the potential application for other purposes (i.e. in the enrichment of extruded products, bakery industry, etc.).

## 3. SUPERCRITICAL FLUID EXTRACTION


With the discovery of supercritical fluids, toxic and ecologically unacceptable organic solvents are being gradually replaced. Supercritical fluid extraction (SFE) is an innovative technology and it represents an excellent alternative to standard extraction procedures using organic solvents. Additional reasons for that are the numerous advantages of supercritical fluids, like better diffusion, lower viscosity, and lower surface tension, which enables them to better penetrate the material used for extracting the targeted substance. In addition, this procedure enables high selectivity and solvent capacity control of the desired component in the supercritical fluid, by changing the pressure and temperature, and simply removing the solvent from the extracts. Furthermore, SC-CO<sub>2</sub> is considered completely safe for industrial application and food processing. This energy efficient process is considered as “clean technology”, because it does not create secondary products that are harmful for the environment and as such is very significant for the food industry (Brunner, 2005; Wang and Weller, 2006; Reverchon and De Marco, 2006; Abbas et al., 2008; Sahena et al., 2009; Temelli, 2009; Cvjetko Bubalo et al., 2015).

In the last few decades, this powerful separation process has drawn an increasing interest in commercial application, particularly due to its technical and environmental advantages compared to the current classical extraction methods using organic solvents. Extracts obtained using SC-CO<sub>2</sub> as the extraction solvent are solvent-free / without any trace of toxic extraction solvents, and are thereby highly valued (Jokić et al., 2014a). CO<sub>2</sub> is an adequate solvent for the extraction of lipophilic non-polar constituents. As such, it has a low affinity to polar compounds. This is one of the main drawbacks of extraction with SC-CO<sub>2</sub>. The low pola-

rity of SC-CO<sub>2</sub> can be overcome by employing polar modifiers (co-solvents, mainly ethanol and methanol) to change the polarity of the supercritical fluid and to increase its solvating power towards the analyte of interest. As such, according to Herrero et al. (2010), the addition of a low percentage of ethanol or methanol (1-10%) to CO<sub>2</sub> expands its extraction rates to include more polar analytes.

### 3.1. HM-SFE System

More than a few hundred commercial plants in the world today are using SFE in different fields (Jokić et al., 2015). The handmade supercritical fluid extraction (HM-SFE) system enables extraction in an inexpensive way. The obtained extraction yields and composition are very similar to those obtained by a commercial SFE system (Jokić et al., 2015; Jokić et al., 2014b). Just like a commercial SFE system, the HM-SFE system is composed of various components that need to be in tune in order to achieve the optimal extraction process. Parameters such as temperature, pressure, and fluid flow rate need to be monitored and controlled in order to enable an efficient and economical extraction process. Furthermore, the system must contain safety features with the possibility of an emergency system shutdown in the event of system failures. When designing the device, it is crucial to pay attention to several factors that make the device safe for work, because supercritical fluids and liquefied gases present a big hazard risk (explosion). Based on our experience in designing the HM-SFE system, we give special importance to the safety of the process. A certificate for working pressure was obtained for each used component, while calculations were carried out for the tube. All parts of the HM-SFE system should be made of stainless steel (AISI 304, AISI 316 Ti) to prevent any corrosion of the material. It is desirable to use certified filters with stainless steel



(1. Compressor; 2. CO<sub>2</sub> tank; 3. Stainless steel coil; 4. Cooling bath; 5. Air driven fluid pump; 6. High pressure needle valves (B-HV); 7. Manometers; 8. Extraction vessel; 9. Separator vessel; 10. Water bath; 11. Centralised system glass fibre heater; 12. Flow meter) (Jokić et al., 2015)

**Figure 1.** The HM-SFE system at the Faculty of Food Technology Osijek

mesh wire, coated with replaceable fine hard filter paper, in order to prevent plugging and possible hazard events.

When designing the extractor, aside from safety, it is very important to make an extractor which is very easy to maintain regarding loading and unloading, cleaning, connecting, and handling. From our experience in designing the HM extractor, we decided to make an extractor with a screw closure system (Jokić et al., 2015). Greater efficiency and continuous work are accomplished by using systems with the multiple parallel-connected extractors. The best closing system for a separator is also a screw type because of the fast extract collecting during the process at any time after depressurising. Due to the fact that it is necessary to heat and cool the fluid, heat exchangers are required in the HM-SFE system. Heat exchangers have the role of conditioning the fluid stream to required temperatures for pumping, extraction, separation, and holding (process storage) (Del Valle et al., 2014). From our experience, the heat exchanger should be placed just before the entrance of the extractor. This enables preheated CO<sub>2</sub> at extraction temperature to enter the extractor. On the HM-SFE system, an air driven non lubricated liquid pump is used to obtain the desirable carbon dioxide pressure. To prevent cavitation and irregularity in the operation of the pump, the head of the pump is additionally cooled through a cooling bath with a stainless steel coil.

The schematic diagram of the designed HM-SFE system (Faculty of Food Technology Osijek, Croatia) is given in **Figure 1** and explained in detail in the previous chapter of our book (Jokić et al., 2015).

#### 4. CONCLUSION

In recent years, SC-CO<sub>2</sub> extraction was demonstrated as a good alternative for processing food industry by-products. There is a great diversity of potential functional products which can be obtained from food by-products. Therefore, the use of food industry by-products has become extremely interesting because they represent cheap and nutritionally valuable raw material, and also have an impact on waste reduction. Modern companies are increasingly investing in research related to the reduction in the amount of waste and in research that would maximise the utilisation of raw materials. By applying SFE technology, it is possible to achieve the "zero waste approach" by ensuring that large amounts of organic waste are not created, which represents a huge environmental and financial problem in almost all branches of the food industry today.

#### References

- Abbas K.A., Mohamed A., Abdulmir A.S., Abas H.A.: A review on supercritical fluid extraction as new analytical method. *American Journal of Biochemistry and Biotechnology* 4: 345–353, 2008.
- Adil I.H., Cetin H.I., Yener M.E., Bayindirli A.: Subcritical (carbon dioxide + ethanol) extraction of polyphenols from apple and peach pomaces, and determination of the antioxidant activities of the extracts. *The Journal of Supercritical Fluids* 43: 55–63, 2007.
- Aladić K.: Optimizacija procesa ekstrakcije konopljinog (*Cannabis Sativa* L.) ulja superkritičnim CO<sub>2</sub> iz pogače nakon hladnog prešanja. *Doktorska disertacija*. Prehrambeno-tehnoški fakultet Osijek, 2015.
- Avelino A., Avelino H.T., Roseiro J.C., Collaco M.T.A.: Saccharification of tomato pomace for the production of biomass. *Bioresource Technology* 61: 159–162, 1997.
- Bak I., Lekli I., Juhasz B., Varga E., Varga B., Gesztelyi R., Szendrei L., Tosaki A.: Isolation and Analysis of Bioactive Constituents of Sour Cherry (*Prunus cerasus*) Seed Kernel: An Emerging Functional Food. *Journal of Medicinal Food* 13: 905–910, 2010.
- Baysal T., Ersus S., Starmans D.A.J.: Supercritical CO<sub>2</sub> extraction of beta carotene and lycopene from tomato paste waste. *Journal of Agricultural and Food Chemistry* 48: 5507–5511, 2000.
- Bernardo-Gil G., Oneto C., Antunes P., Rodriguez M.F., Empis J.M.: Extraction of lipids from cherry seed oil using supercritical carbon dioxide. *European Food Research and Technology* 212: 170–174, 2001.
- Bravi M., Spinoglio F., Verdona N., Adami M., Aliboni A., Andrea A.D., De Santis A., Ferri D.: Improving the extraction of a-tocopherol-enriched oil from grape seeds by supercritical CO<sub>2</sub>. Optimization of the extraction conditions. *Journal of Food Engineering* 78: 488–493, 2007.
- Brunner G.: Supercritical fluids: technology and application to food processing. *Journal of Food Engineering* 67: 21–33, 2005.
- Casas L., Mantell C., Rodríguez M., Martínez de la Ossa E.J., Roldán A., De Ory I., Caro I., Blandino A.: Extraction of resveratrol from the pomace of Palomino fino grapes by supercritical carbon dioxide. *Journal of Food Engineering* 96: 304–308, 2010.
- Cvijetko Bubalo M., Vidović S., Radojčić Redovniković I., Jokić S.: Green solvents for green technologies. *Journal of Chemical Technology & Biotechnology* 90: 1631–1639, 2015.
- Chun Y., John S., Sophia J.X., Yueming J., Dong L.: Effects of supercritical fluid extraction parameters on lycopene yield and antioxidant activity. *Food Chemistry* 113: 1088–1094, 2009.
- De Lucas A., Rincon J., Gracia I.: Influence of operation variables on quality parameters of olive husk oil extracted with CO<sub>2</sub>: Three-step sequential extraction. *Journal of the American Oil Chemists' Society* 80: 181–188, 2003.
- Del Valle, J.M., Núñez, G.A., Aravena, R.I.: Supercritical CO<sub>2</sub> oilseed extraction in multi-vessel plants. 1. Minimization of operational cost. *Journal of Supercritical Fluids* 92: 197–207, 2014.
- Esquivel P., Jimenez V.M.: Functional properties of coffee and coffee by-products. *Food Research International* 46: 488–495, 2012.
- Favati F., Galgano F., Daminano N., Lanzarini G.: Supercritical CO<sub>2</sub> extraction of lycopene and β-carotene from byproducts of the tomato industry. *4th Italian Conference on Supercritical Fluids and their Applications – Proceedings*, 121–128, 1997.
- Giannuzzo A.N., Boggetti H.J., Nazareno M.A., Mishima H.T.: Supercritical fluid extraction of naringin from the peel of *Citrus paradisi*. *Phytochemical Analysis* 14: 221–223, 2003.
- Górnaś P., Rudzińska M., Raczky M., Mišina I., Soliven A., Segliņa D.: Composition of bioactive compounds in kernel oils recovered from sour cherry (*Prunus cerasus* L.) by-products: Impact of the cultivar on potential applications. *Industrial Crops and Products* 82: 44–50, 2016.
- Hartati I.: Hydrotropic extraction of Theobromine from cocoa bean shell. *Momentum* 6: 17–20, 2010.
- Herrero, M., Mendiola, J.A., Cifuentes, A., Ibanez, E.: Supercritical fluid extraction: recent advances and applications. *Journal of Chromatography A* 1217: 2495–2511, 2010.

- Ibáñez E., Palacios J., Señoráns F.J., Santa-María G., Tabera J., Reglero G.: Isolation and separation of tocopherols from olive by-products with supercritical fluids. *Journal of the American Oil Chemists' Society* 77: 187-190, 2000.
- Jerković I., Družić J., Marijanović Z., Gugić M., Jokić S., Roje M.: GC-FID/MS Profiling of Supercritical CO<sub>2</sub> Extracts of Peels from *Citrus aurantium*, *C. sinensis* cv. *Washington navel*, *C. sinensis* cv. *Tarocco* and *C. sinensis* cv. *Doppio Sangugno* from Dubrovnik Area (Croatia). *Natural product communication* 10: 1315-1318, 2015.
- Jokić S., Bijuk M., Aladić K., Bilić M., Molnar M.: Optimization of supercritical CO<sub>2</sub> extraction of grape seed oil using response surface methodology. *International Journal of Food Science and Technology* 51: 403-410, 2016.
- Jokić S., Horvat G., Aladić K.: Design of SFE system using a holistic approach – problems and challenges. In: *Supercritical Fluid Extraction: Technology, Applications and Limitations* (Lindy J, Ed.). Nova Publishers, New York, SAD, pp. 95-122, 2015.
- Jokić S., Vidović S., Aladić K.: Supercritical Fluid Extraction of Edible Oils. In: *Supercritical Fluids: Fundamentals, Properties and Applications* (J. Osborne, Ed.). Nova Publishers, New York, SAD, pp. 205-228, 2014a.
- Jokić S., Moslavac T., Bošnjak A., Aladić K., Rajić M., Bilić M.: Optimization of walnut oil production. *Croatian Journal of Food Science and Technology* 6: 27-35, 2014b.
- Kassama L.S., Shi J., Mittal G.S.: Optimization of supercritical fluid extraction of lycopene from tomato skin with central composite rotatable design model. *Separation and Purification Technology* 60: 278-284, 2008.
- Kesterson J.W., Hendrickson R.: Utilization of Citrus By-Products. *Economic Botany* 12: 164-185, 1958.
- Louli V., Ragoussis N., Magoulas K.: Recovery of phenolic antioxidants from wine industry by-product. *Bioresource Technology* 92: 201-208, 2004.
- Marti M.C.P., Salvador A., Chafer A., Berna A.: Supercritical fluid extraction of resveratrol from grape skin of *Vitis inifera* and determination by HPLC. *Talanta* 54: 735-740, 2001.
- Mezzomo N., Mileo B.R., Friedrich M.T., Martínez J., Ferreira S.R.S.: Supercritical fluid extraction of peach almond oil: Process yield and extract composition. *Bioresource Technology* 101: 5622-5632, 2010.
- Mira B., Blasco M., Berna A., Subirats S.: Supercritical CO<sub>2</sub> extraction of essential oil from orange peel. Effect of operation conditions on the extract composition. *The Journal of Supercritical Fluids* 14: 95-104, 1999.
- Murga R., Ruiz R., Beltran S., Cabezas J.L.: Extraction of natural complex phenols and tannins from grape seeds by using supercritical mixtures of carbon dioxide and alcohol. *Journal of Agricultural and Food Chemistry* 48: 3408-3412, 2000.
- Ollanketo M., Hartonen K., Riekkola M.L., Holm Y., Hiltunen R.: Supercritical carbon dioxide extraction of lycopene in tomato skins. *European Food Research and Technology* 212: 561-565, 2001.
- Özkal S.G., Yener M.E., Bayındırlı L.: Mass transfer modelling of apricot kernel oil extraction with supercritical carbon dioxide. *The Journal of Supercritical Fluids* 35: 119-127, 2005.
- Owusu-Domfeh, K.: The future of cocoa and its by-products in the feeding of livestock. *The Ghana Journal of Agricultural Science* 5: 57-64, 1972.
- O'Shea N., Arendt E.K., Gallagher E.: Dietary fibre and phytochemical characteristics of fruit and vegetable by-products and their recent applications as novel ingredients in food products. *Innovative Food Science and Emerging Technologies* 16: 1-10, 2012.
- Palma M., Taylor L.T.: Extraction of polyphenolic compounds from grape seeds with near critical carbon dioxide. *Journal of Chromatography A* 849: 117-124, 1999.
- Passos C.P., Silva R.M., Da Silva F.A., Coimbra M.A., Silva C.M.: Enhancement of the supercritical fluid extraction of grape seed oil by using enzymatically pre-treated seed. *The Journal of Supercritical Fluids* 48: 225-229, 2009.
- Perretti G.: Supercritical Carbon Dioxide Extraction of Agricultural and Food Processing Wastes and Byproducts. In: *Modern Extraction Techniques* (Turner C., Ed.). ACS Symposium Series; American Chemical Society, Washington, DC, pp. 23-35, 2006.
- Perretti G., Miniati E., Montanari L., Fantozzi P.: Improving the value of rice by-products by SFE. *The Journal of Supercritical Fluids* 26: 63-71, 2003.
- Reverchon E., De Marco I.: Supercritical fluid extraction and fractionation of natural matter. *The Journal of Supercritical Fluids* 38: 146-166, 2006.
- Rossi M.: Supercritical fluid extraction of Cocoa and Cocoa products. In: *Supercritical Fluid Technology in Oil and Lipid Chemistry* (King J.W. & List G.R., Ed.) The American Oil Chemists Society, Champaign, Illinois, pp. 220-224, 1996.
- Rozzi N.L., Singh R.K., Vierling R.A., Watkin B.A.: Supercritical fluid extraction of lycopene from tomato processing byproduct. *Journal of Agricultural and Food Chemistry* 50: 2638-2643, 2002.
- Roy B.C., Goto M., Hirose T.: Temperature and pressure effects on supercritical CO<sub>2</sub> extraction of tomato seed oil. *International Journal of Food Science and Technology* 31:137-141, 1996.
- Sahena F., Zaidul I.S.M., Jinap S., Karim A.A., Abbas K.A., Norulaini N.A.N., Omar A.K.M.: Application of supercritical CO<sub>2</sub> in lipid extraction – A review. *Journal of Food Engineering* 95: 240-253, 2009.
- Sanal I.S., Guvenç A., Salgin U., Mehmetoglu U., Calimli A.: Recycling of apricot pomace by supercritical CO<sub>2</sub> extraction. *The Journal of Supercritical Fluids* 32: 221-230, 2004.
- Schieber A., Stintzing F.C., Carle R.: By-products of plant food processing as a source of functional compounds — Recent developments. *Trends in Food Science & Technology* 12: 401-413, 2001.
- Seabra I.J., Braga M.E.M., Batista M.T., Herminio C.S.: Effect of solvent (CO<sub>2</sub>/ethanol/H<sub>2</sub>O) on the fractionated enhanced solvent extraction of anthocyanins from elderberry pomace. *The Journal of Supercritical Fluids* 54: 145-152, 2010.
- Sorin-Stefan B., Ionescu M., Voicu G., Ungureanu N., Vladut V.: Calculus Elements for Mechanical Presses in Oil Industry. In: *Food Industry* (Dr. Muzzalupo, Ed.). INTECH Open Access Publisher, pp. 471-473, 2013.
- Temelli F.: Perspectives on supercritical fluid processing of fats and oils. *The Journal of Supercritical Fluids* 47: 583-590, 2009.
- Turan S., Topcu A., Karabulut I., Vural H., Hayaloglu A.A.: Fatty acid, triacylglycerol, phytosterol, and tocopherol variations in kernel oil of malatya apricots from Turkey. *The Journal of Agricultural and Food Chemistry* 55: 10787-10794, 2007.
- Voća N.: Proizvodnja toplinske energije iz vinske komine. *Završeno izvješće VIP projekta*. Sveučilište u Zagrebu, Agronomski fakultet, 2010.
- Vagi E., Simandi B., Vasa R.K.P., Daoud H., Kery A., Doleschall F., Nagy B.: Supercritical carbon dioxide extraction of carotenoids, tocopherols and sitosterols from industrial tomato by-products. *The Journal of Supercritical Fluids* 40: 218-226, 2007.
- Wang L., Weller C.L.: Recent advances in extraction of nutraceuticals from plants. *Trends in Food Science & Technology* 17: 300-312, 2006.