



CLINICAL VALUE OF TRANSVAGINAL ULTRASONOGRAPHY IN COMPARISON TO HYSTEROSCOPY WITH HISTOPATHOLOGIC EXAMINATION IN DIAGNOSING ENDOMETRIAL ABNORMALITIES

Hrvojka Soljačić Vraneš¹, Ivka Djaković¹, Zdenko Kraljević¹, Sandra Nakić Radoš²,
Tanja Leniček³ and Krunoslav Kuna¹

¹Department of Gynecology and Obstetrics, Sestre milosrdnice University Hospital Centre, Zagreb, Croatia;

²Department of Psychology, Catholic University of Croatia, Zagreb, Croatia;

³Ljudevit Jurak University Department of Pathology, Sestre milosrdnice University Hospital Centre, Zagreb, Croatia

SUMMARY – The aim of our retrospective study was to compare the performance of transvaginal sonography in relation to histologic diagnosis of samples obtained by hysteroscopy through analysis of data collected over 16 years. Data on suspected formation of endometrial polyp or submucosal fibroid found on ultrasound examination were extracted. The study included a total of 3679 women examined during the 2000-2015 period. All women underwent ultrasound examination preoperatively for better planning the type and scope of operation to be performed. The study included only women with samples for histopathologic analysis collected during the operation. Ultrasound diagnosis of polyps compared with histology showed 89.6% sensitivity and 39.1% specificity. For submucosal myomas, sensitivity was 69.2% and specificity 91.3%. In conclusion, ultrasound is not reliable method for definitive diagnosis but it is an excellent orientation method.

Key words: *Hysteroscopy; Polyps; Ultrasonography; Retrospective studies; Uterine neoplasms; Leiomyoma*

Introduction

Ultrasound has been in use for more than 60 years and nowadays it is the method of choice in uterine pathology evaluation. Transvaginal ultrasound has many advantages in practice. Proximity of tested areas enables high image quality and sharp picture of gynecologic structures. Additional benefits of the method are acquired by pulsed Doppler enabling insight into blood supply of the structures observed. This provides better assessment of normal and pathological structures.

Uterine myomas are the most common tumors in women of reproductive age¹. Myomas appear as echoic or hypoechoic nodules on ultrasound examination. If there are secondary changes in myomas, such as necrosis, calcification, hemorrhage, etc., ultrasound imaging will not be specific. If these secondary changes are located submucosally, differential diagnosis may be difficult because some other uterine changes look alike, i.e. polyps, hyperplasia, blood, mucus, etc. The use of color and pulsed Doppler can differentiate these changes more confidently, although in case of secondary changes such as necrosis or inflammation, low resistance index (RI) can be found as in malignancy².

Endometrial polyps appear as focal or diffuse changes in the uterine lining on ultrasound examination. The easiest visualization of polyps is achieved

Correspondence to: *Ivka Djaković, MD*, Department of Gynecology and Obstetrics, Sestre milosrdnice University Hospital Centre, Vinogradska c. 29, HR-10000 Zagreb, Croatia
E-mail: ivkadj@yahoo.com

Received May 2, 2018, accepted September 11, 2018

when the endometrium is thinnest, that is, immediately after menstrual bleeding in premenopausal women. Endometrial polyp can also be modified by secondary inflammation or necrosis. Examination during menstruation when fluid is present in the uterus or after instillation of contrast into the uterine cavity facilitates visualization of polyps and discernment to the thickened endometrium.

Hysteroscopy dates before ultrasound. The first diagnostic and therapeutic hysteroscopy was performed in the 19th century by Desormeaux³, when hysteroscopic endometrial polyp was treated with silver nitrate. Diagnostic hysteroscopy is used for visualization of the endocervical canal, endometrial cavity and tubal estuary. Operative hysteroscopy involves the use of mechanical, electrosurgical or laser instruments in the treatment of intracavitary pathology.

Hysteroscopy is now a widespread method in the diagnosis and treatment of endometrial abnormalities. The most common complications of hysteroscopic surgery include complications related to anesthesia and analgesia, excessive absorption of ionic liquids, perforation of the uterus, and injury of the surrounding structures. When the monopolar resectoscope is used, thermal injury to the vulva and vagina is possible⁴.

The aim of our retrospective study with 16-year data was to compare the performance of transvaginal sonography in relation to histologic diagnosis of samples obtained by hysteroscopy. Data on suspected formation of endometrial polyp or submucosal myomas found on ultrasound examination were extracted.

Patients and Methods

The study included a total of 3679 women examined during the 2000-2015 period at Sestre milosrdnice University Hospital Centre. All women were admitted for planned hysteroscopic surgery and underwent preoperative ultrasound examination for better planning the type and scope of the operation to be performed. Ultrasound is used to diagnose the presence of myomas or polyps. Endometrial polyps may be visualized on transvaginal ultrasound as nonspecific endometrial thickening or may be identified as focal masses within the endometrium. They can be hypoechoic, isoechoic or echogenic compared to the adjacent myometrium. Leiomyomas are usually well-defined, solid masses. Their echogenicity is usually simi-

lar to that of the myometrium, but sometimes they are hypoechoic or even hyperechoic compared to normal myometrium. They can have calcifications or necrosis. Calcifications are seen as echogenic foci with shadowing, and cystic areas may be seen if there is necrosis or degenerations.

The study included only women with samples for histopathologic analysis collected during the operation. Patients were analyzed in two groups: polyp group and myoma group. Study groups were tested for anthropologic characteristics such as age, menopausal status, marital status and place of residence.

We used χ^2 -test to determine the significance of differences in frequency between the groups. ANOVA (analysis of variance) was used to determine the significance of differences in patient age according to diagnosis and test period. When necessary, post-hoc Tukey HSD test was applied to determine differences among the groups. Online VassarStats calculator (<http://www.vassarstats.net/>) was used to determine sensitivity and specificity of a particular diagnostic test. The level of significance was set at $p < 0.05$ in all tests.

Results

The number of hysteroscopy procedures increased continuously during the study period (χ^2 (15, N=3679)=2331.51, $p < 0.001$).

The most common referral diagnoses were polyps and myomas. The incidence of polyps and myomas significantly increased in the 2000-2015 period (χ^2 (15, N=2945)=1871.34 and χ^2 (15, N=525)=222.18, respectively; $p < 0.001$ both).

The most common types of operation performed were ablation of polyp and ablation of myoma. The overall number of polyp ablations and myoma ablations significantly increased in the 2000-2015 period (χ^2 (15, N=2487)=1468.40 and χ^2 (15, N=525)=233.44, respectively; $p < 0.001$ both) (Fig. 1).

The most common histologic findings were polyps, myomas and normal endometrium, as illustrated in Figure 2. The number of polyps significantly increased in the 2000-2015 period (χ^2 (15, N=2453)=2065.51; $p < 0.001$). The incidence of myomas and normal endometrium (χ^2 (15, N=341)=259.54 and χ^2 (15, N=630)=636.99, $p < 0.001$ both) was significantly changed in the 2000-2015 period.




Fig. 1. Types of operation during the 2000–2015 period.




Fig. 2. Histopathologic findings during the 2000–2015 period.




Fig. 3. Age of women with polyps and myomas during the 2000–2015 period.

Significant correlation was found between the mean age and the diagnosis of polyps and fibroids. During the 2000–2015 period, the age of women diagnosed with polyps decreased by about 5 years, while the age of women diagnosed with myomas increased by about 4 years ($F(14, 2956)=1.74; p<0.05$), (Fig. 3).

Although there were more married women ($\chi^2(3M=2895)=24.07; p<0.01$), the groups did not differ according to marital status (Table 1). Menopausal women had the same incidence of polyps, but significantly fewer myomas ($\chi^2(1, N=2795)=36.25; p<0.001$) (Table 2). The incidence of myomas and polyps was the same in women from urban and rural areas ($\chi^2(1, N=2795)=0.17; p>0.05$) (Table 3).

A total of 2945 women were prediagnosed with polyps. In 2199 of them, diagnosis was confirmed by histopathology (Table 4). In addition, histopathology confirmed the diagnosis of polyps in 254 of 733 wom-

Table 1. Number of women with polyps or myomas according to marital status

	Polyp n (%)	Myoma n (%)
Married	1775 (72.3)	249 (73.0)
Single	325 (13.2)	54 (15.8)
Divorced	87 (3.5)	19 (5.6)
Widow	267 (11.0)	19 (5.6)
Total	2454	341

Table 2. Number of women with polyps or myomas according to menopausal status

		Polyp n (%)	Myoma n (%)
Menopause	Yes	1178 (48.0)	104 (31.7)
	No	1276 (52.0)	237 (68.3)
	Total	2454	341

Table 3. Number of women with polyps or myomas according to residence

		Polyp n (%)	Myoma n (%)
Residence	Rural	746 (30.4)	108 (31.7)
	Urban	1708 (69.6)	233 (68.3)
	Total	2454	341

Table 4. Frequency of polyp prediagnosis compared to definitive histopathology

		Definitive histopathology		Total
		Other	Polyp	
Prediagnosis	Polyp	746	2199	2945
	Other	479	254	733
Total		1225	2453	3678

Table 5. Metrical characteristics of definitive histopathology compared to polyp prediagnosis

	Value	95% confidence interval	
		Lower limit	Upper limit
Prevalence	0.667	0.651	0.682
Sensitivity	0.896	0.884	0.908
Specificity	0.391	0.363	0.419
False-positive	0.252	0.238	0.270
False-negative	0.347	0.312	0.382

Table 6. Frequency of myoma prediagnosis compared to definitive histopathology

		Definitive histopathology		Total
		Other	Myoma	
Prediagnosis	Myoma	289	236	525
	Other	3048	105	3153
Total		3337	341	3678

Table 7. Metrical characteristics of definitive histopathology compared to myoma prediagnosis

	Value	95% confidence interval	
		Lower limit	Upper limit
Prevalence	0.093	0.083	0.103
Sensitivity	0.692	0.640	0.740
Specificity	0.913	0.903	0.923
False-positive	0.550	0.507	0.593
False-negative	0.033	0.027	0.040

en with another referral diagnosis. The sensitivity and specificity for referral diagnosis was 89.6% and 39.1%, respectively (Table 5).

A total of 525 women were prediagnosed with myomas. In 236 of them, the diagnosis was confirmed by

histopathology (Table 6). In addition, histopathology confirmed the diagnosis of myomas in 105 of 3153 women with another referral diagnosis. The sensitivity and specificity for referral diagnosis was 69.2% and 91.3%, respectively (Table 7).

Discussion

Ultrasound is a widespread method that has evolved through sophisticated programs and devices. However, 2D ultrasound is still the most used method⁵. Ultrasound examination is simple, fast, noninvasive and easy to learn, with no side effects, while providing additional information to physical examination. In modern gynecologic routine, ultrasound is irreplaceable.

First attempts at constructing instruments for visualization of intrauterine pathology date back to 1805 when Binzzini constructed a series of concave mirrors to visualize uterine cavity⁶. First hysteroscopy was performed in 1869³. Since then, the procedure has been greatly improved, so that most patients are not hospitalized, even in case of therapeutic procedure⁷.

Complications of hysteroscopy are rare and more often due to therapeutic procedure⁸. Hysteroscopy is used for a wide range of diagnostic and therapeutic algorithms, e.g., assessment of sterility, recurrent miscarriage, uterine anomalies, and diagnosis and treatment of endometrial pathology⁹⁻¹¹. In the last fifteen years, hysteroscopy has been used more frequently for a wide spectrum of indications. Ultrasound is mostly used for detection of suitable cases and proper indications. Although most studies state that endometrial polyps are symptomatic, there are questions of the frequency of hysteroscopy and possibility of ultrasound follow up^{12,13}. The most common diagnoses in hysteroscopy candidates are endometrial polyps and submucosal myomas¹⁴, as well as histology findings¹⁵, as in our study.

Treatment of endometrial polyps and endometrial thickness in postmenopausal women differs. The former should always be treated operatively, whereas the latter require only follow up but the diagnosis is not always possible by ultrasound^{16,17}. So, the diagnosis should be definitive before making this decision. Like other authors¹⁸⁻²⁰, we compared the accuracy of ultrasound diagnosis in 3679 patients.

In our study, the sensitivity and specificity of ultrasound diagnosis of polyps compared with histology was 89.6% and 39.1%, respectively. For submucosal myomas, sensitivity was 69.2% and specificity 91.3%,

suggesting a conclusion that ultrasound is not reliable method for definitive diagnosis but is an excellent orientation method.

Myomas in postmenopausal women are usually asymptomatic and often not treated operatively. However, in premenopausal women, myomas often cause symptoms such as prolonged and heavy bleeding due to cavity deformities, and should be treated operatively.

In premenopausal women without the risk of endometrial hyperplasia or cancer (obesity, Lynch syndrome, polycystic ovary syndrome), polyps can only be followed up if asymptomatic, whereas in postmenopausal women, the risk of malignancy in polyps is higher and hysteroscopic diagnosis and histopathologic verification is recommended²¹.

In conclusion, ultrasound is not reliable method for definitive diagnosis of polyps and myomas but is an excellent orientation method.

References

1. Fleischer AC, Entman SS, Porrath SA, James AE. Sonographic evaluation of uterine malformations and disorders. In: Sanders RC, ed. *The Principles and Practice of Ultrasonography in Obstetrics and Gynecology*. Norwalk: Appleton Century Crofts, 1985: p. 531-45.
2. Kurjak A, Kupešić S, Mirić D. The assessment of benign uterine tumor vascularisation by transvaginal color Doppler. *J Ultrasound Med Biol*. 1992;6:645-9. [http://dx.doi.org/10.1016/0301-5629\(92\)90079-P](http://dx.doi.org/10.1016/0301-5629(92)90079-P)
3. Pantaleoni DC. On endoscopic examination of the cavity of the womb. *Med Press Circ*. 1869;8:26-7.
4. Munro MG, Christianson LA. Complications of hysteroscopic and uterine resectoscopic surgery. *Clin Obstet Gynecol*. 2015;58(4):765-97. <http://dx.doi.org/10.1097/GRF.0000000000000146>
5. Stadtmauer L, Shah A. Gynecologic surgery: preoperative assessment with ultrasound. *Clin Obstet Gynecol*. 2017;60(1): 82-92. <http://dx.doi.org/10.1097/GRF.0000000000000260>
6. Lindeman HJ. The hysteroscopy. In: Van der Pas H, Vancaillie T, eds. *Manual of Hysteroscopy*. Amsterdam-New York-Oxford: Elsevier Science Publishers B.V., 1990; p. 9-12.
7. Dealberti D, Riboni F, Prigione S, Pisani C, Rovetta E, Montella F, Garuti G. New mini-resectoscope: analysis of preliminary quality results in outpatient hysteroscopic polypectomy. *Arch Gynecol Obstet*. 2013;288(2):349-53. <http://dx.doi.org/10.1007/s00404-013-2754-7>
8. Cooper JM, Brady RM. Intraoperative and early postoperative complications of operative hysteroscopy. *Obstet Gynecol Clin North Am*. 2000;27:347-66.
9. El-Matny A, Abou-Salem N, El-Sherbiny W, Saber W. Outpatient hysteroscopy: a routine investigation before assisted reproductive techniques? *Fertil Steril*. 2011;95:272-6. <http://dx.doi.org/10.1016/j.fertnstert.2010.06.033>
10. Zolghardi J, Momtahan M, Aminian K, Ghaffarpasand F, Tavana Z. The value of hysteroscopy in diagnosis of chronic endometritis in patients with unexplained recurrent spontaneous abortion. *Eur J Obstet Gynecol Reprod Biol*. 2011;155:217-20. <http://dx.doi.org/10.1016/j.ejogrb.2010.12.010>
11. Dendrinou S, Grigoriou O, Sakkas EG, Makrakis E, Creatsas G. Hysteroscopy in the evaluation of habitual abortions. *Eur J Contracept Reprod Health Care*. 2008;13:198-200. <http://dx.doi.org/10.1080/13625180801920032>
12. Haimov-Kochman R, Deri-Hasid R, Hamani Y, Voss E. The natural course of endometrial polyps: could they vanish when left untreated? *Fertil Steril*. 2009;92:828. <http://dx.doi.org/10.1016/j.fertnstert.2009.04.054>
13. Lieng M, Istre O, Sandvik L, Qvigstad E. Prevalence, 1-year regression rate, and clinical significance of asymptomatic endometrial polyps: cross-sectional study. *J Minim Invasive Gynecol*. 2009;16:465-71. <http://dx.doi.org/10.1016/j.jmig.2009.04.005>
14. Aslam M, Ijaz L, Tariq S, Shafiqat K, Meher-Un-Nisa, Ashraf R, Kazmi T. Comparison of transvaginal sonography and saline contrast sonohysterography in women with abnormal uterine bleeding: correlation with hysteroscopy and histopathology. *Int J Health Sci*. 2007;1:17-24.
15. Kalampokas T, Gregoriou O, Grigoriadis C, Iavazzo C, Zervakis A, Sofoudis C, Kalampokas E, Botsis D. Comparing transvaginally defined endometrial thickness with hysteroscopic and histopathologic findings in asymptomatic postmenopausal women. *Eur J Gynaecol Oncol*. 2012;33:508-11.
16. Goldstein SR. Significance of incidentally thick endometrial echo on transvaginal ultrasound in postmenopausal women. *Menopause*. 2011;18:434-6. <http://dx.doi.org/10.1097/gme.0b013e31820ad00b>
17. Lieng M, Qvigstad E, Sandvik L, Jorgensen H, Langebrekke A, Istre O. Hysteroscopic resection of symptomatic and asymptomatic endometrial polyps. *J Minim Invasive Gynecol*. 2007; 14:189-94. <http://dx.doi.org/10.1016/j.jmig.2006.09.018>
18. Yantapan A. Comparison of the accuracy of transvaginal sonography and hysteroscopy for the diagnosis of endometrial polyps at Rajavithi Hospital. *J Med Assoc Thai*. 2012;95 Suppl 3:S92-7.
19. Kasraeian M, Asadi N, Ghaffarpasand F, Karimi AA. Value of transvaginal ultrasonography in endometrial evaluation of non-bleeding postmenopausal women. *Climacteric*. 2011;14: 126-31. <http://dx.doi.org/10.3109/13697137.2010.514079>
20. La Sala GB, Blasi I, Gallinelli A, Debbi C, Lopopolo G, Vinci V, Villani MT, Iannotti F. Diagnostic accuracy of sonohysterography and transvaginal sonography as compared with hysteroscopy and endometrial biopsy: a prospective study. *Minerva Gynecol*. 2011;63:421-7.
21. Lieng M, Istre O, Qvigstad E. Treatment of endometrial polyps: a systematic review. *Acta Obstet Gynecol Scand*. 2010;89(8): 992-1002. <http://dx.doi.org/10.3109/00016349.2010.493196>

Sažetak

KLINIČKA VRIJEDNOST TRANSVAGINALNE ULTRASONOGRAFIJE
U USPOREDBI S HISTEROSKOPIJOM I HISTOPATOLOŠKOM ANALIZOM
KOD DIJAGNOZE ENDOMETRALNIH ABNORMALNOSTI

H. Soljačić Vraneš, I. Djaković, Z. Kraljević, S. Nakić Radoš, T. Leniček i K. Kuna

Cilj ovog retrospektivnog istraživanja je bila usporedba pouzdanosti transvaginalnog ultrazvuka u odnosu na histološki nalaz dobiven histeroskopijom kroz analizu podataka skupljenih tijekom 16 godina. Podatci o suspektim endometralnim polipima i submukoznim miomima primijećenim na ultrazvučnom pregledu su izdvojeni. U istraživanje je uključeno 3679 žena pregledanih u razdoblju od 2000. do 2015. godine. Svim ženama je učinjen prijeoperacijski ultrazvučni pregled radi planiranja tipa i opsega zahvata. U istraživanje su uključene samo žene kojima je uzet histološki uzorak tijekom operacije. Ultrazvučno utvrđeni polipi u usporedbi s histološkim nalazom pokazali su osjetljivost od 89,6% i specifičnost od 39,1%. Za submukozne miome utvrđena je osjetljivost od 69,2% i specifičnost od 91,3%. Može se zaključiti kako je ultrazvuk nepouzdana metoda pri donošenju konačne dijagnoze, ali odlična orijentacijska metoda.

Cljučne riječi: *Histeroskopija; Polipi; Ultrazvuk; Retrospektivne studije; Uterus, tumori; Leiomiomi*