

TWO WATER MITE SPECIES (ACARI: HYDRACHNIDIA) FROM KARST SPRINGS NEW FOR THE FAUNA OF CROATIA WITH NOTES ON DISTRIBUTION AND ENVIRONMENTAL PREFERENCES

IVANA POZOJEVIĆ¹, VLADIMIR PEŠIĆ² & SANJA GOTTSTEIN^{1*}

¹Department of Biology, Faculty of Science, University of Zagreb, Rooseveltov trg 6, 10000 Zagreb, Croatia

²Department of Biology, Faculty of Science and Mathematics, University of Montenegro, Montenegro

Pozojević, I., Pešić, V. & Gottstein, S.: Two water mite species (Acari: Hydrachnidia) from karst springs new for the fauna of Croatia with notes on distribution and environmental preferences. Nat. Croat., Vol. 28, No. 2., 415-422, 2019, Zagreb.

One fifth of all described water mite species show clear preferences to spring ecosystems. The main goal of this research was to contribute to the knowledge of water mite assemblages, species richness and abundance in karst springs, with a special emphasis on comparing rheocene and limnocene spring ecosystems. The limnocene spring Modro Oko in the Southern Dalmatia had only one water mite species recorded in the studied period, showing very low abundances, whereas the rheocene spring of Jadro River in the Central Dalmatia showed both higher species richness (ten taxa recorded) and abundance. First records of two water mite species of the family Hygrobatidae from karst springs situated in the coastline area of Croatia are provided and discussed: *Hygrobates setosus* Besseling, 1942 and *Atractides distans* (K. Viets, 1914). Including these two additional records, a total number of 88 water mite species are recorded from Croatia so far.

Key words: Hygrobatidae, karst springs, rheocene, limnocene, new records, Dalmatia

Pozojević, I., Pešić, V. & Gottstein, S.: Dvije vrste vodenih grinja (Acari: Hydrachnidia) iz krških izvora, novih u fauni Hrvatske, s osvrtom na rasprostranjenost i okolišne sklonosti. Nat. Croat., Vol. 28, No. 2., 415-422, 2019, Zagreb.

Čak petina svih opisanih vrsta vodengrinja pokazuje izrazitu povezanost s izvorskim ekosustavima. Glavni cilj ovog rada bio je doprinijeti poznavanju zajednica, bogatstva vrsta i gustoće populacija vodengrinja u krškim izvorima, s posebnim naglaskom na usporedbu limnokrenih i reokrenih ekosustava. U limnokrenom izvoru Modro Oko u južnoj Dalmaciji zabilježena je samo jedna vrsta vodengrinja koja je u istraživanom razdoblju zabilježena s relativno malom gustoćom populacija, dok je u reokrenom izvoru rijeke Jadro u srednjoj Dalmaciji nađeno veće bogatstvo vrsta (10 zabilježenih svojiti) kao i veća gustoća populacija. U ovom radu prikazani su prvi nalazi dviju vrsta vodengrinja porodice Hygrobatidae iz krških izvora priobalne Hrvatske: *Hygrobates setosus* Besseling, 1942 i *Atractides distans* (K. Viets, 1914). S dva dodatna nalaza, u Hrvatskoj je zabilježeno ukupno 88 vrsta vodengrinja.

Ključne riječi: Hygrobatidae, krški izvori, reokreni, limnokreni, novi nalazi, Dalmacija

INTRODUCTION

Mites (Acari) are primarily terrestrial animals (DAVIDS & BELIER, 1979), but some groups like the true water mites (Acari: Hydrachnidia) have invaded aquatic ecosystems (KRANTZ & WALTER, 2009). Since their freshwater invasion, water mites have developed a high diversity - around 6.000 species are described,

* corresponding author: sanja.gottstein@biol.pmf.hr

with experts estimating these numbers to exceed 10.000 worldwide (DAVIDS *et al.*, 2007; GOLDSCHMIDT, 2016). Representatives of this group inhabit most types of freshwater ecosystems (MICCOLI *et al.*, 2013; WIĘCEK *et al.*, 2013), but are often neglected in environmental research due to perceived difficulties in identification (PROCTOR, 2007; GOLDSCHMIDT, 2016). In Europe today, around 1.000 species are described (Watermite.org, 2016), with the research on this group rapidly growing due to recently published comprehensive identification keys (DAVIDS *et al.*, 2007; DI SABATINO *et al.*, 2010; GERECKE *et al.*, 2016). In Croatia, the first checklist was published by PEŠIĆ (2002). Recent research in Croatia focused on the ecology of this group in both lentic and lotic Dinaric karst hydrosystems (POZOJEVIĆ *et al.*, 2018a; 2018b; 2019).

Springs play an important role in water mite assemblages (DI SABATINO *et al.*, 2003; GERECKE *et al.*, 2018). Among all aquatic invertebrates, they represent the highest ratio of crenophilous and crenobiont taxa, with approximately 20% of all species showing preference to spring habitats (GERECKE *et al.*, 2018). In general, karstic springs differ from springs of other typology in a higher degree of heterogeneity in terms of size, discharge characteristics, and water level fluctuations (BONACCI, 1993; SMART & WORTHINGTON 2004; KREŠIĆ, 2010).

The goal of this research is to contribute to the knowledge on the distribution of water mite assemblages, species richness and abundance in karst springs, with a special emphasis on comparing rheocene and limnocene ecosystems.

MATERIAL AND METHODS

The study area is located in the Dinaric Western Balkan ecoregion (ILLIES, 1978) of Croatia. Two karst springs with perennial discharge were selected: a river-forming rheocene, the Jadro spring, and a lake-forming limnocene, the Modro Oko spring (GLAZIER, 2009). The Jadro and Modro Oko springs (Fig. 1) are situated in South and Southeast Croatia, respectively, in a region mostly characterized by the Mediterranean, temperate humid climate with dry and hot summers, and mild and rainy winters (ŠEGOTA & FILIPČIĆ, 2003). The Jadro spring is located in the middle of Dalmatia on the foothills of the Mosor and Kozjak mountains. The spring is situated at an altitude of 35 m a.s.l., and it is part of the Cetina River basin (LOBOREC *et al.*, 2015). The protected freshwater spring Modro Oko is a significant landscape unit situated on the right bank of the Neretva River in Southern Dalmatia at an altitude of 9 m a.s.l.. It is influenced only slightly by seawater ($\leq 4\%$) from the Adriatic, infiltrating through the main body of the Neretva River, a marine lake (Vlaška) and the Crna Rijeka River (JASPRICA & HAFNER, 2005; GOTTSTEIN MATOČEĆ *et al.*, 2006; ŽGANEC *et al.*, 2010).

Each spring was sampled at two separate occasions: Jadro in May 2014 and in January 2015, whereas Modro Oko was sampled in November 2014 and in January 2015. At each sampling occasion, 20 subsamples were collected from each spring, taking all microhabitat types into consideration that made up at least 5% of the total substrate (AQEM Consortium, 2002). The samples from the Modro Oko were taken only within the littoral zone, while the samples from Jadro were

taken from all parts of the spring. The substrate was disturbed up to 5 cm in depth. Benthic macroinvertebrates were collected with a hand net (0.0625 m² area and 200 µm mesh size) and preserved in 96% ethyl alcohol. Water mites from the samples were subsequently isolated and fixed in Koenike's solution. Adult water mite individuals were determined to species or genus level using general keys by DAVIDS *et al.* (2007), DI SABATINO *et al.* (2010) and GERECKE *et al.* (2016). Deutonymphs were determined to genus level using the key of TUZOVSKIJ (1990). All water mite specimens are deposited at the Department of Biology, Faculty of Science, Zagreb, Croatia.




Fig. 1. The location and situation of the two study sites.

The following variables were measured at each spring: air and water temperature, oxygen concentration and saturation (WTW optical oxygen sensor: FDO 925), chemical oxygen demand (COD; potassium permanganate oxidation by Standard Methods - APHA, 2005), conductivity (WTW electrode conductivity measuring cell: TetraCon 925), average current velocity (SonTek FlowTracker), pH (WTW pH electrode: SenTix 940) and alkalinity (titration with 0.1 M HCl with methyl orange used as indicator).

RESULTS AND DISCUSSION

The Jadro spring, because of its rheocrenic nature, is characterized by a considerable current, while in the limnocrene Modro Oko no water current was measurable (Tab. 1). The turbulent water flow in the Jadro rheocrene results in oxygen saturation exceeding 100%, while this value was around 90 % in Modro Oko spring. Between the two respective sampling data, the water temperature

was more stable in the rheocrene (oscillation of 0.1°C), when compared with the limnocrene (oscillation of 1.4°C). The variability of the limnocrene water temperature was influenced by the air temperature because of the slow flow of current and direct insolation, while the riparian shading on the rheocrene was greatly reducing temperature oscillations. The conductivity is in both springs at typical levels for springs with elevated alkalinity due to dissolved calcium carbonate. The sea water infiltration had obviously only a minimum effect on Modro Oko. Both springs are relatively pristine, with little or no inflow of allochthonic organic matter and/or nutrients in the groundwater catchment. Slightly higher levels of organics (expressed by chemical oxygen demand, COD) were measured in the standing water of the limnocrene, when compared with the rheocrene.

Tab. 1. Environmental parameters measured at Jadro and Modro Oko springs, Croatia.

	Jadro Spring		Modro Oko Spring	
Coordinates	N 43°32'30.0"; E 16°31'07.5"		N 43°05'77.0", E 17°51'07.0"	
Spring type	Rheocrene		Limnocrene	
Altitude (m a.s.l.)	17		9	
Sampling date	May 18 th 2014	January 4 th 2015	November 28 th 2014	January 3 rd 2015
Microhabitat coverage	60% phytal (macrophytes); 30% macrolithal; 10% mesolithal	60% microlithal + mesolithal; 40% phytal (macrophytes)	80% phytal (macrophytes) + argyllal*; 20% mesolithal	70% argyllal*; 30% phytal (macrophytes)
Air temperature (°C)	21.5	10.8	15.1	12.3
Water temperature (°C)	12.7	12.6	13.4	12
Oxygen concentration (mg/L)	11.1	10.77	9.17	9.71
Oxygen saturation (%)	104.2	101.7	87.2	90.2
pH	7.9	7.8	7.52	7.41
Conductivity (µS/cm)	375	448	447	474
Alkalinity (mg CaCO ₃ /L)	190	205	220	232.5
Chemical oxygen demand (O ₂ mg/L)	0.67	1.49	4.4	4.65
Current velocity (m/s) - min	0.04	0.48	0	0
Current velocity (m/s) - max	0.24	0.98	0	0
Current velocity (m/s) - average	0.13	0.73	0	0
Water depth at sampling site (cm) - max	60	50	40	70
Water depth at sampling site (cm) - min	25	30	25	5

*argyllal substrate = silt, loam and/or clay

During this research, a total of 102 water mite specimens (92 adults, 7 deutonymphs and 3 larvae) were collected, belonging to 10 species, 7 genera and 6 families (Tab 2). The highest diversity was found at the rheocrene in May 2014 on the mesolithal microhabitat (here also the highest recorded abundance: 200 individuals per square meter). In the limnocrene, during the study period only

Tab. 2. Water mite species and number of specimens recorded at Jadro and Modro Oko springs, Croatia. The calculated abundances per square meter are given in brackets.

Study site:		Jadro Spring				Modro Oko Spring			
		May 18 th 2014	January 4 th 2015	November 28 th 2014	January 3 rd 2015	January 4 th 2015	November 28 th 2014	January 3 rd 2015	
Microhabitat sample:		phytal (macrophytes)	mesolithal	mesolithal + mesolithal	phytal (macrophytes)	phytal (macrophytes) + argyllal	mesolithal	argyllal	phytal (macrophytes)
Family	Species								
Hydryphantidae	<i>Proctia</i> sp.	1 (1.3)							
Hygrobatidae	<i>Atractides</i> sp.	34 (45.3)	5 (13.3)						
	<i>Atractides distans</i> (K.Viets, 1914)		2 (5.3)						
	<i>Atractides gibberpalpis</i> Piersig, 1898	5 (6.7)	1 (2.7)		4 (32)				
	<i>Atractides nodipalpis</i> (Thor, 1899)	4 (5.3)			3 (24)				
	<i>Atractides pumatus</i> (K. Viets, 1920)			2 (2.7)					
	<i>Hygrobatas setosus</i> Besseling, 1942						3 (3.4)	1 (1)	1 (4)
Lebertiidae	<i>Lebertia</i> sp.	4 (5.3)	5 (13.3)		2 (16)				
Oxidae	<i>Oxus setosus</i> (Koenike, 1898)				1 (8)				
Sperchontidae	<i>Sperchon</i> sp.	2 (2.7)	2 (5.3)		1 (8)				
Torrenticolidae	<i>Torrenticola</i> sp.		1 (2.7)		2 (16)				
Hydrachnidia larvae		3 (4)							
Water mite species richness		6	6	2	7	0	1	1	1
Water mite abundance (ind./m ²)		70.7	42.7	4	200	0	3.4	1	4

Hygrobates setosus Besseling, 1942 was found, with low abundances not exceeding 4 individuals per square meter.

The rheocrene was characterized by different rithrobiont taxa such as: *Oxus setosus* (Koenike, 1898), a species usually found in lowland rivers (DI SABATINO *et al.*, 2010), *Atractides distans* (K. Viets, 1914), *A. gibberipalpis* Piersig, 1898, *A. nodipalpis* (Thor, 1899), but also a typical crenobiont species: *A. pennatus* (K. Viets, 1920) (GERECKE, 2003). Following GERECKE *et al.* (2018), rheocrenes should usually be inhabited by assemblages of typical rheocrene species and rithrobionts, whereas limnocrenes should be characterized by crenobionts not specialized to these habitats and ubiquitous species. So far, *Hygrobates marezaensis* Pešić and Dabert, 2017 from Montenegro and Croatia is the only known typical limnocrenobiont (PEŠIĆ *et al.*, 2017). In our study, the rheocrene showed much higher abundances and taxa richness when compared to the limnocrene. *Hygrobates setosus*, the only species found in the Modro Oko, has recently been recorded also in the strong-flowing limnocrene Vitoja in Montenegro (PEŠIĆ *et al.*, 2019). This species has only recently been distinguished from *H. nigromaculatus* Lebert, 1879 by MARTIN *et al.* (2010), and belongs to the *H. nigromaculatus* complex. *Hygrobates nigromaculatus* Lebert, 1879 is usually found in lakes (recorded in Croatia by POZOJEVIĆ *et al.*, 2019), whereas *H. setosus* is in general inhabiting streams.

Since the first checklist of water mites in Croatia (PEŠIĆ, 2002, with 48 species), information on the group was broadened in two occasions in the course of research on water mites of the Balkan Peninsula (PEŠIĆ *et al.*, 2010, 2018). These checklists are missing records from MATONIČKIN & PAVLETIĆ (1959) [*Woolastookia rotundifrons* (K. Viets, 1922)] and the doctoral thesis of LATTINGER (1988) [*Partnunia angusta* (Koenike, 1893), *Paninus torrenticolus* Piersig, 1898 and *Protzia squamosa* Walter, 1908]. Furthermore, POZOJEVIĆ *et al.* (2018a) published records of three water mite species new for the fauna of Croatia from lake Torak and Čikola River, and POZOJEVIĆ *et al.* (2019), in a comprehensive overview for the fauna of reservoirs in the Dinaric ecoregion (ILLIES, 1978), added an additional 19 species. Including the two additional records given here, a total of 88 water mite species is now recorded from Croatia.

ACKNOWLEDGEMENTS

This research was supported by the projects MULTISEK (Multimetric evaluation of crenobiocoenosis) funded by the Environmental Protection and Energy Efficiency Fund (code: 351-01/13-01/108). We are very grateful to two anonymous referees for commenting and correction on an earlier version of the manuscript. Many thanks to Ivica Barač for all of his help during the field trips.

Received July 26, 2019

REFERENCES

- BONACCI, O., 1993: Karst springs hydrographs as indicators of karst aquifers. *Hydrological Sciences Journal* **38**, 51–62.
- DAVIDS, C. & BELIER, R., 1979: Spermatophores and sperm transfer in the water mite *Hydrachna conjecta* Koen. Reflections of the descent of water mites from terrestrial forms. *Acarologia* **21**, 84–90.
- DAVIDS, C., DI SABATINO, A. & GERECKE, R., 2007: Chelicerata: Araneae, Acari I. In GERECKE, R. (ed.) *Süßwasserfauna von Mitteleuropa 7/2-1*. Spektrum Akademischer Verlag, Heidelberg, pp 241–333.
- DI SABATINO, A., CICOLANI, B., & GERECKE, R., 2003: Biodiversity and distribution of water mites (Acari, Hydrachnidia) in spring habitats. *Freshwater Biology* **48**, 2163–2173.
- DI SABATINO, A., GERECKE, R., GLEDHILL, T. & SMIT, H., 2010: Chelicerata: Acari II. In Gerecke R. (ed.) *Süßwasserfauna von Mitteleuropa 7/2-2*. Spektrum, Heidelberg, pp 1–216.
- GERECKE, R., 2003: Water mites of the genus *Atractides* Koch, 1837 (Acari: Parasitengona: Hygrobatidae) in the western Palaearctic region: a revision. *Zoological Journal of the Linnean Society* **138**, 141–378.
- GERECKE, R. (ed.), GLEDHILL, T., PEŠIĆ, V. & SMIT, H., 2016: Chelicerata: Acari III. In GERECKE R. (ed.) *Süßwasserfauna von Mitteleuropa 7/2-3*. Spektrum, Heidelberg, pp 1–417.
- GERECKE, R., MARTIN, P. & GLEDHILL, T., 2018: Water mites (Acari: Parasitengona: Hydrachnidia) as inhabitants of groundwater-influenced habitats - considerations following an update of Limnofauna Europaea. *Limnologica* **69**, 81–93.
- GLAZIER, D.S., 2009: Springs. In LIKENS, G.E. (ed.) *Encyclopedia of Inland Waters*. Elsevier, Oxford, Vol.1, pp 734–755.
- GOLDSCHMIDT, T., 2016: Water mites (Acari, Hydrachnidia): powerful but widely neglected bioindicators – a review. *Neotropical Biodiversity* **2**, 12–25.
- GOTTSTEIN MATOČEC, S., KUZMAN, A., KEROVEC, M., 2006: Life History Traits of the Grass Shrimp *Palaemonetes antennarius* (Decapoda, Palaemonidae) in the Delta of the Neretva River, Croatia. *Limnologica* **36**(1), 42–53.
- ILLIES, J., 1978: *Limnofauna Europaea*. Gustav Fischer Verlag, Stuttgart and New York, pp 1–532.
- JASPRICA, N. & HAFNER, D., 2005: Taxonomic composition and seasonality of diatoms in three Dinaric karstic lakes in Croatia. *Limnologica* **35**, 304–319.
- KRANTZ, G.W., & WALTER, D.E., 2009: *A manual of Acarology*. Texas Tech University Press, Lubbock (TX), third edition, pp 1–807.
- KREŠIĆ, N., 2010: Chapter 2 - Types and classifications of springs. In KREŠIĆ, N. & STEVANOVIĆ, Z. (eds.) *Groundwater Hydrology of Springs. Engineering, Theory, Management, and Sustainability*. Elsevier Inc., Burlington, Oxford. pp 31–85.
- LATTINGER, R. 1988: *Ekološka diferenciranost faune podzemnih voda Medvednice*. [Ecological differentiation of the groundwater fauna of Medvednica] – Doctoral thesis (in Croatian), University of Zagreb, Zagreb, pp 1–448.
- LOBOREC, J., KAPELJ, S., DOGANČIĆ, D. & ŠIRONČIĆ, A.P., 2015: Assessment of Groundwater Vulnerability in Croatian Karstic Aquifer in Jadro and Žrnovnica Springs Catchment Area. In ANDREO, B., CARRASCO, F., DURÁN, J., JIMÉNEZ, P. & LAMOREAUX, J. (eds) *Hydrogeological and Environmental Investigations in Karst Systems*. Environmental Earth Sciences, vol 1. Springer, Berlin, Heidelberg, pp 397–405.
- MARTIN, P., DABERT, M., & DABERT, J., 2010: Molecular evidence for species separation in the water mite *Hygrobatas nigromaculatus* Lebert, 1879 (Acari, Hydrachnidia): evolutionary consequences of the loss of larval parasitism. – *Aquatic Sciences* **72**, 347–360.
- MATONIČKIN, I. & PAVLETIĆ, Z., 1959: Životne zajednice na sedrenim slapovima rijeke Une i na brzacima pritoke Unca [Biocenosis on the travertine cataracts in the river Una and in the rapids of the affluent Unac]. *Musei Macedonici Scientiarum naturalium* **2**, 78–99
- MICCOLI, F.P., LOMBARDO, P. & CICOLANI, B., 2013: Indicator value of lotic water mites (Acari: Hydrachnidia) and their use in macroinvertebrate-based indices for water quality assessment purposes. *Knowledge and Management of Aquatic Ecosystems* **8**, 1–28.
- PEŠIĆ, V., 2002: New records of water mites (Acari, Actinedida) based on the material collected by T. Petkovski from Croatia, including a check-list of species recorded from Croatia. *Natura Croatica* **11**, 447–453.

- PEŠIĆ, V., SMIT, H., GERECKE, R. & DI SABATINO, A., 2010: The water mites (Acari: Hydrachnidia) of the Balkan peninsula, a revised survey with new records and descriptions of five new taxa. *Zootaxa* **2586**, 1–100.
- PEŠIĆ, V., ASADI, M., CIMPEAN, M., DABERT, M., ESEN, Y., GERECKE, R., MARTIN, P., SAVIĆ, A., SMIT, H. & STUR, E., 2017: Six species in one: evidence of cryptic speciation in the *Hygrobatas fluviatilis* complex (Acariformes, Hydrachnidia, Hygrobatidae). *Systematic and Applied Acarology* **22**, 1327–1377.
- PEŠIĆ, V., BAŃKOWSKA, A., GOLDSCHMIDT, T., GRABOWSKI, M., MICHONKI, G. & ZAWAL, A., 2018: Supplement to the checklist of water mites (Acari: Hydrachnidia) from the Balkan peninsula. *Zootaxa* **4394**, 151–184.
- PEŠIĆ, V., SAVIĆ, A., JABLOŃSKA, A., MICHONSKI, G., GRABOWSKI, M., BAŃKOWSKA, A. & ZAWAL, A., 2019: Environmental factors affecting water mite assemblages along eucrenon-hypocrenon gradients in Mediterranean karstic springs. *Experimental and Applied Acarology* **77**, 471–486.
- POZOJEVIĆ, I., BRIGIĆ, A. & GOTTSTEIN, S., 2018: Water mite (Acari: Hydrachnidia) diversity and distribution in undisturbed Dinaric karst springs. *Experimental and Applied Acarology* **76**, 123–138.
- POZOJEVIĆ, I., TERNEJ, I., MIHALJEVIĆ, Z., GOTTSTEIN, S., VUČKOVIĆ, N., DORIĆ, V. & RUMIŠEK, M., 2018: Prey abundance supporting unusual water mite (Acari: Hydrachnidia) community in a sublacustrine spring and tributary river. *Acta Biologica* **25**, 69–75.
- POZOJEVIĆ, I., JURŠIĆ, L., VUČKOVIĆ, N., DORIĆ, V., GOTTSTEIN, S., TERNEJ, I. & MIHALJEVIĆ, Z., 2019: Is the spatial distribution of lentic water mite assemblages (Acari: Hydrachnidia) governed by prey availability? *Experimental and Applied Acarology* **77**(4), 487–510.
- PROCTOR, H.C., 2007: Aquatic mites in assessments of stream invertebrate diversity. In: MORALES-MALACARA, J.B., BEHAN-PELLETIER, V., UECKERMANN, E., et al., (eds) *Acarology XI: proceedings of the International Congress 8–13 September 2002, Mérida, Yucatán, México*. México, Instituto de Biología and Facultad de Ciencias, Universidad Nacional Autónoma de México, Sociedad Latinoamericana de Acarología, pp. 105–117.
- Šegota, T. & FILIPČIĆ, A., 2003: Köppenova podjela klima i hrvatsko nazivlje. *Geoadria* **8**, 17–37
- SMART, C. & WORTHINGTON, S.R.H., 2004: Springs. In GUNN, J. (ed.) *Encyclopedia of caves and karst science*. Taylor & Francis, London, pp. 1495–1505.
- TUZOVSKIJ, P.V., 1990: Key to deutonymphs of water mites. [Opređelitel' deutonymphs vodyanykh kleshchey]. Akademia Nauka UdSSR, Naukau, Moscow, pp. 1–238.
- WATERMITE.ORG, 2016. European Water Mite Research – A website informing on water mite research in Europe. <http://www.watermite.org/>.
- WIĘCEK, M., MARTIN, P. & GABKA, M., 2013: Distribution patterns and environmental correlates of water mites (Hydrachnidia, Acari) in peatland microhabitats. *Experimental and Applied Acarology* **61**, 147–160.
- ŽGANEC, K., GOTTSTEIN, S., GRABOWSKI, M., & PLATVOET, D., 2010: Distribution of the endemic amphipod species *Echinogammarus thoni* (Schäferna, 1922) in watercourses of the Balkan Peninsula. *Lauterbornia* **69**, 41–50.