

Simplex lattice design for optimization of the mass ratio of *Curcuma longa* L., *Curcuma zedoaria* (Christm.) Roscoe and *Curcuma aromatica* Salisb. to maximize curcuminoids content and antioxidant activity

CHAOWALIT MONTON*
PANUPONG CHUANCHOM
PUWADON POPANIT
SUKANYA SETTHARAKSA
PATHAMAPORN PATHOMPAK

*Drug and Herbal Product Research
and Development Center
College of Pharmacy, Rangsit University,
Pathum Thani 12000, Thailand*

The aim of this work was to optimize the mass ratio of three *Curcuma* plants' rhizomes to obtain the highest curcuminoid content and antioxidant activity using the simplex lattice design. The selected *Curcuma* plants were *C. longa*, *C. zedoaria* and *C. aromatica*. The simplex lattice design was applied in the work. The individual curcuminoids (curcumin, demethoxycurcumin and bis-demethoxycurcumin) and total curcuminoid content were determined using high-performance liquid chromatography. *Curcuma longa* alone provided the highest content of bis-demethoxycurcumin and demethoxycurcumin. A mixture of *C. longa* and *C. aromatica* in the mass ratio of 72:28 % provided the highest curcumin content. The results showed that *C. longa* alone exhibited the highest antioxidant activity.

Keywords: *Curcuma*, curcuminoids, HPLC, simplex lattice design, optimization, antioxidant activity

Accepted August 15, 2020
Published online August 26, 2020

Curcuma longa L., *Curcuma zedoaria* (Christm.) Roscoe and *Curcuma aromatica* Salisb. are plants belonging to the Zingiberaceae family. The most extensively used parts of these three plants are the rhizomes. According to the *Thai Herbal Pharmacopoeia* (1), *C. longa* is used as stomachic, carminative, coloring agent and adstringent, whereas *C. zedoaria* is used as stomachic, antidiarrheal and emmenagogue. *C. longa* has several biological and pharmacological activities such as antioxidant, antiinflammatory, antiangiogenic and antiapoptotic (2). Additionally, it also exhibits antihyperlipidemic, anti-Alzheimer, antidepressant, antidiabetic, radioprotective and antimicrobial activities (3). *C. aromatica* exhibits antiinflammatory, wound healing, antitumor, anticancer, mosquito repellent, antiplatelet, antitussive, antioxidant, antimelanogenic and antinephrotoxic activities (4).

All three plants exhibit antioxidant activity, that can protect the skin. Curcuminoids, including curcumin (CUR), demethoxycurcumin (DMC), and bis-demethoxycurcumin (BDMC), are the major chemical constituents in their rhizomes (5). Among these three curcuminoids, CUR is the most abundant and its mechanism of antioxidant activity has been described (6, 7). CUR can defend biomembranes from peroxidative damage by scav-

*Correspondence; e-mail: chaowalit@rsu.ac.th

enging the reactive free radicals. CUR can degrade into several compounds, *i.e.*, *trans*-6-(4'-hydroxy-3'-methoxyphenyl)-2,4-dioxo-5-hexanal, ferulic aldehyde, ferulic acid, ferruloyl methane and vanillin (8). Ferulic acid and vanillin exhibit antioxidant activity as well (9). Another work has also proven that CUR reveals a good antioxidant activity in several *in vitro* assays (10). In addition, DMC and BDMC also exhibit antioxidant activity (11).

The authors attempt is to distinguish the effect of the mass ratio of *C. longa*, *C. zedoaria* and *C. aromatica* rhizome powder on the content of curcuminoids and antioxidant activity, to further develop as an ingredient in the skincare formulations, using the simplex lattice design.

EXPERIMENTAL

Materials

Three standard curcuminoids (purity $\geq 99\%$), *i.e.*, BDMC, DMC and CUR were purchased from Chengdu Biopurify Phytochemicals Ltd., China. DPPH (2,2-diphenyl-1-picrylhydrazyl) was purchased from Sigma-Aldrich Inc., USA. Absolute ethanol was purchased from QR&C, New Zealand. Acetonitrile (HPLC grade) was purchased from Honeywell-Burdick & Jackson, USA. Acetic acid (AR grade) was purchased from Carlo-Erba, France.

Plant samples and extraction procedure

The rhizomes of *C. longa*, *C. zedoaria* and *C. aromatica* were collected from Buached District, Surin Province, Thailand. *C. longa* and *C. aromatica* were collected in November 2017, while *C. zedoaria* was collected in January 2018. The voucher specimens CM-

Table I. Coded and actual value of the model conditions obtained from simplex lattice design

Condition	Coded value			Actual value (mass of dried rhizome powder, g)		
	x_1	x_2	x_3	<i>C. longa</i>	<i>C. zedoaria</i>	<i>C. aromatica</i>
1	1	0	0	30	0	0
2	0	1	0	0	30	0
3	0	0	1	0	0	30
4	0.5	0.5	0	15	15	0
5	0.5	0	0.5	15	0	15
6	0	0.5	0.5	0	15	15
7	0.67	0.17	0.17	20	5	5
8	0.17	0.67	0.17	5	20	5
9	0.17	0.17	0.67	5	5	20
10	0.33	0.33	0.33	10	10	10
11	0.33	0.33	0.33	10	10	10
12	0.33	0.33	0.33	10	10	10

CL001-1-11-2017, CM-CZ001-1-01-2018, and CM-CA002-2-11-2017 were coded for *C. longa*, *C. zedoaria* and *C. aromatica*, resp. (Drug and Herbal Product Research and Development Center, College of Pharmacy, Rangsit University).

Rhizomes of each plant were cleaned, sliced and sun-dried for 3 days. The dried rhizomes were pulverized and passed through a 0.841-mm (20-mesh) sieve. The 30 g of rhizome powder was extracted in an ultrasonic bath for 20 min at different mass ratios (Table I) using 100 mL of absolute ethanol. The extract was separated from the marc after vacuum filtration. The extraction was performed in triplicate; three filtrates were pooled and the solvent was removed by a rotary evaporator.

High-performance liquid chromatography (HPLC) conditions

HPLC method, which was already reported, has been adapted and preliminarily validated (13, 19). Agilent 1260 Infinity HPLC instrument (Agilent Technologies, USA) was used in the analysis. ACE Generix column, 150×4.6 mm, 5 µm (Advanced Chromatography Technologies Ltd, UK) was used. An isocratic system composed of acetonitrile/acetic acid (1 %) in a volume ratio of 55:45, was used at a flow rate of 1 mL min⁻¹. The temperature of the column was set at 30 °C. The injection volume was 10 µL. The response was detected at 425 nm by a photodiode array detector.

HPLC method preliminary validation

HPLC method validation was done according to ICH guidelines (12). The five topics, including linearity and range, selectivity, the limit of detection (*LOD*) and limit of quantitation (*LOQ*), precision and accuracy were demonstrated.

Linearity and range. – A stock solution of each BDMC, DMC and CUR in the concentration of 1000 µg mL⁻¹ was prepared. Methanol was used as the solvent. The mixtures of three standard curcuminoids for each standard were prepared from the stock solutions at five concentration levels (*i.e.*, 10, 25, 50, 75, and 100 µg mL⁻¹). Two sets of mixtures of three standard curcuminoids were prepared. They were filtered using a nylon syringe filter and injected into the HPLC instrument. Each concentration was analyzed in triplicate. The calibration curves for each BDMC, DMC and CUR were constructed. The linear equation, coefficient of determination (*R*²), and range were reported.

Selectivity. – Selectivity was evaluated based on the selectivity ratio and chromatographic resolution. The selectivity ratio was calculated from the capacity factor values of the two peaks.

LOD and LOQ. – The standard solution was diluted with methanol. The *LOD* and *LOQ* were estimated based on the signal-to-noise ratio of 10:1 and 3:1, resp.

Precision. – Three concentration levels of each standard curcuminoid (25, 50, and 75 µg mL⁻¹) were analyzed. The analysis, done within-a-day and on three days, was used to calculate the percent relative standard deviation (RSD). These data were reported as intra-day and inter-day precision, resp.

Accuracy. – The individual three concentration levels of each standard curcuminoids (25, 50 and 75 µg mL⁻¹) were added to the extract (condition 10) solution. Each concentration level was analyzed in triplicate and the accuracy of each concentration level was reported.

Analysis of curcuminoids content

The plant extract was diluted with methanol to a concentration of 200–500 $\mu\text{g mL}^{-1}$. It was filtered using a nylon syringe filter and injected into the HPLC instrument. Each sample was prepared in two sets ($n = 2$) and each set was evaluated in triplicate. The content of each curcuminoid was calculated from the calibration curve and total curcuminoid content was obtained from the summation of three curcuminoids.

Determination of antioxidant activity

Antioxidant activity was determined using the DPPH radical scavenging assay. Extract volumes of 50–1000 $\mu\text{g mL}^{-1}$ were prepared using absolute ethanol as diluent. CUR, DMC, and BDMC were also tested, with a concentration range of 1–1000 $\mu\text{g mL}^{-1}$. The sample (100 μL) was pipetted into a 96-well plate ($n = 3$). 100 μL aliquot of the 200 $\mu\text{mol L}^{-1}$ DPPH ethanolic solution was then added to each well. They were incubated for 30 min in the dark at room temperature. The absorbance of the test sample was measured at 517 nm using a microplate reader (Benchmark Plus, Bio-Rad Laboratories, USA). An equal volume of absolute ethanol and DPPH ethanolic solution was used as a control. The half-maximal inhibitory concentration IC_{50} was reported.

Optimization

A simplex lattice design was used in this work. Design-Expert[®] version 11 was used for the optimization process. Three factors, *i.e.*, the mass ratios of *C. longa*, *C. zedoaria*, and *C. aromatica* rhizome powder were coded as x_1 , x_2 , and x_3 , resp. The value of each factor was between 0 and 100 % and the summation of all factors equals 100 %. Five responses, *i.e.*, BDMC content (y_1), DMC content (y_2), CUR content (y_3), total curcuminoid content (y_4), and IC_{50} value obtained from DPPH radical scavenging assay (y_5) were determined. The contour plot of each response was created. The plots between predicted values *vs.* actual values and residual *vs.* run were produced to confirm the stability and reliability of the estimation. The statistical data, *p*-value of the model and a lack of fit of each response were reported. The desirability function was used to select the optimum conditions – the optimum mass ratio of three *Curcuma* rhizome powders provided the highest curcuminoid content and maximal antioxidant activity (or minimal IC_{50} value).

RESULTS AND DISCUSSION

The method for analysis of curcuminoids in the rhizome of *C. longa*, *C. zedoaria*, and *C. aromatica*, and their mixtures was preliminary validated. The linearity parameters (linear equation and R^2), range and LOD, together with precision and accuracy data, are given in Table II. RSD values of intra- and inter-day precision were 0.1–0.2 and 0.2–3.6 %, resp. In addition, accuracy was 93.4–99.8 % for BDMC, 96.9–102.2 % for DMC, and 99.0–99.2 % for CUR. Fig. 1 shows that BDMC, DMC and CUR were eluted at the retention times of 3.95, 4.30 and 4.70 min, resp. The selectivity ratio of DMC *vs.* BDMC and CUR *vs.* DMC was 1.11 and 1.14, resp., however, for CUR/BDMC it was 1.27. All validation data indicated the acceptable analytical performances of the method.

Table II. Analytical performances of HPLC method

Linear regression parameter					
	Analyte				
	BDMC	DMC	CUR		
Range ($\mu\text{g mL}^{-1}$)	10–100	10–100	10–100		
Linear equation	$y = 161525x - 50686$	$y = 185753x + 15028$	$y = 226712x + 63963$		
R^2	0.9992	0.9995	0.9993		
LOD (ng mL^{-1})	2.5	5.0	5.0		
Precision and accuracy					
Analyte	Concentration ($\mu\text{g mL}^{-1}$)	Precision (RSD, %)		Added concentration ($\mu\text{g mL}^{-1}$)	Accuracy (%)
		Intra-day	Inter-day		
BDMC	25	0.2	2.2	25	93.4
	50	0.2	0.2	50	99.8
	75	0.2	0.2	75	98.5
DMC	25	0.2	3.6	25	96.9
	50	0.1	1.6	50	97.8
	75	0.1	1.2	75	102.2
CUR	25	0.1	0.7	25	99.2
	50	0.1	2.3	50	99.0
	75	0.1	2.5	75	99.2

Chromatographic parameters of the HPLC method are given in Table III.

Both individual and total curcuminoid content in rhizome powders mixture were estimated to observe their variation when different mass ratios of *C. longa*, *C. zedoaria*, and *C. aromatica* were applied. In this case, curcuminoids present in the extract were analyzed and calculated as curcuminoid content in the raw plant material (dried powder of rhizome). Fig. 1 shows the HPLC chromatogram of standard curcuminoids and extracts obtained under the same conditions as shown in Table I. *C. longa* alone (condition 1) mostly contained CUR followed by BDMC and DMC, resp. (Fig. 1a). It contained 1.06 % CUR, 0.63 % DMC and 0.91 % BDMC; total curcuminoids content roughly 2.6 %. Jayaprakasha *et al.* (14) showed *C. longa* to be composed of 1.01–5.65 % CUR, 0.83–3.36 % DMC and 0.42–2.16 % BDMC. In addition, total curcuminoid content was 2.34–9.18 % (14). The slight difference in the content of the curcuminoids was reported by Ali *et al.* (15) who stated 2.1 % CUR, 0.46 % DMC and 0.1 % BDM, whereas Osorio-Tobón *et al.* (16) showed 10.2 % CUR, 5.9–7.0 % DMC and 2.1–6.1 % BDMC in *C. longa*. *C. longa* powder filled in hard gelatin capsule contained high curcuminoids content as previously reported: 6.61–7.67 % CUR, 2.76–3.33 % DMC and 2.64–3.47 % BDMC (17).

C. zedoaria (condition 2) and *C. aromatica* (condition 3) mostly contained DMC (Figs. 1b,c). *C. zedoaria* contained 0.23 % DMC, 0.06 % BDMC and 0.04 % CUR (total curcuminoids




Fig. 1. HPLC chromatograms of: a) mixed standard curcuminoids ($50 \mu\text{g mL}^{-1}$) and extract obtained from: b) condition 1 ($200 \mu\text{g mL}^{-1}$), c) condition 2 ($500 \mu\text{g mL}^{-1}$), d) condition 3 ($500 \mu\text{g mL}^{-1}$) and e) condition 10 ($500 \mu\text{g mL}^{-1}$).

Table III. Chromatographic parameters of HPLC method

Chromatographic parameter	Analyte		
	BDMC	DMC	CUR
t_R (min) ^a	3.95 ± 0.01	4.30 ± 0.00 ₁	4.70 ± 0.00 ₃
Capacity factor (k') ^b	2.07 ± 0.06	2.30 ± 0.00 ₂	2.63 ± 0.06
Tailing factor (T_f) ^b	1.20 ± 0.17	1.37 ± 0.15	1.23 ± 0.25
Number of theoretical plates (N) ^b	2196 ± 94	2252 ± 255	2369 ± 74
Resolution (R_s) ^b	1.50 ± 0.26	1.50 ± 0.26	1.53 ± 0.25
Selectivity (α) ^b	1.11 ± 0.03		1.14 ± 0.03
	–	1.11 ± 0.03	1.27 ± 0.03

Mean ± SD. ^a $n = 3$, ^b $n = 6$.

0.3 %). *C. aromatica* contained 0.43 % DMC, 0.09 % BDMC and 0.08 % CUR (total curcuminoids content 0.6 %), with two times higher total curcuminoids compared with *C. zedoaria*, whereas *C. longa* had approx. 8 and 4 times, resp., higher content of total curcuminoids than the other two species. Paramapojn and Gritsanapan (18) reported that ethanolic extract of *C. zedoaria* rhizome contained 2.73 % CUR, 7.37 % DMC and 1.40 % BDMC, which was close to our results: 1.02 % CUR, 5.98 % DMC and 1.63 % BDMC. Other work of the same colleagues showed that 0.50–0.73 % CUR, 0.23–1.43 % DMC, 0.12–0.44 % BDMC were found in *C. zedoaria* powder (19). *C. zedoaria* consisted of 0–1.5 % CUR as reported by Morishita *et al.* (20). However, no CUR, DMC and BDMC in *C. zedoaria* methanolic extract were reported in a publication by Tohda *et al.* (21) but 0.11 % CUR with no DMC and no BDMC in *C. aromatica* methanolic extract. Morishita *et al.* (20) reported that *C. aromatica* contained 0.05–0.1 % CUR estimated on a dry mass basis. However, sometimes *C. aromatica* may contain more CUR than DMC (22, 23). Fig. 1e shows the mixture of three plants rhizome powders in equal mass ratios (condition 10) which mostly had CUR followed by DMC and BDMC.

Our work documented that CUR exhibited the most potent antioxidant activity compared to DMC and BDMC: IC_{50} values of CUR, DMC and BDMC were 38, 97, and 1,375 $\mu\text{mol L}^{-1}$ (namely, 17.83 ± 1.59 , 32.73 ± 5.07 and $424.10 \pm 20.40 \mu\text{g mL}^{-1}$), resp. All three curcuminoids showed good antioxidant activity. The presence of the methoxy group in CUR could increase antioxidant activity (9), so CUR showed the best antioxidant activity compared with DMC and BDMC. The IC_{50} value of the standard curcuminoids varied in the literature. Abas *et al.* (24) reported the IC_{50} values of CUR, DMC, and BDMC of 31.8, 92.5, and 104.4 $\mu\text{mol L}^{-1}$ (or 11.7, 31.3, and 32.2 $\mu\text{g mL}^{-1}$) (24) with a largely different IC_{50} value of BDMC compared with our result. The low IC_{50} values were reported for CUR by Borra *et al.* (25) of 2.9 $\mu\text{mol L}^{-1}$ (or 1.1 $\mu\text{g mL}^{-1}$) and Sökmen and Khan (26) of 22.8 $\mu\text{mol L}^{-1}$ (or 8.4 $\mu\text{g mL}^{-1}$). On the other hand, high IC_{50} values were also reported: Ak and Gulcin (10) reported that CUR had IC_{50} of 94.6 $\mu\text{mol L}^{-1}$ (or 34.9 $\mu\text{g mL}^{-1}$).

Regarding the plant extracts, extract obtained from the condition 1, which contained *C. longa* alone, exhibited the lowest IC_{50} value ($33.83 \pm 14.05 \mu\text{g mL}^{-1}$, equivalent to $26.01 \pm 1.24 \mu\text{g mL}^{-1}$ total curcuminoids), indicating the most potent antioxidant activity. The IC_{50} of *C. longa* extract was similar to the results of the previous work ($33.5 \mu\text{g mL}^{-1}$) (27). However, low IC_{50} of *C. longa* extract of 9.7 $\mu\text{g mL}^{-1}$ was reported by Zaeoung *et al.* (28). Further-




Fig. 2. Contour plots of the model conditions of: a) BDMC content, b) DMC content, c) CUR content, d) total curcuminoid content, in the plant powder mixtures, and e) IC_{50} value of the extract obtained from the plant powder mixtures.

more, Tanvir *et al.* (29) reported that IC_{50} values of *C. longa* aqueous and ethanolic extracts ranged 5.31–16.55 and 1.08–3.03 $\mu\text{g mL}^{-1}$, resp. In this work, we found that *C. zedoaria* extract obtained from condition 2 and *C. aromatica* extract obtained from condition 3 had IC_{50} values of 54.63 ± 3.37 and 138.72 ± 4.76 $\mu\text{g mL}^{-1}$, resp., which were equivalent to 8.63 ± 0.96 and 7.56 ± 0.17 $\mu\text{g mL}^{-1}$ total curcuminoids, resp. The other samples (extracts obtained from conditions 4–12) contained three plants in different ratios, IC_{50} values ranged from approximately 40 to 200 $\mu\text{g mL}^{-1}$. Our result was different from the previous reports. Souria *et al.* (30) showed that *C. zedoaria* extract had an IC_{50} value of 10.60 $\mu\text{g mL}^{-1}$, whereas Nahak and Sahu (31) reported 40 $\mu\text{g mL}^{-1}$. They also reported an IC_{50} value of *C. aromatica* extract of 120 $\mu\text{g mL}^{-1}$, which was close to the result of Srividya *et al.* (32) of 132.5 $\mu\text{g mL}^{-1}$.

Fig. 2 shows the contour plots of the model conditions of curcuminoids content in dried rhizome powder and IC_{50} value of the extract. The maximum BDMC, DMC and total curcuminoid content in dried rhizome powder were found when *C. longa* was used alone, whereas the minimum was found when *C. zedoaria* was used alone. In the case of CUR in rhizome powder, the maximum value was achieved when *C. longa* was mixed with *C. aromatica* in the mass ratio of 72:28, while the minimum value was found when *C. zedoaria* was mixed with *C. aromatica* in the mass ratio of 89:11. The maximum IC_{50} value of the extract




Fig. 3. Predicted *vs.* actual plots (left) and residual *vs.* run plots (right) of: a) BDMC content, b) DMC content, c) CUR content, d) total curcuminoids content, and e) IC_{50} value.

was found when *C. aromatica* was used alone and the minimum IC_{50} value when *C. longa* was used alone. The results indicated that the most potent antioxidant activity was found when *C. longa* was used alone.

The predicted values obtained from the Design-Expert® were based on the equations shown below. Fig. 3 shows the predicted *vs.* actual value plots and the internally Studentized residuals *vs.* run number plots of the model condition of BDMC, DMC, CUR and total curcuminoids content in rhizome powder and IC_{50} value of the extract. The R^2 between the predicted and actual values of model conditions of BDMC, DMC, CUR and total curcuminoids content in rhizome powder, and IC_{50} value of the extract were 0.8265, 0.4265, 0.9183, 0.7136 and 0.3392, resp. The moderate to high correlation was found for BDMC, CUR and total curcuminoid content, whereas low correlation was found in DMC content and IC_{50} value. The internally Studentized residuals *vs.* run number plots displayed that the data was within the red borderline, indicating data distribution within a 95 % confidence interval (CI). These results could assure the stability of the prediction. The data in Fig. 3 could approve the reliability and stability of the prediction by computer software (33, 34). Table IV shows the *p*-values of the model and the lack of fit of various responses. The *p*-values of experimental models of y_1 , y_3 , and y_4 were less than 0.05 indicating that model terms were significant, whereas the *p*-values of models of y_2 and y_5 were not significant. The lack of fit *p*-values of all responses was more than 0.05 implying that the lack of fit was not significant relative to the pure error. So the experimental design was not significantly affected by the error or coincidence.

Table IV. Statistical data (*p*-value) of the model and lack of fit of each response

Response	<i>p</i> -value	
	Model	Lack of fit
y_1	0.0004*	0.6970
y_2	0.0820	0.7042
y_3	0.0033*	0.8333
y_4	0.0036*	0.6218
y_5	0.1550	0.7096

* Statistical significance

Table V. Predicted value, actual value, percent error, and 95% confidence interval of the prediction

Response	Predicted value	Actual value	Error (%)	95 % CI (lower-upper)
y_1 (% m/m)	1.07	0.91±0.10	-17.6	0.84–1.29
y_2 (% m/m)	0.82	0.63±0.03	-30.2	0.56–1.07
y_3 (% m/m)	1.05	1.06±0.01	0.9	0.67–1.44
y_4 (% m/m)	3.30	2.60±0.12	-26.9	2.43–4.17
y_5 (µg mL ⁻¹)	41.81	33.83±6.60	-23.6	0.00–83.62

CI – confidence interval

The equations used for the estimation of each response are shown below:

$$y_1 = 1.07x_1 + 0.12x_2 + 0.19x_3$$

$$y_2 = 0.82x_1 + 0.35x_2 + 0.57x_3$$

$$y_3 = 1.05x_1 + 0.06x_2 + 0.10x_3 + 1.28x_1x_2 + 2.14x_1x_3 - 0.34x_2x_3$$

$$y_4 = 3.30x_1 + 0.62x_2 + 1.05x_3$$

$$y_5 = 41.81x_1 + 64.08x_2 + 172.34x_3$$

The desirability function could be used for selecting the optimum condition. In this case, the optimum condition was the optimum mass ratio of three *Curcuma* rhizome powders providing the highest curcuminoid content and the maximal antioxidant activity (or minimal IC_{50} value). The desirability value equal to 1 should indicate a fully desired response (35). The desirability value, obtained from the computer software Design-Expert®, showed that the maximum desirability value of 0.978 was reached when *C. longa* was used alone. So, the optimum condition was approached when *C. longa* was used alone. The predicted value, actual value, percent error and range of 95 % CI of the prediction are shown in Table V. When the prediction value was compared with the actual value, the error of the prediction was found approximately 30 % – the lowest percent error was found in y_3 of 0.9 %, and the highest percent error was found in y_2 of –30.2 %. However, the actual value was within the lower and upper limit of the 95 % confidence interval, indicating that the data obtained from the computer software were moderately accurate.

CONCLUSIONS

This work used simplex lattice design to optimize the mass ratio of three *Curcuma* plants' rhizomes to obtain the high curcuminoid content and antioxidant activity. Three plants, *i.e.*, *C. longa*, *C. zedoaria* and *C. aromatica*, were included in the work. The CUR was the major compound found in *C. longa*, whereas DMC was the major compound found in *C. zedoaria* and *C. aromatica*. *C. longa* alone gave the highest content of BDMC and DMC. *C. longa* mixed with *C. aromatica* gave the highest CUR content. The lowest IC_{50} value, obtained from DPPH radical scavenging assay, was found when *C. longa* was used alone. From the optimization process, *C. longa* alone reveals the highest total curcuminoid content and the most potent antioxidant activity compared with *C. zedoaria*, *C. aromatica* and their admixtures. In conclusion, the simplex lattice design could be used to optimize the mass ratio of three *Curcuma* plants rhizomes to obtain the maximized individual and total curcuminoid content as well as antioxidant activity with moderate accuracy.

Acknowledgements. – The authors would like to thank the College of Pharmacy, Rangsit University for facilities support. Three plant samples were authenticated by plant taxonomist, Ajarn Nirun Vipunngun, Department of Pharmacognosy, College of Pharmacy, Rangsit University.

REFERENCES

1. Khamin Chan, in *Thai Herbal Pharmacopoeia 2017*, Department of Medical Sciences, the Agricultural Co-operative Federation of Thailand, Bangkok 2017, pp. 135–142.

2. R. Agarwal, S. K. Gupta, S. Srivastava, P. Agarwal and S. S. Agrawal, Therapeutic potential of *Curcuma longa*, the golden spice of India, in drug discovery for ophthalmic diseases, *Expert Opin. Drug Discov.* **4** (2009) 147–158; <https://doi.org/10.1517/13543770802668117>
3. V. Soleimani, A. Sahebkar and H. Hosseinzadeh, Turmeric (*Curcuma longa*) and its major constituent (curcumin) as nontoxic and safe substances: Review, *Phytother. Res.* **32** (2018) 985–995; <https://doi.org/10.1002/ptr.6054>
4. A. Sikha, A. Harini and L. Hegde Prakash, Pharmacological activities of wild turmeric (*Curcuma aromatica* Salisb): a review, *J. Pharmacogn. Phytochem.* **3** (2015) 1–4.
5. E. Mudge, M. Chan, S. Venkataraman and P. N. Brown, Curcuminoids in turmeric roots and supplements: Method optimization and validation, *Food Anal. Meth.* **9** (2016) 1428–1435; <https://doi.org/10.1007/s12161-015-0326-0>
6. T. Masuda, K. Hidaka, A. Shinohara, T. Maekawa, Y. Takeda and H. Yamaguchi, Chemical studies on antioxidant mechanism of curcuminoid: Analysis of radical reaction products from curcumin, *J. Agric. Food Chem.* **47** (1999) 71–77; <https://doi.org/10.1021/jf9805348>
7. T. Masuda, T. Maekawa, K. Hidaka, H. Bando, Y. Takeda and H. Yamaguchi, Chemical studies on antioxidant mechanism of curcumin: Analysis of oxidative coupling products from curcumin and linoleate, *J. Agric. Food Chem.* **49** (2001) 2539–2547; <https://doi.org/10.1021/jf001442x>
8. K. I. Priyadarsini, The chemistry of curcumin: from extraction to therapeutic agent, *Molecules* **19** (2014) 20091–20112; <https://doi.org/10.3390/molecules191220091>
9. V. P. Menon and A. R. Sudheer, Antioxidant and anti-inflammatory properties of curcumin, *Adv. Exp. Med. Biol.* **595** (2007) 105–125; https://doi.org/10.1007/978-0-387-46401-5_3
10. T. Ak and I. Gulcin, Antioxidant and radical scavenging properties of curcumin, *Chem. Biol. Interact.* **174** (2008) 27–37; <https://doi.org/10.1016/j.cbi.2008.05.003>
11. G. K. Jayaprakasha, L. Jaganmohan Rao and K. K. Sakariah, Antioxidant activities of curcumin, demethoxycurcumin and bisdemethoxycurcumin, *Food Chem.* **98** (2006) 720–724; <https://doi.org/10.1016/j.foodchem.2005.06.037>
12. International Conference on Harmonisation of Technical Requirements for Registration of Pharmaceuticals for Human Use, *ICH Harmonised Tripartite Guideline, Validation of Analytical Procedures: Text and Methodology Q2(R1)*, Current Step 4 version, ICH, Geneva, November 2005; <https://database.ich.org/sites/default/files/Q2%28R1%29%20Guideline.pdf>; last access date May 2, 2019
13. C. Monton, C. Luprasong and L. Charoenchai, Convection combined microwave drying affect quality of volatile oil compositions and quantity of curcuminoids of turmeric raw material, *Rev. Bras. Farmacogn.* **29** (2019) 434–440; <https://doi.org/10.1016/j.bjp.2019.04.006>
14. G. K. Jayaprakasha, L. Jagan Mohan Rao and K. K. Sakariah, Improved HPLC method for the determination of curcumin, demethoxycurcumin, and bisdemethoxycurcumin, *J. Agric. Food Chem.* **50** (2002) 3668–3672; <https://doi.org/10.1021/jf025506a>
15. I. Ali, A. Haque and K. Saleem, Separation and identification of curcuminoids in turmeric powder by HPLC using phenyl column, *Anal. Meth.* **6** (2014) 2526–2536; <https://doi.org/10.1039/c3ay41987h>
16. J. F. Osorio-Tobón, P. I. Carvalho, G. F. Barbero, G. C. Nogueira, M. A. Rostagno and M. A. Meireles, Fast analysis of curcuminoids from turmeric (*Curcuma longa* L.) by high-performance liquid chromatography using a fused-core column, *Food Chem.* **200** (2016) 167–174; <https://doi.org/10.1016/j.foodchem.2016.01.021>
17. C. Monton, L. Charoenchai, J. Suksaeree and L. Sueree, Quantitation of curcuminoid contents, dissolution profile, and volatile oil content of turmeric capsules produced at some secondary government hospitals, *J. Food Drug Anal.* **24** (2016) 493–499; <https://doi.org/10.1016/j.jfda.2016.01.007>
18. S. Paramapojn and W. Gritsanapan, Variation of curcuminoids in ethanolic extract of *Curcuma zedoaria* rhizomes in Thailand by HPLC, *Planta Med.* **73** (2007) P_227; <https://doi.org/10.1055/s-2007-987008>

19. S. Paramapojn and W. Gritsanapan, *Quantitative Analysis of Curcuminoids in Curcuma zedoaria Rhizomes in Thailand by HPLC Method*, in *International Workshop on Medicinal and Aromatic Plants* (Eds. N. Chomchalow and V. Chantrasm), Vol. 1, Chiang Mai (Thailand) 2008; *ISHS Acta Hort.* **786** (2008) 169–174; <https://doi.org/10.17660/ActaHortic.2008.786.18>
20. T. Morishita, H. Yamaguchi, K. Degi, A. Shimizu and H. Nakagawa, Comparison of curcumin contents and antioxidative activities among turmeric species cultivated in Kanto area, *Jpn. J. Crop Sci.* **82** (2013) 56–62; <https://doi.org/10.1626/jcs.82.56>
21. C. Tohda, N. Nakayama, F. Hatanaka and K. Komatsu, Comparison of anti-inflammatory activities of six *Curcuma* rhizomes: A possible curcuminoid-independent pathway mediated by *Curcuma phaeocalis* extract, *Evid. Based Complement. Alternat. Med.* **3** (2006) 255–260; <https://doi.org/10.1093/ecam/nel008>
22. J. Zhang, S. Jinnai, R. Ikeda, M. Wada, S. Hayashida and K. Nakashima, A simple HPLC-fluorescence method for quantitation of curcuminoids and its application to turmeric products, *Anal. Sci.* **25** (2009) 385–388; <https://doi.org/10.2116/analsci.25.385>
23. J. Lee, Y. Jung, J. H. Shin, H. K. Kim, B. C. Moon, D. H. Ryu and G. S. Hwang, Secondary metabolite profiling of *Curcuma* species grown at different locations using GC/TOF and UPLC/Q-TOF MS, *Molecules* **19** (2014) 9535–9551; <https://doi.org/10.3390/molecules19079535>
24. F. Abas, L. S. Hui, S. Ahmad, J. Stanslas, D. A. Israf, K. Shaari and N. H. Lajis, Biological evaluation of curcumin and related diarylheptanoids, *Z. Naturforsch. C* **61** (2006) 625–631; <https://doi.org/10.1515/znc-2006-9-1002>
25. S. K. Borra, P. Gurumurthy, J. Mahendra, K. M. Jayamathi, C. N. Cherian and R. Chand, Antioxidant and free radical scavenging activity of curcumin determined by using different *in vitro* and *ex vivo* models, *J. Med. Plants Res.* **7** (2013) 2680–2690; <https://doi.org/10.5897/JMPR2013.5094>
26. M. Sökmen and M. Akram Khan, The antioxidant activity of some curcuminoids and chalcones, *Inflammopharmacology* **24** (2016) 81–86; <https://doi.org/10.1007/s10787-016-0264-5>
27. F. O. S. Camatari, K. H. Lopes, B. I. Valentim, J. A. Xavier, J. G. da Costa, A. E. G. Santana and M. O. F. Goulart, Antioxidant potential of flours from cereals, tubers, beans and seeds chemical profile of *Curcuma longa* flour, *J. Nutr. Food Sci.* **6** (2016) 1000483; <https://doi.org/10.4172/2155-9600.1000483>
28. S. Zaeoung, A. Plubrukarn and N. Keawpradub, Cytotoxic and free radical scavenging activities of Zingiberaceae rhizomes, *Songklanakarinn J. Sci. Technol.* **27** (2005) 799–812.
29. E. M. Tanvir, M. S. Hossen, M. F. Hossain, R. Afroz, S. H. Gan, M. I. Khalil and N. Karim, Antioxidant properties of popular turmeric (*Curcuma longa*) varieties from Bangladesh, *J. Food Qual.* **2017** (2017) 8471785; <https://doi.org/10.1155/2017/8471785>
30. E. Souria, G. Amin, H. Farsam, H. Jalalizadeh and S. Barezi, Screening of thirteen medicinal plant extracts for antioxidant activity, *Iran. J. Pharm. Res.* **7** (2008) 149–154; <https://doi.org/10.22037/IJPR.2010.758>
31. G. Nahak and R. K. Sahu, Evaluation of antioxidant activity in ethanolic extracts of five curcuma species, *Int. Res. J. Pharm.* **2** (2011) 243–248.
32. A. R. Srividya, P. Dhanabal, Parthkumar Bavadia, V. J. Vishnuvarthan and M. N. S. Kumar, Antioxidant and antidiabetic activity of *Curcuma aromatica*, *Int. J. Res. Ayurveda Pharm.* **3** (2012) 401–405.
33. S. Duangjit, L. M. Mehr, M. Kumpugdee-Vollrath and T. Ngawhirunpat, Role of simplex lattice statistical design in the formulation and optimization of microemulsions for transdermal delivery, *Biol. Pharm. Bull.* **37** (2014) 1948–1957; <https://doi.org/10.1248/bpb.b14-00549>
34. S. Duangjit, Y. Obata, H. Sano, S. Kikuchi, Y. Onuki, P. Opanasopit, T. Ngawhirunpat, Y. Maitani and K. Takayama, Menthosomes, novel ultradeformable vesicles for transdermal drug delivery: optimization and characterization, *Biol. Pharm. Bull.* **35** (2012) 1720–1728; <https://doi.org/10.1248/bpb.b12-00343>
35. M. A. Bezerra, R. E. Santelli, E. P. Oliveira, L. S. Villar and L. A. Escalera, Response surface methodology (RSM) as a tool for optimization in analytical chemistry, *Talanta* **76** (2008) 965–977; <https://doi.org/10.1016/j.talanta.2008.05.019>