

## LAPLACIAN COEFFICIENTS OF TREES

ALI GHALAVAND AND ALI REZA ASHRAFI

ABSTRACT. Let  $G$  be a simple and undirected graph with Laplacian polynomial  $\psi(G, \lambda) = \sum_{k=0}^n (-1)^{n-k} c_k(G) \lambda^k$ . In this paper, exact formulas for the coefficient  $c_{n-4}$  and the number of 4-matchings with respect to the Zagreb indices of a given tree are presented. The chemical trees with first through the fifteenth greatest  $c_{n-4}$ -values are also determined.

### 1. INTRODUCTION

A graph  $G$  consists of two sets  $V = V(G)$  and  $E = E(G)$ . The elements of  $V$  are called the *vertices* of  $G$  and the elements of  $E$  are *edges* of this graph. Each edge is a 2-element subset of vertices  $\{x, y\}$  which is denoted by  $xy$ . A *chemical graph* is a graph in which  $\Delta(G) \leq 4$ , where  $\Delta(G)$  is the maximum degree of vertices in  $G$  and a *tree* is a connected graph without cycles. The vertex degree of  $v \in V(G)$ ,  $deg_G(v)$ , is defined as the number of edges incident to  $v$  and  $N_G(v)$  denotes the set of all vertices adjacent to  $v$ . The distance between two vertices  $x, y \in V(G)$ ,  $d(x, y)$ , is defined as the number of edges in a shortest path connecting them. The summation of all such numbers is called the Wiener index of  $G$  denoted by  $W(G)$ .

For subset  $E'$  of  $E(G)$ , we denote the subgraph of  $G$  obtained by deleting the edges of  $E'$  by  $G - E'$ . If  $E' = \{uv\}$ , then the subgraph  $G - E'$  will be written as  $G - uv$  for short. In addition, for any two nonadjacent vertices  $x$  and  $y$  of  $G$ , let  $G + xy$  be the graph obtained from  $G$  by adding an  $xy$  edge. If two vertices  $x$  and  $y$  are adjacent then we write  $x \sim y$ . The *path* and *star* on  $n$ -vertices are denoted by  $P_n$  and  $S_n$ , respectively. The set of all  $n$ -vertex chemical trees is denoted by  $\mathcal{CT}(n)$ .

Suppose  $\mathcal{G}$  denotes the set of all graphs and  $G, H \in \mathcal{G}$ . If  $V(H) \subseteq V(G)$  and  $E(H) \subseteq E(G)$ , then we say that  $H$  is a *subgraph* of  $G$  and use the notation  $H \subseteq G$ . The number of subgraphs of  $G$  isomorphic to a fixed subgraph  $H$  is denoted by  $\eta(G, H)$ . It is easy to see that  $\eta(G, S_2) = m$ , the number of edges in  $G$ . The number of vertices of degree  $i$  in  $G$  will be denoted by  $n_i = n_i(G)$ .

---

2020 *Mathematics Subject Classification.* 05C31, 05C05.

*Key words and phrases.* Laplacian coefficient, matching, Zagreb index, tree.

It is easy to see that  $\sum_{i=1}^{\Delta(G)} n_i = |V(G)|$ . A map  $Top$  from  $\mathcal{G}$  into the set of all non-negative real numbers is called a *graph invariant* if  $G \cong H$  implies that  $Top(G) = Top(H)$ . *Topological indices* are graph invariants applicable in chemistry.

The graph invariants *Wiener index* [14], *first Zagreb index* and *second Zagreb index* [9], *forgotten topological index* [6] and the *first general Zagreb index* [16], are defined as:

$$\begin{aligned} W(G) &= \sum_{\{u,v\} \subset V(G)} d_G(u,v), \\ M_1(G) &= \sum_{v \in V(G)} deg_G(v)^2, \\ M_2(G) &= \sum_{uv \in E(G)} deg_G(u)deg_G(v), \\ F(G) &= \sum_{v \in V(G)} deg_G(v)^3 = \sum_{uv \in E(G)} [deg_G(u)^2 + deg_G(v)^2], \\ M_1^\alpha(G) &= \sum_{u \in V(G)} deg_G(u)^\alpha, \end{aligned}$$

respectively. Here,  $\alpha \neq 0, 1$  is an arbitrary real number. Furthermore, the first Zagreb index and the forgotten topological index are just the case of  $\alpha = 2, 3$  in the first general Zagreb index, respectively.

The first and second reformulated Zagreb indices of graphs were introduced by Milićević et al. [12]. These graph invariants are edge counterparts of the first and second Zagreb indices, respectively. These numbers can be defined as:

$$\begin{aligned} EM_1(G) &= \sum_{e \sim f} [deg_G(e) + deg_G(f)] = \sum_{e \in E(G)} deg_G(e)^2, \\ EM_2(G) &= \sum_{e \sim f} deg_G(e)deg_G(f). \end{aligned}$$

In this formulas, if  $e = uv$  then  $deg_G(e) = deg_G(u) + deg_G(v) - 2$ . Moreover,  $e \sim f$  means that the edges  $e$  and  $f$  are incident.

Suppose  $G$  is a simple graph with vertex set  $\{v_1, \dots, v_n\}$ . The adjacency matrix of  $G$  is an  $n \times n$  0-1 matrix  $A = (a_{ij})$  such that  $a_{ij}$  is one if and only if there is an edge connecting  $v_i$  and  $v_j$ . The degree matrix,  $D(G)$ , is a square matrix of order  $n$  whose its  $i^{th}$  diagonal entry is equal to  $deg_G(v_i)$  and whose off-diagonal elements are zero. The Laplacian matrix of  $G$  is defined as  $L(G) = D(G) - A(G)$ . The characteristic polynomial of the Laplacian matrix,  $\psi(G, \lambda) = \det(\lambda I_n - L(G))$ , is said to be the Laplacian polynomial of the graph  $G$ . In this paper we write this polynomial in the form of  $\psi(G, \lambda) = \sum_{k=0}^n (-1)^{n-k} c_k(G) \lambda^k$ . It is well-known that  $c_k(G) \geq 0$ , for all  $k$ .

Suppose  $G$  is a simple and undirected graph. The relationship between the coefficients of  $\psi(G, \lambda)$  and the structure of  $G$  was established many years ago by Kel'mans [3, p. 38]. He proved that  $c_k(G) = \sum_{F \in \mathcal{F}_k(G)} \gamma(F)$ , where  $F$  is a spanning forest and the summation goes over the set  $\mathcal{F}_k(G)$  of all spanning forests of  $G$ , possessing exactly  $k$  components and  $\gamma(F)$  is the product of the number of vertices of the components of  $F$ . If  $T$  is an  $n$ -vertex tree, then for  $k \geq 1$ , the elements of  $\mathcal{F}_k(T)$  can be obtained by deleting  $k - 1$  distinct edges from  $T$ . So, it is easy to see that,  $c_1(T) = n$ ,  $c_n(T) = 1$  and  $c_{n-1}(T) = 2(n - 1)$ . Yan et al. [15], proved that  $c_2(T) = W(T)$ . Oliveira et al. [13], obtained closed formulas for the coefficient  $c_{n-2}(T)$  and  $c_{n-3}(T)$  in terms of the number of vertices, the first Zagreb and forgotten indices as  $c_{n-2}(T) = 2n^2 - 5n + 3 - \frac{1}{2}M_1(T)$  and  $c_{n-3}(T) = \frac{1}{3}[4n^3 - 18n^2 + 24n - 10 + F(T) - 3(n - 2)M_1(T)]$ .

A matching  $K$  in a simple graph  $G$  is a set of pairwise non-adjacent edges, that is, no two edges of  $K$  share a common vertex. If  $|K| = k$  then  $K$  is called a  $k$ -matching of  $G$ . The matching polynomial of  $G$  is a generating function for counting the number of  $k$ -matchings in  $G$ . Let  $p(G, k)$  denote the number of  $k$ -matchings in  $G$ . Then the matching polynomial of  $G$  is defined as  $M(G) = \sum_{k \geq 0} (-1)^k p(G, k) x^{n-2k}$ , where  $n = |V(G)|$ . Farrell and Guo [5], established a formula for the number of 3-matchings in terms of the size, degree sequence and number of triangles in given graph  $G$ , and Behmaram [2] continued this work to present a formula for the number of 4-matchings of triangular-free graphs with respect to the number of vertices, edges, degrees and 4-cycles.

## 2. PRELIMINARY RESULTS

The aim of this section is to state some results which are crucial throughout the paper. We encourage the interested readers to consult papers [1, 7] for more details.

The common vertex of two incident edges  $e$  and  $f$  is denoted by  $e \cap f$ . Define the graph invariants  $\alpha(T)$  and  $\beta(T)$  as follows:

$$\alpha(T) = \sum_{u \sim v} \deg_T(u) \deg_T(v) (\deg_T(u) + \deg_T(v)),$$

$$\beta(T) = \sum_{e \sim f} \deg_T(e \cap f) (\deg_T(e) + \deg_T(f)).$$

Suppose  $T$  is a tree. In some of our results we need to have  $\eta(T, H)$  for some special subgraphs of  $T$ . In the following lemma we record some cases which are important in our calculations. The following lemma is a restatement of Lemmas 2.1, 2.2 and 2.3 of [7] in which the number of paths of length 3, 4 and 5 are given.

LEMMA 2.1. *Let  $T$  be an  $n$ -vertex tree. Then,*

$$\eta(T, P_3) = \frac{1}{2}M_1(T) - n + 1,$$

$$\eta(T, P_4) = M_2(T) - M_1(T) + n - 1,$$

$$\eta(T, P_5) = EM_2(T) + EM_1(T) + \frac{3}{2}M_1(T) + \frac{1}{2}M_1^4(T) - \frac{3}{2}F(T) - n + 1 - \beta(T).$$

The number of stars with exactly four and five vertices in a given tree  $T$  are presented in the following lemma which is Lemma 2.2 in [1].

LEMMA 2.2. *Let  $T$  be an  $n$ -vertex graph. Then,*

$$\eta(T, S_4) = \frac{1}{6}F(T) - \frac{1}{2}M_1(T) + \frac{2}{3}m,$$

$$\eta(T, S_5) = \frac{1}{24}M_1^4(T) - \frac{1}{4}F(T) + \frac{11}{24}M_1(T) - \frac{1}{2}m.$$

Let  $T$  be an arbitrary tree and  $T_1, T_2, \dots, T_5$  be graphs depicted in Figure 1. The number of subtrees of  $T$  isomorphic to one of these trees are given in the following lemma. These are restatements of Lemmas 2.3, 2.5., 2.7 and 2.15 in [1].

LEMMA 2.3. *Let  $T$  be an  $n$ -vertex tree. Then we have,*

$$\eta(T, T_1) = n\eta(T, P_4) + 2M_2(T) + F(T) - M_1(T) - 2\eta(T, P_5) - \alpha(T).$$

$$\eta(T, T_2) = \frac{1}{2}\alpha(T) + \frac{5}{2}M_1(T) - 3M_2(T) - \frac{1}{2}F(T) - 2m.$$

$$\begin{aligned} \eta(T, T_3) &= \eta(T, P_3)\left(\frac{1}{2}M_1(T) - n - 3\right) - \frac{5}{4}M_1^4(T) + \frac{11}{2}F(T) + 6M_2(T) \\ &\quad - \frac{33}{4}M_1(T) - 2EM_2(T) + 4m - \alpha(T) + 2\beta(T) - 3EM_1(T). \end{aligned}$$

$$\begin{aligned} \eta(T, T_4) &= \frac{1}{2}\eta(T, P_3)\left((n+1)(n+2) - M_1(T) + 4\right) + \frac{1}{4}(6n+52)M_1(T) \\ &\quad - \frac{1}{4}(2n+36)F(T) + 2M_1^4(T) - (2n+9)M_2(T) + 3EM_2(T) \\ &\quad - 8(n-1) + \frac{5}{2}\alpha(T) - 3\beta(T) + 5EM_1(T). \end{aligned}$$

$$\eta(T, T_5) = (n+2)\eta(T, S_4) - \frac{1}{2}\alpha(T) + \frac{1}{2}F(T) + 3M_2(T) - \frac{1}{6}M_1^4(T) - \frac{4}{3}M_1(T).$$

In [1], the authors proved a useful formula for computing the 4-matching of a tree which is important in our calculations.

THEOREM 2.4. *Let  $T$  be a tree with  $n$  vertices. Then,*

$$\begin{aligned} p(T, 4) &= \frac{1}{24}(n-1)(n^3 + 3n^2 + 22n + 4) - \frac{1}{4}(n^2 + 5n + \frac{27}{6})M_1(T) + \frac{1}{4}M_1(T)^2 \\ &\quad + (n+1)M_2(T) + \frac{1}{6}(2n + \frac{29}{2})F(T) - \frac{21}{24}M_1^4(T) - EM_2(T) \\ &\quad - EM_1(T) + \beta(T) - \alpha(T) - \sum_{\{u,v\} \subset V(T)} \binom{\text{deg}_T(u)}{2} \binom{\text{deg}_T(v)}{2}. \end{aligned}$$

LEMMA 2.5. *Let  $T$  be an  $n$ -vertex tree. Then*

$$\beta(T) - \alpha(T) = M_1^4(T) - 3F(T) + 2M_1(T) - 2M_2(T).$$

PROOF. By definition,

$$\begin{aligned} \beta(T) &= \sum_{e \sim f, e=uv, f=vx} \deg_T(v)(\deg_T(e) + \deg_T(f)) \\ &= \sum_{u \sim v \sim x} \deg_T(v) \left( \deg_T(u) + \deg_T(v) - 2 + \deg_T(v) + \deg_T(x) - 2 \right) \\ &= 2 \sum_{u \sim v \sim x} \deg_T(v)^2 - 4 \sum_{u \sim v \sim x} \deg_T(v) + \sum_{u \sim v \sim x} \deg_T(v)(\deg_T(u) + \deg_T(x)) \\ &= 2 \sum_{v \in V(T)} \binom{\deg_T(v)}{2} \deg_T(v)^2 - 4 \sum_{v \in V(T)} \binom{\deg_T(v)}{2} \deg_T(v) \\ &\quad + \sum_{uv \in E(T)} \deg_T(u) \deg_T(v) (\deg_T(u) + \deg_T(v) - 2) \\ &= \sum_{v \in V(T)} (\deg_T(v)^4 - \deg_T(v)^3) - 2 \sum_{v \in V(T)} (\deg_T(v)^3 - \deg_T(v)^2) \\ &\quad - 2M_2(T) + \alpha(T). \end{aligned}$$

Therefore,  $\beta(T) - \alpha(T) = M_1^4(T) - 3F(T) + 2M_1(T) - 2M_2(T)$ , which completes the proof.  $\square$

LEMMA 2.6. *Let  $T$  be a tree with  $n$  vertices. Then*

$$\begin{aligned} \eta(T, P_5) &= 6n - \frac{1}{4}F(T) - \frac{39}{8}M_1(T) + \frac{1}{2}nM_1(T) - \frac{1}{8}(M_1(T))^2 - \frac{1}{2}n^2 \\ &\quad + \frac{5}{8}M_1^4(T) + EM_2(T) + 3M_2(T) - \frac{11}{2} - \frac{1}{2}EM_1(T) - \beta(T) \\ &\quad + \sum_{\{u,v\} \subset V(T)} \binom{\deg_T(u)}{2} \binom{\deg_T(v)}{2}. \end{aligned}$$

PROOF. By definition,

$$\begin{aligned} \eta(T, P_5) &= \binom{n-1}{4} - \left( \eta(T, T_1) + \eta(T, T_2) + \eta(T, T_3) + \eta(T, T_4) + \eta(T, T_5) \right. \\ &\quad \left. + \eta(T, S_5) + p(T, 4) \right). \end{aligned}$$

Now, we apply Lemmas 2.2, 2.3, Theorem 2.4 and above discussion to deduce that

$$\begin{aligned} \eta(T, P_5) &= 6n - \frac{1}{4}F(T) - \frac{39}{8}M_1(T) + \frac{1}{2}nM_1(T) - \frac{1}{8}(M_1(T))^2 - \frac{1}{2}n^2 \\ &\quad + \frac{5}{8}M_1^4(T) + EM_2(T) + 3M_2(T) - \frac{11}{2} - \frac{1}{2}EM_1(T) - \beta(T) \\ &\quad + \sum_{\{u,v\} \subset V(T)} \binom{\deg_T(u)}{2} \binom{\deg_T(v)}{2}, \end{aligned}$$

proving the lemma.  $\square$

LEMMA 2.7. *Let  $T$  be a tree with  $n$  vertices and  $A(T) = \sum_{\{u,v\} \subset V(T)} \binom{\deg_T(u)}{2} \binom{\deg_T(v)}{2}$ . Then*

$$\begin{aligned} A(T) &= \frac{3}{2}EM_1(T) + \frac{51}{8}M_1(T) - \frac{1}{8}M_1^4(T) - \frac{5}{4}F(T) - 7n + \frac{13}{2} - \frac{1}{2}nM_1(T) \\ &\quad + \frac{1}{8}(M_1(T))^2 + \frac{1}{2}n^2 - 3M_2(T). \end{aligned}$$

PROOF. By two formulas for  $\eta(T, P_5)$  given Lemmas 2.1, 2.6, and a simple calculation we have

$$\begin{aligned} A(T) &= \frac{3}{2}EM_1(T) + \frac{51}{8}M_1(T) - \frac{1}{8}M_1^4(T) - \frac{5}{4}F(T) - 7n + \frac{13}{2} - \frac{1}{2}nM_1(T) \\ &\quad + \frac{1}{8}(M_1(T))^2 + \frac{1}{2}n^2 - 3M_2(T), \end{aligned}$$

proving the lemma.  $\square$

LEMMA 2.8. *Let  $G$  be a graph with  $m$  edges. Then  $EM_1(T) = F(G) + 2M_2(G) - 4M_1(G) + 4m$ .*

PROOF. By definition,

$$\begin{aligned} EM_1(T) &= \sum_{e=uv \in E(G)} \deg_G(e)^2 = \sum_{e=uv \in E(G)} (\deg_G(u) + \deg_G(v) - 2)^2 \\ &= \sum_{e=uv \in E(G)} \left( \deg_G(u)^2 + \deg_G(v)^2 + 2\deg_G(u)\deg_G(v) \right. \\ &\quad \left. - 4(\deg_G(u) + \deg_G(v)) + 4 \right) = F(G) + 2M_2(G) - 4M_1(T) + 4m, \end{aligned}$$

as desired.  $\square$

THEOREM 2.9 (See [1]). *Let  $T$  be a tree with  $n$  vertices. Then*

$$\begin{aligned} c_{n-4}(T) &= (n-1)\left(\frac{16}{24}n^3 - 4n^2 + \frac{348}{24}n - \frac{532}{6}\right) + \frac{17}{8}M_1(T)^2 \\ &+ \left(\frac{4}{6}n - \frac{412}{24}\right)F(T) + \frac{39}{2}EM_1(T) - \frac{108}{48}M_1^4(T) - 40M_2(T) \\ &- \left(n^2 + \frac{7}{2}n - \frac{1920}{24}\right)M_1(T) - 16 \sum_{\{u,v\} \subset V(T)} \binom{\deg_T(u)}{2} \binom{\deg_T(v)}{2}. \end{aligned}$$




FIGURE 1. The graphs  $T_1, \dots, T_5$  and  $S_5$ .

### 3. MAIN RESULTS

Suppose  $T$  is a tree. It is well known that the Laplacian coefficient  $c_{n-2}(T)$  is equal to the Wiener index of  $T$ , while  $c_{n-3}(T)$  is equal to the modified hyper-Wiener index of  $T$ . We refer to [11] for more information on this topic. So, it is natural to think about the coefficient  $c_{n-4}(T)$  and its relationship with some other topological indices of  $T$ .

The following environments are predefined:

THEOREM 3.1. *Let  $T$  be a tree with  $n$  vertices. Then,*

$$\begin{aligned} p(T, 4) &= \frac{1}{24}(n-1)(n^3 + 3n^2 + 10n - 80) + \frac{1}{8}M_1(T)(-2n^2 + M_1(T) - 6n + 36) \\ &+ M_2(T)(n-3) + \frac{1}{6}F(T)(2n-11) + \frac{1}{4}M_1^4(T) - EM_2(T). \end{aligned}$$

PROOF. By Theorem 2.4,

$$\begin{aligned} p(T, 4) &= \frac{1}{24}(n-1)(n^3 + 3n^2 + 22n + 4) - \frac{1}{4}(n^2 + 5n + \frac{27}{6})M_1(T) + \frac{1}{4}M_1(T)^2 \\ &\quad + (n+1)M_2(T) + \frac{1}{6}(2n + \frac{29}{2})F(T) - \frac{21}{24}M_1^4(T) - EM_2(T) - EM_1(T) \\ &\quad + \beta(T) - \alpha(T) - \sum_{\{u,v\} \subset V(T)} \binom{\deg_T(u)}{2} \binom{\deg_T(v)}{2}. \end{aligned}$$

Now, by Lemmas 2.5 and 2.7, we have

$$\begin{aligned} p(T, 4) &= \frac{1}{24}(n-1)(n^3 + 3n^2 + 10n + 160) + \frac{1}{8}M_1(T)(-2n^2 + M_1(T) - 6n - 44) \\ &\quad + M_2(T)(n+2) + \frac{1}{3}F(T)(n+2) + \frac{1}{4}M_1^4(T) - EM_2(T) - \frac{5}{2}EM_1(T), \end{aligned}$$

and by Lemma 2.8,

$$\begin{aligned} p(T, 4) &= \frac{1}{24}(n-1)(n^3 + 3n^2 + 10n - 80) + \frac{1}{8}M_1(T)(-2n^2 + M_1(T) - 6n + 36) \\ &\quad + M_2(T)(n-3) + \frac{1}{6}F(T)(2n-11) + \frac{1}{4}M_1^4(T) - EM_2(T). \end{aligned}$$

This completes the proof.  $\square$

THEOREM 3.2. *Let  $T$  be a tree with  $n$  vertices. Then*

$$\begin{aligned} c_{n-4}(T) &= \frac{1}{6}(n-1)(4n^3 - 24n^2 + 39n - 16) + \frac{1}{3}F(G)(2n-5) \\ &\quad + \frac{1}{8}M_1(T)(-8n^2 + M_1(T) + 36n - 32) - \frac{1}{4}M_1^4(T) - M_2(T). \end{aligned}$$

PROOF. By Lemmas 2.7, 2.8, Theorem 2.9, and simple calculations we have

$$\begin{aligned} c_{n-4}(T) &= \frac{1}{6}(n-1)(4n^3 - 24n^2 + 39n - 16) + \frac{1}{3}F(G)(2n-5) \\ &\quad + \frac{1}{8}M_1(T)(-8n^2 + M_1(T) + 36n - 32) - \frac{1}{4}M_1^4(T) - M_2(T). \end{aligned}$$

Hence the result.  $\square$

A pendant path of a graph  $G$  is a path  $P$ , in which one terminal vertex is of degree at least three, another terminal vertex is a pendant vertex, and all internal vertices (if any exists) are of degree two in  $G$ . It is clear that the number of pendant paths in  $G$  is equal to the number of pendant vertices in  $G$ . An internal path of  $G$  is a path  $I$ , in which two terminal vertices are of degree at least three and each internal vertex (if any exists) is of degree two in  $G$ . We also assume that  $\alpha_i$ ,  $1 \leq i \leq 6$ , are classes of chemical trees presented in Table 1.

**Transformation A.** Suppose  $G$  is a chemical tree with two given pendant paths  $P := v_1v_2 \dots v_k$  and  $Q := u_1u_2 \dots u_l$  such that  $k, l \geq 3$  and  $\deg_G(v_k) = \deg_G(u_l) = 1$ . Define  $G' = G - v_2v_3 + v_3u_l$ .



TABLE 1. Degree distributions of chemical trees with  $2 \leq n_1(T) \leq 5$ .

E.C.	$n_4$	$n_3$	$n_2$	$n_1$	E.C.	$n_4$	$n_3$	$n_2$	$n_1$
$\alpha_1$	0	0	$n-2$	2	$\alpha_4$	1	0	$n-5$	4
$\alpha_2$	0	1	$n-4$	3	$\alpha_5$	1	1	$n-7$	5
$\alpha_3$	0	2	$n-6$	4	$\alpha_6$	0	3	$n-8$	5

LEMMA 3.3. *Let  $G$  and  $G'$  be two chemical trees as described in Transformation A, with  $n$  ( $\geq 4$ ) vertices. Then  $c_{n-4}(G) < c_{n-4}(G')$ .*

PROOF. By definitions of  $G$  and  $G'$ , we have

$$M_1(G) = M_1(G'), \quad F(G) = F(G'), \quad M_1^4(G) = M_1^4(G').$$

Therefore by Theorem 3.2,

$$c_{n-4}(G) - c_{n-4}(G') = M_2(G') - M_2(G) = 2 - \deg_G(v_1).$$

Now,  $\deg_G(v_1) \in \{3, 4\}$  and so,  $c_{n-4}(G) - c_{n-4}(G') < 0$ .  $\square$

**Transformation B.** Suppose  $G$  is a chemical tree with a given internal path  $P_2 := v_1v_2$ . In addition, we assume that  $Q := u_1u_2 \dots u_l$  is a pendant or internal path in  $G$ , such that  $l \geq 4$ . Define  $G' = G - \{v_1v_2, u_1u_2, u_2u_3\} + \{v_1u_2, u_2v_2, u_1u_3\}$ .

LEMMA 3.4. *Let  $G$  and  $G'$  be two chemical trees as described in Transformation B, with  $n$  ( $\geq 8$ ) vertices. Then  $c_{n-4}(G) < c_{n-4}(G')$ .*

PROOF. By definitions of  $G$  and  $G'$ ,  $M_1(G) = M_1(G')$ ,  $F(G) = F(G')$  and  $M_1^4(G) = M_1^4(G')$ . We now apply Theorem 3.2 to deduce that  $c_{n-4}(G) - c_{n-4}(G') = M_2(G') - M_2(G) = 2\deg_G(v_1) + 2\deg_G(v_1) - \deg_G(v_1)\deg_G(v_2) - 4$ . Therefore,  $\deg_G(v_1), \deg_G(v_2) \in \{3, 4\}$  and so  $c_{n-4}(G) - c_{n-4}(G') < 0$ .  $\square$

**Transformation C.** Suppose  $G$  is a chemical tree with a given pendant path  $P_2 := v_1v_2 \dots v_k$  such that  $k \geq 3$  and  $\deg_G(v_k) = 1$ . In addition, we assume that  $Q := u_1u_2 \dots u_l$  is an internal path in  $G$ , such that  $l \geq 3$ . Define  $G' = G - \{v_2v_3, u_1u_2\} + \{u_1v_3, v_ku_2\}$ .

LEMMA 3.5. *Let  $G_1$  and  $G_2$  be two chemical trees as explained in Transformation C, with  $n$  ( $\geq 8$ ) vertices. Then  $c_{n-4}(G) < c_{n-4}(G')$ .*

PROOF. By definitions of  $G$  and  $G'$ ,  $M_1(G) = M_1(G')$ ,  $F(G) = F(G')$  and  $M_1^4(G) = M_1^4(G')$ . Apply Theorem 3.2 to prove that  $c_{n-4}(G) - c_{n-4}(G') = M_2(G') - M_2(G) = 4 + \deg_G(v_1) - [2 + 2\deg_G(v_1)] = 2 - \deg_G(v_1)$ . Since  $\deg_G(v_1) \in \{3, 4\}$ ,  $c_{n-4}(G) - c_{n-4}(G') < 0$ .  $\square$

**Transformation D.** Suppose  $G$  is a chemical tree with two given pendant paths  $P := v_1v_2 \dots v_k$  and  $Q := u_1u_2 \dots u_l$  such that  $\deg_G(v_k) = \deg_G(u_l) = 1$ . Define  $G' = G - v_1v_2 + u_lv_2$ .

Let  $T$  be a tree on  $n$  vertices. Then Gutman and Das in [10] have proved that

$$(3.1) \quad M_1(T) \leq n(n-1),$$

with equality if and only if  $T \cong S_n$ .

**LEMMA 3.6.** *Let  $G$  and  $G'$  be two chemical trees as in Transformation D, with  $n (\geq 8)$  vertices. Then  $c_{n-4}(G) < c_{n-4}(G')$ .*

**PROOF.** By definitions, if  $\deg_G(v_1) = 3$ , then

$$M_1(G) = M_1(G') + 2, \quad F(G) = F(G') + 12, \quad M_1^4(G) = M_1^4(G') + 50.$$

Therefore, by Theorem 3.2 and a simple calculation we have,

$$c_{n-4}(G) - c_{n-4}(G') \geq \frac{1}{2}M_1(G) - 2n^2 + 17n - 41 - M_2(G) + M_2(G').$$

By Equation (3),  $M_1(G) \leq n(n-1)$  and so,

$$c_{n-4}(G) - c_{n-4}(G') \leq \frac{1}{2}(33n - 3n^2) - 41 - M_2(G) + M_2(G').$$

Next by [4, Lemma 2.1],  $M_2(G') \leq M_2(G)$ . This proves that

$$c_{n-4}(G) - c_{n-4}(G') \leq \frac{1}{2}(33n - 3n^2) - 41 < 0.$$

The proof of the case that  $\deg_G(v_1) = 4$ , is similar.  $\square$

**LEMMA 3.7.** [8, Lemma 2.3] *If  $T$  is a chemical tree with  $n$  vertices, then*

$$n_1(T) = 2 + n_3(T) + 2n_4(T) \text{ and } n_2(T) = n - [2 + 2n_3(T) + 3n_4(T)].$$

**LEMMA 3.8.** *There exists a chemical tree of order  $n$  with  $2 \leq n_1(T) \leq 5$ , if and only if  $T$  belongs to one of the equivalence classes (E.C.) given in Table 1.*

**PROOF.** We distinguish the following four cases:

- (1)  $n_1(T) = 2$ .
- (2)  $n_1(T) = 3$ .
- (3)  $n_1(T) = 4$ .
- (4)  $n_1(T) = 5$ .

To prove case (1), let  $n_1(T) = 2$ . Then by Lemma 3.7, there is a tree  $T$  with  $n_1(T) = 2$  if and only if  $n_3(T) + 2n_4(T) = 0$ , if and only if  $n_3(T) = n_4(T) = 0$  if and only if  $n_2(T) = n - 2$  if and only if  $T \in \alpha_1$ . The proofs of the other cases are similar and we omit them.  $\square$

The number of edges connecting vertices of degree  $i$  and  $j$  in a graph  $A$  is denoted by  $m_{i,j}(A)$ . For a positive integer  $n \geq 10$ , we define:

$$B_1 = \{T \in \alpha_5 \mid m_{1,3}(T) = 2, m_{1,4}(T) = 3, m_{2,3}(T) = m_{2,4}(T) = 1, m_{2,2}(T) = n - 8\}.$$

$$B_2 = \{T \in \alpha_6 \mid m_{1,3}(T) = 5, m_{2,3}(T) = 4, \text{ and } m_{2,2}(T) = n - 10\}.$$

By Theorem 3.2, it is easy to see that for each  $T \in B_1$  and  $T' \in B_2$  we have

$$(3.2) \quad c_{n-4}(T) = \frac{1}{6}(2n-9)(2n^3-17n^2+25n+86),$$

$$(3.3) \quad c_{n-4}(T') = \frac{2}{3}n^4 - \frac{26}{3}n^3 + \frac{215}{6}n^2 - \frac{191}{6}n - 63.$$

LEMMA 3.9. *Let  $T$  be a chemical tree with  $n_1(T) \geq 5$ . Then,*

$$c_{n-4}(T) \leq \frac{2}{3}n^4 - \frac{26}{3}n^3 + \frac{215}{6}n^2 - \frac{191}{6}n - 63,$$

*with equality if and only if  $T \in B_2$ .*

PROOF. If  $n_1(T) = 5$ , then Lemmas 3.3, 3.4, 3.5, 3.8, and Equations 3.2, 3.3 give us the result. If  $n_1(T) \geq 6$ , then by repeated application of Transformation  $D$  we obtain a tree, say  $T'$ , such that  $n_1(T') = 5$ , and by Lemma 3.6,  $c_{n-4}(T') > c_{n-4}(T)$ . But  $c_{n-4}(T') \leq \frac{2}{3}n^4 - \frac{26}{3}n^3 + \frac{215}{6}n^2 - \frac{191}{6}n - 63$ , proving the lemma.  $\square$

We now apply Lemma 3.8 and Theorem 3.2, to compute the coefficient  $c_{n-4}$  for all chemical trees with  $n \geq 10$  vertices and  $2 \leq n_1 \leq 4$ .

$$\begin{aligned} A_1 &= \{T \in \alpha_1 \mid m_{1,2}(T) = 2, m_{2,2}(T) = n - 3\}, \\ A_2 &= \{T \in \alpha_2 \mid m_{1,2}(T) = 1, m_{1,3}(T) = 2, m_{2,3}(T) = 1, m_{2,2}(T) = n - 5\}, \\ A_3 &= \{T \in \alpha_2 \mid m_{1,2}(T) = 2, m_{1,3}(T) = 1, m_{2,3}(T) = 2, m_{2,2}(T) = n - 6\}, \\ A_4 &= \{T \in \alpha_2 \mid m_{1,2}(T) = 3, m_{2,3}(T) = 3, m_{2,2}(T) = n - 7\}, \\ A_5 &= \{T \in \alpha_3 \mid m_{1,3}(T) = 4, m_{2,3}(T) = 2, m_{2,2}(T) = n - 7\}, \\ A_6 &= \{T \in \alpha_3 \mid m_{1,2}(T) = 1, m_{1,3}(T) = 3, m_{2,3}(T) = 3, m_{2,2}(T) = n - 8\}, \\ A_7 &= \{T \in \alpha_3 \mid m_{1,2}(T) = 2, m_{1,3}(T) = 2, m_{2,3}(T) = 4, m_{2,2}(T) = n - 9\}, \\ A_8 &= \{T \in \alpha_3 \mid m_{1,2}(T) = 3, m_{1,3}(T) = 1, m_{2,3}(T) = 5, m_{2,2}(T) = n - 10\}, \\ A_9 &= \{T \in \alpha_3 \mid m_{1,2}(T) = 4, m_{2,3}(T) = 6, m_{2,2}(T) = n - 11\}, \\ A_{10} &= \{T \in \alpha_3 \mid m_{1,2}(T) = m_{2,3}(T) = m_{3,3}(T) = 1, m_{1,3}(T) = 3, m_{2,2}(T) = n - 7\}, \\ A_{11} &= \{T \in \alpha_3 \mid m_{1,2}(T) = m_{1,3}(T) = m_{2,3}(T) = 2, m_{3,3}(T) = 1, m_{2,2}(T) = n - 8\}, \\ A_{12} &= \{T \in \alpha_3 \mid m_{1,2}(T) = m_{2,3}(T) = 3, m_{1,3}(T) = m_{3,3}(T) = 1, m_{2,2}(T) = n - 9\}, \\ A_{13} &= \{T \in \alpha_3 \mid m_{1,2}(T) = 4, m_{2,3}(T) = 4, m_{3,3}(T) = 1, m_{2,2}(T) = n - 10\}, \\ A_{14} &= \{T \in \alpha_4 \mid m_{1,2}(T) = 1, m_{1,4}(T) = 3, m_{2,4}(T) = 1, m_{2,2}(T) = n - 6\}, \\ A_{15} &= \{T \in \alpha_4 \mid m_{1,2}(T) = 2, m_{1,4}(T) = 2, m_{2,4}(T) = 2, m_{2,2}(T) = n - 7\}, \\ A_{16} &= \{T \in \alpha_4 \mid m_{1,2}(T) = 3, m_{1,4}(T) = 1, m_{2,4}(T) = 3, m_{2,2}(T) = n - 8\}, \\ A_{17} &= \{T \in \alpha_4 \mid m_{1,2}(T) = 4, m_{2,4}(T) = 4, m_{2,2}(T) = n - 9\}. \end{aligned}$$

Let  $T_i \in A_i$ , for  $i = 1, 2, \dots, 17$ . Then by Theorem 3.2, we have:

$$\begin{aligned}
(3.4) \quad c_{n-4}(T_1) &= \frac{1}{6}(2n-5)(2n-7)(n-3)(n-4), \\
c_{n-4}(T_2) &= \frac{2}{3}n^4 - \frac{26}{3}n^3 + \frac{239}{6}n^2 - \frac{419}{6}n + 25, \\
c_{n-4}(T_3) &= \frac{1}{6}(2n-9)(2n^3 - 17n^2 + 43n - 16), \\
c_{n-4}(T_4) &= \frac{2}{3}n^4 - \frac{26}{3}n^3 + \frac{239}{6}n^2 - \frac{419}{6}n + 23, \\
c_{n-4}(T_5) &= \frac{2}{3}n^4 - \frac{26}{3}n^3 + \frac{227}{6}n^2 - \frac{305}{6}n - 19, \\
c_{n-4}(T_6) &= \frac{2}{3}n^4 - \frac{26}{3}n^3 + \frac{227}{6}n^2 - \frac{305}{6}n - 20, \\
c_{n-4}(T_7) &= c_{n-4}(T_{10}) = \frac{1}{6}(2n-9)(2n^3 - 17n^2 + 37n + 14), \\
c_{n-4}(T_8) &= c_{n-4}(T_{11}) = \frac{2}{3}n^4 - \frac{26}{3}n^3 + \frac{227}{6}n^2 - \frac{305}{6}n - 22, \\
c_{n-4}(T_9) &= c_{n-4}(T_{12}) = \frac{2}{3}n^4 - \frac{26}{3}n^3 + \frac{227}{6}n^2 - \frac{305}{6}n - 23, \\
c_{n-4}(T_{13}) &= \frac{2}{3}n^4 - \frac{26}{3}n^3 + \frac{227}{6}n^2 - \frac{305}{6}n - 24, \\
c_{n-4}(T_{14}) &= \frac{2}{3}n^4 - \frac{26}{3}n^3 + \frac{215}{6}n^2 - \frac{167}{6}n - 87, \\
c_{n-4}(T_{15}) &= \frac{2}{3}n^4 - \frac{26}{3}n^3 + \frac{215}{6}n^2 - \frac{167}{6}n - 89, \\
c_{n-4}(T_{16}) &= \frac{2}{3}n^4 - \frac{26}{3}n^3 + \frac{215}{6}n^2 - \frac{167}{6}n - 91, \\
c_{n-4}(T_{17}) &= \frac{2}{3}n^4 - \frac{26}{3}n^3 + \frac{215}{6}n^2 - \frac{167}{6}n - 93.
\end{aligned}$$

**THEOREM 3.10.** *If  $n \geq 11$ ,  $T_i \in A_i$ , for  $i = 1, 2, \dots, 17$ ,  $T_{18} \in B_2$ , and  $T \in \mathcal{CT}(n) \setminus \{T_1, T_2, \dots, T_{18}\}$ , then  $c_{n-4}(T_1) > c_{n-4}(T_2) > c_{n-4}(T_3) > c_{n-4}(T_4) > c_{n-4}(T_5) > c_{n-4}(T_6) > c_{n-4}(T_7) = c_{n-4}(T_{10}) > c_{n-4}(T_8) = c_{n-4}(T_{11}) > c_{n-4}(T_9) = c_{n-4}(T_{12}) > c_{n-4}(T_{13}) > c_{n-4}(T_{14}) > c_{n-4}(T_{15}) > c_{n-4}(T_{16}) > c_{n-4}(T_{17}) > c_{n-4}(T_{18}) > c_{n-4}(T)$ .*

**PROOF.** By Equations 3.3 and 3.4,  $c_{n-4}(T_1) > c_{n-4}(T_2) > c_{n-4}(T_3) > c_{n-4}(T_4) > c_{n-4}(T_5) > c_{n-4}(T_6) > c_{n-4}(T_7) = c_{n-4}(T_{10}) > c_{n-4}(T_8) = c_{n-4}(T_{11}) > c_{n-4}(T_9) = c_{n-4}(T_{12}) > c_{n-4}(T_{13}) > c_{n-4}(T_{14}) > c_{n-4}(T_{15}) > c_{n-4}(T_{16}) > c_{n-4}(T_{17}) > c_{n-4}(T_{18})$ . Since  $T \notin \{T_1, T_2, \dots, T_{18}\}$ ,  $n_1(T) \geq 5$  and Lemma 3.9, gives the result.  $\square$

**REMARK 3.11.**

1. If  $n = 10$ , then  $c_{n-4}(T_1) > c_{n-4}(T_2) > c_{n-4}(T_3) > c_{n-4}(T_4) > c_{n-4}(T_5) > c_{n-4}(T_6) > c_{n-4}(T_7) = c_{n-4}(T_{10}) > c_{n-4}(T_8) = c_{n-4}(T_{11})$

- $> c_{n-4}(T_{12}) > c_{n-4}(T_{13}) > c_{n-4}(T_{14}) > c_{n-4}(T_{15}) > c_{n-4}(T_{16}) >$   
 $c_{n-4}(T_{17}) > c_{n-4}(T_{18}) > c_{n-4}(T).$
2. If  $n = 9$ , then  $c_{n-4}(T_1) > c_{n-4}(T_2) > c_{n-4}(T_3) > c_{n-4}(T_4) > c_{n-4}(T_5)$   
 $> c_{n-4}(T_6) > c_{n-4}(T_7) = c_{n-4}(T_{10}) > c_{n-4}(T_{11}) > c_{n-4}(T_{12}) >$   
 $c_{n-4}(T_{14}) > c_{n-4}(T_{15}) > c_{n-4}(T_{16}) > c_{n-4}(T_{17}) > c_{n-4}(T).$
  3. If  $n = 8$ , then  $c_{n-4}(T_1) > c_{n-4}(T_2) > c_{n-4}(T_3) > c_{n-4}(T_4) > c_{n-4}(T_5)$   
 $> c_{n-4}(T_6) > c_{n-4}(T_{10}) > c_{n-4}(T_{11}) > c_{n-4}(T_{14}) > c_{n-4}(T_{15}) >$   
 $c_{n-4}(T_{16}) > c_{n-4}(T).$
  4. If  $n = 7$ , then  $c_{n-4}(T_1) > c_{n-4}(T_2) > c_{n-4}(T_3) > c_{n-4}(T_4) > c_{n-4}(T_5)$   
 $> c_{n-4}(T_{10}) > c_{n-4}(T_{14}) > c_{n-4}(T_{15}) > c_{n-4}(T).$
  5. If  $n = 6$ , then  $c_{n-4}(T_1) > c_{n-4}(T_2) > c_{n-4}(T_3) > c_{n-4}(T_{14}) >$   
 $c_{n-4}(T).$
  6. If  $n = 5$ , then  $c_{n-4}(T_1) = c_{n-4}(T_2) = c_{n-4}(S_5).$

## ACKNOWLEDGEMENTS.

The authors are indebted to an anonymous referee that his/her suggestions and helpful remarks led us to improve this paper. The research of the authors is partially supported by the University of Kashan under grant no 364988/169.

## REFERENCES

- [1] A. R. Ashrafi, M. Eliasi and A. Ghalavand, *Laplacian coefficients and Zagreb indices of trees*, Linear Multilinear Algebra **67** (2019), 1736–1749.
- [2] A. Behmaram, *On the number of 4-matchings in graphs*, MATCH Commun. Math. Comput. Chem. **62** (2009), 381–388.
- [3] D. Cvetković, M. Doob and H. Sachs, *Spectra of Graphs – Theory and Application*, Barth, Heidelberg, 1995.
- [4] M. Eliasi and A. Ghalavand, *Trees with the minimal second Zagreb index*, Kragujevac J. Math. **42** (2018), 325–333.
- [5] E. J. Farrell, J. M. Guo and G. M. Constantine, *On matching coefficients*, Discrete Math. **89** (1991), 203–210.
- [6] B. Furtula and I. Gutman, *A forgotten topological index*, J. Math. Chem. **53** (2015), 1184–1190.
- [7] A. Ghalavand, M. Eliasi and A. R. Ashrafi, *Relations between Wiener, hyper-Wiener and some Zagreb type indices*, Kragujevac J. Sci. **41** (2019), 37–42.
- [8] I. Gutman, A. Ghalavand, T. Dehghan-Zadeh and A. R. Ashrafi, *Graphs with smallest forgotten index*, Iranian J. Math. Chem. **8** (2017), 259–273.
- [9] I. Gutman and N. Trinajstić, *Graph theory and molecular orbitals. Total  $\varphi$ -electron energy of alternant hydrocarbons*, Chem. Phys. Lett. **17** (1972), 535–538.
- [10] I. Gutman and K. Ch. Das, *The first Zagreb indices 30 years after*, MATCH Commun. Math. Comput. Chem. **50** (2004), 83–92.
- [11] A. Ilić, *Trees with minimal Laplacian coefficients*, Comput. Math. Appl. **59** (2010), 2776–2783.
- [12] A. Miličević, S. Nikolić and N. Trinajstić, *On reformulated Zagreb indices*, Mol. Divers. **8** (2004), 393–399.

- [13] C. S. Oliveira, N. M. M. de Abreu and S. Jurkewicz, *The characteristic polynomial of the Laplacian of graphs in (a, b)-linear classes*, Linear Algebra Appl. **356** (2002) 113–121.
- [14] H. Wiener, *Structural determination of paraffin boiling points*, J. Am. Chem. Soc. **69** (1947) 17–20.
- [15] W. Yan and Y.-N. Yeh, *Connections between Wiener index and matchings*, J. Math. Chem. **39** (2006) 389–399.
- [16] S. Zhang and H. Zhang, *Unicyclic graphs with the first three smallest and largest first general Zagreb index*, MATCH Commun. Math. Comput. Chem. **55** (2006), 427–438.

## Laplaceovi koeficijenti stabala

*Ali Ghalavand i Ali Reza Ashrafi*

SAŽETAK. Neka je  $G$  jednostavan neusmjereni graf s Laplaceovim polinomom  $\psi(G, \lambda) = \sum_{k=0}^n (-1)^{n-k} c_k(G) \lambda^k$ . U ovom članku, izvedene su egzaktne formule za koeficijent  $c_{n-4}$  te za broj 4-sparivanja s obzirom na zagrebačke indekse danog stabla. Također su određena kemijska stabla koji imaju petnaest najvećih vrijednosti od  $c_{n-4}$ .

Ali Ghalavand  
Department of Pure Mathematics  
Faculty of Mathematical Sciences  
University of Kashan  
Kashan 87317-53153, I. R. Iran  
*E-mail:* [alighalavand@grad.kashanu.ac.ir](mailto:alighalavand@grad.kashanu.ac.ir)

Ali Reza Ashrafi  
Department of Pure Mathematics  
Faculty of Mathematical Sciences  
University of Kashan  
Kashan 87317-53153, I. R. Iran  
*E-mail:* [ashrafi@kashanu.ac.ir](mailto:ashrafi@kashanu.ac.ir)

*Received:* 29.6.2019.

*Revised:* 6.10.2019. ; 10.1.2020.

*Accepted:* 11.2.2020.