

ENVIRONMENT IMPACT ON CHILDREN'S HEALTH

UČINAK OKRUŽENJA NA ZDRAVLJE DJECE

*Carlos António Laranjeira*¹

ABSTRACT

Aim: There has been a growing concern in the last few years with the exposure of children to environmental chemicals. There is little knowledge regarding their effects on developing organisms. The objective of this study was to investigate the recent medical literature on environmental chemical hazards to child health.

Methods: The bibliographic research was conducted using the MEDLINE database, as well as through direct research, limiting the scope to articles published within the past 8 years.

Results: Children have a greater exposure to environmental pollutants than adults, because their metabolic needs and behaviors put them at special risk of contact with chemicals when they breathe, eat, drink or play. Heavy metals, pesticides, persistent organic pollutants and, at home, environmental tobacco smoke have been associated with the increasing number of diseases such as asthma, neurodevelopmental disorders and childhood cancer.

Conclusions: Pediatricians and other health professionals who treat children and adolescents are entrusted with the task of recognizing the hazards that environmental chemical agents may pose.

Key words: Environmental exposure, environmental pollution, chemical compounds

SAŽETAK

Cilj: U proteklih nekoliko godina, u porastu je zabrinutost zbog izloženosti djece kemikalijama koje se nalaze u okruženju. Malo se zna o učincima tih kemikalija na organizme u razvoju. Cilj je ovog istraživanja bio proučiti noviju medicinsku literaturu o opasnostima kemikalija koje se nalaze u okruženju, te njihovu utjecaju na zdravlje djece.

Metode: Proučavanje literature provedeno je korištenjem baze podataka MEDLINE, te neposrednim istraživanjem, pri čemu su uzeti u obzir jedino radovi objavljeni u proteklih osam godina.

Rezultati: Djeca su izložena onečišćivačima koji se nalaze u okruženju od odraslih, jer su zbog svojih metaboličkih potreba i ponašanja izloženi posebnu riziku vezanom uz kontakt s kemikalijama dok dišu, jedu, piju ili se igraju. Teški metali, pesticidi, postojani organski onečišćivači, te kod kuće sekundarni cigaretni dim, dovedeni su u vezu sa sve većim brojem bolesti u djece poput astme, neuroloških razvojnih poremećaja, ili raka.

Zaključak: Pedijatrima i ostalim zdravstvenim djelatnicima koji liječe djecu i adolescente, povjerena je zadaća prepoznavanja opasnosti koje mogu prouzročiti kemijska sredstva koja se nalaze u okruženju.

Ključne riječi: izloženost okruženju, onečišćenost okruženja, kemijski spojevi

INTRODUCTION

Environmental conditions are among the major factors that determine children's health. Unfavorable situations regarding access to drinkable water, sewage disposal and housing conditions contribute to remarkable morbidity and mortality of children in developing countries¹. Moreover, there has been a growing concern all over the world in the last decades with environmental contamination by

¹Escola Superior de Saúde do Campus Universitário Jean Piaget de Viseu, Estrada do Alto do Gaio – Galifonge, 3515776 Lordosa, Portugal

Primljeno: 16. listopada 2006.

Prihvaćeno: 25. siječnja 2007.

Correspondence to: Carlos António Laranjeira, Graduate in Nursing, Doctoral student in Nursing Sciences, Apartado 341, 3501-908 Viseu, Portugal, Mobile phone: +351 918 801 973, e-mail: carloslaranjeira@hotmail.com

chemical agents produced by industrial, mining and agricultural activities, and with their possible relationship with the increase in the prevalence of congenital malformations, asthma, cancer, and neurological and behavioral disorders in children in recent years².

Currently, around 85,000 chemical agents are produced and used in industrialized countries, 2,800 of which are mass-produced (over 500,000 kg a year). Less than 45% of these agents have undergone basic toxicological tests and less than 10% have been investigated as to their toxic effects on developing organisms.

Children are especially vulnerable to the exposure to environmental chemical agents, due to their physiological characteristics: they drink more water, eat more, and breathe in a greater amount of air per kg than adults do. An infant up to the sixth month of life drinks seven times more water, while a preschooler (1 to 5 years old) eats three or four times more per body weight than an average adult. Oxygen supply in a resting infant is twice as high as that which is observed in an adult. On top of that, the hand-to-mouth habit and playing or moving close to the ground contribute to children's greater exposure. Thus, any chemical agents found in water, air, soil and foods are more likely to be ingested by children than by adults³.

Due to the paucity of information about the consequences of such an exposure, a new and challenging field of research has been taking shape: environmental health in pediatrics. This field seeks to shed some light on the effects of environmental exposures to chemical or physical (e.g.: ionizing radiation) agents and on how these agents reach the children, and thus seeks to help with the prevention, diagnosis and management of various diseases that are possibly caused or aggravated by these conditions⁴⁻⁷.

The aim of the present review is to provide information about the susceptibility of children to the exposure to environmental chemical agents, placing special emphasis on the moments of greater vulnerability from intrauterine life to adolescence. It also describes specific aspects of the agents or groups of agents (heavy metals, pesticides, persistent organic pollutants) of greatest interest in our setting and suggests tools, such as environmental history, which can be useful in finding and applying solutions to the problem^{8,9}. Articles published on this subject between 1998 and 2006 were accessed via the Medical Literature, Analysis, and Retrieval System Online (MEDLINE).

EXPOSURE TO CHEMICAL AGENTS AND DEVELOPMENTAL STAGES

The risk of exposure to chemical agents may be present in intrauterine life (fetal growth period). The term placental "barrier" has become completely inadequate after those events involving thalidomide and dietilstilbestrol in the 1950s and 1960s¹⁰.

Besides drugs, lipophilic chemical contaminants such as polycyclic aromatic hydrocarbons (found in cigarette smoke), organic mercury compounds (methylmercury) and ethyl alcohol can easily cross the placenta¹¹.

Some essential processes, such as the central nervous system development, predominantly occur during fetal life and during the first years of life. These processes include differentiation of the cerebral cortex, neuronal migration, synaptogenesis and myelination¹². Fetal vulnerability and the neurotoxicity of some of the afore-mentioned agents during this period have been extensively documented in cases of fetal alcohol syndrome and in severe neurological disorders observed in Iraqi children who had been exposed to methylmercury during their intrauterine life due to their mothers' intake of bread contaminated with mercury fungicides in the 1970s¹³. In the neonatal period, there is a higher risk for the uptake of chemical agents through the skin, since keratinization is only complete around the fifth to eighth day of life. In addition, the body surface/mass ratio is greater in newborn infants than in older children and in adults. For a given exposed area, a newborn infant absorbs three times more than, and a child twice as much as, an adult¹¹.

The metabolic pathways of chemical agents that enter the body are not fully developed at birth and during the first months of life. The capacity to metabolize, detoxify and excrete many chemical agents is different from that observed in adults. In many cases, as in exposures to lead and organophosphates, children are less able to fight these agents due to the fact that they lack or are deficient in enzymes that could metabolize them and eliminate them².

Another important aspect is that occasional difficulties in metabolizing toxic agents may result in the buildup of these agents in storage organs or tissues (e.g.: bones and adipocytes) in case of continued or chronic exposure of children. Given the life expectancy of these children, the consequences of these exposures may be observed only several years later, for instance, possible development of cancer associated with exposure to cigarette smoke com-

ponents, from intrauterine life to the household environment during childhood¹⁴.

Processes caused by biological agents such as bacterial infections or environmental exposure to fungi, or by chemical air pollutants (e.g.: cigarette smoke or products from fuel combustion), certainly interfere with lung development. The proliferation of alveoli and pulmonary capillaries begins in the fetal period and extends to the fifth or eighth years of life, and problems during the normal development of this process may cause permanent sequelae on the morphology and function of the respiratory tract.

Older children (at preschool or school age) interact more constantly with the surrounding environment, due to their nutritional and metabolic requirements, and behavioral characteristics. Besides the household environment, new environments such as leisure areas and school may expose children to chemical contaminants¹⁵. One should recall that, in the so-called developing countries like Brazil, many children live, go to school, and play in places where the soil is highly contaminated by industrial waste, or in areas that are too close to garbage dumps or polluted water sources.

Finally, in adolescents, the quick development of the endocrine system towards sexual maturity is a vulnerable target, making them susceptible to related disorders⁵. Entry into the job market, which sometimes involves activities that are unhealthy and inappropriate for children and adolescents, becomes a possible risk of exposure to toxic substances.

CHEMICAL AGENTS OF INTEREST IN OUR SETTING

Lead

The effects produced by the exposure of children to lead have been well documented for over 100 years¹⁴. With the improved knowledge about the possible effects of lead on child development, blood lead levels currently tolerated are five times lower than those accepted in the 1970s.

The fact that apparently asymptomatic children had low IQ scores, difficulty expressing themselves verbally, attention disorders and behavioral problems was crucial for the establishment of increasingly lower acceptable levels.

The mechanisms through which lead affects the central nervous system functions remain unclear, but its interference in calcium transport and possible effect on the development and maturation of

neuronal connections may lie in the genesis of the observed involvement¹⁴.

In addition to the central nervous system, the urinary tract (kidneys) and the biosynthesis of hemoglobin are the other targets of toxic lead action, and anemia is a common finding in this type of exposure.

Lead pollution persists on a worldwide basis due to its use in metallurgy, manufacture of car batteries, ammunition industries, manufacture of crystals and production of paints and dyes.

Acute lead toxicity is rarely described in children nowadays, and is characterized by severe CNS involvement (encephalopathy), which may develop into seizures followed by coma and death if appropriate life support and specific treatment with chelating agents (e.g.: dimercaprol and edetate calcium disodium) are not available.

Mercury

Mercury is one of the most abundant metals on Earth. In addition to natural sources, it is found in various human activities, such as mineral extraction, coal combustion (thermoelectric plants), electronic industry, paper industry, and pulp bleaching (where mercury is used in the electrolytic process for the procurement of chlorine, being one of the resultant residues).

Metallic mercury, discharged into rivers and oceans, enters the food chain and turns into an organic compound (methylmercury) and is then submitted to biomagnification, where growing concentrations are observed even at the top of the food chain, which comprises large predator fish, which are eventually eaten by humans¹⁵. Organic mercury compounds are highly absorbed via food intake, opposed to metallic mercury, whose uptake is negligible.

Just like lead, organic mercury is toxic to the developing brain¹³. Studies with children exposed to methylmercury *in utero* showed adverse effects on intelligence and poor performance in speech, attention and memory.

Exposures to high levels of mercury, as the ones that occurred in Minamata Bay and in Iraq, cause cerebral palsy and remarkable mental retardation¹⁶.

More recent studies, conducted in the Seychelles, Indian ocean and on the Faroe islands in the North Atlantic, which included children whose mothers had eaten contaminated fish during pregnancy, have yielded controversial results. Whereas in the former one there was no correlation between

the levels of mercury detected in mothers and the cognitive development of concepti, there was an association between mercury levels detected at birth in umbilical cord blood (mean of 22.9 micrograms/liter) and poor cognitive performance at eight years of life in the second study¹³.

Pesticides

Pesticides for agricultural or domestic use, besides their significant morbidity and mortality from acute exposure, especially in developing countries, are a concern in terms of their long-term effects on the health of children exposed through water and soil contamination and food residues.

Organochlorine pesticides (DDT, hexachlorobenzene, aldrin) have a large environmental persistence and, although they were taken out of the market many years ago or been used sparingly nowadays, they can be found in farmlands or in industrial areas where they had been produced. Lactating women can mobilize organochlorine residuals from body stores (fat) and incorporate them into breastmilk. Animal studies have demonstrated the relationship of these agents with carcinogenesis, teratogenesis, immunotoxic effects and as endocrine disrupters¹⁷.

Organophosphate pesticides, such as malathion, chlorpyrifos, diazinon, and carbamates (e.g.: carbaryl) are widely used all over the world for pest control. Their mechanism of action is related to acetylcholinesterase inhibition. The essential role of acetylcholinesterase in the developmental stage (modeling) of neuronal connections in the central nervous system seems to be well established. Therefore, one cannot rule out the possibility that children who are continuously exposed to these agents may develop some problems, with possible neurobehavioral disorders as a result¹⁴.

Persistent organic pollutants (POPs)

Besides organochlorine pesticides, other compounds that are environmentally persistent and build up in storage tissues (e.g.: adipose tissue) have been investigated as to their toxicity to developing organisms. Among these compounds we have polychlorinated biphenyls (PCBs) and the dioxin group, whose best known member is tetrachloro-dibenzodioxin (TCDD). PCBs used to be employed because they made excellent electric and thermal insulating materials, especially for electrical transformers. Although their use was banned, they are still found in the environment due to

industrial residues with occasional water and food contamination.

Dioxins may be present in chlorophenoxy (2-4-dichlorophenoxyacetic acid) herbicides, nowadays in lesser amounts than in the past, but mainly as a by-product of the combustion of materials containing chlorine, as occurs in the incineration of industrial waste or garbage¹³.

PCBs and dioxins have been associated in animals studies with carcinogenesis and with the role of endocrine disrupters, with consequences such as low sperm count and infertility¹⁷.

Tobacco smoke

Tobacco smoke contains a wide range of chemical agents, including carbon monoxide and sulfur dioxide (also found in air pollution emissions in big cities, either from motor vehicles or industrial activities), and extremely toxic compounds, such as hydrogen cyanide, and carcinogenic substances (e.g.: benzopyrene).

The harmful effects of passive exposure of children to cigarette smoke and correlates can be felt *in utero*, extending throughout childhood. The effects of smoking on pregnancy include miscarriage, low birth weight and cognitive disorders. The association of tobacco smoke exposure in the household with recurrent otitis, lower airway infections, and induction or exacerbation of bronchial asthma¹⁴ in children has been well documented. There also seems to be an association between the incidence of cancer in adulthood and exposure to tobacco smoke in childhood¹⁴.

ENVIRONMENTAL EXPOSURE TO CHEMICAL AGENTS AND CHILDHOOD DISEASES

There seems to be a great amount of information and evidence, some of which is based on consistent epidemiological studies, that the occurrence of some diseases or group of diseases has increased in the last decades, especially in northern hemisphere countries^{2,14,16}.

This trend has also been observed in industrialized regions and in big cities in developing countries.

Among the several factors that could explain this increase are the improvement in diagnostic tools (as in the case of childhood cancer), greater exposure to chemical agents (e.g.: air pollutants and pesticides), and to physical agents (e.g.: electromagnetic radiation).

Asthma

Asthma incidence among children aged less than 18 years increased more than twice in the United States² and in other industrialized countries in the last decade. The major triggers of acute asthma attacks are cigarette smoke (at home), volatile organic compounds found in construction materials, cleaning products, adhesives, and pesticides. In the external environment of big cities, ozone (O₃) and sulfur dioxide (SO₂), produced by motor vehicles or by industries and thermoelectric plants, together with particulate material, have a key role in the development of asthma attacks. These chemical agents add up to external biological factors (allergens), such as pollen, fungi, dust mites and others, affecting mainly those children who live under poor housing conditions¹⁴.

Cancer

Despite the reduction in mortality rates and an increase in survival rates, due to diagnostic improvements and anticancer treatments in recent years, the incidence of some types of childhood cancer has surprisingly risen. The incidence of acute lymphoblastic leukemia increased nearly 30% between the 1970s and the 1990s in the United States, rising from 2.8 cases per every 100,000 children to 3.5:100,000.2 Between 1973 and 1994, the incidence of brain cancer increased by 39.6%, with a similar behavior between males and females³. Other types of cancer, found in young adults who had just grown out of adolescence, such as testicular cancer, increased by 68 % in the same time period².

A question that is still open is whether the improvements in diagnosis and case notification could explain such increases or whether the presence of chemical agents in the environment, given the high geographical correlation found in some cases, might be related to the collected epidemiological data¹⁸.

Neurological developmental disorders

It has been estimated that around 6% of live births in the United States may have neurobehavioral disorders, from attention deficit/hyperactivity disorder to autism. The causes have not been clearly established yet, but the association of agents, such as heavy metals (lead, mercury), certain pesticides and persistent organic pollutants (polychlorated biphenyls) with brain injury in children is widely known, as previously described. Neurological developmental disorders probably result from the interaction

between environmental factors and the individual susceptibility of every child, such as genetic predisposition. However, there is no doubt that exposures to high concentrations of agents such as organic mercurials, under certain circumstances (*in utero*), may, in isolation, cause remarkable damage¹⁷.

Endocrine disruption

Endocrine disrupters are natural or exogenous synthetic chemical agents that can mimic or modify the action of hormones¹⁴. Initially, the term was used to describe agents with estrogenic effects, such as estrogens found in plants and in foods of vegetable origin. Currently, the term is also used for disrupters of thyroid function or pancreatic islets (insulin producers) or for those which modify androgenic activity.

Even though few studies have been conducted with humans, some organic chlorinated compounds, such as DDT and chlordecone and polychlorinated biphenyls, revealed estrogenic activity in laboratory studies. Reproductive and fertility disorders were investigated in wild animals, such as birds and reptiles.

In human beings, an association has been made between exposure to endocrine disrupters and the increase in the incidence of hypospadias in boys and the increasingly earlier onset of puberty in girls².

ENVIRONMENTAL HISTORY IN PEDIATRICS

Currently, the training of health professionals and pediatricians does not include environmental health studies, which would properly prepare them for the recognition, treatment and prevention of environmentally related diseases and exposures. Due to biological, psychological, behavioral and social discrepancies between adults and children, there is the necessity for specific education of pediatricians in pediatric environmental health by including such syllabus in the School of Medicine curriculum, in Medical Residency programs and in continued education¹⁹.

The World Health Organization acknowledges that it is necessary to educate and train health professionals for the prevention, diagnosis and treatment of diseases related to environmental risk factors, since these diseases are regarded as an important global public health problem. A series of training courses organized by the WHO have been developed, including the elaboration of teaching materials, information brochures for health professionals and, mainly, the promotion of Environmen-

tal History in Pediatrics as an integrating part of standard medical History^{20,21}.

One of the roles of a pediatrician is to guide and instruct parents and family members on how to reduce or prevent exposures to the most common and potentially harmful chemical substances, but special attention should also be paid to new risks and to the effects produced by prolonged low-dose exposure (chronic exposure)^{22,23}. Many of the currently described and well-documented environmental risks were first discovered by the observation and investigation of some clinicians, for instance, the neurological disorders caused by methylmercury in Minamata, the congenital malformations caused by ethyl alcohol, among others²⁴.

A survey conducted by Kilpatrick et al. with pediatricians in Georgia (USA) about their knowledge of Environmental History revealed that only one in five pediatricians had had a training course in this area, but all of them believed that the information obtained through History could detect harmful exposures²⁵.

Observing the incidence and prevalence of diseases and using such information for interventions is a well-established strategy used in Public Health approaches; the necessity to follow up chronic diseases mediated by environmental factors in children has been increasingly acknowledged. It is believed that at least three clinical situations, as previously described, are correlated or may be exacerbated by the exposure to certain environmental agents: asthma, cancer and neurological development disorders¹⁶.

Information about the environment is important for children's health and should be included in every complete pediatric history^{14,26}.

What is environmental history in Pediatrics? It is a series of basic and concise questions that allow pediatricians to identify possible exposures of children to several chemical, physical and biological environmental factors, according to their specific vulnerabilities at each developmental stage. These questions reflect the regional social aspects of each population, and it is important that environmental history be carried out on a routine basis with all children (symptomatic or not)^{14,10,26,27}.

Environmental history, together with clinical history, allows health professionals to keep track of external and internal environmental conditions, behaviors and risk factors that are important to children's health. Examples include characteristics of the house, school, day care centers and recreation areas; proximity to garbage dumps; urban regions with

heavy traffic and with polluting industries; potential exposure to pesticides or other chemical products, among others²⁸.

Nowadays, there are specialized units that keep track of diseases and other problems related to environmental factors in the USA, Canada, Mexico and Spain, as well as other initiatives in Argentina and Uruguay. Since most health professionals do not have enough knowledge about the effects caused by exposures to chemical and physical agents, these specialized units play a crucial role in the diagnosis and follow-up of patients, provision of information to the population, research, training of health professionals and implementation of preventive measures²⁹.

FINAL CONSIDERATIONS

As we may perceive, pediatricians and other health professionals who treat children and adolescents are entrusted with the task of recognizing the hazards that environmental chemical agents may pose. This task should not be limited to developing skills for the detection of already established health damage or for the implementation of proper preventive measures in the environment shared by children. It is high time pediatricians, along with other professionals, organizations such as environmental organizations, centers for information and toxicological assistance, and non-governmental organizations for the protection of children and of the environment, actively struggle for legislative measures that properly protect our children and adolescents from environmental contamination by chemical agents^{9,29}.

Pediatricians may help promote better environmental health among their patients and in the community where they work in different ways. The inclusion of related topics in usual guidelines (e.g.: vaccination schemes) would be an important step. Measures such as sorting out and recycling garbage, using less pesticide in the household environment, and encouraging the intake of natural foods instead of industrialized ones are some examples of habits that should be promoted^{17,30}. In the community, participation in campaigns for less polluting alternative sources of energy (wind power, solar energy) and for reduced circulation of motor vehicles (good-quality public transportation and implementation of bicycle paths) may contribute to the improvement of environmental conditions in urban centers. Strategies for the preservation of water sources and forests are trends that should be supported by everybody¹⁷.

Finally, at the regional and national levels, pediatricians should engage in and actively advocate health and legal measures that allow the early detection of environmentally related diseases in children and the availability of resources for their management and prevention^{14,17,30}.

REFERENCES

- Bellamy C. Healthy environments for children. *Bull World Health Organ* 2003;81:157.
- Landrigan PJ, Garg A. Chronic effects of toxic environmental exposures on children's health. *J Toxicol Clin Toxicol* 2002;40:449-56.
- Landrigan PJ, Kimmel CA, Correa A, Eskenazi B. Children's health and the environment: public health issues and challenges for risk assessment. *Environ Health Perspect* 2004;112:257-65.
- Cohen Hubal EA, Sheldon LS, Burke JM, McCurdy TR, Berry MR, Rigas ML, et al. Children's exposure assessment: a review of factors influencing children's exposure, and the data available to characterize and assess that exposure. *Environ Health Perspect* 2000;108:475-86.
- Damstra T. Potential effects of certain persistent organic pollutants and endocrine disrupting chemicals on the health of children. *J Toxicol Clin Toxicol* 2002;40:457-65.
- Goldman LR, Koduru S. Chemicals in the environment and developmental toxicity to children: a public health and policy perspective. *Environ Health Perspect* 2000;108(Suppl)3:443-8.
- Makri A, Goveia M, Balbus J, Parkin R. Children's susceptibility to chemicals: a review by developmental stage. *J Toxicol Environ Health B Crit Rev* 2004;7:417-35.
- Balk SJ. Resources for pediatricians. How do I answer questions from parents, patients, teachers, and others? *Pediatr Clin North Am* 2001;48:1099-111, viii.
- Woolf AD. Introduction: children's health and the environment. *J Toxicol Clin Toxicol* 2002;40:447-8.
- Etzel RA, Balk SJ, Reigart JR, Landrigan PJ. Environmental health for practicing pediatricians. *Indian Pediatr* 2003;40:853-60.
- Mazur LJ. Pediatric environmental health. *Curr Probl Pediatr Adolesc Health Care* 2003;33:6-25.
- Bearer CF. Developmental neurotoxicity. Illustration of principles. *Pediatr Clin North Am* 2001;48:1199-213, ix.
- Grigg J. Environmental toxins; their impact on children's health. *Arch Dis Child* 2004;89:244-50.
- Etzel RA, Balk SJ, American Academy of Pediatrics. Committee on Environmental Health. Pediatric environmental health. 2nd ed. Elk Grove Village, IL: American Academy of Pediatrics 2003.
- Gitterman BA, Bearer CF. A developmental approach to pediatric environmental health. *Pediatr Clin North Am* 2001;48:1071-83.
- Woodruff TJ, Axelrad DA, Kyle AD, Nweke O, Miller GG, Hurley BJ. Trends in environmentally related childhood illnesses. *Pediatrics* 2004;113 (4 Suppl):1133-40.
- Children in the new millennium: environmental impact on health. Nairobi: UNEP 2002.
- WHO Regional Office for Europe, European Environmental Agency. Children's Health and environment: a review of evidence. Environmental issue report 29. Copenhagen, EEA 2000.
- McCurdy LE, Roberts J, Rogers B, Love R, Etzel R, Paulson J, et al. Incorporating environmental health into pediatric medical and nursing education. *Environ Health Perspect* 2004;112:1755-60.
- World Health Organization. Children's Environmental Health. Capacity building. 2006 [cited 2006 May 14]. Available from: URL:<http://www.who.int/ceh/capacity/en>.
- World Health Organization. Training package for health care providers. Improving the capacity to diagnose, prevent and manage paediatric diseases linked to the environment. 2006 [cited 2006 May 15]. Available from: URL:<http://www.who.int/ceh/capacity/trainpackage/en/>.
- Brent RL, Weitzman M. The Pediatricians' role and responsibility in educating parents about environmental risks. *Pediatrics* 2004;113:1167-72.
- Davoli CT. Home-sick: the effect of a child's environment on health [editorial]. *Arch Pediatr Adolesc Med* 2002;156:853-4.
- Miller RW. How environmental hazards in childhood have been discovered: carcinogens, teratogens, neurotoxicants, and others. *Pediatrics* 2004;113:945-51.
- Kilpatrick N, Frumkin H, Trowbridge J, Escoffrey C, Geller R, Rubin L, et al. The environmental history in pediatric practice: a study of pediatricians' attitudes, beliefs and practices. *Environ Health Perspect* 2002;110:823-7.
- The National Environmental Educational and Training Foundation (NEETF). Pediatric Environmental History Primer. 2006 [cited 2006 May 14]. Available from: URL:<http://www.neetf.org/Health/primer.pdf>.
- The National Environmental Educational and Training Foundation (NEETF). Position Statement Health Professionals and Environmental Health Education. 2006 [cited 2006 May 14]. Available from: URL:<http://www.neetf.org/Health/PositionStatement2.pdf>.
- World Health Organization. Paediatric environmental health centres. Expertise, information and training on the diagnosis, prevention and management of paediatric diseases linked to environment. 2006 [cited 2006 May 14]. Available from: URL: <http://www.who.int/ceh/capacity/paedehcentres/en>.
- Chance GW. Environmental toxicants and children's health: Cause for concern, time for action. *Paediatr Child Health* 2001;6:731-43.
- Shea KM. Global environmental change and children's health: understanding the challenges and finding solutions. *J Pediatr* 2003;143:149-54.