

Femoral to crural and distal-leg bypasses – five-year-period analysis in University Hospital Centre Zagreb

 **Josip Figl***,
 **Tomislav Meštrović**,
 **Ivan Brižić**,
 **Damir Halužan**,
 **Dino Papeš**,
 **Predrag Pavić**,
 **Andrea Crkvenac Gregorek**,
 **Irena Šnajdar**,
 **Ivan Škorak**

University of Zagreb School of Medicine, University Hospital Centre Zagreb, Zagreb, Croatia

KEYWORDS: critical limb ischemia, distal-leg bypass, femoral to crural bypass.

CITATION: Cardiol Croat. 2021;16(1-2):70. | <https://doi.org/10.15836/ccar2021.70>

***ADDRESS FOR CORRESPONDENCE:** Josip Figl, Klinički bolnički centar Zagreb, Kišpatičeva 12, HR-10000 Zagreb, Croatia. / Phone: +385-1-2388-240 / E-mail: figlmd@gmail.com

ORCID: Josip Figl, <https://orcid.org/0000-0001-6755-8656> • Tomislav Meštrović, <https://orcid.org/0000-0002-3492-3837>
Ivan Brižić, <https://orcid.org/0000-0002-5874-8666> • Damir Halužan, <https://orcid.org/0000-0002-8587-8351>
Dino Papeš, <https://orcid.org/0000-0001-7241-9693> • Predrag Pavić, <https://orcid.org/0000-0002-0117-7663>
Andrea Crkvenac Gregorek, <https://orcid.org/0000-0002-7790-1347> • Irena Šnajdar, <https://orcid.org/0000-0002-4429-1095>
Ivan Škorak, <https://orcid.org/0000-0002-4900-4876>

The goal: To compare the results of most distal femoral bypasses in our institution with results in literature.

Patients and Methods: A single centre retrospective analysis of femoral to crural and distal-leg bypass performed in University Hospital Centre Zagreb between 2014 and 2018.

Results: Primary one-year patency is even a bit better in our hospital than in literature (80% vs 77% respectively) and secondary patency is in the range of the world literature (54% vs 68% respectively).

Conclusion: Surgical revascularization in form of the most distal bypasses, performed in this institution observed, although reserved for a smaller number of patients, is very good and reliable method of treatment, and its results are at the level of world standards and percentages.¹⁻³

RECEIVED:
November 13, 2020

ACCEPTED:
December 18, 2020



LITERATURE

- Wölfle KD, Bruijnen H, Loeprecht H, Rümenapf G, Schweiger H, Grabitz K, et al. Graft patency and clinical outcome of femorodistal arterial reconstruction in diabetic and non-diabetic patients: results of a multicentre comparative analysis. Eur J Vasc Endovasc Surg. 2003 Mar;25(3):229-34. <https://doi.org/10.1053/ejvs.2002.1849>
- van der Zaag ES, Legemate DA, Prins MH, Reekers JA, Jacobs MJ. Angioplasty or bypass for superficial femoral artery disease? A randomised controlled trial. Eur J Vasc Endovasc Surg. 2004 Aug;28(2):132-7. <https://doi.org/10.1016/j.ejvs.2004.04.003>
- Veith FJ, Gupta SK, Ascer E, White-Flores S, Samson RH, Scher LA, Towne JB, Bernhard VM, Bonier P, Flinn WR, et al. Six-year prospective multicenter randomized comparison of autologous saphenous vein and expanded polytetrafluoroethylene grafts in infringuinal arterial reconstructions. J Vasc Surg. 1986 Jan;3(1):104-14. [https://doi.org/10.1016/0741-5214\(86\)90073-X](https://doi.org/10.1016/0741-5214(86)90073-X)