

VIKOR method for multiple criteria group decision making under 2-tuple linguistic neutrosophic environment

Guiwu Wei, Jie Wang, Jianping Lu, Jiang Wu, Cun Wei, Fuad E. Alsaadi & Tasawar Hayat

To cite this article: Guiwu Wei, Jie Wang, Jianping Lu, Jiang Wu, Cun Wei, Fuad E. Alsaadi & Tasawar Hayat (2020) VIKOR method for multiple criteria group decision making under 2-tuple linguistic neutrosophic environment, Economic Research-Ekonomiska Istraživanja, 33:1, 3185-3208, DOI: [10.1080/1331677X.2019.1691036](https://doi.org/10.1080/1331677X.2019.1691036)

To link to this article: <https://doi.org/10.1080/1331677X.2019.1691036>



© 2020 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 25 Nov 2019.



Submit your article to this journal [↗](#)



Article views: 794



View related articles [↗](#)




View Crossmark data [↗](#)



Citing articles: 11 View citing articles [↗](#)

VIKOR method for multiple criteria group decision making under 2-tuple linguistic neutrosophic environment

Guiwu Wei^{a,b} , Jie Wang^a, Jianping Lu^a, Jiang Wu^c, Cun Wei^c, Fuad E. Alsaadi^b and Tasawar Hayat^{d,e}

^aSchool of Business, Sichuan Normal University, Chengdu, P.R. China; ^bCommunications Systems and Networks (CSN) Research Group, Department of Electrical and Computer Engineering, Faculty of Engineering, King Abdulaziz University, Jeddah, Saudi Arabia; ^cSchool of Statistics, Southwestern University of Finance and Economics, Chengdu, P.R. China; ^dDepartment of Mathematics, Quaid-I-Azam University, Islamabad, Pakistan; ^eNonlinear Analysis and Applied Mathematics (NAAM) Research Group, Department of Mathematics, Faculty of Science, King Abdulaziz University, Jeddah, Saudi Arabia

ABSTRACT

In this article, the VIKOR method is proposed to solve the multiple criteria group decision making (MCGDM) with 2-tuple linguistic neutrosophic numbers (2TLNNs). Firstly, the fundamental concepts, operation formulas and distance calculating method of 2TLNNs are introduced. Then some aggregation operators of 2TLNNs are reviewed. Thereafter, the original VIKOR method is extended to 2TLNNs and the calculating steps of VIKOR method with 2TLNNs are proposed. In the proposed method, it's more reasonable and scientific for considering the conflicting criteria. Furthermore, the VIKOR are extended to interval-valued 2-tuple linguistic neutrosophic numbers (IV2TLNNs). Moreover, a numerical example for green supplier selection has been given to illustrate the new method and some comparisons are also conducted to further illustrate advantages of the new method.

ARTICLE HISTORY

Received 1 December 2018
Accepted 14 October 2019

KEYWORDS

Multiple criteria group decision making (MCGDM); 2-tuple linguistic neutrosophic numbers (2TLNNs); interval-valued 2-tuple linguistic neutrosophic numbers (IV2TLNNs); VIKOR method green suppliers selection

JEL CLASIFICACIONES

C43; C61; D81

1. Introduction

In practical decision problems, it's difficult to present the criteria values with real values for the complexity and fuzziness of the alternatives, so it can be more useful and effective to express the criteria values by different kinds of fuzzy numbers, such as intuitionistic fuzzy set (IFSs) (Atanassov, 1986; Li, Gao, & Wei, 2018; Wu, Wei, Gao, & Wei, 2018), Pythagorean fuzzy set (PFSs) (Tang et al., 2019; Tang, Wei, & Gao, 2019a, 2019b; Yager, 2014), q-rung orthopair fuzzy sets (q-ROFSs) (Wang, Gao, Wei, & Wei, 2019; Wang, Wang, Wei, & Wei, 2019; Yager, 2017). The fuzzy set theory which initially introduced by Zadeh (1965) has been proved as a feasible mean in the

CONTACT Guiwu Wei  weiguiwu@163.com

© 2020 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

application of MCGDM (Mahmoudi, Sadi-Nezhad, & Makui, 2016; Sharma, Kumari, & Kar, 2019; Wang, Wei, & Lu, 2018a; Wang, Wang, & Wei, 2019; Wei, 2019; Wei, Wang, Wei, Wei, & Zhang, 2019; Wei, Wang, Wang, Wei, & Zhang, 2019). Atanassov (1986) defined the IFSs which consider the membership degree and the non-membership degree. To depict the indeterminacy membership degree, Smarandache (1999) provided the neutrosophic sets (NSs). Wang, Smarandache, Zhang, and Sunderraman (2010) investigated some theories about single-valued neutrosophic sets (SVNSs) and given the definition of interval neutrosophic sets (INNS). Ye (2018) studied the MADM problems under the hesitant linguistic neutrosophic (HLN) environment. Wang, Tang, and Wei (2018) studied the dual generalized Bonferroni mean (DGBM) aggregation operators under the SVNSs environment and developed some aggregation operators based on the traditional BM operators (Deng, Wei, Gao, & Wang, 2018; Tang & Wei, 2018; Wang, Wei, & Wei, 2018; Wei, 2017; Wei & Zhang, 2019; Xu & Chen, 2011; Zhu & Xu, 2013). Liu and You (2018) proposed some linguistic neutrosophic Hamy mean (LNHM) aggregation operators. Wu, Wu, Zhou, Chen, and Guan (2018) gave the definition of SVN 2-tuple linguistic sets (SVN2TLSs) and proposed some new Hamacher aggregation operators. Ju, Ju, and Wang (2018) extended the SVN2TLSs to interval-valued environment and presented some single-valued neutrosophic interval 2-tuple linguistic Maclaurin symmetric mean (SVN-ITLMSM) operators. Wu, Wang, Wei, and Wei (2018) studied SVNSs with Hamy operators under 2-tuple linguistic varies environment. Wang, Wei et al. (2018) gave the definition of 2-tuple linguistic neutrosophic set (2TLNS) which the truth-membership degree (MD), indeterminacy-membership degree (IMD) and falsity-membership degree (FMD) are depicted by 2-tuple linguistic neutrosophic numbers (2TLNNs). Wang, Wei, and Lu (2018b) developed an extended TODIM model (Gomes & Rangel, 2009; Huang & Wei, 2018; Wang et al., 2018b; Wei, 2018) under 2-tuple linguistic neutrosophic environment and applied the new defined model in safety assessment of a construction project. Wang, Gao, and Wei (2018) studied the Muirhead mean (MM) operator and the dual Muirhead mean (DMM) operator under 2-tuple linguistic neutrosophic environment, then some 2-tuple linguistic neutrosophic Muirhead mean operators were given to deal with green supplier selection. Thereafter, the 2-tuple linguistic neutrosophic set (2TLNS) theory has been broadly used to study MCGDM problems.

For MADM problems, the way to express the assessment information is only one aspect, another vital aspect is selecting best alternative from a given alternative set. In previous document, some traditional decision making model had been applied to MADM problems, such as the EDAS model (Keshavarz Ghorabae, Zavadskas, Olfat, & Turskis, 2015), the MABAC model (Pamucar & Cirovic, 2015), the COPRAS model (Roy, Sharma, Kar, Zavadskas, & Saporauskas, 2019), the TOPSIS model (Chen, 2000; Lai, Liu, & Hwang, 1994), The TODIM model (Gomes & Lima, 1979) and the GRA model (Li & Wei, 2014). As a powerful tool for handling MADM, The VIKOR (Vlsekriterijumska Optimizacija I KOMPromisno Resenje) method (Opricovic & Tzeng, 2004), which owns precious merits of considering the compromise between group utility maximization and individual regret minimization, has been regards as a meaningful tool to apply in many decision making fields in past few years.

Comparing with these above mentioned methods, the VIKOR model has the advantage of taking the compromise between group utility maximization and individual regret minimization into consideration. Du and Liu (2011) extended the traditional VIKOR model to intuitionistic trapezoidal fuzzy environment. Park, Cho, and Kwun (2011) established the interval-valued intuitionistic fuzzy VIKOR model for MADM problems. Qin, Liu, and Pedrycz (2015) proposed an extension of VIKOR model based on interval type-2 fuzzy information. Based on extended hesitant fuzzy linguistic information, Ghadikolaei, Madhoushi, and Divsalar (2018) built new extended VIKOR model for MADM problems. Narayanamoorthy, Geetha, Rakkiyappan, and Joo (2019) developed an extended VIKOR model based on interval-valued intuitionistic hesitant fuzzy entropy for industrial robots selection. Yang, Pang, Shi, and Wang (2018) defined the linguistic hesitant intuitionistic VIKOR model for MADM. Wang, Zhang, Wang, and Li (2018) proposed the projection-based VIKOR model under picture fuzzy environment and applied it for the risk evaluation of construction project. Wu, Xu, Jiang, and Zhong (2019) presented the VIKOR model based on hesitant fuzzy linguistic term sets with possibility distributions.

According to above literature review, we can obtain that the 2-tuple linguistic neutrosophic set (2TLNS) can express the assessment information easily and reasonably, the VIKOR method can consider the conflicting criteria. Thus, to combine these two advantages, we shall propose some extended VIKOR models with 2TLNNs. The structure of our paper is organized as follows. Section 2 introduces the concepts, operation formulas, distance calculating method and some aggregation operators of 2TLNNs. Section 3 extends the original VIKOR method to 2TLNNs and introduce the calculating steps of VIKOR method with 2TLNNs. Section 4 extends the VIKOR method to IV2TLNNs and develops the calculating steps of VIKOR method with IV2TLNNs. Section 5 provides a numerical example for green supplier selection and introduces the comparison between our proposed methods with the existing method. Section 6 gives some summaries of our article.

2. Preliminaries

2.1. 2-tuple linguistic neutrosophic sets

Based on the concepts of 2-tuple linguistic fuzzy set (2TLS) and the fundamental theories of single valued neutrosophic set (SVNS), the 2-tuple linguistic neutrosophic sets (2TLNSs) which firstly defined by Wang, Wei et al. (2018) can be depicted as follows.

Definition 1. (Wang, Wei et al., 2018) Let s_1, s_2, \dots, s_k be a linguistic term set. Any label s_i shows a possible linguistic variable, and $S = \{s_0 = \text{extremelypoor}, s_1 = \text{verypoor}, s_2 = \text{poor}, s_3 = \text{medium}, s_4 = \text{good}, s_5 = \text{verygood}, s_6 = \text{extremelygood}\}$, the 2TLNSs η can be depicted as:

$$\eta = \{(s_\alpha, \phi), (s_\beta, \varphi), (s_\gamma, \gamma)\} \quad (1)$$

Where $\Delta^{-1}(s_\alpha, \phi), \Delta^{-1}(s_\beta, \varphi)$ and $\Delta^{-1}(s_\gamma, \gamma) \in [0, k]$ represent the degree of the truth membership, the indeterminacy membership and the falsity membership which are expressed by 2TLNNs and satisfies the condition $0 \leq \Delta^{-1}(s_\alpha, \phi) + \Delta^{-1}(s_\beta, \varphi) + \Delta^{-1}(s_\gamma, \gamma) \leq 3k$.

Definition 2. (Wang, Wei et al., 2018) Assume there are three 2TLNNs $\eta_1 = \{(s_{\alpha_1}, \phi_1), (s_{\beta_1}, \varphi_1), (s_{\gamma_1}, \gamma_1)\}$, $\eta_2 = \{(s_{\alpha_2}, \phi_2), (s_{\beta_2}, \varphi_2), (s_{\gamma_2}, \gamma_2)\}$ and $\eta = \{(s_\alpha, \phi), (s_\beta, \varphi), (s_\gamma, \gamma)\}$, the operation laws of them can be defined:

$$(1) \eta_1 \oplus \eta_2 = \left\{ \Delta \left(k \left(\frac{\Delta^{-1}(s_{\alpha_1}, \phi_1)}{k} + \frac{\Delta^{-1}(s_{\alpha_2}, \phi_2)}{k} - \frac{\Delta^{-1}(s_{\alpha_1}, \phi_1)}{k} \cdot \frac{\Delta^{-1}(s_{\alpha_2}, \phi_2)}{k} \right) \right), \right. \\ \left. \Delta \left(k \left(\frac{\Delta^{-1}(s_{\beta_1}, \varphi_1)}{k} \cdot \frac{\Delta^{-1}(s_{\beta_2}, \varphi_2)}{k} \right) \right), \Delta \left(k \left(\frac{\Delta^{-1}(s_{\gamma_1}, \gamma_1)}{k} \cdot \frac{\Delta^{-1}(s_{\gamma_2}, \gamma_2)}{k} \right) \right) \right\};$$

$$(2) \eta_1 \otimes \eta_2 = \left\{ \Delta \left(k \left(\frac{\Delta^{-1}(s_{\alpha_1}, \phi_1)}{k} \cdot \frac{\Delta^{-1}(s_{\alpha_2}, \phi_2)}{k} \right) \right), \right. \\ \left. \Delta \left(k \left(\frac{\Delta^{-1}(s_{\beta_1}, \varphi_1)}{k} + \frac{\Delta^{-1}(s_{\beta_2}, \varphi_2)}{k} - \frac{\Delta^{-1}(s_{\beta_1}, \varphi_1)}{k} \cdot \frac{\Delta^{-1}(s_{\beta_2}, \varphi_2)}{k} \right) \right), \right. \\ \left. \Delta \left(k \left(\frac{\Delta^{-1}(s_{\gamma_1}, \gamma_1)}{k} + \frac{\Delta^{-1}(s_{\gamma_2}, \gamma_2)}{k} - \frac{\Delta^{-1}(s_{\gamma_1}, \gamma_1)}{k} \cdot \frac{\Delta^{-1}(s_{\gamma_2}, \gamma_2)}{k} \right) \right) \right\};$$

$$(3) \lambda \eta = \left\{ \Delta \left(k \left(1 - \left(1 - \frac{\Delta^{-1}(s_\alpha, \phi)}{k} \right)^\lambda \right) \right), \Delta \left(k \left(\frac{\Delta^{-1}(s_\beta, \varphi)}{k} \right)^\lambda \right), \Delta \left(k \left(\frac{\Delta^{-1}(s_\gamma, \gamma)}{k} \right)^\lambda \right) \right\}, \lambda > 0;$$

$$(4) \eta^\lambda = \left\{ \Delta \left(k \left(\frac{\Delta^{-1}(s_\alpha, \phi)}{k} \right)^\lambda \right), \Delta \left(k \left(1 - \left(1 - \frac{\Delta^{-1}(s_\beta, \varphi)}{k} \right)^\lambda \right) \right), \Delta \left(k \left(1 - \left(1 - \frac{\Delta^{-1}(s_\gamma, \gamma)}{k} \right)^\lambda \right) \right) \right\}, \lambda > 0.$$

According to the Definition 2, it's clear that the operation laws have the following properties.

$$(1) \eta_1 \oplus \eta_2 = \eta_2 \oplus \eta_1, \eta_1 \otimes \eta_2 = \eta_2 \otimes \eta_1, \left((\eta_1)^{\lambda_1} \right)^{\lambda_2} = (\eta_1)^{\lambda_1 \lambda_2}; \tag{2}$$

$$(2) \lambda(\eta_1 \oplus \eta_2) = \lambda \eta_1 \oplus \lambda \eta_2, (\eta_1 \otimes \eta_2)^\lambda = (\eta_1)^\lambda \otimes (\eta_2)^\lambda; \tag{3}$$

$$(3) \lambda_1 \eta_1 \oplus \lambda_2 \eta_1 = (\lambda_1 + \lambda_2) \eta_1, (\eta_1)^{\lambda_1} \otimes (\eta_1)^{\lambda_2} = (\eta_1)^{(\lambda_1 + \lambda_2)}. \tag{4}$$

Definition 3. (Wang, Wei et al., 2018) Let $\eta = \{(s_\alpha, \phi), (s_\beta, \varphi), (s_\gamma, \gamma)\}$ be a 2TLNN, the score and accuracy functions of η can be expressed:

$$s(\eta) = \frac{\left(2k + \Delta^{-1}(s_\alpha, \phi) - \Delta^{-1}(s_\beta, \varphi) - \Delta^{-1}(s_\gamma, \gamma)\right)}{3k}, s(\eta) \in [0, 1] \tag{5}$$

$$h(\eta) = \Delta^{-1}(s_\alpha, \phi) - \Delta^{-1}(s_\gamma, \gamma), h(\eta) \in [-k, k] \tag{6}$$

For two 2TLNNs η_1 and η_2 , based on the Definition 3, then

- (1) if $s(\eta_1) < s(\eta_2)$, then $\eta_1 < \eta_2$;
- (2) if $s(\eta_1) > s(\eta_2)$, then $\eta_1 > \eta_2$;
- (3) if $s(\eta_1) = s(\eta_2)$, $h(\eta_1) < h(\eta_2)$, then $\eta_1 < \eta_2$;
- (4) if $s(\eta_1) = s(\eta_2)$, $h(\eta_1) > h(\eta_2)$, then $\eta_1 > \eta_2$;
- (5) if $s(\eta_1) = s(\eta_2)$, $h(\eta_1) = h(\eta_2)$, then $\eta_1 = \eta_2$.

2.2. The normalized Hamming distance

Definition 4. Let $\eta_1 = \{(s_{\alpha_1}, \phi_1), (s_{\beta_1}, \varphi_1), (s_{\gamma_1}, \gamma_1)\}$ and $\eta_2 = \{(s_{\alpha_2}, \phi_2), (s_{\beta_2}, \varphi_2), (s_{\gamma_2}, \gamma_2)\}$ be two 2TLNNs, then we can get the normalized Hamming distance:

$$d(\eta_1, \eta_2) = \frac{1}{3k} \left(|\Delta^{-1}(s_{\alpha_1}, \phi_1) - \Delta^{-1}(s_{\alpha_2}, \phi_2)| + |\Delta^{-1}(s_{\beta_1}, \varphi_1) - \Delta^{-1}(s_{\beta_2}, \varphi_2)| + |\Delta^{-1}(s_{\gamma_1}, \gamma_1) - \Delta^{-1}(s_{\gamma_2}, \gamma_2)| \right) \tag{7}$$

Theorem 1. Assume there are three 2TLNNs $\eta_1 = \{(s_{\alpha_1}, \phi_1), (s_{\beta_1}, \varphi_1), (s_{\gamma_1}, \gamma_1)\}$, $\eta_2 = \{(s_{\alpha_2}, \phi_2), (s_{\beta_2}, \varphi_2), (s_{\gamma_2}, \gamma_2)\}$ and $\eta_3 = \{(s_{\alpha_3}, \phi_3), (s_{\beta_3}, \varphi_3), (s_{\gamma_3}, \gamma_3)\}$, the Hamming distance d has the following properties:

- (P1) $0 \leq d(\eta_1, \eta_2) \leq 1$;
- (P2) if $d(\eta_1, \eta_2) = 0$, then $\eta_1 = \eta_2$;
- (P3) $d(\eta_1, \eta_2) = d(\eta_2, \eta_1)$;
- (P4) $d(\eta_1, \eta_2) + d(\eta_2, \eta_3) \geq d(\eta_1, \eta_3)$.

Proof. (P1) $0 \leq d(\eta_1, \eta_2) \leq 1$

Since $\Delta^{-1}(s_{\alpha_1}, \phi_1), \Delta^{-1}(s_{\alpha_2}, \phi_2) \in [0, k]$, then $0 \leq |\Delta^{-1}(s_{\alpha_1}, \phi_1) - \Delta^{-1}(s_{\alpha_2}, \phi_2)| \leq k$, similarly we can get $0 \leq |\Delta^{-1}(s_{\beta_1}, \varphi_1) - \Delta^{-1}(s_{\beta_2}, \varphi_2)| \leq k, 0 \leq |\Delta^{-1}(s_{\gamma_1}, \gamma_1) - \Delta^{-1}(s_{\gamma_2}, \gamma_2)| \leq k$, then $0 \leq |\Delta^{-1}(s_{\alpha_1}, \phi_1) - \Delta^{-1}(s_{\alpha_2}, \phi_2)| + |\Delta^{-1}(s_{\beta_1}, \varphi_1) - \Delta^{-1}(s_{\beta_2}, \varphi_2)| + |\Delta^{-1}(s_{\gamma_1}, \gamma_1) - \Delta^{-1}(s_{\gamma_2}, \gamma_2)| \leq 3k$, So $0 \leq \left(|\Delta^{-1}(s_{\alpha_1}, \phi_1) - \Delta^{-1}(s_{\alpha_2}, \phi_2)| + |\Delta^{-1}(s_{\beta_1}, \varphi_1) - \Delta^{-1}(s_{\beta_2}, \varphi_2)| + |\Delta^{-1}(s_{\gamma_1}, \gamma_1) - \Delta^{-1}(s_{\gamma_2}, \gamma_2)| \right) \leq 3k$.

Therefore $0 \leq d(\eta_1, \eta_2) \leq 1$, the proof is completed.

- (P2) if $d(\eta_1, \eta_2) = 0$, then $\eta_1 = \eta_2$

$$\begin{aligned}
 d(\eta_1, \eta_2) &= \frac{1}{3k} \left(|\Delta^{-1}(s_{\alpha_1}, \phi_1) - \Delta^{-1}(s_{\alpha_2}, \phi_2)| + |\Delta^{-1}(s_{\beta_1}, \varphi_1) - \Delta^{-1}(s_{\beta_2}, \varphi_2)| + |\Delta^{-1}(s_{\gamma_1}, \gamma_1) - \Delta^{-1}(s_{\gamma_2}, \gamma_2)| \right) = 0 \\
 &\Rightarrow \left(|\Delta^{-1}(s_{\alpha_1}, \phi_1) - \Delta^{-1}(s_{\alpha_2}, \phi_2)| = 0, |\Delta^{-1}(s_{\beta_1}, \varphi_1) - \Delta^{-1}(s_{\beta_2}, \varphi_2)| = 0, |\Delta^{-1}(s_{\gamma_1}, \gamma_1) - \Delta^{-1}(s_{\gamma_2}, \gamma_2)| = 0 \right) \\
 &\Rightarrow \left(\Delta^{-1}(s_{\alpha_1}, \phi_1) = \Delta^{-1}(s_{\alpha_2}, \phi_2), \Delta^{-1}(s_{\beta_1}, \varphi_1) = \Delta^{-1}(s_{\beta_2}, \varphi_2), \Delta^{-1}(s_{\gamma_1}, \gamma_1) = \Delta^{-1}(s_{\gamma_2}, \gamma_2) \right)
 \end{aligned}$$

That means $\eta_1 = \eta_2$, so (P2) if $d(\eta_1, \eta_2) = 0$, then $\eta_1 = \eta_2$ is right.

(P3) $d(\eta_1, \eta_2) = d(\eta_2, \eta_1)$

$$\begin{aligned}
 d(\eta_1, \eta_2) &= \frac{1}{3k} \left(|\Delta^{-1}(s_{\alpha_1}, \phi_1) - \Delta^{-1}(s_{\alpha_2}, \phi_2)| + |\Delta^{-1}(s_{\beta_1}, \varphi_1) - \Delta^{-1}(s_{\beta_2}, \varphi_2)| + |\Delta^{-1}(s_{\gamma_1}, \gamma_1) - \Delta^{-1}(s_{\gamma_2}, \gamma_2)| \right) \\
 &= \frac{1}{3k} \left(|\Delta^{-1}(s_{\alpha_2}, \phi_2) - \Delta^{-1}(s_{\alpha_1}, \phi_1)| + |\Delta^{-1}(s_{\beta_2}, \varphi_2) - \Delta^{-1}(s_{\beta_1}, \varphi_1)| + |\Delta^{-1}(s_{\gamma_2}, \gamma_2) - \Delta^{-1}(s_{\gamma_1}, \gamma_1)| \right) = d(\eta_2, \eta_1)
 \end{aligned}$$

So we complete the proof. (P3) $d(\eta_1, \eta_2) = d(\eta_2, \eta_1)$ is hold.

(P4) $d(\eta_1, \eta_2) + d(\eta_2, \eta_3) \geq d(\eta_1, \eta_3)$

$$\begin{aligned}
 d(\eta_1, \eta_2) &= \frac{1}{3k} \left(|\Delta^{-1}(s_{\alpha_1}, \phi_1) - \Delta^{-1}(s_{\alpha_3}, \phi_3)| + |\Delta^{-1}(s_{\beta_1}, \varphi_1) - \Delta^{-1}(s_{\beta_3}, \varphi_3)| \right. \\
 &\quad \left. + |\Delta^{-1}(s_{\gamma_1}, \gamma_1) - \Delta^{-1}(s_{\gamma_3}, \gamma_3)| \right) \\
 &= \frac{1}{3k} \left(|\Delta^{-1}(s_{\alpha_1}, \phi_1) - \Delta^{-1}(s_{\alpha_2}, \phi_2) + \Delta^{-1}(s_{\alpha_2}, \phi_2) - \Delta^{-1}(s_{\alpha_3}, \phi_3)| \right. \\
 &\quad \left. + |\Delta^{-1}(s_{\beta_1}, \varphi_1) - \Delta^{-1}(s_{\beta_2}, \varphi_2) + \Delta^{-1}(s_{\beta_2}, \varphi_2) - \Delta^{-1}(s_{\beta_3}, \varphi_3)| \right. \\
 &\quad \left. + |\Delta^{-1}(s_{\gamma_1}, \gamma_1) - \Delta^{-1}(s_{\gamma_2}, \gamma_2) + \Delta^{-1}(s_{\gamma_2}, \gamma_2) - \Delta^{-1}(s_{\gamma_3}, \gamma_3)| \right) \\
 &\leq \frac{1}{3k} \left(|\Delta^{-1}(s_{\alpha_1}, \phi_1) - \Delta^{-1}(s_{\alpha_2}, \phi_2)| + |\Delta^{-1}(s_{\alpha_2}, \phi_2) - \Delta^{-1}(s_{\alpha_3}, \phi_3)| \right. \\
 &\quad \left. + |\Delta^{-1}(s_{\beta_1}, \varphi_1) - \Delta^{-1}(s_{\beta_2}, \varphi_2)| + |\Delta^{-1}(s_{\beta_2}, \varphi_2) - \Delta^{-1}(s_{\beta_3}, \varphi_3)| \right. \\
 &\quad \left. + |\Delta^{-1}(s_{\gamma_1}, \gamma_1) - \Delta^{-1}(s_{\gamma_2}, \gamma_2)| + |\Delta^{-1}(s_{\gamma_2}, \gamma_2) - \Delta^{-1}(s_{\gamma_3}, \gamma_3)| \right) \\
 &= d(\eta_1, \eta_2) + d(\eta_2, \eta_3)
 \end{aligned}$$

2.3. The aggregation operators of 2TLNNs

Definition 5. (Wang, Wei et al., 2018). Let $\eta_j = \{(s_{\alpha_j}, \phi_j), (s_{\beta_j}, \varphi_j), (s_{\gamma_j}, \gamma_j)\}$ ($j = 1, 2, \dots, n$) be a group of 2TLNNs, then the 2TLNNWA and 2TLNNWG operators are defined as follows.

$$2TLNNWA(\eta_1, \eta_2, \dots, \eta_n) = \omega_1 \eta_1 \oplus \omega_2 \eta_2 \cdots \oplus \omega_n \eta_n = \bigoplus_{j=1}^n \omega_j \eta_j \tag{8}$$

and

$$2TLNNWG(\eta_1, \eta_2, \dots, \eta_n) = (\eta_1)^{\omega_1} \otimes (\eta_2)^{\omega_2} \cdots \otimes (\eta_n)^{\omega_n} = \bigotimes_{j=1}^n (\eta_j)^{\omega_j} \tag{9}$$

where ω_j is weighting vector of $\eta_j, j = 1, 2, \dots, n$. which satisfies $0 \leq \omega_j \leq 1, \sum_{j=1}^n \omega_j = 1$.

Theorem 2. (Wang, Wei et al., 2018) Let $\eta_j = \{(s_{\alpha_j}, \phi_j), (s_{\beta_j}, \varphi_j), (s_{\gamma_j}, \gamma_j)\}$ ($j = 1, 2, \dots, n$) be a group of 2TLNNs, then the operation results by 2TLNNWA and 2TLNNWG operators are also a 2TLNN where

$$\begin{aligned}
 2TLNNWA(\eta_1, \eta_2, \dots, \eta_n) &= \bigoplus_{j=1}^n \omega_j \eta_j \\
 &= \left\langle \begin{aligned} &\Delta \left(k \left(1 - \prod_{j=1}^n \left(1 - \frac{\Delta^{-1}(s_{\alpha_j}, \phi_j)}{k} \right)^{\omega_j} \right) \right), \Delta \left(k \prod_{j=1}^n \left(\frac{\Delta^{-1}(s_{\beta_j}, \varphi_j)}{k} \right)^{\omega_j} \right), \\ &\Delta \left(k \prod_{j=1}^n \left(\frac{\Delta^{-1}(s_{\gamma_j}, \gamma_j)}{k} \right)^{\omega_j} \right). \end{aligned} \right\rangle \quad (10)
 \end{aligned}$$

and

$$\begin{aligned}
 2TLNNWG(\eta_1, \eta_2, \dots, \eta_n) &= \bigotimes_{j=1}^n (\eta_j)^{\omega_j} \\
 &= \left\langle \begin{aligned} &\Delta \left(k \prod_{j=1}^n \left(\frac{\Delta^{-1}(s_{\alpha_j}, \phi_j)}{k} \right)^{\omega_j} \right), \Delta \left(k \left(1 - \prod_{j=1}^n \left(1 - \frac{\Delta^{-1}(s_{\beta_j}, \varphi_j)}{k} \right)^{\omega_j} \right) \right), \\ &\Delta \left(k \left(1 - \prod_{j=1}^n \left(1 - \frac{\Delta^{-1}(s_{\gamma_j}, \gamma_j)}{k} \right)^{\omega_j} \right) \right). \end{aligned} \right\rangle \quad (11)
 \end{aligned}$$

3. The VIKOR model for 2TLNNs MCGDM problems

Assume that $\{A_1, A_2, \dots, A_m\}$ be a group of alternatives, $\{D_1, D_2, \dots, D_t\}$ be a list of experts with weighting vector be $\{v_1, v_2, \dots, v_t\}$, and $\{G_1, G_2, \dots, G_n\}$ be a list of criteria with weighting vector be $\{\omega_1, \omega_2, \dots, \omega_n\}$, thereby satisfying $\omega_i \in [0, 1]$, $v_i \in [0, 1]$ and $\sum_{i=1}^n \omega_i = 1$, $\sum_{i=1}^t v_i = 1$. Construct the evaluation matrix $\eta^\lambda = [\eta_{ij}^\lambda]_{m \times n}$, $i = 1, 2, \dots, m, j = 1, 2, \dots, n, \lambda = 1, 2, \dots, t$, where $\eta_{ij}^\lambda = \{(s_{\alpha_{ij}}, \phi_{ij})^\lambda, (s_{\beta_{ij}}, \varphi_{ij})^\lambda, (s_{\gamma_{ij}}, \gamma_{ij})^\lambda\}$ means the estimate results of the alternative A_i ($i = 1, 2, \dots, m$) based on the criterion G_j ($j = 1, 2, \dots, n$) by expert D^λ ($\lambda = 1, 2, \dots, t$). $\Delta^{-1}(s_{\alpha_{ij}}, \phi_{ij})^\lambda \in [0, k]$ denotes the degree of truth-membership (TMD), $\Delta^{-1}(s_{\beta_{ij}}, \varphi_{ij})^\lambda \in [0, k]$ denotes the degree of indeterminacy-membership (IMD) and $\Delta^{-1}(s_{\gamma_{ij}}, \gamma_{ij})^\lambda \in [0, k]$ denotes the degree of falsity-membership (FMD) $0 \leq \Delta^{-1}(s_{\alpha_{ij}}, \phi_{ij})^\lambda + \Delta^{-1}(s_{\beta_{ij}}, \varphi_{ij})^\lambda + \Delta^{-1}(s_{\gamma_{ij}}, \gamma_{ij})^\lambda \leq 3k$, ($i = 1, 2, \dots, m, j = 1, 2, \dots, n, \lambda = 1, 2, \dots, t$).

Consider both the 2TLNNs theories and the traditional VIKOR model; we try to propose the VIKOR method with 2TLNNs to study MCGDM problems effectively. The method can be depicted as follows:

Step 1. Construct the decision matrix $\eta^\lambda = [\eta_{ij}^\lambda]_{m \times n}$, and utilize overall values of $\eta^\lambda = [\eta_{ij}^\lambda]_{m \times n}$ to $\eta = [\eta_{ij}]_{m \times n}$ by using equal (10) or (11);

Step 2. Compute the positive ideal solution (PIS) A^+ and the negative ideal solution (NIS) A^- ;

$$A^+ = \left\{ \Delta^{-1}(s_{\alpha_j}, \phi_j)^+, \Delta^{-1}(s_{\beta_j}, \varphi_j)^+, \Delta^{-1}(s_{\gamma_j}, \gamma_j)^+ \right\} (j = 1, 2, \dots, n) \tag{12}$$

$$A = \left\{ \Delta^{-1}(s_{\alpha_j}, \phi_j)^-, \Delta^{-1}(s_{\beta_j}, \varphi_j)^-, \Delta^{-1}(s_{\gamma_j}, \gamma_j)^- \right\} (j = 1, 2, \dots, n) \tag{13}$$

For benefit attribute

$$\begin{aligned} A^+ &= \left\{ \Delta^{-1}(s_{\alpha_j}, \phi_j)^+, \Delta^{-1}(s_{\beta_j}, \varphi_j)^+, \Delta^{-1}(s_{\gamma_j}, \gamma_j)^+ \right\} \\ &= \left\{ \max_i \left(\Delta^{-1}(s_{\alpha_j}, \phi_j) \right), \min_i \left(\Delta^{-1}(s_{\beta_j}, \varphi_j) \right), \min_i \left(\Delta^{-1}(s_{\gamma_j}, \gamma_j) \right) \right\} \end{aligned} \tag{14}$$

$$\begin{aligned} A^- &= \left\{ \Delta^{-1}(s_{\alpha_j}, \phi_j)^-, \Delta^{-1}(s_{\beta_j}, \varphi_j)^-, \Delta^{-1}(s_{\gamma_j}, \gamma_j)^- \right\} \\ &= \left\{ \min_i \left(\Delta^{-1}(s_{\alpha_j}, \phi_j) \right), \max_i \left(\Delta^{-1}(s_{\beta_j}, \varphi_j) \right), \max_i \left(\Delta^{-1}(s_{\gamma_j}, \gamma_j) \right) \right\} \end{aligned} \tag{15}$$

For cost attribute

$$\begin{aligned} A^+ &= \left\{ \Delta^{-1}(s_{\alpha_j}, \phi_j)^+, \Delta^{-1}(s_{\beta_j}, \varphi_j)^+, \Delta^{-1}(s_{\gamma_j}, \gamma_j)^+ \right\} \\ &= \left\{ \min_i \left(\Delta^{-1}(s_{\alpha_j}, \phi_j) \right), \max_i \left(\Delta^{-1}(s_{\beta_j}, \varphi_j) \right), \max_i \left(\Delta^{-1}(s_{\gamma_j}, \gamma_j) \right) \right\} \end{aligned} \tag{16}$$

$$\begin{aligned} A^- &= \left\{ \Delta^{-1}(s_{\alpha_j}, \phi_j)^-, \Delta^{-1}(s_{\beta_j}, \varphi_j)^-, \Delta^{-1}(s_{\gamma_j}, \gamma_j)^- \right\} \\ &= \left\{ \max_i \left(\Delta^{-1}(s_{\alpha_j}, \phi_j) \right), \min_i \left(\Delta^{-1}(s_{\beta_j}, \varphi_j) \right), \min_i \left(\Delta^{-1}(s_{\gamma_j}, \gamma_j) \right) \right\} \end{aligned} \tag{17}$$

Step 3. Based on the Equation (7) and the attribute weighting vector ω_j , we can calculate the values of τ_i and ψ_i which express the average and the worst group scores of η_i .

$$\tau_i = \sum_{j=1}^n \omega_j \frac{d \left(\begin{array}{c} \left\{ \Delta^{-1}(s_{\alpha_j}, \phi_j)^+, \Delta^{-1}(s_{\beta_j}, \varphi_j)^+, \Delta^{-1}(s_{\gamma_j}, \gamma_j)^+ \right\}, \\ \left\{ \Delta^{-1}(s_{\alpha_j}, \phi_j), \Delta^{-1}(s_{\beta_j}, \varphi_j), \Delta^{-1}(s_{\gamma_j}, \gamma_j) \right\} \end{array} \right)}{d \left(\begin{array}{c} \left\{ \Delta^{-1}(s_{\alpha_j}, \phi_j)^+, \Delta^{-1}(s_{\beta_j}, \varphi_j)^+, \Delta^{-1}(s_{\gamma_j}, \gamma_j)^+ \right\}, \\ \left\{ \Delta^{-1}(s_{\alpha_j}, \phi_j)^-, \Delta^{-1}(s_{\beta_j}, \varphi_j)^-, \Delta^{-1}(s_{\gamma_j}, \gamma_j)^- \right\} \end{array} \right)} \tag{18}$$

$$\psi_i = \max_j \left\{ \omega_j \frac{d \left(\begin{matrix} \{ \Delta^{-1}(s_{x_j}, \phi_j)^+, \Delta^{-1}(s_{\beta_j}, \varphi_j)^+, \Delta^{-1}(s_{\gamma_j}, \gamma_j)^+ \}, \\ \{ \Delta^{-1}(s_{x_j}, \phi_j), \Delta^{-1}(s_{\beta_j}, \varphi_j), \Delta^{-1}(s_{\gamma_j}, \gamma_j) \} \end{matrix} \right)}{d \left(\begin{matrix} \{ \Delta^{-1}(s_{x_j}, \phi_j)^+, \Delta^{-1}(s_{\beta_j}, \varphi_j)^+, \Delta^{-1}(s_{\gamma_j}, \gamma_j)^+ \}, \\ \{ \Delta^{-1}(s_{x_j}, \phi_j)^-, \Delta^{-1}(s_{\beta_j}, \varphi_j)^-, \Delta^{-1}(s_{\gamma_j}, \gamma_j)^- \} \end{matrix} \right)} \right\} \quad (19)$$

where d is the normalized Hamming distance and $0 \leq \omega_j \leq 1$ means the weight of attributes which satisfies $\sum_{i=1}^n \omega_i = 1$.

Step 4. Compute the values of Q_i based on the results of τ_i and ψ_i , the calculating formula is characteristic as follows.

$$Q_i = \rho \frac{(\tau_i - \tau^+)}{(\tau^- - \tau^+)} + (1 - \rho) \frac{(\psi_i - \psi^+)}{(\psi^- - \psi^+)} \quad (20)$$

where

$$\tau^+ = \min_i \tau_i, \tau^- = \max_i \tau_i \quad (21)$$

$$\psi^+ = \min_i \psi_i, \psi^- = \max_i \psi_i \quad (22)$$

where ρ means the coefficient of decision making strategic. $\rho > 0.5$ depicts “the maximum group utility”, $\rho = 0.5$ depicts equality and $\rho < 0.5$ depicts the minimum regret.

Step 5. To choose the best alternative by rank the values of Q_i , the alternative with minimum value is the best choice.

4. The VIKOR method for IV2TLNNs MCDM problems

4.1. The IV2TLNSs

To solve MCDM problems more effectively, we extend the 2TLNSs to interval-valued environment to propose the IV2TLNSs as follows.

Definition 6. Let s_1, s_2, \dots, s_k be a linguistic term set. Any label s_i shows a possible linguistic variable, the IV2TLNSs $\tilde{\eta}$ can be depicted as:

$$\tilde{\eta} = \left\{ [(s_x, \phi)^L, (s_x, \phi)^U], [(s_\beta, \varphi)^L, (s_\beta, \varphi)^U], [(s_\gamma, \gamma)^L, (s_\gamma, \gamma)^U] \right\} \quad (23)$$

where $[\Delta^{-1}(s_x, \phi)^L, \Delta^{-1}(s_x, \phi)^U]$, $[\Delta^{-1}(s_\beta, \varphi)^L, \Delta^{-1}(s_\beta, \varphi)^U]$ and $[\Delta^{-1}(s_\gamma, \gamma)^L, \Delta^{-1}(s_\gamma, \gamma)^U] \in [0, k]$ represent the degree of the truth membership, the indeterminacy membership

and the falsity membership which are expressed by IV2TLNNs and satisfies the condition $0 \leq \Delta^{-1}(s_{\alpha}, \phi)^U + \Delta^{-1}(s_{\beta}, \varphi)^U + \Delta^{-1}(s_{\gamma}, \gamma)^U \leq 3k$.

Definition 7. Let $\tilde{\eta} = \{[(s_{\alpha}, \phi)^L, (s_{\alpha}, \phi)^U], [(s_{\beta}, \varphi)^L, (s_{\beta}, \varphi)^U], [(s_{\gamma}, \gamma)^L, (s_{\gamma}, \gamma)^U]\}$ be an IV2TLNN, the score and accuracy functions of $\tilde{\eta}$ can be expressed:

$$s(\tilde{\eta}) = \frac{1}{6k} \left\{ \begin{aligned} & \left(2k + \Delta^{-1}(s_{\alpha}, \phi)^L - \Delta^{-1}(s_{\beta}, \varphi)^L - \Delta^{-1}(s_{\gamma}, \gamma)^L \right) \\ & + \left(2k + \Delta^{-1}(s_{\alpha}, \phi)^U - \Delta^{-1}(s_{\beta}, \varphi)^U - \Delta^{-1}(s_{\gamma}, \gamma)^U \right) \end{aligned} \right\}, s(\tilde{\eta}) \in [0, 1] \quad (24)$$

$$h(\tilde{\eta}) = \frac{\left(\Delta^{-1}(s_{\alpha}, \phi)^L - \Delta^{-1}(s_{\gamma}, \gamma)^L \right) + \left(\Delta^{-1}(s_{\alpha}, \phi)^U - \Delta^{-1}(s_{\gamma}, \gamma)^U \right)}{2}, h(\tilde{\eta}) \in [-k, k] \quad (25)$$

For two IV2TLNNs $\tilde{\eta}_1$ and $\tilde{\eta}_2$, based on the Definition 7, then

- (1) if $s(\tilde{\eta}_1) < s(\tilde{\eta}_2)$, then $\tilde{\eta}_1 < \tilde{\eta}_2$;
- (2) if $s(\tilde{\eta}_1) > s(\tilde{\eta}_2)$, then $\tilde{\eta}_1 > \tilde{\eta}_2$;
- (3) if $s(\tilde{\eta}_1) = s(\tilde{\eta}_2)$, $h(\tilde{\eta}_1) < h(\tilde{\eta}_2)$, then $\tilde{\eta}_1 < \tilde{\eta}_2$;
- (4) if $s(\tilde{\eta}_1) = s(\tilde{\eta}_2)$, $h(\tilde{\eta}_1) > h(\tilde{\eta}_2)$, then $\tilde{\eta}_1 > \tilde{\eta}_2$;
- (5) if $s(\tilde{\eta}_1) = s(\tilde{\eta}_2)$, $h(\tilde{\eta}_1) = h(\tilde{\eta}_2)$, then $\tilde{\eta}_1 = \tilde{\eta}_2$.

Definition 8. Let $\tilde{\eta}_1 = \{[(s_{\alpha_1}, \phi_1)^L, (s_{\alpha_1}, \phi_1)^U], [(s_{\beta_1}, \varphi_1)^L, (s_{\beta_1}, \varphi_1)^U], [(s_{\gamma_1}, \gamma_1)^L, (s_{\gamma_1}, \gamma_1)^U]\}$ and $\tilde{\eta}_2 = \{[(s_{\alpha_2}, \phi_2)^L, (s_{\alpha_2}, \phi_2)^U], [(s_{\beta_2}, \varphi_2)^L, (s_{\beta_2}, \varphi_2)^U], [(s_{\gamma_2}, \gamma_2)^L, (s_{\gamma_2}, \gamma_2)^U]\}$ be two IV2TLNNs, then we can get the normalized Hamming distance:

$$d(\tilde{\eta}_1, \tilde{\eta}_2) = \frac{1}{6k} \left(\begin{aligned} & \left| \Delta^{-1}(s_{\alpha_1}, \phi_1)^L - \Delta^{-1}(s_{\alpha_2}, \phi_2)^L \right| + \left| \Delta^{-1}(s_{\beta_1}, \varphi_1)^L - \Delta^{-1}(s_{\beta_2}, \varphi_2)^L \right| \\ & + \left| \Delta^{-1}(s_{\gamma_1}, \gamma_1)^L - \Delta^{-1}(s_{\gamma_2}, \gamma_2)^L \right| + \left| \Delta^{-1}(s_{\alpha_1}, \phi_1)^U - \Delta^{-1}(s_{\alpha_2}, \phi_2)^U \right| \\ & + \left| \Delta^{-1}(s_{\beta_1}, \varphi_1)^U - \Delta^{-1}(s_{\beta_2}, \varphi_2)^U \right| + \left| \Delta^{-1}(s_{\gamma_1}, \gamma_1)^U - \Delta^{-1}(s_{\gamma_2}, \gamma_2)^U \right| \end{aligned} \right) \quad (26)$$

Theorem 3. Let $\tilde{\eta}_1 = \{[(s_{\alpha_1}, \phi_1)^L, (s_{\alpha_1}, \phi_1)^U], [(s_{\beta_1}, \varphi_1)^L, (s_{\beta_1}, \varphi_1)^U], [(s_{\gamma_1}, \gamma_1)^L, (s_{\gamma_1}, \gamma_1)^U]\}$, $\tilde{\eta}_2 = \{[(s_{\alpha_2}, \phi_2)^L, (s_{\alpha_2}, \phi_2)^U], [(s_{\beta_2}, \varphi_2)^L, (s_{\beta_2}, \varphi_2)^U], [(s_{\gamma_2}, \gamma_2)^L, (s_{\gamma_2}, \gamma_2)^U]\}$ and $\tilde{\eta}_3 = \{[(s_{\alpha_3}, \phi_3)^L, (s_{\alpha_3}, \phi_3)^U], [(s_{\beta_3}, \varphi_3)^L, (s_{\beta_3}, \varphi_3)^U], [(s_{\gamma_3}, \gamma_3)^L, (s_{\gamma_3}, \gamma_3)^U]\}$, the Hamming distance d also has the following properties:

- (P1) $0 \leq d(\tilde{\eta}_1, \tilde{\eta}_2) \leq 1$; (P2) if $d(\tilde{\eta}_1, \tilde{\eta}_2) = 0$, then $\tilde{\eta}_1 = \tilde{\eta}_2$;
 (P3) $d(\tilde{\eta}_1, \tilde{\eta}_2) = d(\tilde{\eta}_2, \tilde{\eta}_1)$; (P4) $d(\tilde{\eta}_1, \tilde{\eta}_2) + d(\tilde{\eta}_2, \tilde{\eta}_3) \geq d(\tilde{\eta}_1, \tilde{\eta}_3)$.

4.2. The aggregation operators of IV2TLNNs

Definition 9. Let $\tilde{\eta}_j = \{[(s_{\alpha_j}, \phi_j)^L, (s_{\alpha_j}, \phi_j)^U], [(s_{\beta_j}, \varphi_j)^L, (s_{\beta_j}, \varphi_j)^U], [(s_{\gamma_j}, \gamma_j)^L, (s_{\gamma_j}, \gamma_j)^U]\}$ be a group of IV2TLNNs, then the IV2TLNNWA and IV2TLNNWG operators can be defined as follows.

$$\begin{aligned}
 & \text{IV2TLNNA}(\tilde{\eta}_1, \tilde{\eta}_2, \dots, \tilde{\eta}_n) = \bigoplus_{j=1}^n \omega_j \tilde{\eta}_j \\
 & = \left\{ \begin{aligned} & \left[\Delta \left(k \left(1 - \prod_{j=1}^n \left(1 - \frac{\Delta^{-1}(s_{\alpha_j}, \phi_j)^L}{k} \right)^{w_j} \right) \right), \Delta \left(k \left(1 - \prod_{j=1}^n \left(1 - \frac{\Delta^{-1}(s_{\alpha_j}, \phi_j)^U}{k} \right)^{w_j} \right) \right) \right], \\ & \left[\Delta \left(k \prod_{j=1}^n \left(\frac{\Delta^{-1}(s_{\beta_j}, \varphi_j)^L}{k} \right)^{w_j} \right), \Delta \left(k \prod_{j=1}^n \left(\frac{\Delta^{-1}(s_{\beta_j}, \varphi_j)^U}{k} \right)^{w_j} \right) \right], \\ & \left[\Delta \left(k \prod_{j=1}^n \left(\frac{\Delta^{-1}(s_{\gamma_j}, \gamma_j)^L}{k} \right)^{w_j} \right), \Delta \left(k \prod_{j=1}^n \left(\frac{\Delta^{-1}(s_{\gamma_j}, \gamma_j)^U}{k} \right)^{w_j} \right) \right]. \end{aligned} \right. \quad (27)
 \end{aligned}$$

and

$$\begin{aligned}
 & \text{IV2TLNNG}(\tilde{\eta}_1, \tilde{\eta}_2, \dots, \tilde{\eta}_n) = \bigotimes_{j=1}^n (\tilde{\eta}_j)^{\omega_j} \\
 & = \left\{ \begin{aligned} & \left[\Delta \left(k \prod_{j=1}^n \left(\frac{\Delta^{-1}(s_{\alpha_j}, \phi_j)^L}{k} \right)^{w_j} \right), \Delta \left(k \prod_{j=1}^n \left(\frac{\Delta^{-1}(s_{\alpha_j}, \phi_j)^U}{k} \right)^{w_j} \right) \right], \\ & \left[\Delta \left(k \left(1 - \prod_{j=1}^n \left(1 - \frac{\Delta^{-1}(s_{\beta_j}, \varphi_j)^L}{k} \right)^{w_j} \right) \right), \Delta \left(k \left(1 - \prod_{j=1}^n \left(1 - \frac{\Delta^{-1}(s_{\beta_j}, \varphi_j)^U}{k} \right)^{w_j} \right) \right) \right], \\ & \left[\Delta \left(k \left(1 - \prod_{j=1}^n \left(1 - \frac{\Delta^{-1}(s_{\gamma_j}, \gamma_j)^L}{k} \right)^{w_j} \right) \right), \Delta \left(k \left(1 - \prod_{j=1}^n \left(1 - \frac{\Delta^{-1}(s_{\gamma_j}, \gamma_j)^U}{k} \right)^{w_j} \right) \right) \right]. \end{aligned} \right. \quad (28)
 \end{aligned}$$

4.3. Computing steps for MCGDM problems with IV2TLNNS

Assume that $\{A_1, A_2, \dots, A_m\}$ be a group of alternatives and $\{G_1, G_2, \dots, G_n\}$ be a list of criteria with weighting vector be $\{\omega_1, \omega_2, \dots, \omega_n\}$, thereby satisfying $\omega_i \in [0, 1]$ and $\sum_{i=1}^n \omega_i = 1$. Construct the evaluation matrix $\tilde{\eta} = [\tilde{\eta}_{ij}]_{m \times n}$, $i = 1, 2, \dots, m, j = 1, 2, \dots, n$ where $\tilde{\eta}_{ij} = \{[(s_{\alpha_{ij}}, \phi_{ij})^L, (s_{\alpha_{ij}}, \phi_{ij})^U], [(s_{\beta_{ij}}, \varphi_{ij})^L, (s_{\beta_{ij}}, \varphi_{ij})^U], [(s_{\gamma_{ij}}, \gamma_{ij})^L, (s_{\gamma_{ij}}, \gamma_{ij})^U]\}$ means the estimate results of the alternative $A_i (i = 1, 2, \dots, m)$ based on the criterion $G_j (j = 1, 2, \dots, n)$. The calculating steps also can be depicted as follows:

Step 1. Construct the decision matrix $\tilde{\eta} = [\tilde{\eta}_{ij}]_{m \times n}$;

Step 2. Compute the positive ideal solution A^+ and the negative ideal solution A^- ;

$$A^+ = \left\{ \begin{aligned} & \left[\left(\Delta^{-1}(s_{\alpha_j}, \phi_j)^L \right)^+, \left(\Delta^{-1}(s_{\alpha_j}, \phi_j)^U \right)^+ \right], \\ & \left[\left(\Delta^{-1}(s_{\beta_j}, \varphi_j)^L \right)^+, \left(\Delta^{-1}(s_{\beta_j}, \varphi_j)^U \right)^+ \right], \\ & \left[\left(\Delta^{-1}(s_{\gamma_j}, \gamma_j)^L \right)^+, \left(\Delta^{-1}(s_{\gamma_j}, \gamma_j)^U \right)^+ \right] \end{aligned} \right\} (j = 1, 2, \dots, n) \quad (29)$$

$$A^- = \left\{ \begin{array}{l} \left[\left(\Delta^{-1}(s_{\alpha_j}, \phi_j)^L \right)^-, \left(\Delta^{-1}(s_{\alpha_j}, \phi_j)^U \right)^- \right], \\ \left[\left(\Delta^{-1}(s_{\beta_j}, \varphi_j)^L \right)^-, \left(\Delta^{-1}(s_{\beta_j}, \varphi_j)^U \right)^- \right], \\ \left[\left(\Delta^{-1}(s_{\gamma_j}, \gamma_j)^L \right)^-, \left(\Delta^{-1}(s_{\gamma_j}, \gamma_j)^U \right)^- \right] \end{array} \right\} \quad (j = 1, 2, \dots, n) \quad (30)$$

For benefit attribute

$$\left\{ \begin{array}{l} \left[\left(\Delta^{-1}(s_{\alpha_j}, \phi_j)^L \right)^+, \left(\Delta^{-1}(s_{\alpha_j}, \phi_j)^U \right)^+ \right], \\ \left[\left(\Delta^{-1}(s_{\beta_j}, \varphi_j)^L \right)^+, \left(\Delta^{-1}(s_{\beta_j}, \varphi_j)^U \right)^+ \right], \\ \left[\left(\Delta^{-1}(s_{\gamma_j}, \gamma_j)^L \right)^+, \left(\Delta^{-1}(s_{\gamma_j}, \gamma_j)^U \right)^+ \right] \end{array} \right\} = \left\{ \begin{array}{l} \left[\max_i \left(\Delta^{-1}(s_{\alpha_j}, \phi_j)^L \right), \max_i \left(\Delta^{-1}(s_{\alpha_j}, \phi_j)^U \right) \right], \\ \left[\min_i \left(\Delta^{-1}(s_{\beta_j}, \varphi_j)^L \right), \min_i \left(\Delta^{-1}(s_{\beta_j}, \varphi_j)^U \right) \right], \\ \left[\min_i \left(\Delta^{-1}(s_{\gamma_j}, \gamma_j)^L \right), \min_i \left(\Delta^{-1}(s_{\gamma_j}, \gamma_j)^U \right) \right] \end{array} \right\} \quad (31)$$

$$\left\{ \begin{array}{l} \left[\left(\Delta^{-1}(s_{\alpha_j}, \phi_j)^L \right)^-, \left(\Delta^{-1}(s_{\alpha_j}, \phi_j)^U \right)^- \right], \\ \left[\left(\Delta^{-1}(s_{\beta_j}, \varphi_j)^L \right)^-, \left(\Delta^{-1}(s_{\beta_j}, \varphi_j)^U \right)^- \right], \\ \left[\left(\Delta^{-1}(s_{\gamma_j}, \gamma_j)^L \right)^-, \left(\Delta^{-1}(s_{\gamma_j}, \gamma_j)^U \right)^- \right] \end{array} \right\} = \left\{ \begin{array}{l} \left[\min_i \left(\Delta^{-1}(s_{\alpha_j}, \phi_j)^L \right), \min_i \left(\Delta^{-1}(s_{\alpha_j}, \phi_j)^U \right) \right], \\ \left[\max_i \left(\Delta^{-1}(s_{\beta_j}, \varphi_j)^L \right), \max_i \left(\Delta^{-1}(s_{\beta_j}, \varphi_j)^U \right) \right], \\ \left[\max_i \left(\Delta^{-1}(s_{\gamma_j}, \gamma_j)^L \right), \max_i \left(\Delta^{-1}(s_{\gamma_j}, \gamma_j)^U \right) \right] \end{array} \right\} \quad (32)$$

For cost attribute

$$\left\{ \begin{array}{l} \left[\left(\Delta^{-1}(s_{\alpha_j}, \phi_j)^L \right)^+, \left(\Delta^{-1}(s_{\alpha_j}, \phi_j)^U \right)^+ \right], \\ \left[\left(\Delta^{-1}(s_{\beta_j}, \varphi_j)^L \right)^+, \left(\Delta^{-1}(s_{\beta_j}, \varphi_j)^U \right)^+ \right], \\ \left[\left(\Delta^{-1}(s_{\gamma_j}, \gamma_j)^L \right)^+, \left(\Delta^{-1}(s_{\gamma_j}, \gamma_j)^U \right)^+ \right] \end{array} \right\} = \left\{ \begin{array}{l} \left[\min_i \left(\Delta^{-1}(s_{\alpha_j}, \phi_j)^L \right), \min_i \left(\Delta^{-1}(s_{\alpha_j}, \phi_j)^U \right) \right], \\ \left[\max_i \left(\Delta^{-1}(s_{\beta_j}, \varphi_j)^L \right), \max_i \left(\Delta^{-1}(s_{\beta_j}, \varphi_j)^U \right) \right], \\ \left[\max_i \left(\Delta^{-1}(s_{\gamma_j}, \gamma_j)^L \right), \max_i \left(\Delta^{-1}(s_{\gamma_j}, \gamma_j)^U \right) \right] \end{array} \right\} \quad (33)$$

$$\left\{ \begin{array}{l} \left[\left(\Delta^{-1}(s_{\alpha_j}, \phi_j)^L \right)^-, \left(\Delta^{-1}(s_{\alpha_j}, \phi_j)^U \right)^- \right], \\ \left[\left(\Delta^{-1}(s_{\beta_j}, \varphi_j)^L \right)^-, \left(\Delta^{-1}(s_{\beta_j}, \varphi_j)^U \right)^- \right], \\ \left[\left(\Delta^{-1}(s_{\gamma_j}, \gamma_j)^L \right)^-, \left(\Delta^{-1}(s_{\gamma_j}, \gamma_j)^U \right)^- \right] \end{array} \right\} = \left\{ \begin{array}{l} \left[\max_i \left(\Delta^{-1}(s_{\alpha_j}, \phi_j)^L \right), \max_i \left(\Delta^{-1}(s_{\alpha_j}, \phi_j)^U \right) \right], \\ \left[\min_i \left(\Delta^{-1}(s_{\beta_j}, \varphi_j)^L \right), \min_i \left(\Delta^{-1}(s_{\beta_j}, \varphi_j)^U \right) \right], \\ \left[\min_i \left(\Delta^{-1}(s_{\gamma_j}, \gamma_j)^L \right), \min_i \left(\Delta^{-1}(s_{\gamma_j}, \gamma_j)^U \right) \right] \end{array} \right\} \quad (34)$$

Step 3. Based on the Equation (26) and the attribute weighting vector ω_j , we can calculate the values of $\tilde{\tau}_i$ and $\tilde{\psi}_i$ which express the average and the worst group scores of $\tilde{\eta}_i$.

$$\tilde{\tau}_i = \sum_{j=1}^n \omega_j \frac{d \left(\left\{ \left[\begin{array}{l} (\Delta^{-1}(s_{z_j}, \phi_j)^L)^+, (\Delta^{-1}(s_{z_j}, \phi_j)^U)^+ \\ (\Delta^{-1}(s_{\beta_j}, \varphi_j)^L)^+, (\Delta^{-1}(s_{\beta_j}, \varphi_j)^U)^+ \\ (\Delta^{-1}(s_{\gamma_j}, \gamma_j)^L)^+, (\Delta^{-1}(s_{\gamma_j}, \gamma_j)^U)^+ \end{array} \right] \right\}, \left\{ \left[\begin{array}{l} \Delta^{-1}(s_{z_j}, \phi_j)^L, \Delta^{-1}(s_{z_j}, \phi_j)^U \\ \Delta^{-1}(s_{\beta_j}, \varphi_j)^L, \Delta^{-1}(s_{\beta_j}, \varphi_j)^U \\ \Delta^{-1}(s_{\gamma_j}, \gamma_j)^L, \Delta^{-1}(s_{\gamma_j}, \gamma_j)^U \end{array} \right] \right\} \right)}{d \left(\left\{ \left[\begin{array}{l} (\Delta^{-1}(s_{z_j}, \phi_j)^L)^+, (\Delta^{-1}(s_{z_j}, \phi_j)^U)^+ \\ (\Delta^{-1}(s_{\beta_j}, \varphi_j)^L)^+, (\Delta^{-1}(s_{\beta_j}, \varphi_j)^U)^+ \\ (\Delta^{-1}(s_{\gamma_j}, \gamma_j)^L)^+, (\Delta^{-1}(s_{\gamma_j}, \gamma_j)^U)^+ \end{array} \right] \right\}, \left\{ \left[\begin{array}{l} (\Delta^{-1}(s_{z_j}, \phi_j)^L)^-, (\Delta^{-1}(s_{z_j}, \phi_j)^U)^- \\ (\Delta^{-1}(s_{\beta_j}, \varphi_j)^L)^-, (\Delta^{-1}(s_{\beta_j}, \varphi_j)^U)^- \\ (\Delta^{-1}(s_{\gamma_j}, \gamma_j)^L)^-, (\Delta^{-1}(s_{\gamma_j}, \gamma_j)^U)^- \end{array} \right] \right\} \right)} \quad (35)$$

$$\tilde{\psi}_i = \max_j \left\{ \omega_j \frac{d \left(\left\{ \left[\begin{array}{l} (\Delta^{-1}(s_{z_j}, \phi_j)^L)^+, (\Delta^{-1}(s_{z_j}, \phi_j)^U)^+ \\ (\Delta^{-1}(s_{\beta_j}, \varphi_j)^L)^+, (\Delta^{-1}(s_{\beta_j}, \varphi_j)^U)^+ \\ (\Delta^{-1}(s_{\gamma_j}, \gamma_j)^L)^+, (\Delta^{-1}(s_{\gamma_j}, \gamma_j)^U)^+ \end{array} \right] \right\}, \left\{ \left[\begin{array}{l} \Delta^{-1}(s_{z_j}, \phi_j)^L, \Delta^{-1}(s_{z_j}, \phi_j)^U \\ \Delta^{-1}(s_{\beta_j}, \varphi_j)^L, \Delta^{-1}(s_{\beta_j}, \varphi_j)^U \\ \Delta^{-1}(s_{\gamma_j}, \gamma_j)^L, \Delta^{-1}(s_{\gamma_j}, \gamma_j)^U \end{array} \right] \right\} \right)}{d \left(\left\{ \left[\begin{array}{l} (\Delta^{-1}(s_{z_j}, \phi_j)^L)^+, (\Delta^{-1}(s_{z_j}, \phi_j)^U)^+ \\ (\Delta^{-1}(s_{\beta_j}, \varphi_j)^L)^+, (\Delta^{-1}(s_{\beta_j}, \varphi_j)^U)^+ \\ (\Delta^{-1}(s_{\gamma_j}, \gamma_j)^L)^+, (\Delta^{-1}(s_{\gamma_j}, \gamma_j)^U)^+ \end{array} \right] \right\}, \left\{ \left[\begin{array}{l} (\Delta^{-1}(s_{z_j}, \phi_j)^L)^-, (\Delta^{-1}(s_{z_j}, \phi_j)^U)^- \\ (\Delta^{-1}(s_{\beta_j}, \varphi_j)^L)^-, (\Delta^{-1}(s_{\beta_j}, \varphi_j)^U)^- \\ (\Delta^{-1}(s_{\gamma_j}, \gamma_j)^L)^-, (\Delta^{-1}(s_{\gamma_j}, \gamma_j)^U)^- \end{array} \right] \right\} \right)} \right\} \quad (36)$$

where d is the normalized Hamming distance and $0 \leq \omega_j \leq 1$ means the weight of attributes which satisfies $\sum_{i=1}^n \omega_i = 1$.

Step 4. Compute the values of Q_i based on the results of $\tilde{\tau}_i$ and $\tilde{\psi}_i$, the calculating formula is characteristic as follows.

$$Q_i = \rho \frac{(\tilde{\tau}_i - \tilde{\tau}^+)}{(\tilde{\tau}^- - \tilde{\tau}^+)} + (1 - \rho) \frac{(\tilde{\psi}_i - \tilde{\psi}^+)}{(\tilde{\psi}^- - \tilde{\psi}^+)} \quad (37)$$

where

$$\tilde{\tau}^+ = \min_i \tilde{\tau}_i, \tau^- = \max_i \tilde{\tau}_i \quad (38)$$

$$\tilde{\psi}^+ = \min_i \tilde{\psi}_i, \tilde{\psi}^- = \max_i \tilde{\psi}_i \quad (39)$$

where ρ means the coefficient of decision making strategic. $\rho > 0.5$ depicts “the maximum group utility”, $\rho = 0.5$ depicts equality and $\rho < 0.5$ depicts the minimum regret.

Step 5. To choose the best alternative by rank the values of Q_i , the alternative with minimum value is the best choice.

5. The numerical example

5.1. Numerical for 2TLNNs MCGDM problems

After China’s entering into the WTO, the economy has developed rapidly and it has held a high rate of economic growth. But the stamina of development is confronted with severe challenges: On the one hand, the international economic situation is continuously changing and many enterprises in China are limited by international green barriers; On the other hand, while enjoying great economic development achievements, people also realized that our country’s environment and resources are becoming more and more serious. While China’s economic development is growing at a high speed, the ecological environment and natural resources have been seriously injured and the contradiction between natural resource environment and social economic development has become increasingly obvious. Under the background of people’s urgent need for environmental protection and healthy living, many enterprises in China are aware of the necessity and importance of green health and low carbon environmental protection for the survival and development of enterprises. Green suppliers selection is a classical MADM problem. In this chapter, we provide a numerical example to select best green suppliers selection by using VIKOR method with 2TLNNs. Assume that five possible green suppliers $A_i(i = 1, 2, 3, 4, 5)$ to be selected and four criteria to assess these green suppliers: ① G_1 is the product quality factor; ② G_2 is environmental factors; ③ G_3 is delivery factor; ④ G_4 is price factor. The five possible green suppliers $A_i(i = 1, 2, 3, 4, 5)$ are to be evaluated with 2TLNNs with the four criteria by three experts (criteria weight $\omega = (0.32, 0.13, 0.35, 0.20)$, experts weight $\nu = (0.25, 0.35, 0.40)$.), which are given in Tables 1–3.

Step 1. Utilize overall values of $\eta^\lambda = [\eta_{ij}^\lambda]_{m \times n}$ to $\eta = [\eta_{ij}]_{m \times n}$ by using 2TLNNWA operator, the aggregation results are listed in Table 4.

Table 1. 2TLNNs evaluation matrix by the first expert.

	G_1	G_2	G_3	G_4
A_1	$\{(s_4,0), (s_2,0), (s_1,0)\}$	$\{(s_5,0), (s_3,0), (s_2,0)\}$	$\{(s_4,0), (s_1,0), (s_1,0)\}$	$\{(s_3,0), (s_2,0), (s_2,0)\}$
A_2	$\{(s_5,0), (s_4,0), (s_4,0)\}$	$\{(s_3,0), (s_4,0), (s_2,0)\}$	$\{(s_2,0), (s_1,0), (s_3,0)\}$	$\{(s_3,0), (s_1,0), (s_2,0)\}$
A_3	$\{(s_5,0), (s_4,0), (s_3,0)\}$	$\{(s_2,0), (s_4,0), (s_5,0)\}$	$\{(s_3,0), (s_3,0), (s_4,0)\}$	$\{(s_2,0), (s_1,0), (s_4,0)\}$
A_4	$\{(s_3,0), (s_2,0), (s_3,0)\}$	$\{(s_4,0), (s_3,0), (s_2,0)\}$	$\{(s_3,0), (s_3,0), (s_4,0)\}$	$\{(s_2,0), (s_1,0), (s_1,0)\}$
A_5	$\{(s_1,0), (s_4,0), (s_5,0)\}$	$\{(s_2,0), (s_3,0), (s_1,0)\}$	$\{(s_3,0), (s_4,0), (s_5,0)\}$	$\{(s_2,0), (s_4,0), (s_3,0)\}$

Table 2. 2TLNNs evaluation matrix by the second expert.

	G_1	G_2	G_3	G_4
η_1	$\{(s_5,0), (s_1,0), (s_2,0)\}$	$\{(s_4,0), (s_3,0), (s_1,0)\}$	$\{(s_4,0), (s_2,0), (s_1,0)\}$	$\{(s_5,0), (s_1,0), (s_2,0)\}$
η_2	$\{(s_4,0), (s_3,0), (s_3,0)\}$	$\{(s_3,0), (s_2,0), (s_4,0)\}$	$\{(s_2,0), (s_1,0), (s_3,0)\}$	$\{(s_5,0), (s_4,0), (s_2,0)\}$
η_3	$\{(s_3,0), (s_4,0), (s_3,0)\}$	$\{(s_2,0), (s_4,0), (s_5,0)\}$	$\{(s_5,0), (s_1,0), (s_2,0)\}$	$\{(s_2,0), (s_1,0), (s_2,0)\}$
η_4	$\{(s_4,0), (s_5,0), (s_4,0)\}$	$\{(s_2,0), (s_3,0), (s_2,0)\}$	$\{(s_3,0), (s_3,0), (s_4,0)\}$	$\{(s_4,0), (s_4,0), (s_5,0)\}$
η_5	$\{(s_2,0), (s_4,0), (s_5,0)\}$	$\{(s_3,0), (s_1,0), (s_5,0)\}$	$\{(s_2,0), (s_3,0), (s_4,0)\}$	$\{(s_2,0), (s_1,0), (s_3,0)\}$

Table 3. 2TLNNs evaluation matrix by the third expert.

	G ₁	G ₂	G ₃	G ₄
A ₁	{{(s ₅ ,0), (s ₁ ,0), (s ₁ ,0)}}	{{(s ₄ ,0), (s ₁ ,0), (s ₂ ,0)}}	{{(s ₃ ,0), (s ₃ ,0), (s ₁ ,0)}}	{{(s ₄ ,0), (s ₂ ,0), (s ₂ ,0)}}
A ₂	{{(s ₅ ,0), (s ₄ ,0), (s ₅ ,0)}}	{{(s ₃ ,0), (s ₂ ,0), (s ₁ ,0)}}	{{(s ₂ ,0), (s ₁ ,0), (s ₄ ,0)}}	{{(s ₄ ,0), (s ₅ ,0), (s ₃ ,0)}}
A ₃	{{(s ₂ ,0), (s ₁ ,0), (s ₄ ,0)}}	{{(s ₅ ,0), (s ₁ ,0), (s ₃ ,0)}}	{{(s ₄ ,0), (s ₃ ,0), (s ₄ ,0)}}	{{(s ₅ ,0), (s ₂ ,0), (s ₃ ,0)}}
A ₄	{{(s ₂ ,0), (s ₂ ,0), (s ₃ ,0)}}	{{(s ₄ ,0), (s ₁ ,0), (s ₂ ,0)}}	{{(s ₅ ,0), (s ₃ ,0), (s ₂ ,0)}}	{{(s ₁ ,0), (s ₄ ,0), (s ₅ ,0)}}
A ₅	{{(s ₁ ,0), (s ₄ ,0), (s ₅ ,0)}}	{{(s ₂ ,0), (s ₄ ,0), (s ₄ ,0)}}	{{(s ₃ ,0), (s ₄ ,0), (s ₃ ,0)}}	{{(s ₂ ,0), (s ₄ ,0), (s ₄ ,0)}}

Table 4. The aggregation values by 2TLNNWA operator.

	G ₁	G ₂
A ₁	{{(s ₅ , -0.1892), (s ₁ , 0.1892), (s ₁ , 0.2746)}}	{{(s ₄ , 0.3182), (s ₂ , -0.0668), (s ₂ , -0.4308)}}
A ₂	{{(s ₅ , -0.2746), (s ₄ , -0.3831), (s ₄ , -0.0455)}}	{{(s ₃ , 0.0000), (s ₂ , 0.3784), (s ₂ , -0.0681)}}
A ₃	{{(s ₃ , 0.4425), (s ₂ , 0.2974), (s ₃ , 0.3659)}}	{{(s ₄ , -0.2974), (s ₂ , 0.2974), (s ₄ , 0.0760)}}
A ₄	{{(s ₃ , 0.0794), (s ₃ , -0.2438), (s ₃ , 0.3178)}}	{{(s ₃ , 0.4509), (s ₂ , -0.0668), (s ₂ , 0.0000)}}
A ₅	{{(s ₁ , 0.3756), (s ₄ , 0.0000), (s ₅ , 0.0000)}}	{{(s ₂ , 0.3831), (s ₂ , 0.2914), (s ₃ , 0.0582)}}

	G ₃	G ₄
A ₁	{{(s ₄ , -0.3522), (s ₂ , -0.0221), (s ₁ , 0.0000)}}	{{(s ₄ , 0.2634), (s ₂ , -0.4308), (s ₂ , 0.0000)}}
A ₂	{{(s ₂ , 0.0000), (s ₁ , 0.0000), (s ₃ , 0.3659)}}	{{(s ₄ , 0.2634), (s ₃ , 0.0925), (s ₂ , 0.3522)}}
A ₃	{{(s ₄ , 0.2634), (s ₂ , 0.0423), (s ₃ , 0.1383)}}	{{(s ₄ , -0.2974), (s ₁ , 0.3195), (s ₃ , -0.2028)}}
A ₄	{{(s ₄ , 0.0668), (s ₃ , 0.0000), (s ₃ , 0.0314)}}	{{(s ₃ , -0.4313), (s ₃ , -0.1716), (s ₃ , 0.3437)}}
A ₅	{{(s ₃ , -0.3178), (s ₄ , -0.3831), (s ₄ , -0.2303)}}	{{(s ₂ , 0.0000), (s ₂ , 0.4623), (s ₃ , 0.3659)}}

Step 2. Compute the values of A⁺ (PIS) and A⁻ (NIS), for all attributes are benefit and based on the formula (16) and (17), we can obtain the (PIS) A⁺ and (NIS)A⁻ as follows.

$$A^+ = \left\{ \begin{array}{l} \{(s_5, -0.1892), (s_1, 0.1892), (s_1, 0.2746)\}, \\ \{(s_4, 0.3182), (s_2, -0.0668), (s_2, -0.4308)\}, \\ \{(s_4, 0.2634), (s_1, 0.0000), (s_1, 0.0000)\}, \\ \{(s_4, 0.2634), (s_1, 0.3195), (s_2, 0.0000)\} \end{array} \right\}$$

$$A^- = \left\{ \begin{array}{l} \{(s_1, 0.3756), (s_4, 0.0000), (s_5, 0.0000)\}, \\ \{(s_2, 0.3831), (s_2, 0.3784), (s_4, 0.0760)\}, \\ \{(s_2, 0.0000), (s_4, -0.3831), (s_4, -0.2303)\}, \\ \{(s_2, 0.0000), (s_3, 0.0925), (s_3, 0.3659)\} \end{array} \right\}$$

Step 3. Based on the Equation (7) and the attribute weighting vector ω_j, calculate the values of τ_i and ψ_i.

$$\tau_1 = 0.0821, \tau_2 = 0.5137, \tau_3 = 0.4351, \tau_4 = 0.5677, \tau_5 = 0.9161,$$

$$\psi_1 = 0.0729, \psi_2 = 0.2118, \psi_3 = 0.1466, \psi_4 = 0.1934, \psi_5 = 0.3200.$$

Table 5. Ordering by the VIKOR method with 2TLNNs.

Parameter	Q_1	Q_2	Q_3	Q_4	Q_5	Ordering
$\rho = 0.0$	0.0000	0.5621	0.2982	0.4878	1.0000	$Q_1 > Q_3 > Q_4 > Q_2 > Q_5$
$\rho = 0.1$	0.0000	0.5576	0.3107	0.4973	1.0000	$Q_1 > Q_3 > Q_4 > Q_2 > Q_5$
$\rho = 0.3$	0.0000	0.5487	0.3357	0.5162	1.0000	$Q_1 > Q_3 > Q_4 > Q_2 > Q_5$
$\rho = 0.5$	0.0000	0.5398	0.3607	0.5351	1.0000	$Q_1 > Q_3 > Q_4 > Q_2 > Q_5$
$\rho = 0.7$	0.0000	0.5309	0.3858	0.5540	1.0000	$Q_1 > Q_3 > Q_4 > Q_2 > Q_5$
$\rho = 1.0$	0.0000	0.5175	0.4233	0.5823	1.0000	$Q_1 > Q_3 > Q_4 > Q_2 > Q_5$

Step 4. Compute the values of Q_i based on the results of τ_i and ψ_i , the calculating values are listed as follows. (Let $\rho = 0.4$)

$$Q_1 = 0.0000, Q_2 = 0.5442, Q_3 = 0.3482, Q_4 = 0.5256, Q_5 = 1.0000.$$

Step 5. To choose the best alternative by rank the values of Q_i , the ranking of Q_i is $Q_1 > Q_3 > Q_4 > Q_2 > Q_5$, and the best choice is η_1 .

By altering the parameter ρ , we can derive the following results which listed in Table 5.

From Table 5, we can easily find that the ordering of alternatives are same, which indicates our developed method has the robustness and can be applied to deal with practical decision making problems.

5.2. Comparative analyses

In this section, we compare our proposed VIKOR method under 2TLNNs with the 2TLNNWA and 2TLNNWG operators defined by Wang, Wei et al. (2018). Based on the values of Table 4 and attributes weighting vector $\omega = (0.32, 0.13, 0.35, 0.20)^T$, we can compute overall value η_i by 2TLNNWA and 2TLNNWG operators.

We can get calculating results η_i by 2TLNNWA operator:

$$\begin{aligned} \eta_1 &= \{(s_4, 0.2963), (s_2, -0.4001), (s_1, 0.3163)\} \\ \eta_2 &= \{(s_4, -0.2615), (s_2, 0.1166), (s_3, 0.0692)\} \\ \eta_3 &= \{(s_4, -0.1558), (s_2, -0.0267), (s_3, 0.2448)\} \\ \eta_4 &= \{(s_3, 0.4351), (s_3, -0.2747), (s_3, 0.0146)\} \\ \eta_5 &= \{(s_2, 0.1265), (s_3, 0.2595), (s_4, -0.0744)\} \end{aligned}$$

We can get calculating results η_i by 2TLNNWG operator:

$$\begin{aligned} \eta_1 &= \{(s_4, 0.2030), (s_2, -0.3488), (s_1, 0.3771)\} \\ \eta_2 &= \{(s_3, 0.2297), (s_3, -0.3938), (s_3, 0.2565)\} \\ \eta_3 &= \{(s_4, -0.1997), (s_2, 0.0282), (s_3, 0.2931)\} \\ \eta_4 &= \{(s_3, 0.3222), (s_3, -0.2358), (s_3, 0.0783)\} \\ \eta_5 &= \{(s_2, 0.0115), (s_3, 0.4173), (s_4, 0.1509)\} \end{aligned}$$

Table 6. Alternative scores $s(\eta_i)$ by 2TLNNA and 2TLNNWG operators.

2TLNNA operator	2TLNNWG operator
$s(\eta_1) = 0.7433, s(\eta_2) = 0.5863, s(\eta_3) = 0.5903,$ $s(\eta_4) = 0.5386, s(\eta_5) = 0.3856.$	$s(\eta_1) = 0.7319, s(\eta_2) = 0.5204, s(\eta_3) = 0.5822, s(\eta_4) =$ $0.5267, s(\eta_5) = 0.3580.$

Table 7. Rank of Alternatives by 2TLNNA and 2TLNNWG operators.

	order
2TLNNA	$A_1 > A_3 > A_2 > A_4 > A_5$
2TLNNWG	$A_1 > A_3 > A_4 > A_2 > A_5$
2TLNNs VIKOR	$A_1 > A_3 > A_4 > A_2 > A_5$

Then, we calculate the alternative scores $s(\eta_i)$ by score functions of 2TLNNs which are listed in Table 6.

The ranking of alternatives by 2TLNNA and 2TLNNWG operators are listed in Table 7.

Compare the values of our proposed VIKOR method under 2TLNNs with 2TLNNA and 2TLNNWG operators, the results are slightly different in ranking of alternatives and the best alternatives are same, VIKOR method with 2TLNNs can consider the conflicting attributes and can be more reasonable and scientific in the application of MCGDM problems.

5.3. Numerical case for MCDM problems with IV2TLNNs

In this chapter, if the evaluation values of five green suppliers are depicted by IV2TLNNs, then we can study the MCDM problems by using the VIKOR method with IV2TLNNs, the decision matrix are listed in Table 8 (attribute weighting vector $\omega = (0.4, 0.2, 0.1, 0.1, 0.3)^T$).

Step 1. Construct the decision matrix (See Table 8)

Step 2. Compute the values of A^+ (PIS) and A^- (NIS), for all attributes are benefit and based on the formula (18) and (19), we can obtain the (PIS) A^+ and (NIS) A^- as follows.

$$A^+ = \left\{ \begin{array}{l} \{[(s_4, 0), (s_6, 0)], [(s_1, 0), (s_2, 0)], [(s_2, 0), (s_3, 0)]\}, \\ \{[(s_3, 0), (s_5, 0)], [(s_2, 0), (s_3, 0)], [(s_1, 0), (s_2, 0)]\}, \\ \{[(s_5, 0), (s_6, 0)], [(s_1, 0), (s_3, 0)], [(s_1, 0), (s_2, 0)]\}, \\ \{[(s_5, 0), (s_6, 0)], [(s_2, 0), (s_3, 0)], [(s_1, 0), (s_2, 0)]\} \end{array} \right\}$$

$$A^- = \left\{ \begin{array}{l} \{[(s_1, 0), (s_2, 0)], [(s_4, 0), (s_5, 0)], [(s_5, 0), (s_6, 0)]\}, \\ \{[(s_1, 0), (s_3, 0)], [(s_5, 0), (s_6, 0)], [(s_4, 0), (s_5, 0)]\}, \\ \{[(s_1, 0), (s_2, 0)], [(s_4, 0), (s_5, 0)], [(s_3, 0), (s_4, 0)]\}, \\ \{[(s_1, 0), (s_2, 0)], [(s_4, 0), (s_6, 0)], [(s_2, 0), (s_3, 0)]\} \end{array} \right\}$$

Table 8. IV2TLNNs evaluation matrix.

	G_1	G_2
A_1	$\{(s_4,0),(s_5,0)\},\{(s_1,0),(s_2,0)\},\{(s_2,0),(s_3,0)\}$	$\{(s_3,0),(s_5,0)\},\{(s_2,0),(s_3,0)\},\{(s_1,0),(s_2,0)\}$
A_2	$\{(s_1,0),(s_2,0)\},\{(s_2,0),(s_3,0)\},\{(s_3,0),(s_4,0)\}$	$\{(s_1,0),(s_3,0)\},\{(s_2,0),(s_4,0)\},\{(s_3,0),(s_5,0)\}$
A_3	$\{(s_4,0),(s_5,0)\},\{(s_3,0),(s_4,0)\},\{(s_4,0),(s_5,0)\}$	$\{(s_2,0),(s_3,0)\},\{(s_2,0),(s_4,0)\},\{(s_1,0),(s_2,0)\}$
A_4	$\{(s_4,0),(s_6,0)\},\{(s_2,0),(s_4,0)\},\{(s_5,0),(s_6,0)\}$	$\{(s_3,0),(s_4,0)\},\{(s_2,0),(s_3,0)\},\{(s_1,0),(s_2,0)\}$
A_5	$\{(s_1,0),(s_2,0)\},\{(s_4,0),(s_5,0)\},\{(s_5,0),(s_6,0)\}$	$\{(s_2,0),(s_3,0)\},\{(s_5,0),(s_6,0)\},\{(s_4,0),(s_5,0)\}$
	G_3	G_4
A_1	$\{(s_5,0),(s_6,0)\},\{(s_1,0),(s_3,0)\},\{(s_2,0),(s_3,0)\}$	$\{(s_5,0),(s_6,0)\},\{(s_3,0),(s_4,0)\},\{(s_1,0),(s_2,0)\}$
A_2	$\{(s_5,0),(s_6,0)\},\{(s_4,0),(s_5,0)\},\{(s_1,0),(s_2,0)\}$	$\{(s_3,0),(s_4,0)\},\{(s_2,0),(s_3,0)\},\{(s_1,0),(s_2,0)\}$
A_3	$\{(s_4,0),(s_5,0)\},\{(s_3,0),(s_4,0)\},\{(s_2,0),(s_3,0)\}$	$\{(s_1,0),(s_2,0)\},\{(s_4,0),(s_5,0)\},\{(s_2,0),(s_3,0)\}$
A_4	$\{(s_3,0),(s_4,0)\},\{(s_4,0),(s_5,0)\},\{(s_2,0),(s_3,0)\}$	$\{(s_1,0),(s_2,0)\},\{(s_2,0),(s_3,0)\},\{(s_1,0),(s_2,0)\}$
A_5	$\{(s_1,0),(s_3,0)\},\{(s_2,0),(s_5,0)\},\{(s_3,0),(s_4,0)\}$	$\{(s_2,0),(s_3,0)\},\{(s_4,0),(s_6,0)\},\{(s_2,0),(s_3,0)\}$

Step 3. Based on the Equation (11) and the attribute weighting vector ω_j , calculate the values of $\tilde{\tau}_i$ and $\tilde{\psi}_i$.

$$\tilde{\tau}_1 = 0.0736, \tilde{\tau}_2 = 0.4678, \tilde{\tau}_3 = 0.5632, \tilde{\tau}_4 = 0.4307, \tilde{\tau}_5 = 0.9350,$$

$$\tilde{\psi}_1 = 0.0400, \tilde{\psi}_2 = 0.2316, \tilde{\psi}_3 = 0.2800, \tilde{\psi}_4 = 0.1895, \tilde{\psi}_5 = 0.4000.$$

Step 4. Compute the values of Q_i based on the results of $\tilde{\tau}_i$ and $\tilde{\psi}_i$, the calculating values are listed as follows. (Let $\rho = 0.4$)

$$Q_1 = 0.0000, Q_2 = 0.5024, Q_3 = 0.6274, Q_4 = 0.4150, Q_5 = 1.0000.$$

Step 5. To choose the best alternative by rank the values of Q_i , the ranking of Q_i is $Q_1 > Q_4 > Q_2 > Q_3 > Q_5$, and the best choice is $\tilde{\eta}_1$.

By altering the parameter ρ , we can derive the following results which listed in Table 9.

From Table 9, we can easily find that the ordering of alternatives are same, which indicates our developed method has the robustness and can be applied to deal with practical decision making problems.

5.4. Comparative analyses

In this section, we compare our proposed the extend VIKOR method under IV2TLNNs with the IV2TLNNWA and IV2TLNNWG operators. Based on the values of Table 4 and attributes weighting vector $\omega = (0.4, 0.2, 0.1, 0.3)^T$, we can utilize overall $\tilde{\eta}_{ij}$ by IV2TLNNWA and IV2TLNNWG operators.

Table 9. Ordering by the VIKOR method with IV2TLNNs.

Parameter	Q_1	Q_2	Q_3	Q_4	Q_5	Ordering
$\rho = 0.0$	0.0000	0.5322	0.6667	0.4152	1.0000	$Q_1 > Q_4 > Q_2 > Q_3 > Q_5$
$\rho = 0.1$	0.0000	0.5247	0.6568	0.4151	1.0000	$Q_1 > Q_4 > Q_2 > Q_3 > Q_5$
$\rho = 0.3$	0.0000	0.5098	0.6372	0.4150	1.0000	$Q_1 > Q_4 > Q_2 > Q_3 > Q_5$
$\rho = 0.5$	0.0000	0.4949	0.6175	0.4149	1.0000	$Q_1 > Q_4 > Q_2 > Q_3 > Q_5$
$\rho = 0.7$	0.0000	0.4800	0.5979	0.4148	1.0000	$Q_1 > Q_4 > Q_2 > Q_3 > Q_5$
$\rho = 1.0$	0.0000	0.4577	0.5684	0.4146	1.0000	$Q_1 > Q_4 > Q_2 > Q_3 > Q_5$

We can get calculating results $\tilde{\eta}_i$ by IV2TLNNWA operator:

$$\begin{aligned} \tilde{\eta}_1 &= \{[(s_4, 0.3562), (s_6, 0.0000)], [(s_2, -0.4029), (s_3, -0.2192)], [(s_1, 0.4142), (s_2, 0.4495)]\} \\ \tilde{\eta}_2 &= \{[(s_2, 0.3481), (s_6, 0.0000)], [(s_2, 0.1435), (s_3, 0.3442)], [(s_2, -0.0668), (s_3, 0.1698)]\} \\ \tilde{\eta}_3 &= \{[(s_3, -0.0243), (s_4, 0.1118)], [(s_3, 0.0157), (s_4, 0.2769)], [(s_2, 0.2974), (s_3, 0.3935)]\} \\ \tilde{\eta}_4 &= \{[(s_3, 0.0267), (s_6, 0.0000)], [(s_2, 0.1435), (s_4, -0.4577)], [(s_2, 0.0403), (s_3, 0.2321)]\} \\ \tilde{\eta}_5 &= \{[(s_2, -0.4721), (s_3, -0.3659)], [(s_4, -0.0975), (s_5, 0.4772)], [(s_3, 0.4516), (s_5, -0.4877)]\} \end{aligned}$$

We can get calculating results $\tilde{\eta}_i$ by IV2TLNNWG operator:

$$\begin{aligned} \tilde{\eta}_1 &= \{[(s_4, 0.1289), (s_5, 0.3783)], [(s_2, -0.1024), (s_3, 0.0196)], [(s_2, -0.4721), (s_3, -0.4641)]\} \\ \tilde{\eta}_2 &= \{[(s_2, -0.3668), (s_3, -0.0196)], [(s_2, 0.2679), (s_4, -0.4785)], [(s_2, 0.3199), (s_4, -0.2974)]\} \\ \tilde{\eta}_3 &= \{[(s_2, 0.2974), (s_3, 0.4294)], [(s_3, 0.1863), (s_4, 0.3755)], [(s_3, -0.1698), (s_4, -0.0477)]\} \\ \tilde{\eta}_4 &= \{[(s_2, 0.4208), (s_4, -0.1789)], [(s_2, 0.2679), (s_4, -0.2855)], [(s_3, 0.4314), (s_6, 0.0000)]\} \\ \tilde{\eta}_5 &= \{[(s_1, 0.4142), (s_3, -0.4492)], [(s_4, 0.1339), (s_6, 0.0000)], [(s_4, 0.0567), (s_6, 0.0000)]\} \end{aligned}$$

Calculating the alternative scores $s(\tilde{\eta}_i)$ by score functions of IV2TLNNs which listed in Table 10.

The ranking of alternatives by IV2TLNNWA and IV2TLNNWG operators are listed in Table 11.

Compare the values of our proposed VIKOR method under IV2TLNNs with IV2TLNNWA and IV2TLNNWG operators, the results are slightly different in ranking of alternatives and the best alternatives are same, IV2TLNNs VIKOR method can consider the conflicting attributes and can be more reasonable and scientific in the application of MCGDM problems.

5.5. Discussion

Based on above two numerical examples, we can easily find our proposed methods can express more fuzzy information and apply broadly situations in real MCGDM problems. Based on 2-tuple linguistic neutrosophic fuzzy set (2TLNS) and traditional VIKOR method, we develop the 2-tuple linguistic neutrosophic VIKOR method and the interval-valued 2-tuple linguistic neutrosophic VIKOR method; our research results can be more suitable for MCGDM problems than single-valued neutrosophic VIKOR method depicted in literature (Huang, Wei, & Wei, 2017). For the single-valued neutrosophic VIKOR method can't deal with MCGDM problems which the assessment results are depicted with 2TLNNs.

Table 10. Alternative scores $s(\tilde{\eta}_i)$ by IV2TLNNA and IV2TLNNWG operators.

IV2TLNNA operator	IV2TLNNWG operator
$s(\tilde{\eta}_1) = 0.7254, s(\tilde{\eta}_2) = 0.6044, s(\tilde{\eta}_3) = 0.5029,$ $s(\tilde{\eta}_4) = 0.6130, s(\tilde{\eta}_5) = 0.3005.$	$s(\tilde{\eta}_1) = 0.6813, s(\tilde{\eta}_2) = 0.4667, s(\tilde{\eta}_3) = 0.4273,$ $s(\tilde{\eta}_4) = 0.4119, s(\tilde{\eta}_5) = 0.2160.$

Table 11. Rank of Alternatives by IV2TLNNA and IV2TLNNWG operators.

	Order
IV2TLNNA	$A_1 > A_4 > A_2 > A_3 > A_5$
IV2TLNNWG	$A_1 > A_2 > A_3 > A_4 > A_5$
IV2TLNNs VIKOR	$A_1 > A_4 > A_2 > A_3 > A_5$

Furthermore, in complicated decision-making environment, the decision maker's risk attitude is an important factor to think about. the VIKOR methods, which consider the compromise between group utility maximization and individual regret minimization, can make this come true by altering the parameters whereas other decision making ways such as the 2TLNNA operator, the 2TLNNWG operator, the IV2TLNNA operator and the IV2TLNNWG operator don't have the ability that dynamic adjust to the parameter according to the decision maker's risk attitude, so it is difficult to solve the risk multiple attribute decision making in real practice.

6. Conclusion

The 2-tuple linguistic neutrosophic set (2TLNS), which is the generalized form of 2-tuple linguistic set (2TLS) and single-valued neutrosophic set (SVNS), can express the assessment information more easily and reasonably. The VIKOR method, which can consider the compromise between group utility maximization and individual regret minimization, can derive more accuracy decision making results. In this paper, based on traditional VIKOR method and the 2-tuple linguistic neutrosophic set, we develop the 2-tuple linguistic neutrosophic VIKOR method. Furthermore, we extend the 2TLNSs to interval-valued environment and propose the VIKOR method with IV2TLNNs. Moreover, a numerical example for green supplier selection has been proposed to illustrate the new method and some comparisons are also conducted to further illustrate advantages of the new method. In the future, our proposed VIKOR method with 2TLNNs and VIKOR method with IV2TLNNs can be applied to the risk analysis (Wei, Qin, Li, Zhu, & Wei, 2019; Wei, Yu, Liu, & Cao, 2018), the MCGDM problems (Hashemi, Mousavi, Zavadskas, Chalekaee, & Turskis, 2018; Yazdani, Zarate, Coulibaly, & Zavadskas, 2017; Zavadskas, Turskis, Vilutienė, & Lepkova, 2017) and many other uncertain and fuzzy environments (Deng & Gao, 2019; Gao, 2018; Gupta, Mehrlawat, & Grover, 2019; Li & Lu, 2019; Lu & Wei, 2019; Wang, Gao, & Lu, 2019; Wang, 2019; Wu, Gao, & Wei, 2019; Wu, Wang, & Gao, 2019; Xian, Chai, & Guo, 2019).

Disclosure statement

No potential conflict of interest was reported by the authors.

Funding

The work was supported by the National Natural Science Foundation of China under Grant No. 71571128 and the Humanities and Social Sciences Foundation of Ministry of Education of the People's Republic of China (14XJCZH002).

ORCID

Guiwu Wei  <http://orcid.org/0000-0001-9074-2005>

References

- Atanassov, K. T. (1986). Intuitionistic fuzzy sets. *Fuzzy Sets and Systems*, 20(1), 87–96. doi:10.1016/S0165-0114(86)80034-3
- Chen, C. T. (2000). Extensions of the TOPSIS for group decision-making under fuzzy environment. *Fuzzy Sets and Systems*, 114(1), 1–9. doi:10.1016/S0165-0114(97)00377-1
- Deng, X. M., & Gao, H. (2019). TODIM method for multiple attribute decision making with 2-tuple linguistic Pythagorean fuzzy information. *Journal of Intelligent & Fuzzy Systems*, 37, 1769–1780. doi:10.3233/JIFS-179240
- Deng, X. M., Wei, G. W., Gao, H., & Wang, J. (2018). Models for safety assessment of construction project with some 2-tuple linguistic Pythagorean fuzzy Bonferroni mean operators. *IEEE Access*, 6, 52105–52137. doi:10.1109/ACCESS.2018.2869414
- Du, Y., & Liu, P. D. (2011). Extended fuzzy VIKOR method with intuitionistic trapezoidal fuzzy numbers. *Information—An International Interdisciplinary Journal*, 14, 2575–2583.
- Gao, H. (2018). Pythagorean fuzzy Hamacher prioritized aggregation operators in multiple attribute decision making. *Journal of Intelligent & Fuzzy Systems*, 35, 2229–2245. doi:10.3233/JIFS-172262
- Ghadikolaei, A. S., Madhoushi, M., & Divsalar, M. (2018). Extension of the VIKOR method for group decision making with extended hesitant fuzzy linguistic information. *Neural Computing and Applications*, 30, 3589–3602. doi:10.1007/s00521-017-2944-5
- Gomes, L., & Lima, M. (1979). TODIM: Basics and application to multicriteria ranking of projects with environmental impacts. *Foundations of Computing and Decision Sciences*, 16, 113–127.
- Gomes, L., & Rangel, L. A. D. (2009). An application of the TODIM method to the multicriteria rental evaluation of residential properties. *European Journal of Operational Research*, 193, 204–211.
- Gupta, P., Mehlawat, M. K., & Grover, N. (2019). A generalized TOPSIS method for intuitionistic fuzzy multiple attribute group decision making considering different scenarios of attributes weight information. *International Journal of Fuzzy Systems*, 21(2), 369–387. doi:10.1007/s40815-018-0563-7
- Hashemi, H., Mousavi, S. M., Zavadskas, E. K., Chalekaee, A., & Turskis, Z. (2018). A new group decision model based on grey-intuitionistic fuzzy-ELECTRE and VIKOR for contractor assessment problem. *Sustainability*, 10, 1635. doi:10.3390/su10051635
- Huang, Y. H., & Wei, G. W. (2018). TODIM method for Pythagorean 2-tuple linguistic multiple attribute decision making. *Journal of Intelligent & Fuzzy Systems*, 35, 901–915. doi:10.3233/JIFS-171636
- Huang, Y. H., Wei, G. W., & Wei, C. (2017). VIKOR method for interval neutrosophic multiple attribute group decision-making. *Information*, 8(4), 144. doi:10.3390/info8040144
- Ju, D. W., Ju, Y. B., & Wang, A. H. (2018). Multiple attribute group decision making based on Maclaurin symmetric mean operator under single-valued neutrosophic interval 2-tuple linguistic environment. *Journal of Intelligent & Fuzzy Systems*, 34, 2579–2595. doi:10.3233/JIFS-17496

- Keshavarz Ghorabae, M., Zavadskas, E. K., Olfat, L., & Turskis, Z. (2015). Multi-criteria inventory classification using a new method of evaluation based on distance from average solution (EDAS). *Informatica*, 26(3), 435–451. doi:10.15388/Informatica.2015.57
- Lai, Y. J., Liu, T. Y., & Hwang, C. L. (1994). TOPSIS for MODM. *European Journal of Operational Research*, 76(3), 486–500. doi:10.1016/0377-2217(94)90282-8
- Li, X. Y., & Wei, G. W. (2014). GRA method for multiple criteria group decision making with incomplete weight information under hesitant fuzzy setting. *Journal of Intelligent & Fuzzy Systems*, 27, 1095–1105.
- Li, Z. X., Gao, H., & Wei, G. W. (2018). Methods for multiple attribute group decision making based on intuitionistic fuzzy Dombi Hamy mean operators. *Symmetry*, 10(11), 574. doi:10.3390/sym10110574
- Li, Z. X., & Lu, M. (2019). Some novel similarity and distance and measures of Pythagorean fuzzy sets and their applications. *Journal of Intelligent & Fuzzy Systems*, 37, 1781–1799. doi:10.3233/JIFS-179241
- Liu, P. D., & You, X. L. (2018). Some linguistic neutrosophic Hamy mean operators and their application to multi-attribute group decision making. *PLoS One*, 13, e0193027. doi:10.1371/journal.pone.0193027
- Lu, J. P., & Wei, C. (2019). TODIM method for performance appraisal on social-integration-based rural reconstruction with interval-valued intuitionistic fuzzy information. *Journal of Intelligent & Fuzzy Systems*, 37, 1731–1740. doi:10.3233/JIFS-179236
- Mahmoudi, A., Sadi-Nezhad, S., & Makui, A. (2016). An extended fuzzy VIKOR for group decision-making based on fuzzy distance to supplier selection. *Scientia Iranica*, 23(4), 1879–1892. doi:10.24200/sci.2016.3934
- Narayanamoorthy, S., Geetha, S., Rakkiyappan, R., & Joo, Y. H. (2019). Interval-valued intuitionistic hesitant fuzzy entropy based VIKOR method for industrial robots selection. *Expert Systems with Applications*, 121, 28–37. doi:10.1016/j.eswa.2018.12.015
- Opricovic, S., & Tzeng, G. H. (2004). Compromise solution by MCDM methods: A comparative analysis of VIKOR and TOPSIS. *European Journal of Operational Research*, 156(2), 445–455. doi:10.1016/S0377-2217(03)00020-1
- Pamucar, D., & Cirovic, G. (2015). The selection of transport and handling resources in logistics centers using multi-attributive border approximation area comparison (MABAC). *Expert Systems with Applications*, 42, 3016–3028.
- Park, J. H., Cho, H. J., & Kwun, Y. C. (2011). Extension of the VIKOR method for group decision making with interval-valued intuitionistic fuzzy information. *Fuzzy Optimization and Decision Making*, 10(3), 233–253. doi:10.1007/s10700-011-9102-9
- Qin, J. D., Liu, X. W., & Pedrycz, W. (2015). An extended VIKOR method based on prospect theory for multiple attribute decision making under interval type-2 fuzzy environment. *Knowledge-Based Systems*, 86, 116–130. doi:10.1016/j.knosys.2015.05.025
- Roy, J., Sharma, H. K., Kar, S., Zavadskas, E. K., & Saparauskas, J. (2019). An extended COPRAS model for multi-criteria decision-making problems and its application in web-based hotel evaluation and selection. *Economic Research-Ekonomska Istraživanja*, 32(1), 219–253. doi:10.1080/1331677X.2018.1543054
- Sharma, H. K., Kumari, K., & Kar, S. (2019). Short-term forecasting of air passengers based on the hybrid rough set and the double exponential smoothing model. *Intelligent Automation and Soft Computing*, 25, 1–14. doi:10.31209/2018.100000036
- Smarandache, F. (1999). *A unifying field in logics. Neutrosophy: Neutrosophic probability, set and logic*. Rehoboth, NM: American Research Press.
- Tang, M., Wang, J., Lu, J. P., Wei, G. W., Wei, C., & Wei, Y. (2019). Dual hesitant Pythagorean fuzzy Heronian mean operators in multiple attribute decision making. *Mathematics*, 7, 344.
- Tang, X. Y., & Wei, G. W. (2018). Models for green supplier selection in green supply chain management with Pythagorean 2-tuple linguistic information. *IEEE Access*, 6, 18042–18060. doi:10.1109/ACCESS.2018.2817551

- Tang, X. Y., Wei, G. W., & Gao, H. (2019a). Models for multiple attribute decision making with interval-valued Pythagorean fuzzy Muirhead mean operators and their application to green suppliers selection. *Informatica*, 30(1), 153–186. doi:10.15388/Informatica.2019.202
- Tang, X. Y., Wei, G. W., & Gao, H. (2019b). Pythagorean fuzzy Muirhead mean operators in multiple attribute decision making for evaluating of emerging technology commercialization. *Economic Research-Ekonomska Istraživanja*, 32(1), 1667–1696. doi:10.1080/1331677X.2019.1638808
- Wang, H., Smarandache, F., Zhang, Y. Q., & Sunderraman, R. (2010). Single valued neutrosophic sets. *Multispace Multistruct*, 4, 410–413.
- Wang, J., Gao, H., & Lu, M. (2019). Approaches to strategic supplier selection under interval neutrosophic environment. *Journal of Intelligent & Fuzzy Systems*, 37, 1707–1730. doi:10.3233/JIFS-179235
- Wang, J., Gao, H., & Wei, G. (2018). Some 2-tuple linguistic neutrosophic number Muirhead mean operators and their applications to multiple attribute decision making. *Journal of Experimental & Theoretical Artificial Intelligence*, 31, 409–439. doi:10.1080/0952813X.2018.1552320
- Wang, J., Gao, H., Wei, G. W., & Wei, Y. (2019). Methods for multiple-attribute group decision making with q-rung interval-valued orthopair fuzzy information and their applications to the selection of green suppliers. *Symmetry*, 11(1), 56. doi:10.3390/sym11010056
- Wang, J., Tang, X. Y., & Wei, G. W. (2018). Models for multiple attribute decision-making with dual generalized single-valued neutrosophic Bonferroni mean operators. *Algorithms*, 11, 2. doi:10.3390/a11010002
- Wang, J., Wei, G. W., & Lu, M. (2018a). An extended VIKOR method for multiple criteria group decision making with triangular fuzzy neutrosophic numbers. *Symmetry*, 10(10), 497. doi:10.3390/sym10100497
- Wang, J., Wei, G. W., & Lu, M. (2018b). TODIM method for multiple attribute group decision making under 2-tuple linguistic neutrosophic environment. *Symmetry*, 10, 486. doi:10.3390/sym10100486
- Wang, J., Wei, G. W., & Wei, Y. (2018). Models for green supplier selection with some 2-tuple linguistic neutrosophic number Bonferroni mean operators. *Symmetry*, 10(5), 131. doi:10.3390/sym10050131
- Wang, L., Zhang, H. Y., Wang, J. Q., & Li, L. (2018). Picture fuzzy normalized projection-based VIKOR method for the risk evaluation of construction project. *Applied Soft Computing*, 64, 216–226.
- Wang, P., Wang, J., & Wei, G. W. (2019). EDAS method for multiple criteria group decision making under 2-tuple linguistic neutrosophic environment. *Journal of Intelligent & Fuzzy Systems*, 37, 1597–1608. doi:10.3233/JIFS-179223
- Wang, P., Wang, J., Wei, G. W., & Wei, C. (2019). Similarity measures of q-rung orthopair fuzzy sets based on cosine function and their applications. *Mathematics*, 7, 340.
- Wang, R. (2019). Research on the application of the financial investment risk appraisal models with some interval number Muirhead mean operators. *Journal of Intelligent & Fuzzy Systems*, 37, 1741–1752.
- Wei, G. W. (2017). Picture uncertain linguistic Bonferroni mean operators and their application to multiple attribute decision making. *Kybernetes*, 46(10), 1777–1800. doi:10.1108/K-01-2017-0025
- Wei, G. W. (2018). TODIM method for picture fuzzy multiple attribute decision making. *Informatica*, 29(3), 555–566. doi:10.15388/Informatica.2018.181
- Wei, G. W. (2019). Pythagorean fuzzy Hamacher power aggregation operators in multiple attribute decision making. *Fundamenta Informaticae*, 166(1), 57–85. doi:10.3233/FI-2019-1794
- Wei, G. W., Wang, J., Wei, C., Wei, Y., & Zhang, Y. (2019). Dual hesitant Pythagorean fuzzy Hamy mean operators in multiple attribute decision making. *IEEE Access*, 7, 86697–86716.
- Wei, G. W., Wang, R., Wang, J., Wei, C., & Zhang, Y. (2019). Methods for evaluating the technological innovation capability for the high-tech enterprises with generalized interval neutrosophic number Bonferroni mean operators. *IEEE Access*, 7, 86473–86492. doi:10.1109/ACCESS.2019.2925702

- Wei, G. W., & Zhang, Z. P. (2019). Some single-valued neutrosophic Bonferroni power aggregation operators in multiple attribute decision making. *Journal of Ambient Intelligence and Humanized Computing*, 10(3), 863–882.
- Wei, Y., Qin, S., Li, X., Zhu, S., & Wei, G. (2019). Oil price fluctuation, stock market and macroeconomic fundamentals: Evidence from China before and after the financial crisis. *Finance Research Letters*, 30, 23–29.
- Wei, Y., Yu, Q., Liu, J., & Cao, Y. (2018). Hot money and China's stock market volatility: Further evidence using the GARCH-MIDAS model. *Physica A: Statistical Mechanics and Its Applications*, 492, 923–930.
- Wu, L. P., Gao, H., & Wei, C. (2019). VIKOR method for financing risk assessment of rural tourism projects under interval-valued intuitionistic fuzzy environment. *Journal of Intelligent & Fuzzy Systems*, 37, 2001–2008. doi:10.3233/JIFS-179262
- Wu, L. P., Wang, J., & Gao, H. (2019). Models for competitiveness evaluation of tourist destination with some interval-valued intuitionistic fuzzy Hamy mean operators. *Journal of Intelligent & Fuzzy Systems*, 36, 5693–5709.
- Wu, L. P., Wei, G. W., Gao, H., & Wei, Y. (2018). Some interval-valued intuitionistic fuzzy Dombi Hamy mean operators and their application for evaluating the elderly tourism service quality in tourism destination. *Mathematics*, 6(12), 294. doi:10.3390/math6120294
- Wu, Q., Wu, P., Zhou, L. G., Chen, H. Y., & Guan, X. J. (2018). Some new Hamacher aggregation operators under single-valued neutrosophic 2-tuple linguistic environment and their applications to multi-attribute group decision making. *Computers & Industrial Engineering*, 116, 144–162. doi:10.1016/j.cie.2017.12.024
- Wu, S. J., Wang, J., Wei, G. W., & Wei, Y. (2018). Research on construction engineering project risk assessment with some 2-tuple linguistic neutrosophic Hamy mean operators. *Sustainability*, 10, 1536.
- Wu, Z. B., Xu, J. P., Jiang, X. L., & Zhong, L. (2019). Two MAGDM models based on hesitant fuzzy linguistic term sets with possibility distributions: VIKOR and TOPSIS. *Information Sciences*, 473, 101–120.
- Xian, S. D., Chai, J. H., & Guo, H. L. (2019). Z Linguistic-induced ordered weighted averaging operator for multiple attribute group decision-making. *International Journal of Intelligent Systems*, 34(2), 271–296.
- Xu, Z. S., & Chen, Q. (2011). A multi-criteria decision making procedure based on interval-valued intuitionistic fuzzy bonferroni means. *Journal of Systems Science and Systems Engineering*, 20(2), 217–228.
- Yager, R. R. (2014). Pythagorean membership grades in multicriteria decision making. *IEEE Transactions on Fuzzy Systems*, 22(4), 958–965. doi:10.1109/TFUZZ.2013.2278989
- Yager, R. R. (2017). Generalized orthopair fuzzy sets. *IEEE Transactions on Fuzzy Systems*, 25(5), 1222–1230. doi:10.1109/TFUZZ.2016.2604005
- Yang, W., Pang, Y. F., Shi, J. R., & Wang, C. J. (2018). Linguistic hesitant intuitionistic fuzzy decision-making method based on VIKOR. *Neural Computing & Applications*, 29, 613–626.
- Yazdani, M., Zarate, P., Coulibaly, A., & Zavadskas, E. K. (2017). A group decision making support system in logistics and supply chain management. *Expert Systems with Applications*, 88, 376–392.
- Ye, J. (2018). Multiple attribute decision-making methods based on the expected value and the similarity measure of hesitant neutrosophic linguistic numbers. *Cognitive Computation*, 10(3), 454–463. doi:10.1007/s12559-017-9535-8
- Zadeh, L. A. (1965). Fuzzy sets. In *Information and control* (Vol. 8, pp. 338–356). doi:10.1016/S0019-9958(65)90241-X
- Zavadskas, E. K., Turskis, Z., Vilitienė, T., & Lepkova, N. (2017). Integrated group fuzzy multi-criteria model: Case of facilities management strategy selection. *Expert Systems with Applications*, 82, 317–331.
- Zhu, B., & Xu, Z. S. (2013). Hesitant fuzzy Bonferroni means for multi-criteria decision making. *Journal of the Operational Research Society*, 64(12), 1831–1840. doi:10.1057/jors.2013.7