

## Multiple Primary Melanomas in a Young Patient

A 45-year-old HIV-negative Caucasian man with no reported past medical history was referred to our Department with a large (7 cm in diameter) oozing nodule on the occipital region of the scalp with spontaneous periodical bloody or purulent discharge. The lesion had appeared over a period of six months, had an irregular color, non-specific dermoscopic features, and resembled squamous cell carcinoma. The physical examination revealed three more atypical melanocytic lesions (on the abdomen, back, and buccal mucosa), and multiple swollen occipital, postauricular, as well as superficial and deep cervical lymph nodes. After clinical evaluation, the patient reported having another *in situ* melanoma (submammary region) excised 7 years ago. All the lesions were excised and sent for histopathologic examination, which was compatible with primary cutaneous melanoma. Total body computed tomography revealed the presence of multiple visceral metastases, and the patient was referred to an oncologist. He did not consent to proceed to genetic testing.



**Figure 1.** A large (7 cm in diameter), well demarcated, non-tender, predominantly flesh-colored nodule with purulent discharge, closely resembling squamous cell carcinoma.



**Figure 2.** During surgical excision, large black-colored areas were observed at the base of the nodule.



**Figure 3.** A primary superficially spreading melanoma was also excised from the patient's abdominal area.



**Figure 4.** A primary nodular melanoma on the patient's back.



**Figure 5.** The fourth primary melanoma was located on the mucosa of the hard palate.

**Georgia Kyriakou<sup>1</sup>, Efthymia Gialeli<sup>1</sup>,  
Eleftheria Vryzaki<sup>1</sup>, Stauros Balasis<sup>2</sup>,  
Sophia Georgiou<sup>1</sup>**

<sup>1</sup>*Department of Dermatology, University General  
Hospital of Patras, Rion, Greece*

<sup>2</sup>*Department of Orthopaedics, University General  
Hospital of Patras, Rion, Greece*

**Corresponding author:**

Georgia Kyriakou, MD, MsC, cPhD.  
Department of Dermatology, University General  
Hospital of Patras  
University General Hospital of Patras  
Rion 265 04  
Greece  
*geo\_kyr@yahoo.gr*

The occurrence of multiple primary melanomas (MPM) is a rare but recognized phenomenon. The estimated incidence of a second primary tumor ranges from 0.2% to 8.7% in large retrospective reviews. While 63% to 88% of patients with MPM are reported to have two primary tumors, the occurrence of more than four primary melanomas is considered exceedingly rare (1-2). Whether the presence of multiple primary melanomas is a function of increased genetic susceptibility of the individual, consistent exposure to a common exogenous promoter of malignancy, or a combination of these two factors remains to be elucidated. These patients should undergo intensive dermatologic screening for the rest of their lives and should consider genetic testing.

**References:**

1. Buljan M, Tomić Sremec N, Sremec J, Tomas D, Crnarić I, Šitum M. Disease progression in cases of multiple primary melanoma. *Acta Dermatovenerol Croat.* 2015;23:260-4.
2. Claesson M, Holmström P, Hallberg S, Gillstedt M, Gonzalez H, Wennberg AM, *et al.* Multiple primary melanomas: a common occurrence in Western Sweden. *Acta Derm Venereol.* 2017;97:715-9.

Received: April 25, 2019

Accepted: November 16, 2020