

# Traditional CVD Risk Factors and Socio-Economic Deprivation in Roma Minority Population of Croatia

Hrvojka Zeljko<sup>1</sup>, Tatjana Škarić-Jurić<sup>2</sup>, Nina Smolej Narančić<sup>2</sup>, Marijana Peričić Salihović<sup>2</sup>, Irena Martinović Klarić<sup>2</sup>, Maja Barbalić<sup>2</sup>, Lovorka Barać Lauc<sup>2</sup> and Branka Jančićjević<sup>2</sup>

<sup>1</sup> General Hospital »Sveti Duh«, Zagreb

<sup>2</sup> Institute for Anthropological Research, Zagreb

## ABSTRACT

Researches into health inequalities consistently show disadvantages in health status, morbidity and mortality for various ethnic minority groups. Current knowledge about prevention of cardiovascular diseases (CVD) mainly derives from studies carried out in populations of European origin while the evidences involving Roma population are scarce. Roma, an ethnic minority of northern Indian origin, live in many countries throughout the world and are well known for preserved traditions and resistance to assimilation. They are most often marginalized economically, spatially, politically and in terms of culture. In order to assess the health status and health-related lifestyle attributes, a multidisciplinary anthropological and epidemiological community-based study was carried out including a total of 423 members of the Bayash Roma minority population living in two regions of Croatia (144 men and 279 women, aged 18–84 yrs). Hypertension (HT) was found in 24.8% Bayash Roma (21.5% men and 26.5% women) using standard diagnostic criteria (i.e. BP  $\geq$  140/90 mm Hg or taking antihypertensive therapy). The prevalence increases from 5.9% in the age group 18–34 yrs; 35.0% in the age group 35–64 yrs and 51.4% in the age group 65+ yrs. The prevalence of hypertension in the Bayash Roma is almost half of the magnitude of what is usually reported for the general population of Croatia. It is also lower when compared with other European populations and this finding is not due to comparative younger average age of the Bayash sample. The significant association of hypertension with age and BMI was confirmed in this study and the importance of non-traditional SES-related CVD risk factors was highlighted. Smoking is a part of traditional Roma life-style and with 70% of smokers almost the entire population is equally exposed to this risk factor in their family environment. Since homogeneously distributed, this risk factor did not show to be a significant predictor of hypertension. The extent to which hypertension is influenced by traditional CVD risk factors as well as by some SES indicators was also assessed using a forward stepwise method of the multivariate logistic analysis. Each risk factor was explored as quantitative variable as well as qualitative one using various cut-offs. The best model showed to be the one having age and BMI presented as quantitative variables and sex, region, smoking status, income and schooling years as categorical ones; with cut-off 3 for number of income sources and 8 for the number of schooling years. In spite of the low prevalence of hypertension, the presented results are showing that Bayash Roma are bearing a high CVD risk factors load. We expect that with westernization of their life-style and along with increase of the economic power, the proportion of CVD in population of Bayash Roma will also increase. Therefore, it is important to recognize the need for early cardiovascular disease risk factors prevention in this minority population.

**Key words:** hypertension, prevalence, CVD, risk factors, SES, poverty, Roma, Croatia

## Introduction

It is well known that cardiovascular diseases (CVD) are estimated to be the leading cause of death and disability-adjusted life years<sup>1</sup>. A number of epidemiological

studies have demonstrated that lifestyle greatly influences the cardiovascular risk pattern of populations and that modernization is associated with increased inci-

dence of non-communicable diseases, such as hypertension, obesity, dyslipidemia and diabetes mellitus.

Along with still growing evidence coming from epidemiological studies carried out world-wide that continually confirm the importance of classical CVD risk factors, the knowledge of the relationship between CVD risk and poverty expands only recently.

Socioeconomic status (SES) is consistently among the most fundamental determinants of health<sup>2,3</sup>. Influence of SES on cardiovascular disease could be viewed as the combined effects of social disparities in health-related behaviors, environmental conditions and the contact and delivery of health care<sup>3,4</sup>.

Research into health inequalities has shown consistent disadvantages in health status, morbidity and mortality for various ethnic minority groups<sup>5,6</sup>. Current knowledge about prevention of coronary heart disease and other cardiovascular disease mainly derives from studies carried out in populations of European origin<sup>7</sup> while the evidence involving Roma population is scarce<sup>6,8-12</sup>.

Roma, an ethnic minority of northern Indian origin, live in many countries throughout the world and are well known for preserved traditions and resistance to assimilation. They are most often marginalized economically, spatially, politically and in terms of culture. Croatia has a substantial Roma population, the exact size estimated to be over 30 000 (about 1% of the total population of Croatia)<sup>13</sup>. The Bayash are one of many Romani branches who, between 14<sup>th</sup> and 19<sup>th</sup> centuries lived in historical Romanian states Wallachia and Moldavia, where they were kept as slaves. After 1856, when the slavery in Romania was finally abolished, larger Bayash groups immigrated to Croatia<sup>14</sup>.

From the genetic-epidemiological perspective, Roma population might be considered as a particularly vulnerable group for developing common chronic diseases. Na-

mely, with the increase in life expectancy, the prevalence of chronic disease in Roma population may rise even at a higher rate than in other populations since bearing additional both genetic and environmental risks. Those risks are connected with the following characteristics of Roma populations:

a) Low socio-economic status and cultural specificities that shape lifestyle patterns compromising health-related behaviors;

b) High degree of reproductive isolation that enhances the possibility of increased frequencies of homozygous genotypes as the consequences of drift and inbreeding<sup>10</sup>.

In order to assess the health status and health-related lifestyle attributes of the Roma minority population living in Croatia, a multidisciplinary anthropological and epidemiological community-based study was carried out in 2005/06 in Bayash settlements in Baranja and Medimurje.

Goals of the present study:

- To report the preliminary results on blood pressure levels and prevalence of hypertension in Bayash Roma minority population of Croatia and to compare it to surrounding populations;
- To assess the importance of some traditional CVD risk factors (BMI, smoking) in this population;
- To investigate the possibility of the association of blood pressure values with some indicators of socio-economic status (SES indicators of poverty) in this population.

## Participants and Methods

The dataset used in this study is a subset of the extensive material collected in field studies carried out in 2005 to 2006 in Bayash settlements in Baranja and Medimurje regions of Croatia (Figure 1). The study was held in cooperation with the Office for National Minorities of the Government of the Republic of Croatia and the Croatian National Institute of Public Health.

All examinees voluntarily participated in the study and before signing the informed consent were informed about the goals, methods and expectations of the project. The study protocol included measurement of various health-related biometric traits and an extensive interview (more thorough information can be found in the previous report<sup>6</sup>).

A total of 423 members of the Bayash population living in Baranja and Medimurje (144 men and 279 women, aged 18–84 yrs) were included in the study. Whereas the gender structure of two regional samples is similar, the age structure differ significantly reflecting voluntary participation. The comparability of two Bayash sub-populations was achieved by presenting additionally the prevalence of hypertension within separate age groups: 18–34; 35–64 and 65+ years' olds.

Blood pressure measurements were performed in the sitting position through the use of a mercury sphygmo-



Figure 1. Map of Croatia

manometer by an experienced and certified examiner. The second measurement was taken as valid due to the fact that participants had been sitting for 10–15 minutes before the second measurement.

The hypertensive group («cases») incorporates participants who declared themselves as hypertensives (indicating previously diagnosed hypertension by personal physician) and those in whom the measured systolic blood pressure values were equal or higher than 140 mmHg or diastolic blood pressure measures were equal or higher than 90 mmHg.

Short anthropometry was undertaken following standard IBP Protocol<sup>15</sup>. Body mass index (BMI) was calculated as weight (kg) / stature(m)<sup>2</sup>. BMI equal or higher than 25 was used as an obesity criterion.

Smoking status was assessed qualitatively – as the proportion of smokers – and quantitatively using estimate »pack per year«. »Pack per year« was calculated as the number of cigarettes smoked daily multiplied by years of smoking and divided by 20.

The socio-economic status was represented by two variables – the total number of income sources (ranging in the sample from zero to three) and years of education (ranging in the sample from zero to twelve). Due to the staggering unemployment rate – e.g., only 7 persons reported to have permanent jobs – the economic status was roughly estimated using the total number of financial resources. Namely, the main financial inputs in Bayash population are social welfare support allowance (83.6%) and child allowance (47.0%), while a smaller proportion of the population depends on retirement allowance and salaries coming from occasional or permanent jobs<sup>6</sup>.

The education level was presented as a quantitative variable »years of schooling« and two qualitative ones: the proportion of persons who ever attended school and the proportion of persons who finished elementary education (8 and more years of schooling). According to the recently published data<sup>6</sup>, the proportion of Bayash who never attended school amounts to 33.3% and the average schooling years for those 66.7% who were enrolled in education system amounts to 5.3 years. The overall average participation of the Bayash in formal education system is 3.6 years and only 10% completed 8-year elementary education, 4.4% were enrolled in secondary education whereas none of the examinees reported university level education.

### *Statistical analyses*

Descriptive statistics was shown in absolute numbers and frequencies for the qualitative variables, and means and standard deviations for the quantitative ones. Differences in qualitative variables (sex, region, hypertensive status) were tested using Fisher's exact test and differences in quantitative variables by t-test. The blood pressure values were age-adjusted by means of multiple regression analysis and standardized residuals were used in relating blood pressure values with the number of income sources. The association of hypertension with CVD

risk factors was assessed by logistic regression analysis. All analyses were performed by SPSS 10.0 statistical package for Windows (SPSS Inc., Chicago, IL, USA), with statistical significance set at  $p \leq 0.05$ .

## **Results**

Descriptive statistics for age, systolic and diastolic blood pressure, body mass index, smoking status, schooling years and the number of income sources of the Bayash sample is presented in Table 1.

Significant differences among two sexes are found for age, intensity of smoking, schooling years and the number of income sources. In this sample females were older than males (42.3 yrs. vs. 38.9 yrs.). While the proportion of smokers in two sexes is similar, males are heavier smokers compared to females (24.9 vs. 19.8 pack/year). A higher proportion of men attend school and their schooling lasts on the average 1.3 years longer than that of women. Men also have significantly more income resources than women.

The prevalence of hypertension in Bayash sample is 24.8% (M: 21.5%, F: 26.5%). As presented in Table 2, according to the age group the prevalence increases from 5.9% in the age group 18–34 yrs; 35.0% in the age group 35–64 yrs to 51.4% in the age group 65+ yrs. In neither age group the difference between males and females was found to be significant.

Hypertension was more prevalent in the Bayash from Baranja (31.2%) than those from Međimurje (14.7%). But the comparison of hypertension prevalence within age groups indicated that the observed regional difference could be attributed to the age differences between two samples.

The relation between measured hypertension (mHT) and previously diagnosed hypertension (dHT) in Bayash population is given in Table 3. In both sub-samples the majority of persons belong to non-hypertensive group (68.9% in Baranja and 85.4% in Međimurje). The »undiagnosed hypertension« is present in 19.6% of the Bayash in Baranja and 12.9% in Međimurje. The »unsuccessfully treated hypertension« is present in 8.9% Bayash in Baranja and 1.2% in Međimurje, while the lowest percentages refer to »successfully treated hypertension« (2.7% in Baranja and 0.6% in Međimurje).

The association of hypertension with CVD risk factors is presented by comparing the prevalence of the investigated CVD risk factors between hypertensives (cases) and normotensives (controls) (Table 4). The prevalence of each CVD risk factor – i.e. smoking, overweight, less than 3 sources of income, less than 8 years of education and never attending school – in cases and controls is presented for men and women separately. The possible influence of age was attenuated by considering the narrower age group (35–64 yrs) across all the samples.

Among five examined indicators, the prevalence of smoking was the only one that did not differ between hypertensive and normotensive persons. And this is equally

**TABLE 1.**  
DESCRIPTIVE STATISTICS FOR CHARACTERISTICS RELATED TO HYPERTENSION (AGE STRUCTURE, BLOOD PRESSURE, BODY MASS INDEX, SMOKING STATUS, EDUCATION AND INCOME) IN BAYASH POPULATION OF CROATIA. SEX DIFFERENCES WERE TESTED USING T-TEST FOR QUANTITATIVE AND FISHER'S EXACT TEST FOR QUALITATIVE VARIABLES

	Total (N=423)	Men (N=144)	Women (N=279)	p (M vs.F)
Age (mean, SD)	41.1 (15.2)	38.9 (15.1)	42.3 (15.1)	0.029
18-34 yrs (n, %)	169 (40.0)	64 (44.4)	105 (37.6)	
35-64 yrs (n, %)	217 (51.3)	69 (47.9)	148 (53.0)	
65+ yrs (n, %)	37 (8.7)	11 (7.6)	26 (9.3)	
Blood pressure				
Systolic (mean, SD)	120.2 (25.8)	119.3 (18.5)	120.7 (28.9)	0.598
Diastolic (mean, SD)	75.8 (13.5)	76.9 (11.4)	75.2 (14.5)	0.221
Hypertension (n, %)	105 (24.8)	31 (21.5)	74 (26.5)	0.260
BMI (mean, SD)	25.5 (5.7)	25.3 (5.0)	25.6 (6.0)	0.595
BMI >25 (n, %)	202 (48.6)	71 (50.0)	131 (47.8)	0.672
Smokers (n, %)	297 (70.4)	104 (72.2)	193 (69.4)	0.551
Pack/year (mean, SD)	21.6 (18.0)	24.9 (21.9)	19.8 (15.3)	0.013
Schooling years: 0-12 y (mean, SD)	3.5 (3.3)	4.4 (3.4)	3.1 (3.2)	0.000
Ever attended school (n, %)	267 (66.4)	111 (80.4)	156 (59.1)	0.000
Schooling years 8+ (n, %)	64 (15.9)	25 (18.1)	39 (14.8)	0.384
Income sources (mean, SD)	1.6 (0.8)	1.8 (0.9)	1.5 (0.7)	0.001
Income sources <3 (n, %)	347 (86.5)	103 (75.2)	244 (92.4)	0.000

**TABLE 2.**  
PREVALENCE OF HYPERTENSION IN BAYASH ROMA BY SEX, AGE AND REGION (TESTED BY CHI<sup>2</sup>-TEST)

Age group	Total sample %	Men %	Women %	p; Men vs. Women	Baranja %	Medimurje %	p; Baranja vs. Medimurje
18-34	5.9	9.4	3.8	0.137	4.2	7.1	0.428
35-64	35.0	30.4	37.2	0.333	39.0	25.4	0.057
65+	51.4	36.4	57.7	0.235	51.4	50.0	0.969
Total	24.8	21.5	26.5	0.260	31.2	14.7	0.000

**TABLE 3.**  
RELATION BETWEEN PREVIOUSLY DIAGNOSED HYPERTENSION (DHT) AND ELEVATED BLOOD PRESSURE VALUES (»MEASURED HYPERTENSION«; MHT) AND IN BAYASH POPULATION

Hypertension status	Definition	Baranja		Medimurje	
		N	%	N	%
Non-hypertensive	No dHT and no mHT	179	68.9	139	85.3
Undiagnosed hypertension	No dHT but mHT	51	19.6	21	12.9
Successfully treated hypertension	dHT but no mHT	7	2.7	1	0.6
Unsuccessfully treated hypertension	dHT and mHT	23	8.9	2	1.2
Total		260	100	163	100

true at the level of the total sample as it is for each investigated sex and age group.

The association of hypertension with BMI and with income sources showed to be significant for both sexes in the age 35-64 yrs as well as in the total sample. The proportion of persons having BMI  $\geq 25$  is higher in cases (M: 85.7%, F: 77.4%) than in controls (M: 54.4%; F: 47.3%).

The difference in proportion of persons having less than 3 income sources amounted to 95.2 vs. 70.2% in men and 98.0 vs. 87.5% in women for cases and controls respectively.

For two estimates of education – less than 8 schooling years and proportion of persons who never attended school – the difference was significant only at the total



**TABLE 4.**  
PREVALENCE OF POTENTIAL RISK FACTORS FOR HYPERTENSION IN HYPERTENSIVE (HT) AND NON-HYPERTENSIVE GROUP OF BAYASH ROMA BY SEX AND AGE (TESTED BY CHI<sup>2</sup>-TEST)

	Men			Women		
	HT (%)	non-HT (%)	p; HT vs. Non-HT	HT (%)	non-HT (%)	p; HT vs. Non-HT
<b>Smokers</b>						
35-64	66.7	72.9	0.599	69.1	76.1	0.353
Total	64.5	74.3	0.280	64.9	71.1	0.320
<b>Overweight persons</b>						
35-64	85.7	54.4	0.013	77.4	47.3	0.000
Total	74.2	43.2	0.002	77.5	37.4	0.000
<b>Income – less than 3 sources</b>						
35-64	95.2	70.2	0.021	98.0	87.5	0.033
Total	90.3	70.8	0.027	98.6	90.3	0.025
<b>Schooling years – less than 8</b>						
35-64	95.2	85.1	0.231	94.0	86.2	0.160
Total	96.8	77.6	0.015	95.6	81.6	0.005
<b>Schooling years – never attended school</b>						
35-64	28.6	21.3	0.512	54.0	49.4	0.606
Total	32.3	15.9	0.043	59.4	34.4	0.000

sample level. Since the difference between cases and controls was not significant within the age group 35-64 yrs, it implies that the connection of the level of education to hypertension could be attributed mainly to the connection of both education measures to age.

The relation of systolic and diastolic blood pressure to the number of income sources is also presented on Figure 2 for men and women separately. The blood pressure values are age-adjusted by means of multiple regression analysis and standardized residuals are presented for each income category. The tendency of blood pressure, irrespective of age, to be higher among those having no income and in those having a smaller number of income sources is visible in both sexes.

The extent to which hypertension is influenced by the years of education and income was also assessed using the forward stepwise method of the multivariate logistic analysis. Smoking status, BMI, length of education, and source of income, age, sex and region were considered as independent variables (Table 5).

In the course of the analysis different models were tested that considered various combinations of risk factors treated as a quantitative variable or qualitative variables with different cut-offs. The socio-economic status was represented by the total number of income sources (0-3) and years of education (0-12). Those two variables showed to be influential when categorized as low income 0-2 sources vs. 3 sources and less than 8 years of schooling (0-7) vs. 8+ (8-12) finished years of schooling (i.e. finished elementary school). The number of income sources was equally significant predictor when represented as a quantitative variable.

The two best models both included the age and BMI (as quantitative variables) and sex, region, smoking sta-




Figure 2. Systolic and diastolic blood pressure in relation to number of income sources in Bayash Roma (sex and age adjusted blood pressure data).

**TABLE 5.**  
MULTIVARIATE SCREENING FOR PREDICTOR VARIABLES FOR  
HYPERTENSION BY LOGISTIC REGRESSION

Predictor variable	Hypertension
Region (Baranja = referent)	1.035 (0.514–2.085); p=0.923
Sex (male = referent)	1.269 (0.703–2.293); p=0.423
Smoker (no = referent)	0.922 (0.494–1.721); p=0.799
Education (0–7 yrs = referent)	3.466 (1.057–11.367); p=0.040
Age (years)	1.056 (1.033–1.080); p<0.001
Body mass index (kg/m <sup>2</sup> )	1.166 (1.107–1.229); p<0.001
Income (none = referent)	1.733 (1.120–2.682); p=0.014

Predictor variable	Hypertension
Region (Baranja = referent)	1.254 (0.638–2.462); p=0.511
Sex (male = referent)	1.321 (0.728–2.395); p=0.359
Smoker (no = referent)	0.886 (0.474–1.657); p=0.706
Education (0–7 yrs = referent)	3.682 (1.135–11.942); p=0.030
Age (yrs)	1.059 (1.037–1.083); p<0.001
Body mass index (kg/m <sup>2</sup> )	1.178 (1.116–1.243); p<0.001
Income (0–2 sources = referent)	5.549 (1.627–18.930); p=0.006

Coding: 1 – normal blood pressure; 2 – hypertension. Abbreviations: OR – odds ratio; CI – confidence intervals

tus, schooling years (as categorical ones) (Table 5). In the first model the income was included as a quantitative variable (with 4 categories) while in the second model it was treated dichotomously (0–2 vs. 3). Age and BMI, expectedly, showed to be the significant predictors for hypertension (at  $p < 0.001$  level), while sex, region and smoking status were not.

The hypertension was associated with a lower number of schooling years (OR=3.466, 95%CI=1.057–11.367,  $p=0.040$ ) and fewer sources of income (OR=1.733, 95%CI=1.120–2.682,  $p=0.014$ ). The risk for being hypertensive is increased for each year of life by OR = 1.056 (95%CI=1.033–1.080,  $p < 0.001$ ) and for each unit of BMI by OR = 1.166 (95%CI=1.107–1.229,  $p < 0.001$ ).

The second best model is the one using cut-off 3 for the number of income source while the other variables are the same as in the previous model. Here, the risk for hypertension is 5.5 x higher in persons who have 0–2 sources of income compared to those who have 3 different sources of income (OR=5.549, 95%CI=1.627–18.930,

$p=0.006$ ). Both presented regression models explain 40% of variance of the dependant variable and are equally acceptable.

## Discussion

The Bayash Roma of Croatia are a socio-economically marginalized minority population characterized by staggering high unemployment rate and high rate of illiteracy. Their society still sustains a system of internal moral norms and social rules. In our recent study<sup>6</sup> it was shown that the majority of Bayash population in Croatia lives in large groups, completely segregated from the rest of the population, in the outskirts of villages (e.g. Village of Krušanec in Međimurje or Torjanci in Baranja) or small towns (e.g. settlements of Zlatnica and Groblje in Darda or Rupa in Beli Manastir) with non-existent or unsatisfactory essential facilities, such as sewage, waterworks, garbage collection, or roads. The Bayash families sometimes cannot afford connection to existing public utilities<sup>6</sup>. Poverty and cultural specificities of the Bayash are the major reasons for the persistence of traditional lifestyle.

There is a strong cultural basis for obesity, tobacco use, fatty diet, and inbreeding present in this population – which all predispose them to hypertension and other CVD risk factors. The Roma, worldwide, are among the populations that are experiencing a rapid transition from traditional lifestyle, characterized by high levels of physical activity and a diet with relatively low energy, to a more sedentary lifestyle with an excess of caloric intake<sup>16</sup>.

Arterial hypertension is a complex disorder that represents one of the most important modifiable risk factors for myocardial infarction, stroke, end-stage renal disease and peripheral vascular disease<sup>17</sup>. The other modifiable CVD risk factors include dyslipidemia, unhealthy diet, physical inactivity and stress while non-modifiable risks are advanced age, gender, ethnicity, and familial susceptibility. The Interheart study<sup>18</sup> estimate that nine potentially modifiable risk factors contribute to more than 90% of CHD worldwide, and more than 95% of patients who died from CHD had at least one of the four traditional risk factors (smoking, hypertension, hyperlipidemia and diabetes)<sup>18</sup>.

The significant association of hypertension with traditional CVD risk factors – age and BMI – was confirmed in the present study, while no association with smoking habits was found in this population. Smoking is a part of traditional Roma life-style and with 70% of smokers in the population<sup>6</sup> it can be assumed that almost the entire population is exposed to this risk factor in their family environment. Therefore, due to homogenously distributed risk from smoking, this factor did not show to be a significant predictor of hypertension.

The here presented results showed no differences in the prevalence of hypertension among regions and between sexes. The fact that the previous report<sup>6</sup> indicated that those differences existed, requires further explana-

**TABLE 6.**  
PREVALENCE OF HYPERTENSION FOR THE AGE GROUP 35–64 YRS IN CROATIAN BAYASH ROMA COMPARED TO SEVERAL EUROPEAN AND AMERICAN POPULATIONS<sup>23,25</sup>.

Population (sorted by increasing order of prevalence)	Total (%)	Men (%)	Women (%)	M – W
Canada <sup>25</sup>	27.4	31.0	23.8	7.2
North America <sup>25</sup>	27.6	30.4	24.8	5.6
USA <sup>25</sup>	27.8	29.8	25.8	4.0
Bayash (present study)	35.0	30.4	37.2	–6.8
Italy <sup>25</sup>	37.7	44.8	30.6	14.2
Sweden <sup>25</sup>	38.4	44.8	32.0	12.8
England <sup>25</sup>	41.7	46.9	36.5	10.4
Croatia <sup>23</sup>	42.8	46.7	39.1	7.6
Europe <sup>25</sup>	44.2	49.7	38.6	11.1
Spain <sup>25</sup>	46.8	49.0	44.6	4.4
Finland <sup>25</sup>	48.7	55.7	41.6	14.1
Germany <sup>25</sup>	55.3	60.2	50.3	9.9

The same standard criteria for hypertension were used in all studies (ie BP  $\geq$ 140/90 mm Hg or treatment with antihypertensive medication)

tion. It lies in the discrepancy of the definition of hypertension applied in the two studies. According to the first study, hypertensives were all those who had a diagnosed hypertension (who are under medication), while in the present study the definition was expanded to all those with elevated measured blood pressure. When these broader criteria were applied, all regional and sex differences disappeared.

The prevalence of cardiovascular disease as well as morbidity and mortality due to other non-communicable diseases are increasing in transitional and developing countries. Croatia, with CVD mortality falling within the range 35%–65% of total mortality, is considered to be at the third stage of epidemiological transition. The CVD epidemic (predominantly ischemic heart disease and cerebrovascular disease) in transition countries is due to the increased rates of hypertension, obesity, smoking and sedentary lifestyle<sup>18</sup>. The decrease of CVD prevalence in developed countries is a well documented phenomenon. In accordance with this pattern are e.g. findings of significantly lower prevalence of hypertension (28 %; M: 29.8%; F: 25.8%) in USA<sup>19</sup>, as an economically developed country, compared to Croatia (37.5%; M: 35.2; F: 39.7%)<sup>19</sup>.

In economically developed countries the inverse relationship exists between the socio-economic status and the presence of CVD risk factors and CVD mortality<sup>20</sup>. For example, in 58 American Gypsies<sup>10</sup> the prevalence of hypertension is as high as 73%. Other contrasting examples of this relation are the »first nations« of Canada<sup>21</sup> and urban Swedes<sup>22</sup>.

In spite of the above mention assumptions, the Bayash population in Croatia is characterized by a low prevalence of hypertension. Namely, we found that Bayash Roma had a much lower prevalence of arterial hypertension (24.8%) when compared with the general population of Croatia (27.7–44.2%)<sup>19,23–24</sup>. The analysis by age groups allow us to conclude that the lower prevalence of hypertension in Bayash than within the general population of Croatia<sup>23</sup> is not due to comparative younger average age of the Bayash sample (Figure 3). The comparison of hypertension prevalence in various countries within age group 35–64 yrs. and applying the same criteria for hypertension revealed that the prevalence in Bayash is the lowest among European populations<sup>25</sup> and it falls near the levels found in USA and in Canada<sup>25</sup> (Table 6).

The expected connection of the low socio-economic status with the increase of cardiovascular disease risk factors is (smilingly) in sharp contrast to the findings of the present study. The vivid explanation for the obtained relations could be found in the work of Ezeamama et al. (2006)<sup>20</sup>, who, within the context of socio-economic development in the Samoan archipelago, demonstrated the positive associations between the socio-economic status and traditional CVD risk factors at the early stages of economic development. Those authors also suggest that this relation changes along with the economic development of the population (Figure 4)<sup>20</sup>, with high incidence of traditional CVD risk factors characterizing the developing societies while their low incidence of CVD risk fac-




Figure 3. Comparison of the prevalence of hypertension in Bayash Roma and in general population of Croatia<sup>23</sup> by sex and age.




Figure 4. Theoretical association between SES and CVD risk factors by level of economic development according to Ezeamama et al. (2006)<sup>20</sup>. Copyright permission will be requested from Soc. Sci. Med./

tors is a mark of economically developed societies as well as it is for the opposite – the traditional undeveloped societies.

In case the above-described association between socio-economic status and cardiovascular disease risk factors by the level of economic development is universally applicable, we could hypothesize that the Bayash Roma population of Croatia – who in spite of their low socio-economic status and the presence of other risk factors for CVD are not developing CVD – are socio-economically still in a »pre-developing stage«. If this assumption is valid, we can expect that along with the westernization of their life-style and the increase of their economic power the increase in prevalence of hypertension in Bayash Roma of Croatia will occur. However, it is interesting that within the here presented Roma sample the »developed« model could be observed – with lower blood pressure values among the richer Roma.

Considering the relation between measured hypertension (mHT) and previously diagnosed hypertension (dHT) in Bayash population, it should be pointed out that as long as hypertension is not a prevalent health problem for Bayash Roma population, the given rates of unsuccessfully treated hypertension are not alarming. But it has to be kept in mind that presently as much as 74% of persons with elevated BP do not have diagnosed hypertension. And if the prevalence of HT increases (which we expect to happen) the proportion of successfully treated persons in this population will be minute if the health care and health awareness remain at the present-day level. Bearing in mind the chronic nature of car-

diovascular disease and its associated long-term disability, mortality and financial cost, it is crucial that effective early cardiovascular disease risk prevention be applied to Roma minority population in Croatia.

Summing up we can state that although the prevalence of hypertension in Croatia is within the range of other European populations, the Bayash Roma minority population is characterized by its lower frequencies. This interesting finding surely calls for attention and various environmental as well as genetic hypotheses (i.e. the impact of genetic isolation and inbreeding) should be tested more thoroughly in the future.

In stressing out the importance of CVD prevention, it is important to take into consideration the necessity for the socially marginalized and economically deprived groups not only to be included socio-economically into modern society but for a parallel the public health policy based on preventive programs should be developed.

## Conclusions

The prevalence of hypertension in the Bayash Roma is almost half the magnitude of what is usually reported for the general population of Croatia. The prevalence of hypertension in Bayash showed no significant difference between the region of Baranja and Međimurje. The significant association of hypertension with age and BMI was confirmed and the importance of non-traditional SES-related CVD risk factors was highlighted.

Although the relation of blood pressure level and SES indicators within this population was negative (as is commonly found in developed populations), low prevalence of hypertension along with poverty, persistence of traditional lifestyle and with the presence of other risk factors for CVD – requires that this finding be taken with precaution. Namely, we expect that in the future – with westernization of their life-style and the increase of the economic power – the prevalence of hypertension and other CVD in the population of Bayash Roma in Croatia will increase. Therefore, it is important to recognize the need for the early cardiovascular disease risk factors prevention in this minority population.

## Acknowledgements

This research was supported by the grants of the Ministry of Science, Education and Sports of the Republic of Croatia (grants: 196-1962766-2747 to NSN and 196-1962766-2763 to BJ) as well as by the Wenner-Gren Foundation (grant: 7349 to IMK).

## REFERENCES

1. MURRAY, C. J. L., A. D. LOPEZ (Eds.): The global burden of disease: a comprehensive assessment of mortality and disability from disease, injuries and risk factors in 1990 and projected to 2020. (Harvard School of Public Health, Boston, 1996). — 2. DAVEY SMITH, G., M. EGGER, *BMJ*, 307 (1993) 1085. — 3. KAPLAN, G. A., J. E. KEIL, *Circulation*, 88 (1993) 1973. — 4. KAPLAN, G. A., J. W. LYNCH, *Prev. Med.*, 29

(pt2) (1999) (pt2) S30-S35. — 5. KRIEGER, N., *Am. J. Public Health*, 95 (2005) 2155. — 6. ŠKARIĆ-JURIĆ, T., I. MARTINOVIĆ KLARIĆ, N. SMOLEJ NARANČIĆ, S. DRMIĆ, M. PERIČIĆ SALIHOVIĆ, L. BARAČ LAUC, J. MILIČIĆ, M. BARBALIĆ, M. ZAJEC, B. JANIČIJEVIĆ, *Croat. Med. J.*, 48 (2007) 708. — 7. YUSUF, S., S. REDDY, S. OUNPUU, S. ANAND, *Circulation*, 104 (2001) 2746. — 8. ZAJC, M., N. SMOLEJ NA-



- RANČIĆ, T. ŠKARIĆ-JURIĆ, J. MILIČIĆ, M. BARBALIĆ, K. MELJANAC SALOPEK, I. MARTINOVIĆ KLARIĆ, B. JANIČIJEVIĆ, Coll. Antropol., 30 (2006) 783. — 9. ŠKARIĆ-JURIĆ, T., M. ZAJC, N. SMOLEJ NARANČIĆ, M. BARBALIĆ, M. PERIČIĆ SALIHOVIĆ, L. BARAČ LAUC, Coll. Antropol., 30 (2006) 761. — 10. THOMAS, J. D., M. M. DOUCETTE, D. C. THOMAS, J. D. STOECKLE, Lancet, 2(8555) (1987) 377. — 11. KRAJCOVICOVA-KUDLACKOVA, M., P. BLAZICEK, E. GINTER, V. SPUSTOVA, Bratisl. Lek. Listy, 103 (2002) 459. — 12. DOLINSKA, S., M. KUDLACKOVA, E. GINTER, Bratisl. Lek. Listy, 108 (2007) 207. — 13. GOVERNMENT OF THE REPUBLIC OF CROATIA: National Programme for Roma: Action Plan for the Decade of Roma Inclusion 2005-2015. Available from URL: [www.vlada.hr/nacionalniprogramromi/](http://www.vlada.hr/nacionalniprogramromi/). — 14. FRASER, A.: The Gypsies. (Blackwell Publishers, Oxford, 1992). — 15. WEINER, J. S., J. A. LOURIE: Practical human biology. (Academic Press, New York, 1981). — 16. VOZAROVA DE COURTEN, B., M. DE COURTEN, R. L. HANSON, A. ZAHORAKOVA, H. P. EGYENES, P. A. TATARANNI, P. H. BENNETT, J. VOZAR, Diabetes Res. Clin. Pract., 62 (2003) 95. — 17. BRANDY, K., N. D. DICKSON, J. BLACKLEDGE, M. H. IHAB, Am. J. Med. Sciences, 332 (2006) 211. — 18. CAREVIĆ, V., M. RUMBOLDT, Z. RUMBOLDT, Acta Med. Croatica, 61 (2007) 299. — 19. JELAKOVIĆ, B.: Epidemiologija arteriske hipertenzije u Hrvatskoj. In Croat. (Predavanje: Simpozij: Prostorna distribucija populacijskih kardiovaskularnih rizika u Hrvatskoj, Akademija medicinskih znanosti Hrvatska, Zagreb, 2005). — 20. EZEAMAMA, A. E., S. VIALI, J. TUTELE, S. T. McGARVEY, Soc. Sci. Med., 63 (2006) 2533. — 21. ANAND, S. S., S. YUSUF, R. JACOBS, Lancet, 358 (2001) 1147. — 22. ENGSTROM, G., I. JERNTRÖP, H. PESSAH-RASMUSSEN, Stroke, 32 (2001) 1098. — 23. ERCEG, M., V. HRABAK-ŽERJAVIĆ, U. IVIČEVIĆ, A. HEMIK: Hrvatska zdravstvena anketa: Povišen krvni tlak. In Croat. (Predavanje: Simpozij: Prostorna distribucija populacijskih kardiovaskularnih rizika u Hrvatskoj, Akademija medicinskih znanosti Hrvatska, Zagreb, 2005). — 24. TUREK, S., I. RUDAN, N. SMOLEJ-NARANČIĆ, L. SZIROVICZA, M. ČUBRILO-TUREK, V. ŽERJAVIĆ-HRABAK, A. RAK-KAIĆ, D. VRHOVSKI-HEBRANG, Ž. PREBEG, M. LJUBIČIĆ, B. JANIČIJEVIĆ, P. RUDAN, Coll. Antropol., 25 (2001) 77. — 25. WOLF-MAIER, K., R. S. COOPER, J. R. BANEGAS, S. GIAMPAOLI, H.-W. HENSE, M. JOFFRES, M. KASTARINEN, N. POULTER, P. PRIMATESTA, F. RODRIGUEZ-AR-TALEJO, B. STEGMAYR, M. THAMM, J. TUOMILEHTO, D. VANUZZO, F. VESCIO, JAMA, 289 (2003) 2363.

*T. Škarić-Jurić*

*Institute for Anthropological Research, Gajeva 32, 10000 Zagreb, Croatia  
e-mail: tanja@inantro.hr*