



Review

DOI: 10.17508/CJFST.2022.14.1.04

Benefits of using edible insects as alternative protein source: A Review

 Nura Abdullahi^{1*}, Enerst Chukwusoro Igwe², Munir Abba Dandago¹

¹Department of Food Science and Technology, Kano University of Science and Technology, Wudil, P.M.B 3244, Kano State, Nigeria

²Department of Food Science and Technology, Nnamdi Azikiwe University, PMB 5025, Awka, Anambra State, Nigeria

ARTICLE INFO

Article history:

Received: January 11, 2021

Accepted: July 15, 2021

Keywords:

entomophagy
protein substitute
meat alternative
insect-eating
unconventional protein

ABSTRACT

There is an urgent need for alternative protein sources due to the rapid population growth, climate change, environmental degradation by pollution, food-fuel competition and the reduction in arable land for agricultural use. Conventional livestock production is also deleterious to the environment due to the production of greenhouse gasses and ammonia. This article provides insights into the potentials of edible insects as novel food ingredients. The manuscript provides concise explanations for the need of embracing additional protein sources, edible insects consumption and their nutritional benefits and environmental and economic advantages of using edible insects as food. Literature was gathered through an online search on the Science Direct database and Google Scholar, relevant papers published between January 2002 and November 2020 were cited. Edible insects are good source of essential nutrients. They are rich in proteins and essential amino acids, contain good quality lipids and significant amounts of important minerals. They are potential source of proteins for humans and animals. They can play an important role in global food security by providing essential nutrients to the increasing global population. This can only be achieved when more attention is given to their production and processing. Creating awareness among new consumers on their nutritional and environmental benefits and the development of food products with appealing sensory properties will surely improve their acceptance as food.

Introduction

The practice of insect-eating, also known as entomophagy, is as old as the history of humans (Sun-Waterhouse et al., 2016; Cartay et al., 2020; Das et al., 2019). Insects were a vital source of food to early men and nowadays they play an imperative role in the nutrition of Latin America, Africa, and Asia (Srivastava et al., 2009; Fernandez-Cassi et al., 2019). Edible insects are among the most neglected and underutilized forest resources (Muimba-Kankolongo et al., 2015) with great potentials in providing nutritional security and income at local and international levels (Padulosi et al., 2018).

Edible insects are excellent source of essential nutrients and bioactive compounds (Sun-Waterhouse et al., 2016), they are rich in proteins with good amino acids composition, contain mono and polyunsaturated fatty acids, significant amounts of indispensable minerals (Zielińska et al., 2015) and petty amounts of cholesterol (Giordano et al., 2018). Protein, fat, mineral, and vitamin contents of edible insects are similar to those of meat (Baiano, 2020). Insect farming can protect the environment and provide proteins for pets, livestock, poultry, aquaculture, and humans (Tomberlin et al., 2015; Govorushko, 2019). In addition to the provision of nutrients, edible insects are also natural renewable resources that degraded organic waste (Das et al., 2019).

*Corresponding author E-mail: nurafst@gmail.com

There is a growing interest in the development and marketing of novel food due to the increasing demand for protein-rich foods and the obligatory need to diversify protein sources (Pippinato et al., 2020). Passion to reconsider ignored food sources and incorporate them into modern recipes has attracted the interest of researchers, chefs, and businessmen (Testa et al., 2016). There is also a rising demand for conventional protein substitutes that are green and cost-effective to meet up the diverse nutritional, sensory, and safety requirements (Fasolin et al., 2019).

The need for additional sources of protein

There is a rising demand for conventional protein substitutes that are green and cost-effective to meet up the diverse nutritional, sensory, and safety requirements (Fasolin et al., 2019). Livestock production is among the major anthropogenic activities that lead to eco-degradation and global warming (Abbasi et al., 2016). Rapid climate changes, environmental degradation by pollution, diminishing resources, and accelerated population growth also cause a serious setback to the global food supply (Dermody and Chatterjee, 2016). The production of conventional proteins is associated with many environmental problems. Intake reduction of about 90% of the current consumption is required to curtail the sustainability issues related to meat production (Videbæk and Grunert, 2020). Providing sufficient amounts of protein to the exponentially growing world population, projected to be 9.7 billion by 2050, will be a serious challenge (Tomberlin et al., 2015; Gallo and Federico, 2018). Moreover, income and population growth in some countries cause a significant increase in the consumption of animal protein (Gallo and Federico, 2018). An increase of 60-70% in animal product consumption is expected (Makkar et al., 2014).

Conventional sources will not be enough to sustain the fast-growing population (Rao, 2016). Alternative sources of proteins are required to complement the projected shortfall (Megido et al., 2014; Tang et al., 2019) and deal with the rising costs of animal proteins, environmental pressures (van Huis et al., 2013), the decline in the availability of arable land used for agricultural purposes (van Huis, 2015; Nadeau et al., 2015) and food-fuel competition (Makkar et al., 2014). An exponential increase in food production is needed (van Huis et al., 2013). A production increase of at least 60% is required to meet up with the demand (New, 2013). There is a need to investigate and explore other sources of protein in addition to traditional ones (Poelaert et al., 2018; Patel et al., 2019). This can be achieved through innovative

changes in the food supply (Dicke, 2018). The high pressure on protein and other essential resources hasten the interest of using insects as food and feed (Gjerris et al., 2016). Insects are an important source of food to the diversity of animal species including humans (Srivastava et al., 2009). Edible insects are among the leading alternative protein sources that are receiving adequate attention (Mancini et al., 2019). Nutrients deficit can be solved by carefully studying local insects and chose the species with similar nutrients profiles (Nadeau et al., 2015).

Developing countries are battling food insecurity due to an increase in population and a decrease in food production (Gahukar, 2011). The majority of the people in these countries are suffering from hunger and undernutrition, and food-related diseases are affecting more than 50% of their population (Ramos-Elorduy, 2009). The risk of micronutrient deficiency is increasing in these countries because the most of the staple foods consumed are deficient in some essential nutrients including essential amino acids (Lategan, 2019). Most of these countries have abundant wild insects that can solve or minimize these problems, but taboos associated with edible-insects, their seasonal and localized nature prevent their consumption globally year-round (Nantanga and Amakali, 2020). Edibleinsects can be a part of the global food recipes when these hurdles are overcome (Akhtar and Isman, 2018; Patel, 2019; Caporizzi et al., 2019; Skrivervik, 2020).

Insect consumption

Entomophagy is an old tradition that is still active in many traditions (Yen, 2015a). The dietary pattern of any community depends on their surrounding ecosystem and resources available within their premises (Ramos-Elorduy, 2009). Insects are consumed comfortably and habitually in societies that accepted their consumption and are viewed as an important source of nutrition and medicine (Costa-Neto and Dunkel, 2016). Even children were encouraged to consume locally processed insects in some localities (Bose, 2020). The effects of lifestyle changes on entomophagy are quite insignificant in some parts of the world as the tradition remains the same over time (Dube et al., 2013). Eating insects as food is determined by culture, religious practice (van Huis et al., 2013; Niassy et al., 2016; Ruby and Rozin, 2019b), dietary needs (Yen, 2015b), location, and availability (Chakravorty, 2014). Insects are predominantly consumed more in tropical countries, because they occur naturally in mass (van Huis, 2015). Insects are favoured food in some parts of Latin America, West Africa and Southeast Asia (Ruby et al.,

2015). They are also an important diet to Australian indigenes, Arabs in the Middle East, and South and Central Americans (Chung, 2008). Insects are consumed more in Thailand, China, Japan, Korea, and Indonesia than in any other nation (Chung, 2008). Thailand's insect industry is well developed with more than 20,000 registered insect farming enterprises (New, 2013).

Entomophagy is more common in rural areas (Chung et al., 2002) and the consumption of species can be restricted to a tribe or community (Niassy et al., 2016). In some parts of Africa, more than 90% of a community can participate in entomophagy (Dube et al., 2013). Central African region is the leading consumer of insects on the continent (Kelemu et al., 2015). Insects are consumed more by women and children than adult men in Africa (Akullo et al., 2017). A different inclination may likely emerge in the Western countries as men are more interested in entomophagy than women in Denmark (Videbæk and Grunert, 2020), America (Ruby and Rozin, 2019), and Italy (Tuccillo et al., 2020).

Insects that are consumed by humans are those that can easily gather in a huge quantity and do not have any unpleasant odour (Srivastava et al., 2009). Insects are consumed as a whole, as processed powder or paste, as extracts such as protein isolate (Williams et al., 2016), raw, cooked, or as ingredients in food preparation (Gahukar, 2011; Ayensu et al., 2019; Imathiu, 2020). They are consumed as either snacks or side meal (Akullo et al., 2017), while in some cultures insects, and other arthropods, are consumed as a staple (Srivastava et al., 2009). In Africa, insects are mostly consumed as complimentary food, and as the main dish in very few cases (Niassy et al., 2016). In India, raw insects are consumed (after trimming wings and legs) with processed fish (Shantibala et al., 2012), while bamboo worms and honey bees are processed and incorporated into condiments such as curry powder (Shantibala et al., 2012). Termites are boiled, blended with spices, mold into a ball, and cooked in banana leaves to serve as a main dish in Uganda (Akullo et al., 2017). In Northeast India, soft-bodied insects are consumed raw or processed into a condiment, while hard-bodied insects are consumed either roasted or fried (Shantibala et al., 2012).

House cricket (*Acheta domesticus*) is the most reported edible insect in the literature. Other commonly consumed insects in the world are grasshoppers, locusts beetles, termites, and caterpillars (Cartay et al., 2020; van Huis, 2020a). Different parts of insects at various stages of development are consumed as food. It could be eggs, larvae or nymphs, pupae, or adults (Chung, 2008). Insects are consumed by many as just traditionally inherited diets without

knowing their nutritional potentials (Akullo et al., 2017). Non-native insect consumers prefer to eat insect products such as flour than eating whole insects (Ruby et al., 2015). People interested in tasting novel foods will accept new food, such as insects, quite easily unlike people with the opposite opinion (Hartmann and Bearth, 2019).

Benefits of using insects as food

More attention was given to entomophagy in recent years due to identified nutritional and environmental benefits (Castro-López et al., 2020). Eating insects is encouraging due to the number of benefits including better nutrient composition, feed conversion efficiency, organic waste conversion, fewer requirements for water, and low greenhouse gas emission (Niassy and Ekesi, 2016). Insects are used in human medicine, as an aphrodisiac, animal feed, and bait during fishing, also as an agent for biodegradation (Niassy et al., 2016). Insects have a shorter production duration and their feed conversion rate is higher than that of higher animals (Megido et al., 2014; Fernandez-Cassi et al., 2019). They are poikilothermic, they do not use the energy generated through metabolism for body temperature control (Alexander et al., 2019). They have higher fecundity and require small space when compared with conventional livestock (Ayensu et al., 2019). The edible insect has a higher protein value and production efficiency when compared with conventional livestock (Dermody and Chatterjee, 2016). Insects eat inedible biomass of plants and produce high-quality proteins with all essential amino acids (Tong et al., 2011). Insects such as termites, grasshoppers, caterpillars, weevils, and houseflies are by weight richer in protein than beef, chicken, pork, or lamb (Srivastava et al., 2009). Entomophagy is expected to be among the key solutions to global food shortage through the provision of a high amounts of good quality proteins, fats, vitamins and minerals (Tang et al., 2019).

Nutritional and health benefits

The use of insects as food by various communities contributed to food security in many ways (Müller et al., 2016). The nutritional value of edible insects will promote their consumption and they may appear on the menu of many eateries in the future (Bednářová et al., 2013). Edible insects can play an important role in ensuring food and feed security through the development of safe and adequate rearing and processing methods that will preserve their nutritive value (Rumpold and Schlüter, 2015). In doing so, a

balance must be maintained between safety and nutritional qualities (Grabowski and Klein, 2017). Growing location, feed composition, killing and processing methods significantly affect the nutritional compositions of edible insects (Table 1). Any of these factors can cause noticeable variations in nutritional composition, even within species. Sex is another factor that affects the nutritional composition of edible insects. Kulma et al. (2019) reported variation in proximate composition and chitin contents between male and female house cricket, such that the male cricket contains more proteins and chitin, while female contains more lipids. The protein value of insects is affected by the rearing condition and harvesting age (Morales-Ramos et al., 2018). The design of an insect rearing facility can also have effects on protein quality (Berggren et al., 2018). Compositions of insect nutrients depend on the rearing method, feed composition (Chinarak et al., 2020) and their ability to convert feed into body mass (Straub et al., 2019). Growing season significantly affects protein, fat, and mineral contents, location and season affect vitamin B₁₂ content, location affects mineral contents within the same season (Ssepuuya et al., 2019). Gut microbiota composition and eating patterns also determine vitamin B₁₂ content (Schmidt et al., 2019). Zou et al. (2017) reported significant variations in the lipid contents of different strains of silkworm pupae. Nutrients in edible insects can promote good health (Ayensu et al., 2019). Some species provide health-promoting constituents (Dicke, 2018). Edible insects are seen as an alternative to conventional animal proteins in terms of their nutritional qualities (Roos and van Huis, 2017). The nutritional content of many insects is similar to that of conventional livestock (Kelemu et al., 2015; Lillford and Hermansson, 2020) and in some cases is superior to conventional livestock, milk, and eggs (Nadeau et al., 2015). Insect proteins are cheap protein substitute that can be used to fight malnutrition among the vulnerable groups in developing and underdeveloped nations (Ayensu et al., 2019). The likelihoods of using insects as conventional protein alternative depend greatly on understanding their nutritional profile (Payne et al., 2016) and confirming the claimed health benefits using acceptable health researches including human trial (Roos and van Huis, 2017). Most edible insects contain more proteins and ash (minerals) than conventional protein sources, but possess lower levels of essential amino acids (Table 1). Edible insects can provide carbohydrates, proteins, fats, minerals, and vitamins (Gahukar, 2011; Ayensu et al., 2019). The proximate profile of most insects is better than that of other meats (Gere et al., 2019). The chitin in the edible insect with exoskeleton can be a

good source of fiber (Koko and Mariod, 2020). The most common fiber in edible insects is N-acetylglucosamine (chitin), located at the exoskeleton and also the main component of the insect body (Caporizzi et al., 2019).

Certain insects and insect products such as honey and pollen are used as both food and medicine (Chung, 2008). Various species of insects contain bioactive compounds with potential health benefits that can contribute to combating global health challenges (Roos and van Huis, 2017). The essential *Lachnospiraceae* bacteria, responsible for fiber digestion, is among the natural biota found in house crickets (Frigerio et al., 2020). Chemicals produced by insects for self-defense can be used in the production of antimicrobial and anticancer drugs (Srivastava et al., 2009). Insects have enormous potentials in providing essential nutrients and active substances not only to humans but also to livestock (Tang et al., 2019). Livestock fed with insects tends to have better growth and amazing sensory qualities as reported in chicken by Waithanji et al. (2019).

Most edible insects have favourable nutrient contents, some can possess a low amount of iodine, vitamin A and some essential amino acids (Lategan, 2019). Insects, such as silk moth, can contain a large amount of proteins with all essential amino acids, numerous essential minerals, vitamins, and fatty acids (Tong et al., 2011). *Gryllus assimilis* and *Zophobas morio* are rich in palmitic, oleic, linoleic, and essential minerals (calcium, magnesium, iron, and zinc), they have protein contents of 65.52% and 46.80% and lipid contents of 21.80% and 43.64%, respectively (Araújo et al., 2019).

Edible insects can provide well-balanced proteins by converting organic waste (Payne et al., 2016). Edible insects are loaded with essential nutrients, they are rich in protein and can provide a well-balanced diet rich in essential amino acids since the protein value is very similar to that of meat and fish (Das et al., 2019). They are an excellent source of essential amino acids with easy digestibility and can be used to tackle chronic malnutrition when large production is promoted (Cartay et al., 2020). Insect protein has good digestibility in its raw form and after processing, and its amino acid profile meets the human dietary requirements (Poelaert et al., 2018). Edible insects contain more proteins than plant protein sources, including soybean (Rumpold and Schlüter, 2015) and they meet the WHO recommendation for amino acids (Tang et al., 2019). Termites have more than 35% of proteins, while crickets, grasshoppers, and locusts can contain more than 60% of proteins (Rumpold and Schlüter, 2015). Silkworm powder is more nutritious than snail, fish, chicken, beef, and pork (Tong et al.,

2011). Crickets, honeybees, and palm weevil larvae were reported to be nutritionally superior to meat (Payne et al., 2016). González-Escobar et al. (2018) associated the higher protein value of insects with the abundant indigenous nitrogen-fixing bacteria.

The concentration of essential amino acids in many edible insects meets FAO/WHO standards (Oibiokpa et al., 2018). The most common limiting amino acids in insects are methionine (Zeece, 2020), cysteine, tryptophan (Tang et al., 2019), and lysine (Testa et al., 2016), but can contain up to 46 to 96% of essential amino acids (Testa et al., 2016). Tryptophan is limited in long-horned grasshopper (Ssepuuya et al., 2019), locust, and cricket; lysine in the scarab beetle; leucine in the silkworm (Köhler et al., 2019); lysine, methionine, and cysteine in yellow mealworms (Severini et al., 2018); methionine and cysteine in house cricket (Kulma et al., 2019); methionine in beetle and cricket (Manditsera et al., 2019); phenylalanine in cockroaches (Kulma et al., 2020), and methionine and lysine in black soldier fly (Surendra et al., 2020). Termite, moth caterpillar, and grasshopper possess low levels of sulfur-containing amino acids (methionine and cysteine) and isoleucine (Oibiokpa et al., 2018).

Insect larvae usually contain more fat than conventional meat and insect, in general, contains healthier lipids, because their polyunsaturated fatty acids content is much more superior when compared with that of conventional meat (Table 1). The higher fat content in the larvae of some insects provides attractive flavour and facilitates edibility (Tang et al., 2019). Insects contain a significant amount of unsaturated fatty acid and can be used as a source of polyunsaturated fatty acids in the landlocked countries with limited access to fish (Koko and Mariod, 2020). The lipid of silkworm pupae contains more than 70 % polyunsaturated fatty acid which is rich in α -linolenic acid (Zou et al., 2017). Omega-3 and omega-6 fatty acid contents of mealworms are comparable to that of fish (van Huis et al., 2013). Feeding insects with fish waste increases their polyunsaturated fatty acid content. An increase in polyunsaturated fatty acid content was observed in black soldier fly larvae fed with fish waste (Barroso et al., 2019). The omega-3 and omega-6 fatty acid ratio in *Eulepida mashona* and *Henicus whellani* is well-balanced (Manditsera et al., 2019). *Imbrasia* caterpillars have appreciable levels of omega-3 and omega-6 fatty acids that can prevent several diseases, *Imbrasia truncata* and *I. epimethea* contains healthy lipids and all essential amino acids at the levels higher than that recommended by FAO (Mba et al., 2019). Oil extracted from *Tenebrio molitor* and *Pachymerus nucleorum* larvae noticeably

reduced cholesterol and glucose levels in the experimental rat (Alves et al., 2019).

Edible insects are rich in minerals such as copper, iron, magnesium, manganese, phosphorous, selenium, and zinc (Rao, 2016). Entomophagy can be a cheap means for combating iron and zinc deficiencies in developing countries (Christensen et al., 2006). Incorporation of the edible insect into cereal-based food increases iron consumption and facilitates growth in infants (Ayensu et al., 2019). The calcium, sodium, manganese, and zinc contents of black soldier fly larvae meal are higher than that of soybean meal and other plant protein sources (Fisher et al., 2020). The consumption of 50 g of wild *Eulepida mashona* and *Henicus whellani* can supply 30% of daily recommended levels of zinc and 50% of daily iron requirement (Manditsera et al., 2019). Earthworm, house cricket, and mealworm larvae have excellent vitamin content (Gere et al., 2019). Cockroach, mealworm, cricket, and grasshopper contain a significant amount of vitamin B₁₂ (Schmidt et al., 2019).

Environmental benefits

Protein production using insects is more friendly to the environment when compared to the resource intense conventional livestock farming (Stull et al., 2018). Insect-based foods were perceived to have lower environmental impact than beef products (Kusch and Fiebelkorn, 2019a). Entomophagy is expected to reduce agricultural land footprint and significantly curtail food waste when universally accepted (Alexander et al., 2017). One of the reasons for promoting insect-eating is the concern about the sustainability of the global ecosystem (Waltner-Toews and Houle, 2017). The integration of insect protein into the human diet is a sustainable alternative to conventional animal proteins (Megido et al., 2014), because conventional animal production puts too much pressure on the environment (Kusch and Fiebelkorn, 2019b; Weindl et al., 2020a) to the level that threatens the integrity of the ecosystem (Weindl et al., 2020b). Promoting insect consumption as an alternative source of proteins will reduce the environmental impact associated with conventional livestock production (Yen, 2015a).

Major environmental challenges such as deforestation and global warming can be minimized when insects are incorporated into diets (Jensen and Lieberoth, 2019). When compared with conventional livestock, insects require less space (Rumpold and Schlüter, 2015) and insect farming generates lower greenhouse gas and ammonia than livestock farming (Oonincx et al., 2010; Testa et al., 2016).

Table 1. Proximate values, Minerals, total essential amino acids (EAA) and total polyunsaturated fatty acids (PUFAs) in commonly consumed edible insect species

Insect	Location/ Habitant	Feed	Killing method	Protein	Lipids	Ash	Fiber	Fe	Ca	P	Cu	Zn	EAA	PUFAs	Reference
Farmed species															
Crickets															
<i>Acheta domestica</i>	Thailand		Obtained as ready-to-eat, deep-fried	53.90				5.81	176.90	762	3.00	21.80	37.0		(Köhler et al., 2019)
	Thailand	Fed with grains and vegetable mix	Obtained as whole insect powder	48.48	28.75	3.50	7.64						50.60	36.17	(Osimani et al., 2018)
	Netherlands		Obtained as whole insect powder	64.4	23.7	4.9	6.9	7.7	103.6	1008	3.4	24.4			(Ribeiro et al., 2019)
<i>Gryllodes sigillatus</i>	Netherlands		Obtained as whole insect powder	65.3	23.5	4.2	7.1	4.7	117.3	782	4.9	16.8			(Ribeiro et al., 2019)
Cockroaches															
		Dog pellets, staled bread													
<i>Blaberus craniifer</i>	Czech Republic	bread, apple, carrot, and lettuce leaves	24 hrs Fasting and freezing	66.60	24.30	1.40	7.40								(Kulma et al., 2020)
		Dog pellets, staled bread													
<i>Zophobas morio</i> larvae	Czech Republic	apple, carrot, and lettuce leaves	24 hrs Fasting and freezing	48.10	34.00	0.7	5.3								(Kulma et al., 2020)
<i>Nauphoeta cinerea</i>	Brazil	fruit and vegetable remains	Boiled water	59.78	21.27	5.11	6.85								(de Oliveira et al., 2017)
<i>Tenebrio molitor</i> larvae	France	wheat bran, cereal, premix and NaCl	freezing (-18 °C)	65.50	24.50										(Azagoh et al., 2016)
Bombay locust (<i>Paungia succinata</i> L.)	Thailand		Obtained as ready-to-eat, deep-fried	36.31				3.45	65.60	267	1.69	8.22	33.2		(Köhler et al., 2019)
Scarab beetle (<i>Holotrichia</i> sp.)	Thailand		Obtained as ready-to-eat, deep-fried	28.92				9.10	75.10	260	1.08	8.80	36.7		(Köhler et al., 2019)
Mulberry silkworm (<i>Bombyx mori</i> L.)	Thailand		Obtained as ready-to-eat, steamed	50.20				3.18	107.60	526	0.94	15.50	43.9		(Köhler et al., 2019)
Sago palm weevil (<i>Rhynchophorus ferrugineus</i>) larvae	Thailand	Fed with sago palm trunk	Blanching for 5 minutes	18.0-28.5	52.4-60.1	2.4-2.9	3.8-4.5	2.63	43.3	494	1.03	8.33			(Chinarak et al., 2020)

Insect	Location/ Habitant	Feed	Killing method	Protein % Dry matter	Lipids Mg/100g	Ash g/100g	Fiber g/100g	Fe	Ca	P	Cu	Zn	AAA	PUFAs	Reference
Laboratory grown															
Mealworms (<i>Tenebrio molitor</i>)	Belgium	wheat flour, brewer's yeast, and wheat bran	24 hrs Fasting and freezing (-18 °C)	43.00	40.90	3.40								22.9	(Megidoa et al., 2018)
	Germany	wheat bran	Freezing (-21 °C)	52.00	26.75		6.10						40.8		(Kröncke et al., 2018)
Wild species															
Beetle (<i>Eulepida mashona</i>)	Zimbabwe		Degutting and freezing	54.47	7.98	4.69		25-54				16-21	28.8	24-35	(Mandisera et al., 2019)
Cricket (<i>Henicus whellani</i>)	Zimbabwe		Degutting and freezing	65.75	7.68	4.99		30-55				16-21	27.4	42-47	(Mandisera et al., 2019)
long-horned grasshopper (<i>Tetigonia viridissima</i>)	Kampala, Uganda		No information	34.2-45.8	42.2-54.3	1.79- 2.72	3.93- 5.34	48.6	70	517	1.91	11.24	49.4	63.5-64.7	(Ssepuuya et al., 2019)
Short-horned grasshopper (<i>Chondactis rosea</i>)	Arunachal Pradesh, India		Freezing	68.88	7.88	4.16	12.38	7.81	300		3.62	10.83	24.9		(Chakravorty et al., 2014)
Mole cricket, (<i>Brachytrapes orientalis</i>)	Arunachal Pradesh, India		Freezing	65.74	6.33	4.33	8.75	18.66	76.28		1.53	8.50	21.3		(Chakravorty et al., 2014)
Meadow grasshopper (<i>Chorthippus parallelus Zetterstedt</i>)	Local grassland Belgium		Fasting for 6 hours and freezing (-20 °C)	69.00	7.88	4.00									(Paul et al., 2016)
African palm weevil (<i>Rhynchophorus phoenicis</i>) larvae	Yaoundé, Cameroon		Freezing (-20 °C)	27.50	71.62	2.18								4.0	(Mba et al., 2017)
Arboreal ant (<i>Oecophylla smaragdina</i>)	Papumpare district, India		Freezing	55.28	14.99	2.59	19.84		74.67		0.85	18.97		8.19	(Chakravorty et al., 2016)
Conventional proteins															
Whole cow milk				24.4	33.5	5.1		0.15	118	96.56	0.03	0.33	49.08	0.138	
Egg (poultry, whole raw)				56.5	39.0	3.4		1.82	49.44	185	0.07	1.23	59.41	1.152	
Chicken				58.5	38.0	3.4		1.27	20.52	199	0.25	1.77	56.53	0.661	
Beef				56.9	40.3	2.6		2.22	6.50	201	0.07	4.64	57.9	1.008	(Longvah et al., 2017)
Pork				47.0	50.8	2.0		0.91	9.95	157	0.23	2.08	53.7	2.129	
Catfish				69.7	27.4	4.1		0.82	21.99	157	0.03	0.71	60.13	0.978	
Soybean				37.7	21.0	5.0	22.8	8.29	239	483	1.29	4.01	40.76	11.279	

Insects have a higher feed conversion efficiency and require less water and land (Rao, 2016).

Putrefied wastes from food processing can be valorized by insects. Barroso et al. (2019) demonstrate how fish waste can be converted into polyunsaturated fatty acids when used as insect feed, though, Skrivervik (Skrivervik, 2020) reported that most of the commercial insect farmers are not using food waste as insect feed. Black soldier fly can efficiently convert waste into a high-value protein with qualities similar to that of soy meal and fishmeal (Surendra et al., 2020). Despite the little negative environmental impacts associated with insect farming, the utilization of their by-products can have additional benefits to the environment. Ulmer et al. (2020) reported that using honeybee drone brood as a novel source of protein can minimize the environmental impacts of beekeeping by 8%.

Berggren et al. (2019) opposed the sustainability claims associated with edible-insects production. According to them, many ecological factors were not given appropriate considerations. These factors include the choice of right species, integrating insects into the agricultural system, their wastes disposal, and risk of mass harvesting to the local ecosystem.

Economic benefits

Edible insects have higher market potentials despite inadequate regulation and little value addition in their business (Odongo et al., 2018). The global market of edible insects was worth 55 million United States Dollars (USD) in 2017 and a gross addition of 43.5% is expected by 2024 (Koko and Mariod, 2020). The predicted increase in the demand for animal protein for both food and feed makes insect farming a lucrative and sustainable business that can promote income generation and provide employment (Niassy and Ekesi, 2016). The profitability of insect farming depends on their ability to eat from cheap, readily available, and unutilized organic wastes (van Huis, 2020b).

The low production cost, which is virtually zero in some developing countries, promotes edibleinsects consumption (Tang et al., 2019). Commercializing insect farming is important when looking at their nutritional and economical value (Srivastava et al., 2009). The production of insect protein is cheaper and more sustainable than the production of conventional proteins (Hartmann and Bearth, 2019), as conventional livestock production depends to a great extent on resources that are also suitable for human consumption (Dicke, 2018). Conventional animal protein is becoming more expensive due to obstacles impeding the development of the livestock industry in

developing countries. Continuous increase in the price of animal feed, fishmeal, soybeans, and grains intensifies the difficulties in accessing animal protein by many individuals (Niassy and Ekesi, 2016). Insect farming will provide a suitable alternative as many edible species can grow mainly on organic waste (Dicke, 2018), although, associating edible insects with waste utilization, upsurges their rejection (Rumpold and Langen, 2020). Insect farming also requires less technical and capital expenditure for rearing and harvesting equipment when compared with conventional livestock farming (Testa et al., 2016).

Insects provide rich proteins that are relatively affordable unlike meat proteins (Srivastava et al., 2009). Edible insects are becoming a fashionable solution to poverty by the role they play in many food systems (Tang et al., 2019). Insect gathering can offer jobs and incomes at both households and industrial scales of processing (van Huis et al., 2013) as insects are easily collected, processed, and sold with minimal technical and capital expenditure (Rao, 2016). The insect industry is expected to be prosperous through the development of products extracted from insects. Additives such as food colours can be extracted from insects (Tang et al., 2019).

Devastating insect pests can be exploited to reduce damage caused by these insects to food crops and to minimize the use of insecticides in the agricultural field (Flores et al., 2015). Edible insects such as locusts can cause devastating damage to crops (van Huis, 2020a). Eating insect pests such as grasshopper will reduce their population and can contribute to economic development (Chung et al., 2002).

Recommendation and conclusion

Recommendation

1. Alternative sources of protein are necessary since traditional rearing of beef, poultry and pork is not sustainable as the price of grains and, subsequently, that of meat are expected to raise in the decades to come.
2. As many people are willing to accept insects as food, research needs to be intensified in their production to develop the value chain.
3. A worldwide campaign on consumption of edible insects should be launched and emphasis should be given to countries with acute food shortages. Women should be engaged in the entomophagy campaign as they decide the types of food to be consumed by the family.
4. Since the allergy status of many edible insects is not fully understood, insect-based foods should be

consumed with caution. A clear understanding of edible insect allergic status will improve their acceptability. Individuals allergic to crustacean and other seafood must be cautious with insect-based foods. The possibility of blocking allergic reactions through the modification of processing conditions should be studied.

Conclusions

Edible insects can contribute to global food security when more attention is given to their production and processing. Accepting insects as novel food ingredient will improve their consumption and open more research areas on their safety, nutritional and health benefits and functionality. Insects gathering can provide jobs and incomes, save food crops, improve their safety, and reduce the cost of their production. Insects can also provide important industrial raw materials such as food supplements and other additives. Edible insects are excellent source of important nutrients. Most edible insects contain more proteins and ash (minerals) than conventional protein sources, they are good source of essential amino acids, fatty acids, and indispensable minerals. Insects larvae usually contain more fat than conventional meat and insects, in general, contain healthier lipids due to their good polyunsaturated fatty acids content. Accepting insects as food will improve the nutritional status of many, especially in developing countries, that are suffering from malnutrition and hunger.

Author Contributions:

Nura Abdullahi: Conceptualization, literature search, original draft preparation.

Ernest Chukwusoro Igwe: Supervision, redesign, writing-reviewing, literature search, and editing.

Munir Abba Dandago: Supervision, literature search, writing-review, and editing.

Funding: None

References

- Abbasi, T., Abbasi, T., Abbasi, S. A. (2016). Reducing the global environmental impact of livestock production: The minilivestock option. *Journal of Cleaner Production* 112, 1754–1766. <https://doi.org/10.1016/j.jclepro.2015.02.094>
- Akhtar, Y., Isman, M. B. (2018). Insects as an Alternative Protein Source. In *Proteins in Food Processing*; R. Y. Yada (Ed.): *Second Edition* (2nd ed., pp. 263–288). Elsevier Ltd. <https://doi.org/10.1016/B978-0-08-100722-8.00011-5>
- Akullo, J., Obaa, B. B., Acai, J. O., Nakimbugwe, D., Agea, J. G. (2017). Knowledge, attitudes and practices on edible insects in Lango sub-region, northern Uganda. *Journal of Insects as Food and Feed* 3(2), 73–81. <https://doi.org/10.3920/JIFF2016.0033>
- Alexander, P., Brown, C., Arneith, A., Dias, C., Finnigan, J., Moran, D., Rounsevell, M. D. A. (2017). Could consumption of insects, cultured meat or imitation meat reduce global agricultural land use? *Global Food Security* 15, 22–32. <https://doi.org/10.1016/j.gfs.2017.04.001>
- Alexander, P., Brown, C., Dias, C., Moran, D., Rounsevell, M. D. A. (2019). Sustainable Proteins Production. In *Proteins: Sustainable Source, Processing and Applications*; C. Galanakis (Ed.), (1st ed., pp. 1–39). <https://doi.org/10.1016/b978-0-12-816695-6.00001-5>
- Alves, A. V., Freitas de Lima, F., Granzotti da Silva, T., Oliveira, V. S. de, Kassuya, C. A. L., Sanjinez-Argandoña, E. J. (2019). Safety evaluation of the oils extracted from edible insects (*Tenebrio molitor* and *Pachymerus nucleorum*) as novel food for humans. *Regulatory Toxicology and Pharmacology* 102, 90–94. <https://doi.org/10.1016/j.yrtph.2019.01.013>
- Araújo, R. R. S., Benfica, T. A. dos S. R., Ferraz, V. P., Moreira Santos, E. (2019). Nutritional composition of insects *Gryllus assimilis* and *Zophobas morio*: Potential foods harvested in Brazil. *Journal of Food Composition and Analysis* 76, 22–26. <https://doi.org/10.1016/j.jfca.2018.11.005>
- Ayensu, J., Annan, R. A., Edusei, A., Lutterrodt, H. (2019). Beyond nutrients, health effects of entomophagy: a systematic review. *Nutrition and Food Science* 49(1), 2–17. <https://doi.org/10.1108/NFS-02-2018-0046>
- Azagoh, C., Ducept, F., Garcia, R., Rakotozafy, L., Cuvelier, M. E., Keller, S., ... Mezdour, S. (2016). Extraction and physicochemical characterization of *Tenebrio molitor* proteins. *Food Research International* 88, 24–31. <https://doi.org/10.1016/j.foodres.2016.06.010>
- Baiano, A. (2020). Edible insects: An overview on nutritional characteristics, safety, farming, production technologies, regulatory framework, and socio-economic and ethical implications. *Trends in Food Science and Technology* 100, 35–50. <https://doi.org/10.1016/j.tifs.2020.03.040>
- Barroso, F. G., Sánchez-Muros, M. J., Rincón, M. Á., Rodríguez-Rodríguez, M., Fabrikov, D., Morote, E., Guil-Guerrero, J. L. (2019). Production of n-3-rich insects by bioaccumulation of fishery waste. *Journal of Food Composition and Analysis* 82, 103237. <https://doi.org/10.1016/j.jfca.2019.103237>
- Bednárová, M., Borkovcová, M., Mleek, J., Rop, O., Zeman, L. (2013). Edible insects - Species suitable for entomophagy under condition of Czech Republic. *Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis* 61(3), 587–593. <https://doi.org/10.11118/actaun201361030587>
- Berggren, Å., Jansson, A., Low, M. (2019). Approaching Ecological Sustainability in the Emerging Insects-as-Food Industry. *Trends in Ecology and Evolution* 34(2), 132–138. <https://doi.org/10.1016/j.tree.2018.11.005>

- Berggren, Jansson, A., Low, M. (2018). Using current systems to inform rearing facility design in the insect-as-food industry. *Journal of Insects as Food and Feed* 4(3), 167–170. <https://doi.org/10.3920/JIFF2017.0076>
- Bose, P. (2020). *Forest foods for tribals in selected regions of India and their sustainability*. In Nutritional and Health Aspects of Food in South Asian Countries; J. Prakash, V. Waisundara, V. Prakash, (Eds.). Elsevier Inc. <https://doi.org/10.1016/b978-0-12-820011-7.00005-8>
- Caporizzi, R., Derossi, A., Severini, C. (2019). Cereal-Based and Insect-Enriched Printable Food: From Formulation to Post-Processing Treatments. Status and Perspectives. In Fundamentals of 3D Food Printing and Applications F. C. Godoi, B. R. Bhandari, S. Prakash, & M. Zhang (Eds.), (pp. 93–116). Elsevier Inc. <https://doi.org/10.1016/b978-0-12-814564-7.00004-3>
- Cartay, R., Dimitroi, V., Feldman, M. (2020). An Insect Bad for Agriculture but Good for Human Consumption: The Case of *Rhynchophorus palmarum*: A Social Science Perspective. In Edible Insects; H. Mikkola (Ed.), (1st ed., pp. 1–18). IntechOpen. <https://doi.org/http://dx.doi.org/10.5772/intechopen.87165>
- Castro-López, C., Santiago-López, L., Vallejo-Cordoba, B., González-Córdova, A. F., Liceaga, A. M., García, H. S., Hernández-Mendoza, A. (2020). An insight to fermented edible insects: A global perspective and prospective. *Food Research International* 137, 109750. <https://doi.org/10.1016/j.foodres.2020.109750>
- Chakravorty, J. (2014). Diversity of Edible Insects and Practices of Entomophagy in India: An Overview. *Journal of Biodiversity, Bioprospecting and Development* 01(03), 1–6. <https://doi.org/10.4172/2376-0214.1000124>
- Chakravorty, J., Ghosh, S., Jung, C., Meyer-Rochow, V. B. (2014). Nutritional composition of *Chondacris rosea* and *Brachytrupes orientalis*: Two common insects used as food by tribes of Arunachal Pradesh, India. *Journal of Asia-Pacific Entomology* 17(3), 407–415. <https://doi.org/10.1016/j.aspen.2014.03.007>
- Chakravorty, J., Ghosh, S., Megu, K., Jung, C., Meyer-Rochow, V. B. (2016). Nutritional and anti-nutritional composition of *Oecophylla smaragdina* (Hymenoptera: Formicidae) and *Odontotermes* sp. (Isoptera: Termitidae): Two preferred edible insects of Arunachal Pradesh, India. *Journal of Asia-Pacific Entomology* 19(3), 711–720. <https://doi.org/10.1016/j.aspen.2016.07.001>
- Chinarak, K., Chaijan, M., Panpipat, W. (2020). Farm-raised sago palm weevil (*Rhynchophorus ferrugineus*) larvae: Potential and challenges for promising source of nutrients. *Journal of Food Composition and Analysis* 92, 103542. <https://doi.org/10.1016/j.jfca.2020.103542>
- Christensen, D. L., Orech, F. O., Mungai, M. N., Larsen, T., Friis, H., Aagaard-Hansen, J. (2006). Entomophagy among the Luo of Kenya: A potential mineral source? *International Journal of Food Sciences and Nutrition* 57(3), 198–203. <https://doi.org/10.1080/09637480600738252>
- Chung, A. Y., Khen, C. V., Unchi, S., Momin, B. (2002). Edible insects and entomophagy in Sabah, Malaysia. *Malayan Nature Journal* 56(2), 131–44.
- Chung, A. Y. C. (2008). An Overview of Edible Insects and Entomophagy in Borneo Arthur. In *UN-FAO Workshop on Forest Insects as Food: Humans Bite Back* (pp. 19–21). Chiang Mai, Thailand.
- Costa-Neto, E. M., Dunkel, F. V. (2016). Insects as Food: History, Culture, and Modern Use around the World. In *Insects as Sustainable Food Ingredients*; A. T. Dossey, J. A. Morales-Ramos, & M. G. Rojas (Eds.), (pp. 29–60). Elsevier Inc. <https://doi.org/10.1016/b978-0-12-802856-8.00002-8>
- Das, K., Bardoloi, S., Mazid, S. (2019). A study on the prevalence of entomophagy among the Koch-Rajbongshis of North Salmara subdivision of Bongaigaon district. *International Journal of Basic and Applied Research* 9(3), 382–388.
- de Oliveira, L. M., da Silva Lucas, A. J., Cadaval, C. L., Mellado, M. S. (2017). Bread enriched with flour from cinereous cockroach (*Nauphoeta cinerea*). *Innovative Food Science and Emerging Technologies* 44, 30–35. <https://doi.org/10.1016/j.ifset.2017.08.015>
- Dermody, J., Chatterjee, I. (2016). Food Glorious Food, Fried Bugs and Mustard! Exploring the Radical Idea of Entomophagy in Advancing Sustainable Consumption to Protect the Planet. In *Academy of Marketing Annual Conference, Newcastle* (pp. 1–10). Newcastle.
- Dicke, M. (2018). Insects as feed and the Sustainable Development Goals. *Journal of Insects as Food and Feed* 4(3), 147–156. <https://doi.org/10.3920/JIFF2018.0003>
- Dube, S., Dlamini, N. R., Mafunga, A., Mukai, M., Dhlamini, Z. (2013). A Survey on Entomophagy Prevalence in Zimbabwe. *African Journal of Food, Agriculture, Nutrition and Development* 13(1), 7242–7253. <https://doi.org/10.1016/j.bpapap.2013.06.007>
- Fasolin, L. H., Pereira, R. N., Pinheiro, A. C., Martins, J. T., Andrade, C. C. P., Ramos, O. L., Vicente, A. A. (2019). Emergent food proteins – Towards sustainability, health and innovation. *Food Research International* 125, 108586. <https://doi.org/10.1016/j.foodres.2019.108586>
- Fernandez-Cassi, X., Supeanu, A., Vaga, M., Jansson, A., Boqvist, S., Vagsholm, I. (2019). The house cricket (*Acheta domesticus*) as a novel food: A risk profile. *Journal of Insects as Food and Feed* 5(2), 137–157. <https://doi.org/10.3920/JIFF2018.0021>
- Fisher, H. J., Collins, S. A., Hanson, C., Mason, B., Colombo, S. M., Anderson, D. M. (2020). Black soldier fly larvae meal as a protein source in low fish meal diets for Atlantic salmon (*Salmo salar*). *Aquaculture* 521, 1–9. <https://doi.org/10.1016/j.aquaculture.2020.734978>
- Flores, C. R., Ponce-Reyes, R., Rojas-García, F. (2015). Exploiting a pest insect species *Sphenarium*

- purpurascens for human consumption: Ecological, social, and economic repercussions. *Journal of Insects as Food and Feed* 1(1), 75–84. <https://doi.org/10.3920/JIFF2014.0013>
- Frigerio, J., Agostinetto, G., Galimberti, A., De Mattia, F., Labra, M., Bruno, A. (2020). Tasting the differences: Microbiota analysis of different insect-based novel food. *Food Research International* 137, 1–10. <https://doi.org/10.1016/j.foodres.2020.109426>
- Gahukar, R. T. (2011). Entomophagy and human food security. *International Journal of Tropical Insect Science* 31(3), 129–144. <https://doi.org/10.1017/S1742758411000257>
- Gallo, M., Federico, N. (2018). Novel Foods: Insects - Technology. In *Encyclopedia of Food Security and Sustainability* (pp. 1–5). Elsevier. <https://doi.org/10.1016/B978-0-12-812687-5.22133-3>
- Gere, A., Radványi, D., Héberger, K. (2019). Which insect species can best be proposed for human consumption? *Innovative Food Science and Emerging Technologies* 52, 358–367. <https://doi.org/10.1016/j.ifset.2019.01.016>
- Giordano, S., Clodoveo, M. L., Gennaro, B. De, Corbo, F. (2018). Factors determining neophobia and neophilia with regard to new technologies applied to the food sector: A systematic review. *International Journal of Gastronomy and Food Science* 11, 1–19. <https://doi.org/10.1016/j.ijgfs.2017.10.001>
- Gjerris, M., Gamborg, C., Röcklinsberg, H. (2016). Ethical aspects of insect production for food and feed. *Journal of Insects as Food and Feed* 2(2), 101–110. <https://doi.org/10.3920/JIFF2015.0097>
- González-Escobar, J. L., Grajales-Lagunes, A., Smoliński, A., Chagolla-López, A., De León-Rodríguez, A., Barba de la Rosa, A. P. (2018). Microbiota of edible *Liometopum apiculatum* ant larvae reveals potential functions related to their nutritional value. *Food Research International* 109, 497–505. <https://doi.org/10.1016/j.foodres.2018.04.049>
- Govorushko, S. (2019). Global status of insects as food and feed source: A review. *Trends in Food Science and Technology* 91, 436–445. <https://doi.org/10.1016/j.tifs.2019.07.032>
- Grabowski, N. T., Klein, G. (2017). Microbiological analysis of raw edible insects. *Journal of Insects as Food and Feed* 3(1), 7–14. <https://doi.org/10.3920/JIFF2016.0004>
- Hartmann, C., Bearth, A. (2019). Bugs on the Menu: Drivers and Barriers of Consumer Acceptance of Insects as Food. In *Edible Insects in the Food Sector*; G. Sogari, C. Mora, & D. Menozzi (Eds.), (First, pp. 45–55). Springer Nature Switzerland. https://doi.org/https://doi.org/10.1007/978-3-030-22522-3_4_45
- Imathiu, S. (2020). Benefits and food safety concerns associated with consumption of edible insects. *NFS Journal* 18, 1–11. <https://doi.org/10.1016/j.nfs.2019.11.002>
- Jensen, N. H., Lieberoth, A. (2019). We will eat disgusting foods together – Evidence of the normative basis of Western entomophagy-disgust from an insect tasting. *Food Quality and Preference* 72, 109–115. <https://doi.org/10.1016/j.foodqual.2018.08.012>
- Kelemu, S., Niassy, S., Torto, B., Fiaboe, K., Affognon, H., Tonnang, H., ... Ekesi, S. (2015). African edible insects for food and feed: Inventory, diversity, commonalities and contribution to food security. *Journal of Insects as Food and Feed* 1(2), 103–119. <https://doi.org/10.3920/JIFF2014.0016>
- Köhler, R., Kariuki, L., Lambert, C., Biesalski, H. K. (2019). Protein, amino acid and mineral composition of some edible insects from Thailand. *Journal of Asia-Pacific Entomology* 22(1), 372–378. <https://doi.org/10.1016/j.aspen.2019.02.002>
- Koko, M. Y. F., Mariod, A. A. (2020). Sensory Quality of Edible Insects. In *African Edible Insects as Alternative Source of Food, Oil, Protein and Bioactive Components*; A. Adam Mariod (Ed.), (First, pp. 115–122). Springer Nature Switzerland. https://doi.org/https://doi.org/10.1007/978-3-030-32952-5_7
- Kröncke, N., Bösch, V., Woyzichowski, J., Demtröder, S., Benning, R. (2018). Comparison of suitable drying processes for mealworms (*Tenebrio molitor*). *Innovative Food Science and Emerging Technologies* 50, 20–25. <https://doi.org/10.1016/j.ifset.2018.10.009>
- Kulma, M., Kouřimská, L., Homolková, D., Božik, M., Plachý, V., Vrabec, V., ... Plachý, V. (2020). Effect of developmental stage on the nutritional value of edible insects. A case study with *Blaberus craniifer* and *Zophobas morio*. *Journal of Food Composition and Analysis* 92, 103570. <https://doi.org/10.1016/j.jfca.2020.103570>
- Kulma, M., Kouřimská, L., Plachý, V., Božik, M., Adámková, A., Vrabec, V. (2019). Effect of sex on the nutritional value of house cricket, *Acheta domestica* L. *Food Chemistry* 272, 267–272. <https://doi.org/10.1016/j.foodchem.2018.08.049>
- Kusch, S., Fiebelkorn, F. (2019a). Environmental impact judgments of meat, vegetarian, and insect burgers: Unifying the negative footprint illusion and quantity insensitivity. *Food Quality and Preference* 78, 103731. <https://doi.org/10.1016/j.foodqual.2019.103731>
- Kusch, S., Fiebelkorn, F. (2019b). Environmental impact judgments of meat, vegetarian, and insect burgers: Unifying the negative footprint illusion and quantity insensitivity. *Food Quality and Preference* 78, 1–10. <https://doi.org/10.1016/j.foodqual.2019.103731>
- Lategan, A. (2019). *An assessment of the potential of edible insect consumption in reducing human nutritional deficiencies in South Africa while considering food and nutrition security aspects*. Stellenbosch University. Retrieved from <https://scholar.sun.ac.za/handle/10019.1/106070>
- Lillford, P., Hermansson, A. M. (2020). Global missions and the critical needs of food science and technology. *Trends in Food Science and Technology*. <https://doi.org/10.1016/j.tifs.2020.04.009>
- Longvah, T., Ananthan, R., Bhaskarachary, K., Venkaiah,

- K. (2017). *Indian Food Composition Tables*. Hyderabad, India: National Institute of Nutrition, Indian Council of Medical Research.
- Makkar, H. P. S., Tran, G., Heuzé, V., Ankers, P. (2014). State-of-the-art on use of insects as animal feed. *Animal Feed Science and Technology* 197, 1–33. <https://doi.org/10.1016/j.anifeedsci.2014.07.008>
- Mancini, S., Moruzzo, R., Riccioli, F., Paci, G. (2019). European consumers' readiness to adopt insects as food. A review. *Food Research International* 122, 661–678. <https://doi.org/10.1016/j.foodres.2019.01.041>
- Manditsera, F. A., Luning, P. A., Fogliano, V., Lakemond, C. M. M. (2019). The contribution of wild harvested edible insects (Eulepida mashona and Henicus whellani) to nutrition security in Zimbabwe. *Journal of Food Composition and Analysis* 75, 17–25. <https://doi.org/10.1016/j.jfca.2018.09.013>
- Mba, A. R. F., Kansci, G., Viau, M., Hafnaoui, N., Meynier, A., Demmano, G., Genot, C. (2017). Lipid and amino acid profiles support the potential of Rhynchophorus phoenicis larvae for human nutrition. *Journal of Food Composition and Analysis* 60, 64–73. <https://doi.org/10.1016/j.jfca.2017.03.016>
- Mba, A. R. F., Kansci, G., Viau, M., Rougerie, R., Genot, C. (2019). Edible caterpillars of Imbrasia truncata and Imbrasia epimethea contain lipids and proteins of high potential for nutrition. *Journal of Food Composition and Analysis* 79, 70–79. <https://doi.org/10.1016/j.jfca.2019.03.002>
- Megido, C. R., Sablon, L., Geuens, M., Brostaux, Y., Alabi, T., Blecker, C., ... Francis, F. (2014). Edible insects acceptance by belgian consumers: Promising attitude for entomophagy development. *Journal of Sensory Studies* 29, 14–20. <https://doi.org/10.1111/joss.12077>
- Megido, C. R., Poelaert, C., Ernens, M., Liotta, M., Blecker, C., Danthine, S., ... Francis, F. (2018). Effect of household cooking techniques on the microbiological load and the nutritional quality of mealworms (*Tenebrio molitor* L. 1758), 106, 503–508. <https://doi.org/10.1016/j.foodres.2018.01.002>
- Morales-Ramos, J. A., Rojas, M. G., Dossey, A. T. (2018). Age-dependent food utilisation of *Acheta domesticus* (Orthoptera: Gryllidae) in small groups at two temperatures. *Journal of Insects as Food and Feed* 4(1), 51–60. <https://doi.org/10.3920/JIFF2017.0062>
- Muimba-Kankolongo, A., Ng'andwe, P., Mwitwa, J., Banda, M. K. (2015). Non-Wood Forest Products, Markets, and Trade. In *Forest Policy, Economics, and Markets in Zambia*; P. Ng'andwe, J. Mwitwa, & A. Muimba-Kankolong (Eds.), (1st ed., pp. 67–104). Elsevier Inc. <https://doi.org/10.1016/B978-0-12-804090-4.00003-3>
- Müller, A., Evans, J., Payne, C. L. R., Roberts, R. (2016). Entomophagy and power. *Journal of Insects as Food and Feed* 2(2), 121–136. <https://doi.org/10.3920/JIFF2016.0010>
- Nadeau, L., Nadeau, I., Franklin, F., Dunkel, F. (2015). The Potential for Entomophagy to Address Undernutrition. *Ecology of Food and Nutrition* 54(3), 200–208. <https://doi.org/10.1080/03670244.2014.930032>
- Nantanga, K. K. M., Amakali, T. (2020). Diversification of mopane caterpillars (*Gonimbrasia belina*) edible forms for improved livelihoods and food security. *Journal of Arid Environments* 177, 104148. <https://doi.org/10.1016/j.jaridenv.2020.104148>
- New, T. R. (2013). Two major contributions to understanding entomophagy. *Journal of Insect Conservation* 17, 857–858. <https://doi.org/10.1007/s10841-013-9580-x>
- Niassy, S., Affognon, H. D., Fiaboe, K. K. M., Akutse, K. S., Tanga, C. M., Ekesi, S. (2016). Some key elements on entomophagy in Africa: Culture, gender and belief. *Journal of Insects as Food and Feed* 2(3), 139–144. <https://doi.org/10.3920/JIFF2015.0084>
- Niassy, S., Ekesi, S. (2016). Contribution to the knowledge of entomophagy in Africa. *Journal of Insects as Food and Feed* 2(3), 137–138. <https://doi.org/10.3920/JIFF2016.x003>
- Odongo, W., Okia, C. A., Nalika, N., Nzabamwita, P. H., Ndimubandi, J., Nyeko, P. (2018). Marketing of edible insects in Lake Victoria basin: The case of Uganda and Burundi. *Journal of Insects as Food and Feed* 4(4), 285–293. <https://doi.org/10.3920/JIFF2017.0071>
- Obiokpa, F. I., Akanya, H. O., Jigam, A. A., Saidu, A. N., Egwim, E. C. (2018). Protein quality of four indigenous edible insect species in Nigeria. *Food Science and Human Wellness* 7, 175–183. <https://doi.org/10.1016/j.fshw.2018.05.003>
- Ooninx, D. G. A. B., van Itterbeek, J., Heetkamp, M. J. W., van den Brand, H., van Loon, J. J. A., van Huis, A. (2010). An exploration on greenhouse gas and ammonia production by insect species suitable for animal or human consumption. *PLoS ONE* 5(12), 1–7. <https://doi.org/10.1371/journal.pone.0014445>
- Osimani, A., Milanović, V., Cardinali, F., Roncolini, A., Garofalo, C., Clementi, F., ... Aquilanti, L. (2018). Bread enriched with cricket powder (*Acheta domesticus*): A technological, microbiological and nutritional evaluation. *Innovative Food Science and Emerging Technologies* 48, 150–163. <https://doi.org/10.1016/j.ifset.2018.06.007>
- Padulosi, S., Cawthorn, D. M., Meldrum, G., Flore, R., Halloran, A., Mattei, F. (2018). Leveraging neglected and underutilized plant, fungi, and animal species for more nutrition sensitive and sustainable food systems. *Encyclopedia of Food Security and Sustainability* 3, 361–370. <https://doi.org/10.1016/B978-0-08-100596-5.21552-7>
- Patel, S. (2019). Insects as a Source of Sustainable Proteins. In *Proteins: Sustainable Source, Processing and Applications*; C. Galanakis (Ed.), (pp. 41–61). Elsevier Inc. <https://doi.org/10.1016/b978-0-12-816695-6.00002-7>
- Patel, S., Suleria, H. A. R., Rauf, A. (2019). Edible insects as innovative foods: Nutritional and functional assessments. *Trends in Food Science and Technology* 86, 352–359. <https://doi.org/10.1016/j.tifs.2019.02.033>
- Paul, A., Frederich, M., Uyttenbroeck, R., Malik, P.,

- Filocco, S., Richel, A., ... Danthine, S. (2016). Nutritional composition and rearing potential of the meadow grasshopper (*Chorthippus parallelus* Zetterstedt). *Journal of Asia-Pacific Entomology* 19(4), 1111–1116. <https://doi.org/10.1016/j.aspen.2016.09.012>
- Payne, C., Dobermann, D., Forkes, A., House, J., Josephs, J., McBride, A., ... Soares, S. (2016). Insects as food and feed: European perspectives on recent research and future priorities. *Journal of Insects as Food and Feed* 2(4), 269–276. <https://doi.org/10.3920/JIFF2016.0011>
- Pippinato, L., Gasco, L., Di Vita, G., Mancuso, T. (2020). Current scenario in the European edible-insect industry: a preliminary study. *Journal of Insects as Food and Feed* 6(1), 1–12. <https://doi.org/10.3920/jiff2020.0008>
- Poelaert, C., Francis, F., Alabi, T., Caparros Megido, R., Crahay, B., Bindelle, J., Beckers, Y. (2018). Protein value of two insects, subjected to various heat treatments, using growing rats and the protein digestibility-corrected amino acid score. *Journal of Insects as Food and Feed* 4(2), 77–87. <https://doi.org/10.3920/JIFF2017.0003>
- Ramos-Elorduy, J. (2009). Anthro-entomophagy: Cultures, evolution and sustainability. *Entomological Research* 39, 271–288. <https://doi.org/10.1111/j.1748-5967.2009.00238.x>
- Rao, M. S. (2016). Scope of Insect Farming and Entomophagy. In *Reshaping Agriculture and Nutrition Linkages for Food and Nutrition Security*; K. Sreedevi, S. R. Nagarjuna, K. Pushpanjali, K. Nagasree, G. Nirmala, & S. Raju (Eds.), (pp. 136–142). Hyderabad, India: Central Research Institute for Dryland Agriculture. Retrieved from <http://crida.in>
- Ribeiro, J. C., Lima, R. C., Maia, M. R. G., Almeida, A. A., Fonseca, A. J. M., Cabrita, A. R. J., Cunha, L. M. (2019). Impact of defatting freeze-dried edible crickets (*Acheta domesticus* and *Gryllobasys sigillatus*) on the nutritive value, overall liking and sensory profile of cereal bars. *Lwt-Food Science and Technology* 113, 108335. <https://doi.org/10.1016/j.lwt.2019.108335>
- Roos, N., van Huis, A. (2017). Consuming insects: Are there health benefits? *Journal of Insects as Food and Feed* 3(4), 225–229. <https://doi.org/10.3920/JIFF2017.x007>
- Ruby, M. B., Rozin, P. (2019). Disgust, sushi consumption, and other predictors of acceptance of insects as food by Americans and Indians. *Food Quality and Preference* 74, 155–162. <https://doi.org/10.1016/j.foodqual.2019.01.013>
- Ruby, M. B., Rozin, P., Chan, C. (2015). Determinants of willingness to eat insects in the USA and India. *Journal of Insects as Food and Feed* 1(3), 215–225. <https://doi.org/10.3920/JIFF2015.0029>
- Rumpold, B. A., Langen, N. (2020). Consumer acceptance of edible insects in an organic waste-based bioeconomy. *Current Opinion in Green and Sustainable Chemistry* 23, 80–84. <https://doi.org/10.1016/j.cogsc.2020.03.007>
- Rumpold, B. A., Schlüter, O. (2015). Insect-based protein sources and their potential for human consumption: Nutritional composition and processing. *Animal Frontiers* 5(2), 20–24. <https://doi.org/10.2527/af.2015-0015>
- Schmidt, A., Call, L. M., Macheiner, L., Mayer, H. K. (2019). Determination of vitamin B 12 in four edible insect species by immunoaffinity and ultra-high performance liquid chromatography. *Food Chemistry* 281, 124–129. <https://doi.org/10.1016/j.foodchem.2018.12.039>
- Severini, C., Azzollini, D., Albenzio, M., Derossi, A. (2018). On printability, quality and nutritional properties of 3D printed cereal based snacks enriched with edible insects. *Food Research International* 106, 666–676. <https://doi.org/10.1016/j.foodres.2018.01.034>
- Shantibala, T., Lokeshwari, R. K., Sharma, H. D. (2012). Entomophagy practices among the ethnic communities of Manipur, North-east India. *International Journal of Integrative Sciences, Innovation and Technology* 1(5), 13–20. Retrieved from www.ijit.webs.com
- Skrivervik, E. (2020). Insects' contribution to the bioeconomy and the reduction of food waste. *Heliyon* 6, e03934. <https://doi.org/10.1016/j.heliyon.2020.e03934>
- Srivastava, S. K., Babu, N., Pandey, H. (2009). Traditional insect bioprospecting - As human food and medicine. *Indian Journal of Traditional Knowledge* 8(4), 485–494.
- Ssepuuya, G., Smets, R., Nakimbugwe, D., Borght, M. Van Der, Claes, J. (2019). Nutrient composition of the long-horned grasshopper *Ruspolia diffrrens* Serville: Effect of swarming season and sourcing geographical area. *Food Chemistry* 301, 125305. <https://doi.org/10.1016/j.foodchem.2019.125305>
- Straub, P., Tanga, C. M., Osuga, I., Windisch, W., Subramanian, S. (2019). Experimental feeding studies with crickets and locusts on the use of feed mixtures composed of storable feed materials commonly used in livestock production. *Animal Feed Science and Technology* 255, 114215. <https://doi.org/10.1016/j.anifeedsci.2019.114215>
- Stull, V. J., Wamulume, M., Mwalukanga, M. I., Banda, A., Bergmans, R. S., Bell, M. M. (2018). “We like insects here”: entomophagy and society in a Zambian village. *Agriculture and Human Values* 35(4), 867–883. <https://doi.org/10.1007/s10460-018-9878-0>
- Sun-Waterhouse, D., Waterhouse, G. I. N., You, L., Zhang, J., Liu, Y., Ma, L., ... Dong, Y. (2016). Transforming insect biomass into consumer wellness foods: A review. *Food Research International* 89, 129–151. <https://doi.org/10.1016/j.foodres.2016.10.001>
- Surendra, K. C., Tomberlin, J. K., van Huis, A., Cammack, J. A., Heckmann, L. H. L., Khanal, S. K. (2020). Rethinking organic wastes bioconversion: Evaluating the potential of the black soldier fly (*Hermetia illucens* (L.) (BSF)). *Waste Management* 117, 58–80. <https://doi.org/10.1016/j.wasman.2020.07.050>
- Tang, C., Yang, D., Liao, H., Sun, H., Liu, C., Wei, L., Li, F. (2019). Edible insects as a food source: a review.

- Food Production, Processing and Nutrition* 1(1), 1–13. <https://doi.org/10.1186/s43014-019-0008-1>
- Testa, M., Stillo, M., Maffei, G., Andriolo, V., Gardois, P., Zotti, C. M. (2016). Ugly but tasty: A systematic review of possible human and animal health risks related to entomophagy. *Critical Reviews in Food Science and Nutrition* 57, 3747–3759. <https://doi.org/10.1080/10408398.2016.1162766>
- Tomberlin, J. K., van Huis, A., Benbow, M. E., Jordan, H., Astuti, D. A., Azzollini, D., ... Zheng, L. (2015). Protecting the environment through insect farming as a means to produce protein for use as livestock, poultry, and aquaculture feed. *Journal of Insects as Food and Feed* 1(4), 307–309. <https://doi.org/10.3920/JIFF2015.0098>
- Tong, L., Yu, X., Liu, H. (2011). Insect food for astronauts: gas exchange in silkworms fed on mulberry and lettuce and the nutritional value of these insects for human consumption during deep space flights. *Bulletin of Entomological Research* 101, 613–622. <https://doi.org/10.1017/s0007485311000228>
- Tuccillo, F., Marino, M. G., Torri, L. (2020). Italian consumers' attitudes towards entomophagy: Influence of human factors and properties of insects and insect-based food. *Food Research International* 137, 1–10. <https://doi.org/10.1016/j.foodres.2020.109619>
- Ulmer, M., Smetana, S., Heinz, V. (2020). Utilizing honeybee drone brood as a protein source for food products: Life cycle assessment of apiculture in Germany. *Resources, Conservation and Recycling* 154, 1–14. <https://doi.org/10.1016/j.resconrec.2019.104576>
- van Huis, A. (2015). Edible insects contributing to food security? *Agriculture and Food Security* 4(1), 1–9. <https://doi.org/10.1186/s40066-015-0041-5>
- van Huis, A. (2020a). Importance of Insects as Food in Africa. In African Edible Insects as Alternative Source of Food, Oil, Protein and Bioactive Components; Abdalbasit Adam Mariod (Ed.), (first, pp. 1–17). Springer Nature Switzerland.
- van Huis, A. (2020b). Insects as food and feed, a new emerging agricultural sector: A review. *Journal of Insects as Food and Feed* 6(1), 27–44. <https://doi.org/10.3920/JIFF2019.0017>
- van Huis, A., Itterbeeck, J. Van, Klunder, H., Mertens, E., Halloran, A., Muir, G., Vantomme, P. (2013). *Edible insects: future prospects for food and feed security*. Rome: Food and Agriculture Organization of the United Nations. Retrieved from <https://library.wur.nl/WebQuery/wurpubs/fulltext/258042>
- Videbæk, P. N., Grunert, K. G. (2020). Disgusting or delicious? Examining attitudinal ambivalence towards entomophagy among Danish consumers. *Food Quality and Preference* 83, 103913. <https://doi.org/10.1016/j.foodqual.2020.103913>
- Waithanji, E., Affognon D., H., King'ori, S., Diiro, G., Nakimbugwe, D., Fiaboe, K. K. M. (2019). Insects as feed: Gendered knowledge attitudes and practices among poultry and Pond Fish farmers in Kenya. *NJAS - Wageningen Journal of Life Sciences* 100312. <https://doi.org/10.1016/j.njas.2019.100312>
- Waltner-Toews, D., Houle, K. (2017). Biophilia on the Dinner Plate: a Conversation about Ethics and Entomophagy. *Food Ethics* 1(2), 157–171. <https://doi.org/10.1007/s41055-017-0015-3>
- Weindl, I., Ost, M., Wiedmer, P., Schreiner, M., Neugart, S., Klopsch, R., ... Klaus, S. (2020a). Sustainable food protein supply reconciling human and ecosystem health: A Leibniz Position. *Global Food Security* 25, 1–10. <https://doi.org/10.1016/j.gfs.2020.100367>
- Weindl, I., Ost, M., Wiedmer, P., Schreiner, M., Neugart, S., Klopsch, R., ... Klaus, S. (2020b). Sustainable food protein supply reconciling human and ecosystem health: A Leibniz Position. *Global Food Security* 25, 100367. <https://doi.org/10.1016/j.gfs.2020.100367>
- Williams, J. P., Williams, J. R., Kirabo, A., Chester, D., Peterson, M. (2016). Nutrient Content and Health Benefits of Insects. In *Insects as Sustainable Food Ingredients*; A. T. Dossey, J. A. Morales-Ramos, & M. G. Rojas (Eds.), (pp. 61–84). Elsevier Inc. <https://doi.org/10.1016/B978-0-12-802856-8/00003-X>
- Yen, A. L. (2015a). Foreword: Why a journal of insects as food and feed? *Journal of Insects as Food and Feed* 1(1), 1–2. <https://doi.org/10.3920/JIFF2015.x001>
- Yen, A. L. (2015b). Insects as food and feed in the Asia Pacific region: Current perspectives and future directions. *Journal of Insects as Food and Feed* 1(1), 33–55. <https://doi.org/10.3920/JIFF2014.0017>
- Zeece, M. (2020). Food systems and future directions. In *Introduction to the Chemistry of Food* (p. 389). Academic Press. <https://doi.org/10.1016/b978-0-12-809434-1.00009-8>
- Zielińska, E., Baraniak, B., Karaś, M., Rybczyńska, K., Jakubczyk, A. (2015). Selected species of edible insects as a source of nutrient composition. *Food Research International*. <https://doi.org/10.1016/j.foodres.2015.09.008>
- Zou, Y. X., Hu, T. G., Shi, Y., Liu, J., Mu, L. X., Xiao, Y., Liao, S. T. (2017). Establishment of a model to evaluate the nutritional quality of *Bombyx mori* Linnaeus (Lepidoptera, Bombycidae) pupae lipid based on principal components. *Journal of Asia-Pacific Entomology* 20(4), 1364–1371. <https://doi.org/10.1016/j.aspen.2017.05.012>