

## SIMILITUDE EXCEPTIONAL THETA CORRESPONDENCES

PETAR BAKIĆ, WEE TECK GAN AND GORDAN SAVIN

*In honor of Marko Tadić  
from whom we have learned so much*

ABSTRACT. We describe a systematic way of constructing dual pairs of similitude groups. We study the theta correspondences arising in this way and prove that Howe duality holds for the similitude dual pair if and only if it holds for the original reductive dual pair used in our construction. Several examples of exceptional correspondences are discussed.

### 1. INTRODUCTION

Let  $F$  be a local field and suppose  $(G, H)$  is a reductive dual pair in a simple linear algebraic group  $\mathcal{E}$  over  $F$ . Hence,  $G$  and  $H$  are subgroups of  $\mathcal{E}$  which are mutual centralizers of each other. We thus have a homomorphism of algebraic groups

$$i : G \times H \longrightarrow \mathcal{E}.$$

If  $\Omega$  is a minimal representation of  $\mathcal{E}(F)$ , we may pull back  $\Omega$  to the product group  $G(F) \times H(F)$  and study the resulting branching problem: this is the usual setup of theta correspondence.

However, the map  $i$  is frequently not injective. Typically, one has (multiplicative type) central subgroups

$$Z \hookrightarrow G \quad \text{and} \quad Z \hookrightarrow H$$

which are identified under  $i$ . More precisely, one has

$$\text{Ker}(i) = Z^\nabla = \{(z, z^{-1}) : z \in Z\},$$

so that one has an injection

$$(G \times H)/Z^\nabla \hookrightarrow \mathcal{E}.$$

---

2020 *Mathematics Subject Classification.* 11S90, 17A75, 17C40.  
*Key words and phrases.* theta correspondences, similitude groups.

This already occurs in the setting of the classical theta correspondence for the symplectic-orthogonal dual pair

$$i : \mathrm{O}(V) \times \mathrm{Sp}(W) \longrightarrow \mathrm{Sp}(V \otimes W)$$

whose kernel is the diagonally embedded  $Z = \mu_2$ .

In this short note, we explain how the non-injectivity of  $i$  can be exploited to extend the theta correspondence for  $G(F) \times H(F)$  to the setting of similitude groups associated to  $G$  and  $H$ . Indeed, as we shall see, this non-injectivity is the reason for the existence of such a theory.

Here is a brief summary of the content of the paper. In §2, we associate to a dual pair  $G \times H \longrightarrow \mathcal{E}$  as above, and some additional data, a similitude dual pair  $\tilde{G} \times \tilde{H}$ . After introducing the associated similitude version of theta correspondence in §3 and demonstrating some simple properties of this similitude theta correspondence in §4, we show in §5 that the Howe duality theorem holds for  $G \times H$  if and only if the analogous theorem holds for  $\tilde{G} \times \tilde{H}$ ; this is Theorem 5.1 and should be considered the main result of this paper. In §6, we illustrate our theory for a few families of dual pairs in exceptional groups. Just as classical dual pairs in symplectic groups can be associated to a pair of Hermitian and skew-Hermitian spaces (via the tensor product of the two spaces), we highlight three families of dual pairs in exceptional groups which can be obtained from pairs of exceptional algebraic structures. These are a pair of composition algebra and Jordan algebra, a pair of twisted composition algebras or a pair of Jordan pairs. For each of these, we discuss whether a theory of similitude theta correspondence exists and if so, what the relevant similitude groups are. In §7, we discuss how the useful notion of seesaw duality can be extended to the similitude setting, using the families of examples discussed in §6 as illustration. Finally, §8 discusses the similitude theta correspondences in the global setting and §9 collects together some basic results in Clifford theory that are used in the paper.

## 2. Similitude Groups

Let us first introduce the notion and construction of the relevant similitude groups. We will continue to work in the context of the introduction.

**2.1. Initial data.** The initial data needed for the construction of similitude groups is described in the following hypotheses:

- (a) the commutative group  $Z$  can be embedded into an induced torus:

$$j : Z \hookrightarrow T := \mathrm{Res}_{K/F} \mathbb{G}_m$$

where  $K$  is an étale  $F$ -algebra of finite rank. Here, and elsewhere,  $\mathrm{Res}$  denotes the standard Weil restriction functor.

(b) One has

$$S := T/j(Z) \hookrightarrow \text{Res}_{E/F}(\mathbb{G}_m),$$

for some  $E/F$ .

We shall fix these data in what follows.

**2.2. Examples.** Let us give some examples:

(1) if  $Z = \prod_{i=1}^r \mu_{n_i}$ , one takes embeddings  $\mu_{n_i} \hookrightarrow \mathbb{G}_m$ , so that one has

$$1 \longrightarrow Z \longrightarrow T = \prod_{i=1}^r \mathbb{G}_m \longrightarrow T = \prod_{i=1}^r \mathbb{G}_m \longrightarrow 1$$

where the map  $T \rightarrow T$  is  $z \mapsto z^{n_i}$  on the  $i$ -th coordinate.

(2) if

$$Z = \text{Res}_{K/F}^1(\mathbb{G}_m) := \text{Ker}(N_{K/F} : \text{Res}_{K/F}(\mathbb{G}_m) \rightarrow \mathbb{G}_m),$$

one takes  $T = \text{Res}_{K/F}\mathbb{G}_m$  and one has

$$1 \longrightarrow Z \longrightarrow T \xrightarrow{N_{K/F}} \mathbb{G}_m \longrightarrow 1$$

(3) if

$$Z = \text{Res}_{K/F}^1(\mu_n) := \text{Ker}(N_{K/F} : \text{Res}_{K/F}(\mu_n) \rightarrow \mathbb{G}_m),$$

one takes  $T = \text{Res}_{K/F}\mathbb{G}_m$  and one has

$$1 \longrightarrow Z \longrightarrow T \xrightarrow{N_{K/F} \times [n]} \mathbb{G}_m \times \text{Res}_{K/F}(\mathbb{G}_m).$$

Note that in this last example, the sequence above is not exact on the right, as the image of the last arrow is a codimension 1 torus. Consider however the special case when

$$|n - [K : F]| = 1.$$

In this case, it turns out that the composite

$$T/Z \xrightarrow{N_{K/F} \times [n]} \mathbb{G}_m \times \text{Res}_{K/F}(\mathbb{G}_m) \xrightarrow{(t,x) \mapsto tx^{-1}} \text{Res}_{K/F}(\mathbb{G}_m) = T$$

is an isomorphism of algebraic tori. Indeed, if  $x \in \text{Res}_{K/F}(\mathbb{G}_m)$  lies in the kernel of the above composite map, so that

$$N_{K/F}(x) = x^n,$$

then on taking norms on both sides, we conclude that

$$N_{K/F}(x)^{[K:F]} = N_{K/F}(x)^n,$$

and hence  $x^n = N_{K/F}(x) = 1$ , i.e.  $x \in \text{Res}_{K/F}^1(\mu_n)$ . In the examples we shall consider later, the rather peculiar condition  $|n - [K : F]| = 1$  will turn out to be satisfied.

**2.3. Similitude groups.** Now we make the following definition:

DEFINITION 2.1. *The similitude group  $\tilde{G}$  associated to  $G$  and the data in (a) and (b) above is:*

$$\tilde{G} := (G \times T)/Z^\nabla.$$

The associated similitude homomorphism is the map

$$\text{sim}_G : \tilde{G} \rightarrow T/j(Z) \hookrightarrow \text{Res}_{E/F}\mathbb{G}_m.$$

given by the second projection and (b).

Then observe that one has:

$$1 \longrightarrow G \longrightarrow \tilde{G} \xrightarrow{\text{sim}_G} \text{Res}_{E/F}\mathbb{G}_m$$

and

$$1 \longrightarrow T \longrightarrow \tilde{G} \longrightarrow G/Z \longrightarrow 1$$

Likewise, we have the similitude group and similitude character

$$\text{sim}_H : \tilde{H} := (H \times T)/Z^\nabla \longrightarrow \text{Res}_{E/F}(\mathbb{G}_m)$$

associated to  $H$  and the data in (a) and (b), as well as analogs of the two exact sequences above.

**2.4. Surjectivity of similitude factor.** Recalling that  $S = T/j(Z)$ , we have a commutative diagram of short exact sequences of algebraic groups:

$$\begin{array}{ccccccc} 1 & \longrightarrow & Z & \xrightarrow{j} & T & \longrightarrow & S \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \parallel \\ 1 & \longrightarrow & G & \longrightarrow & \tilde{G} & \xrightarrow{\text{sim}_G} & S \longrightarrow 1. \end{array}$$

On taking  $F$ -points, we obtain the following commutative diagram with exact rows:

$$\begin{array}{ccccc} T(F) & \longrightarrow & S(F) & \xrightarrow{\alpha} & H^1(F, Z) \\ \downarrow & & \parallel & & \downarrow \beta \\ \tilde{G}(F) & \xrightarrow{\text{sim}_G} & S(F) & \longrightarrow & H^1(F, G) \end{array}$$

From this, we see that the similitude map  $\text{sim}_G : \tilde{G}(F) \rightarrow S(F)$  is surjective if and only if the map  $S(F) \rightarrow H^1(F, G)$  is 0. Since the latter map factors as:

$$S(F) \xrightarrow{\alpha} H^1(F, Z) \xrightarrow{\beta} H^1(F, G),$$

we obtain:

PROPOSITION 2.2. *The map  $\text{sim}_G : \tilde{G}(F) \rightarrow S(F)$  is surjective if one of the following conditions holds:*

- $\alpha : S(F) \rightarrow H^1(F, Z)$  is 0;
- $\beta : H^1(F, Z) \rightarrow H^1(F, G)$  is 0.

Observe that the first condition in the proposition depends only on the initial data (a) and (b) in §2.1, whereas the second condition depends only on the pair  $Z \subset G$ .

**2.5. The group  $\tilde{J}^{\text{sim}}$ .** Going back to dual pairs, let us set

$$\tilde{J}^{\text{sim}} = (\tilde{G} \times \tilde{H})^{\text{sim}} = \{(g, h) \in \tilde{G} \times \tilde{H} : \text{sim}_G(g) \cdot \text{sim}_H(h) = 1\}.$$

Note that one has exact sequences:

$$1 \longrightarrow G(F) \longrightarrow \tilde{J}^{\text{sim}}(F) \longrightarrow \tilde{H}(F)$$

and

$$1 \longrightarrow H(F) \longrightarrow \tilde{J}^{\text{sim}}(F) \longrightarrow \tilde{G}(F)$$

One denotes the image of the last arrow in each sequence above by  $\tilde{H}(F)^+$  and  $\tilde{G}(F)^+$  respectively. Then one has the following containments with finite index:

$$T(F) \cdot G(F) \subset \tilde{G}(F)^+ \quad \text{and} \quad T(F) \cdot H(F) \subset \tilde{H}(F)^+.$$

Observe moreover that

$$\tilde{G}(F)^+ = \tilde{G}(F) \quad \text{if and only if} \quad \text{sim}_G(\tilde{G}(F)) \subseteq \text{sim}_H(\tilde{H}(F))$$

and

$$\tilde{H}(F)^+ = \tilde{H}(F) \quad \text{if and only if} \quad \text{sim}_H(\tilde{H}(F)) \subseteq \text{sim}_G(\tilde{G}(F)).$$

In particular,  $\tilde{G}(F)^+ = \tilde{G}(F)$  if  $\text{sim}_H : \tilde{H}(F) \rightarrow S(F)$  is surjective.

**DEFINITION 2.3.** *We shall call the pair of groups  $(\tilde{G}(F)^+, \tilde{H}(F)^+)$  constructed above a similitude dual pair.*

### 3. Similitude Theta Correspondence

We now consider the theory of theta correspondence. Suppose that  $\Omega$  is a minimal representation of  $\mathcal{E}(F)$  [GS]. If  $\pi \in \text{Irr}(G(F))$ , then the big theta lift of  $\pi$  to  $H(F)$  is by definition

$$\Theta(\pi) = (\Omega \otimes \pi^\vee)_{G(F)},$$

i.e. the largest quotient of  $\Omega \otimes \pi^\vee$  on which  $G(F)$  acts trivially (here  $\pi^\vee$  denotes the contragredient of  $\pi$ ). This is a smooth representation of  $H(F)$ . As is customary, we let  $\theta(\pi)$  denote the maximal semisimple quotient of  $\Theta(\pi)$ . Likewise, one has a smooth  $G(F)$ -representation  $\Theta(\sigma)$  (and  $\theta(\sigma)$ ) for any  $\sigma \in \text{Irr}(H(F))$ .

**DEFINITION 3.1.** *We say that the dual pair  $(G, H)$  satisfies the Howe duality property, or the Howe duality theorem holds for  $(G, H)$ , if  $\Theta(\pi)$  is of finite length with a unique irreducible quotient (if nonzero) for any  $\pi \in \text{Irr}(G(F))$  and likewise for  $\Theta(\sigma)$  for any  $\sigma \in \text{Irr}(H(F))$ .*

We would now like to extend the theta correspondence for  $(G(F), H(F))$  to the setting of the similitude dual pair  $(\tilde{G}(F)^+, \tilde{H}(F)^+)$ . For this, we observe that

$$T^\nabla = \{(t, t^{-1}) : t \in T\} \subset \tilde{J}^{\text{sim}}$$

is a central subgroup. Setting

$$J^{\text{sim}} = \tilde{J}^{\text{sim}}/T^\nabla = ((\tilde{G} \times \tilde{H})^{\text{sim}})/T^\nabla,$$

one sees that the natural inclusion gives an isomorphism

$$(G \times H)/Z^\nabla \simeq J^{\text{sim}} = ((\tilde{G} \times \tilde{H})^{\text{sim}})/T^\nabla.$$

In particular, one has a natural homomorphism

$$\iota : \tilde{J}^{\text{sim}} \twoheadrightarrow J^{\text{sim}} \hookrightarrow \mathcal{E}.$$

On taking  $F$ -points and noting that  $T$  is an induced torus, so that  $H^1(F, T)$  is trivial, we see that

$$((G \times H)/Z^\nabla)(F) = J^{\text{sim}}(F) = (\tilde{G}(F) \times \tilde{H}(F))^{\text{sim}}/T^\nabla(F),$$

so that one has

$$\iota : \tilde{J}^{\text{sim}}(F) \twoheadrightarrow J^{\text{sim}}(F) \hookrightarrow \mathcal{E}(F).$$

In particular, any representation of  $\mathcal{E}(F)$  can be pulled back via  $\iota$  to

$$\tilde{J}^{\text{sim}}(F) = (\tilde{G}(F) \times \tilde{H}(F))^{\text{sim}}.$$

Now one has the following short exact sequences:

$$1 \longrightarrow G(F) \longrightarrow \tilde{J}^{\text{sim}}(F) \longrightarrow \tilde{H}(F)^+ \longrightarrow 1$$

and

$$1 \longrightarrow H(F) \longrightarrow \tilde{J}^{\text{sim}}(F) \longrightarrow \tilde{G}(F)^+ \longrightarrow 1.$$

Given an irreducible representation  $\tilde{\pi}$  of  $\tilde{G}(F)^+$ , we may regard  $\tilde{\pi}^\vee$  as a representation of  $\tilde{J}^{\text{sim}}(F)$  via the natural surjection  $\tilde{J}^{\text{sim}}(F) \rightarrow \tilde{G}(F)^+$  given in the second short exact sequence above. Noting that  $G(F)$  is a normal subgroup of  $\tilde{J}^{\text{sim}}(F)$  by the first short exact sequence above, we set

$$\Theta(\tilde{\pi}) := (\Omega \otimes \tilde{\pi}^\vee)_{G(F)},$$

so that  $\Theta(\tilde{\pi})$  is a smooth representation of  $\tilde{H}(F)^+$ . Likewise for  $\tilde{\sigma} \in \text{Irr}(\tilde{H}(F)^+)$ , one has

$$\Theta(\tilde{\sigma}) = (\Omega \otimes \tilde{\sigma}^\vee)_{H(F)}$$

which is a smooth representation of  $\tilde{G}(F)^+$ .

One can rephrase the above definitions slightly by introducing the similitude minimal representation. Set

$$\tilde{\Omega} = \text{ind}_{\tilde{J}^{\text{sim}}(F)}^{\tilde{J}(F)^+} \Omega$$

where

$$\tilde{J} = \tilde{G} \times \tilde{H}$$

and

$$\tilde{J}(F)^+ = \tilde{G}(F)^+ \times \tilde{H}(F)^+.$$

Then  $\tilde{\Omega}$  is a representation of the similitude dual pair  $\tilde{G}(F)^+ \times \tilde{H}(F)^+$  and we can define  $\Theta(\tilde{\pi})$  and  $\Theta(\tilde{\sigma})$  in the usual way. We leave it to the reader to verify that the two definitions are equivalent.

As in the “isometry” case, we can define the notion of the Howe duality property holding for  $(\tilde{G}(F)^+, \tilde{H}(F)^+)$ :

**DEFINITION 3.2.** *We say that the similitude dual pair  $(\tilde{G}(F)^+, \tilde{H}(F)^+)$  satisfies the Howe duality property if  $\Theta(\tilde{\pi})$  has finite length and unique irreducible quotient (if nonzero) for any  $\tilde{\pi} \in \text{Irr}(\tilde{G}(F)^+)$ , and likewise for  $\Theta(\tilde{\sigma})$  with  $\tilde{\sigma} \in \text{Irr}(\tilde{H}(F)^+)$ .*

#### 4. Central Characters

Let us record some simple properties of the similitude theta correspondence.

**LEMMA 4.1 (Central characters).** *Suppose that  $\pi \in \text{Irr}(\tilde{G}(F)^+)$  has  $T(F)$ -central character  $\chi$ . Then the smooth  $\tilde{H}(F)^+$ -module  $\Theta(\tilde{\pi})$  has  $T(F)$ -central character  $\chi$ .*

**PROOF.** This is because  $T^\nabla(F) \subset \tilde{J}^{\text{sim}}(F)$  acts trivially on  $\Omega$  and so  $T(F) \subset \tilde{H}(F)^+$  acts on

$$\Theta(\tilde{\pi}) = (\Omega \otimes \tilde{\pi}^\vee)_{G(F)}$$

in the same way as  $T(F) \subset \tilde{G}(F)^+$  acts on  $\tilde{\pi}$ .  $\square$

One can also decompose the  $\tilde{J}(F)^+$ -module  $\tilde{\Omega}$  according to characters of the central subgroup  $T(F) \times T(F)/T^\nabla(F)$ . Each character of  $T(F) \times T(F)/T^\nabla(F)$  is of the form  $\chi \otimes \chi$ , where  $\chi$  is a character of  $T(F)$ . We can consider the  $\chi$ -isotypic quotient  $\tilde{\Omega}_{T,\chi}$  of  $\tilde{\Omega}$ , which can be described as follows.

One has an intermediate group

$$\tilde{J}^{\text{sim}}(F) \subset \tilde{J}^{\text{sim}}(F) \cdot (T(F) \times T(F)) \subset \tilde{J}(F)^+.$$

We may extend the  $\tilde{J}^{\text{sim}}(F)$ -module  $\Omega$  to the intermediate group  $\tilde{J}^{\text{sim}}(F) \cdot (T(F) \times T(F))$  by letting  $T(F) \times T(F)$  acts by  $\chi \otimes \chi$ . This extension is well-defined because  $\chi \otimes \chi$  is trivial on the intersection

$$\tilde{J}^{\text{sim}}(F) \cap (T(F) \times T(F)) = T^\nabla(F).$$

We denote this extended representation by  $\Omega_\chi$ . Then one has

$$\tilde{\Omega}_{T,\chi} \simeq \text{ind}_{\tilde{J}^{\text{sim}}(F) \cdot (T(F) \times T(F))}^{\tilde{J}(F)^+} \Omega_\chi.$$

It will sometimes be convenient to restrict to a fixed  $T(F)$ -central character  $\chi$ , in which case one will be working with  $\tilde{\Omega}_{T,\chi}$ .

## 5. The Howe Duality Property

In this section, we shall show:

**THEOREM 5.1.** *The Howe duality property holds for the dual pair  $(G(F), H(F))$  if and only if the Howe duality property holds for the similitude dual pair  $(\tilde{G}(F)^+, \tilde{H}(F)^+)$ .*

**5.1. Some lemmas.** We shall show this in a series of lemmas, beginning with the following observation:

**LEMMA 5.2.** *Let  $V$  be a not-necessarily-smooth representation of  $\tilde{G}(F)^+$  on which the central subgroup  $T(F)$  acts by a character  $\chi$ . Then:*

(i) *A vector  $v \in V$  is smooth with respect to  $\tilde{G}(F)^+$  if and only if it is smooth with respect to  $G(F)$ .*

(ii) *If  $V^\infty$  denotes the subspace of smooth vectors of  $V$  (with respect to  $\tilde{G}(F)^+$  or  $G(F)$ ), then  $V^\infty$  has finite length as a  $\tilde{G}(F)^+$ -module if and only if it has finite length as a  $G(F)$ -module.*

(iii) *If  $V^\infty$  is an irreducible smooth  $\tilde{G}(F)^+$ -module, then  $V^\infty$  is semisimple as a  $G(F)$ -module. Indeed,*

$$V^\infty|_{G(F)} \simeq m \cdot \bigoplus_{i=1}^k V_i$$

where the  $V_i$ 's are irreducible smooth  $G(F)$ -modules which are pairwise inequivalent. Moreover,  $\tilde{G}(F)^+$  permutes the isomorphism classes of the  $V_i$ 's transitively.

(iv) *Suppose that  $V = V^\infty$  is of finite length (as  $\tilde{G}(F)^+$ -module or  $G(F)$ -module). Let  $\text{co}\tilde{\text{soc}}(V)$  be the cosocle (or maximal semisimple quotient) of  $V$  as a  $\tilde{G}(F)^+$ -module and likewise let  $\text{cosoc}(V)$  be the cosocle of  $V$  as a  $G(F)$ -module. Then the natural map*

$$\text{cosoc}(V) \longrightarrow \text{co}\tilde{\text{soc}}(V)$$

*is an isomorphism of  $G(F)$ -modules.*

**PROOF.** All these follow readily from the fact that  $T(F) \cdot G(F)$  is an open subgroup of finite index in  $\tilde{G}(F)^+$  and the Clifford theory recounted in Section 9 below. More precisely,

- (i) follows from the fact that an open compact subgroup of  $G(F) \cdot T(F)$  is an open compact subgroup of  $\tilde{G}(F)^+$ ;
- (ii) follows from Proposition 9.1(1);



- (iii) follows from Proposition 9.1(2) and Proposition 9.4;
- (iv) follows from Proposition 9.1(3).

□

Next we have:

LEMMA 5.3. *Let  $\tilde{\pi} \in \text{Irr}(\tilde{G}(F)^+)$  and suppose (as in Lemma 5.2(iii)) that*

$$\tilde{\pi}|_{G(F)} \simeq m \cdot \bigoplus_{i=1}^k \pi_i$$

for some  $\pi_i \in \text{Irr}(G(F))$ . Then we have:

$$(5.1) \quad \Theta(\tilde{\pi})|_{H(F)} = m \cdot \bigoplus_i \Theta(\pi_i).$$

In particular,  $\Theta(\tilde{\pi})$  is nonzero if and only if  $\Theta(\pi_i) \neq 0$  for some (equivalently for all)  $i$ .

PROOF. We compute:

$$\begin{aligned} \Theta(\tilde{\pi})|_{H(F)} &= (\Omega \otimes \tilde{\pi}^\vee)_{G(F)} \quad (\text{by definition}) \\ &= \left( \Omega \otimes \left( m \cdot \bigoplus_i \pi_i^\vee \right) \right)_{G(F)} \\ &= m \cdot \bigoplus_i (\Omega \otimes \pi_i^\vee)_{G(F)} \\ &= m \cdot \bigoplus_i \Theta(\pi_i). \end{aligned}$$

Here, we have used Lemma 5.2(i) which ensures that the contragredient of  $\tilde{\pi}$  as a representation of  $\tilde{G}(F)$  is the same as its contragredient as a representation of  $G(F)$ , so that

$$\tilde{\pi}^\vee|_{G(F)} \simeq m \cdot \bigoplus_i \pi_i^\vee.$$

This completes the proof of the lemma.

□

By Lemma 5.3 and Lemma 5.2, we deduce:

COROLLARY 5.4.

$\Theta(\tilde{\pi})$  has finite length as an  $\tilde{H}(F)^+$ -module

if and only if

$\Theta(\pi_i)$  has finite length as an  $H(F)$ -module for all  $i$ .

Assuming this finiteness condition holds, one has:

$$\Theta(\tilde{\pi}) = m \cdot \bigoplus_i \Theta(\pi_i)$$

and

$$(5.2) \quad \theta(\tilde{\pi}) = m \cdot \bigoplus_{i \in I} \theta(\pi_i)$$

as  $H(F)$ -modules.

**5.2. Proof of Theorem 5.1.** We can now prove Theorem 5.1. Suppose first that the Howe duality property holds for the pair  $(G, H)$ . By symmetry, for any  $\tilde{\pi} \in \text{Irr}(\tilde{G}(F)^+)$ , we need to show that if  $\Theta(\tilde{\pi})$  is nonzero, then it has finite length and  $\theta(\tilde{\pi})$  is irreducible.

Suppose that

$$\tilde{\pi}|_{G(F)} \simeq m \cdot \bigoplus_{i \in I} \pi_i$$

for  $\pi_i \in \text{Irr}(G(F))$ . Then by Corollary 5.4,

$$\Theta(\tilde{\pi})|_{H(F)} = m \cdot \bigoplus_i \Theta(\pi_i).$$

By hypothesis,  $\Theta(\pi_i)$  has finite length for each  $i$ , and hence so does  $\Theta(\tilde{\pi})$  as an  $\tilde{H}(F)$ -module by Corollary 5.4. Moreover, by Lemma 5.3,  $\Theta(\pi_i)$  is nonzero for each  $i$  and  $\sigma_i := \theta(\pi_i)$  is irreducible by hypothesis.

Let  $\tilde{\sigma} \subset \theta(\tilde{\pi})$  be an irreducible summand. Since (by (5.2))

$$\theta(\tilde{\pi})|_{H(F)} = m \cdot \bigoplus_{i \in I} \theta(\pi_i) = m \cdot \bigoplus_{i \in I} \sigma_i,$$

we see that

$$\tilde{\sigma}|_{H(F)} \simeq n \cdot \bigoplus_i \sigma_i \quad \text{for some } 0 < n \leq m.$$

If  $m = 1$ , then  $\theta(\tilde{\pi}) = \tilde{\sigma}$  and we would have been done already. For general  $m$ , we argue as follows. Switching the roles of  $G$  and  $H$  in the above argument, it follows by Corollary 5.4 that

$$\theta(\tilde{\sigma})|_{G(F)} = n \cdot \bigoplus_i \theta(\sigma_i).$$

By hypothesis, each  $\theta(\sigma_i)$  is irreducible and hence  $\theta(\sigma_i) = \pi_i$ . But since  $\tilde{\pi} \subset \theta(\tilde{\sigma})$ , we deduce that  $m \leq n$ . Hence, we must have  $m = n$ , so that  $\theta(\tilde{\pi}) = \tilde{\sigma}$  is irreducible.

Conversely, suppose now that the Howe duality property holds for the similitude pair  $(\tilde{G}(F)^+, \tilde{H}(F)^+)$ . By symmetry, take any  $\pi \in \text{Irr}(G(F))$  and choose  $\tilde{\pi} \in \text{Irr}(\tilde{G}(F)^+)$  such that  $\pi \subset \tilde{\pi}$ . Suppose

$$\tilde{\pi}|_{G(F)} \simeq m \bigoplus_{i \in I} \pi_i$$

with  $\pi_i \in \text{Irr}(G(F))$  for each  $i$ , and  $\pi = \pi_{i_0}$  for some  $i_0 \in I$ .

By Cor. 5.4, we have

$$\Theta(\tilde{\pi})|_{H(F)} = m \cdot \bigoplus_i \Theta(\pi_i).$$

Since we are assuming that  $\Theta(\pi) \neq 0$ , it follows by Lemma 5.3 that  $\Theta(\tilde{\pi})$  is nonzero. By hypothesis and Cor. 5.4,  $\Theta(\tilde{\pi})$  has finite length as a  $H(F)$ -module, and hence so does  $\Theta(\pi_i)$  for each  $i$ .

Consider  $\tilde{\sigma} := \theta(\tilde{\pi})$ , which is irreducible by hypothesis. By (5.2), we see that

$$(5.3) \quad \tilde{\sigma}|_{H(F)} \simeq m \cdot \bigoplus_{i \in I} \theta(\pi_i),$$

with each  $\theta(\pi_i)$  semisimple of finite length. On writing each  $\theta(\pi_i)$  as a sum of irreducible summands, we have:

$$(5.4) \quad \tilde{\sigma}|_{H(F)} \simeq n \bigoplus_{j \in J} \sigma_j$$

with

$$\text{non-isomorphic } \sigma_j \in \text{Irr}(H(F)), \quad |I| \leq |J| \quad \text{and } n \geq m.$$

Then applying (5.2) with the roles of  $G$  and  $H$  exchanged, one has

$$m \cdot \bigoplus_{i \in I} \pi_i = \tilde{\pi}|_{G(F)} = \theta(\tilde{\sigma})|_{G(F)} = n \cdot \bigoplus_{j \in J} \theta(\sigma_j),$$

with  $\theta(\sigma_j) \neq 0$ . This implies that

$$m = n, \quad |I| = |J| \quad \text{and } \theta(\sigma_j) = \pi_j,$$

after fixing some bijection of  $I$  with  $J$ . In particular, going back to (5.3) and (5.4), we deduce that  $\theta(\pi_i) = \sigma_i$  is irreducible for each  $i \in I$ .

This completes the proof of Theorem 5.1.

## 6. Examples

We give some interesting examples of similitude dual pairs and similitude theta correspondences.

**6.1. Classical dual pairs.** We begin by revisiting the case of classical dual pairs.

- (Symplectic-orthogonal) For a quadratic space  $V$  and a symplectic vector space  $W$ , we have

$$\iota : H \times G = O(V) \times \mathrm{Sp}(W) \longrightarrow \mathrm{Sp}(V \otimes W).$$

Then

$$\mathrm{Ker}(\iota) = \mu_2^\nabla.$$

Hence, we have  $Z = \mu_2$  and may take  $T = \mathbb{G}_m$ , so that

$$\tilde{H} = GO(V) \quad \text{and} \quad \tilde{G} = \mathrm{GSp}(W)$$

are the usual similitude groups. Then one has

$$\iota : \tilde{J}^{\mathrm{sim}} = (GO(V) \times \mathrm{GSp}(W))^{\mathrm{sim}} \longrightarrow \mathrm{Sp}(V \otimes W)$$

extending  $i$ .

Recall however that the Weil representation  $\Omega$  is not a representation of  $\mathrm{Sp}(V \otimes W)$  but rather of its metaplectic cover  $\mathrm{Mp}(V \otimes W)$ . In the classical theta correspondence, one needs to first construct liftings of  $\iota$  to  $\mathrm{Mp}(V \otimes W)$ :

$$\tilde{\iota} : O(V) \times \mathrm{Sp}(W) \longrightarrow \mathrm{Mp}(V \otimes W),$$

before one can restrict the Weil representation to  $O(V) \times \mathrm{Sp}(W)$ . Such splittings exist if  $\dim V$  is even and have been systematically constructed by Kudla [Ku]. Likewise, to obtain a similitude theta correspondence, we would need to extend  $\tilde{\iota}$  to:

$$\tilde{\iota} : (GO(V) \times \mathrm{GSp}(W))^{\mathrm{sim}} \longrightarrow \mathrm{Mp}(V \otimes W).$$

Such extensions have been constructed by B. Roberts [R]. This accounts for the main complexity in the theory of similitude theta correspondences for symplectic-orthogonal dual pairs.

- (Unitary) If  $V$  is a Hermitian space and  $W$  a skew-Hermitian space relative to a quadratic extension  $E/F$ , one has

$$\iota : U(V) \times U(W) \longrightarrow \mathrm{Sp}(\mathrm{Res}_{E/F}(V \otimes_E W))$$

with

$$\mathrm{Ker}(\iota) = (E^1)^\nabla$$

where we have written  $E^1$  for  $\mathrm{Res}_{E/F}^1(\mathbb{G}_m)$ . In this case, we take  $T = \mathrm{Res}_{E/F}\mathbb{G}_m$  to get the usual similitude groups

$$\tilde{G} = \mathrm{GU}(V) \quad \text{and} \quad \tilde{H} = \mathrm{GU}(W).$$

As in the symplectic-orthogonal case, one needs to construct a lifting

$$\tilde{\iota} : \tilde{J}^{\mathrm{sim}} = (\mathrm{GU}(V) \times \mathrm{GU}(W))^{\mathrm{sim}} \longrightarrow \mathrm{Mp}(\mathrm{Res}_{E/F}(V \otimes_E W))$$

before one can consider the similitude theta correspondence. However, the unitary case is somewhat better than the symplectic-orthogonal one.

More precisely, note that

$$\iota : \tilde{J}^{\text{sim}} = (\text{GU}(V) \times \text{GU}(W))^{\text{sim}} \longrightarrow \text{Sp}(V \otimes W)$$

has image contained in  $\text{U}(V \otimes_E W)$ . However, the metaplectic covering is split over  $\text{U}(V \otimes_E W)$ ! By fixing such a splitting, we thus obtain

$$\tilde{J}^{\text{sim}} \longrightarrow \text{U}(V \otimes_E W) \longrightarrow \text{Mp}(V \otimes_E W).$$

Thus, there is no need to do extra work beyond that needed for splitting the isometry dual pairs.

**6.2. Exceptional dual pairs.** Given the somewhat sporadic nature of the geometry of exceptional groups, it is not surprising that it is harder to formulate a uniform theory of dual pairs in exceptional groups. Nonetheless, we shall attempt to do so for a few families of such dual pairs, by realizing the dual pairs as

$$G \times H = \text{Aut}(A) \times \text{Aut}(B)$$

for two algebraic structures  $A$  and  $B$ .

**6.3. Composition algebras and cubic norm structures.** We refer the reader to [KMRT, Chap. VIII, §33] for the notion of composition algebras and [KMRT, Chap. 9, §37 and §38] for the notion of cubic norm structures (also known as Freudenthal-Jordan algebras). Let  $C$  be a composition algebra over  $F$ , so that  $\dim C = 1, 2, 4$  or  $8$ . Let  $J$  be a nontrivial cubic norm structure over  $F$ , so that  $J$  has dimension  $3, 6, 9, 15$  or  $27$ . Then one has an isometry dual pair [MS, Ru]

$$\text{Aut}(C) \times \text{Aut}(J) \hookrightarrow \mathcal{E}$$

where  $\mathcal{E}$  is a certain ambient adjoint group. We enumerate the most interesting (split) cases, where  $C$  or  $J$  have the maximal dimensions.

- For  $\dim C = 8$ , one has  $\text{Aut}(C) = G_2$  and as  $J$  varies, one has the dual pairs

$$G_2 \times \text{Aut}(J) \subset \mathcal{E}$$

with  $\text{Aut}(J)$  and  $\mathcal{E}$  given in the following table.

$\dim J$	3	6	9	15	27
$\text{Aut}(J)$	$S_3$	$\text{SO}_3$	$\text{PGL}_3 \rtimes \mathbb{Z}/2\mathbb{Z}$	$\text{PGSp}_6$	$F_4$
$\mathcal{E}$	$\text{PGSO}_8 \rtimes S_3$	$F_4$	$E_6 \rtimes \mathbb{Z}/2\mathbb{Z}$	$E_7$	$E_8$

- For  $\dim J = 27$ , one has  $\text{Aut}(J) = F_4$  and as  $C$  varies, one has the dual pairs

$$F_4 \times \text{Aut}(C) \subset \mathcal{E}$$

with  $\text{Aut}(C)$  and  $\mathcal{E}$  given in the following table.

$\dim C$	2	4	8
$\text{Aut}(C)$	$\mathbb{Z}/2\mathbb{Z}$	$\text{PGL}_2$	$G_2$
$\mathcal{E}$	$E_6 \rtimes \mathbb{Z}/2\mathbb{Z}$	$E_7$	$E_8$

Observe that the map from  $\text{Aut}(C) \times \text{Aut}(J)$  to  $\mathcal{E}$  is injective. Thus, in such cases, we do not have a theory of similitude theta correspondence.

**6.4. Twisted composition algebras.** Let  $E$  be an étale cubic  $F$ -algebra. Then one has the notion of twisted composition algebras with respect to  $E/F$ ; the reader can consult [KMRT, Chap. 8, §36] for this. One way such a twisted composition algebra arises is to start with a composition  $F$ -algebra  $C$  and consider  $C^b = C \otimes_F E$ . Then a construction in [KMRT, §36C, Pg 499] equips  $C^b$  with the structure of a twisted composition algebra, built out of the composition algebra structure on  $C$ . In fact, if  $C^b$  is any twisted composition algebra, then  $\dim_E(C^b) = 1, 2, 4$  or  $8$  [KMRT, Cor. 36.4, Pg 492].

Let  $C_1^b$  and  $C_2^b$  be two twisted composition algebras relative to  $E/F$ . Then one has a dual pair [GS2]

$$i : \text{Aut}_E(C_1^b) \times \text{Aut}_E(C_2^b) \longrightarrow \mathcal{E}$$

where  $\mathcal{E}$  is some ambient adjoint group. We enumerate the most interesting case with  $\dim_E C_1^b$  maximal. Then one has  $\text{Aut}_E(C_1^b) = \text{Spin}_8^E$ , a simply-connected quasi-split group of type  $D_4$  determined by  $E$ . As  $C_2^b$  varies (we set  $C^b = C_2^b$  to simplify notation), one obtains the dual pair

$$\text{Spin}_8^E \times \text{Aut}_E(C^b) \longrightarrow \mathcal{E}$$

with  $\text{Aut}(C^b)$  and  $\mathcal{E}$  given in the following table (we assume  $E = F^3$  so that the groups are split), where  $\text{Res}$  stands for  $\text{Res}_{E/F}$ :

$\dim_E C^b$	1	2	4	8
$\text{Aut}(C^b)$	$\text{Res}(\mu_2)/\mu_2$	$\text{Res}(\mathbb{G}_m)/\mathbb{G}_m \rtimes \mathbb{Z}/2\mathbb{Z}$	$\text{Res}(\text{SL}_2)/\mu_2$	$\text{Spin}_8^E$
$\mathcal{E}$	$F_4$	$E_6 \rtimes \mathbb{Z}/2\mathbb{Z}$	$E_7$	$E_8$

Observe that the group  $\text{Aut}_E(C^b)$  has center isomorphic to

$$Z = \text{Res}_{E/F}(\mu_2)/\mu_2 \simeq \text{Res}_{E/F}^1(\mu_2)$$

where the last isomorphism is given by

$$x \mapsto N_{E/F}(x)/x =: x^\#.$$

Moreover, the map

$$i : \text{Spin}_8^E \times \text{Aut}(C^b) \longrightarrow \mathcal{E}$$

has

$$\text{Ker}(i) = Z^\nabla \simeq \text{Res}_{E/F}^1(\mu_2).$$

Thus, in this case, we have a theory of similitude theta correspondence which we shall now explicate.

We first pick the data (a) and (b) in §2.1 as in Example (3) of §2.2. Namely, we take  $T = \text{Res}_{E/F}(\mathbb{G}_m)$  and consider the natural embedding

$$j : \text{Res}_{E/F}^1(\mu_2) \hookrightarrow T = \text{Res}_{E/F}(\mathbb{G}_m).$$

As in §2.2, we then have the embedding

$$T/j(Z) \xrightarrow{N_{E/F} \times [2]} \mathbb{G}_m \times \text{Res}_{E/F}(\mathbb{G}_m)$$

whose cokernel has dimension 1. In this special case, since  $3 - 2 = 1$ , it turns out that

$$T/j(Z) \simeq \text{Res}_{E/F}(\mathbb{G}_m) = T$$

via the map

$$x \mapsto N_{E/F}(x)/x^2 = x^\# / x.$$

Now let us consider the similitude groups

$$\tilde{G} = (\text{Spin}_8^E \times T)/Z^\nabla \xrightarrow{pr_2} T/j(Z) \xrightarrow{x \mapsto x^\# / x} T = \text{Res}_{E/F}(\mathbb{G}_m)$$

and

$$\tilde{H} = (\text{Aut}_E(C^b) \times T)/Z^\nabla \xrightarrow{pr_2} T/j(Z) \xrightarrow{x \mapsto x^\# / x} T = \text{Res}_{E/F}(\mathbb{G}_m),$$

Hence, the similitude maps  $\text{sim}_G$  and  $\text{sim}_H$  on the group of  $F$ -rational points both take value in  $T(F) = E^\times$ .

We would like to know what the groups  $\tilde{G}(F)^+$  and  $\tilde{H}(F)^+$  are. If  $F$  is nonarchimedean, then  $H^1(F, \text{Spin}_8^E) = 0$  (because  $\text{Spin}_8^E$  is simply connected semisimple; see [Kn]) and by Proposition 2.2, we see that the map  $\text{sim}_G : \tilde{G}(F) \rightarrow E^\times$  is surjective. Hence  $\tilde{H}(F)^+ = \tilde{H}(F)$  when  $F$  is nonarchimedean. The same is true for  $\tilde{G}(F)^+$  if  $\dim_E C^b = 8$ . We shall now consider the question of whether  $\text{sim}_H : \tilde{H}(F) \rightarrow E^\times$  is surjective when  $\dim_E C^b = 2$  or  $4$ , ignoring the somewhat degenerate case when  $\dim_E C^b = 1$ . For this, it is convenient to give an alternative description of  $\tilde{H}$  for which it is easier to describe  $\tilde{H}(F)$ .

- When  $\dim_E C^b = 4$ , the natural inclusion

$$\text{Res}_{E/F}(\text{SL}_2 \times \mathbb{G}_m) \hookrightarrow \text{Res}_{E/F}(\text{GL}_2 \times \mathbb{G}_m)$$

descends to an isomorphism

$$\tilde{H} \longrightarrow \text{Res}_{E/F}(\text{GL}_2 \times \mathbb{G}_m) / \iota(\text{Res}(\mathbb{G}_m))$$

where

$$\iota(t) = (t, (t^{-1})^\#).$$

Then the group  $\tilde{H}(F)$  is given by

$$\tilde{H}(F) = (\text{GL}_2(E) \times E^\times) / \iota(E^\times)$$

and the similitude character is

$$\text{sim}_H(h, x) = \det(h) \cdot \frac{x^\#}{x}.$$

From this, it is clear that  $\text{sim}_H$  is surjective, so that  $\tilde{G}(F)^+ = \tilde{G}(F)$ .

- When  $\dim_E C^\flat = 2$  and recalling that  $T = \text{Res}_{E/F}(\mathbb{G}_m)$ , we have an isomorphism:

$$\begin{aligned} \tilde{H}^0 &= (T \times T)/(\mathbb{G}_m \times \text{Res}_{E/F}(\mu_2)) \\ &\simeq \downarrow (t_1, t_2) \mapsto (t_1, t_1^{-1}, t_2) \\ &(T \times T \times T)/(\mathbb{G}_m \times T) \end{aligned}$$

where

- (i) the embedding

$$\mathbb{G}_m \times \text{Res}_{E/F}(\mu_2) \longrightarrow T \times T$$

in the first line is given by

$$(a, z) \mapsto (az, (z^{-1})^\#),$$

- (ii) the embedding

$$\mathbb{G}_m \times T \longrightarrow T \times T \times T$$

in the second line is given by

$$(a, t) \mapsto (at, a^{-1}t, (t^{-1})^\#).$$

With this alternative description of  $\tilde{H}^0$ , all tori involved are induced tori, so that

$$\tilde{H}^0(F) \simeq (E^\times \times E^\times \times E^\times)/(F^\times \times E^\times)$$

where  $F^\times \times E^\times$  is embedded into  $(E^\times)^3$  as in (ii) above. In this incarnation, the similitude character is given by

$$\text{sim}_H(x, y, z) = x \cdot y \cdot \frac{z^\#}{z} \in E^\times.$$

In particular,  $\text{sim}_H$  is surjective onto  $E^\times$ , so that  $\tilde{G}(F)^+ = \tilde{G}(F)$  in this case as well.

In particular, let us consider the case where the central character is trivial. Then we obtain for example the similitude dual pairs of adjoint type:

$$\text{PGSO}_8^E(F) \times \begin{cases} \text{PGSO}_8^E(F) \text{ in } E_8; \\ \text{PGL}_2(E) \text{ in } E_7; \\ (E^\times/F^\times) \rtimes \mathbb{Z}/2\mathbb{Z} \text{ in } E_6 \rtimes \mathbb{Z}/2\mathbb{Z}. \end{cases}$$



**6.5. Jordan pairs.** We refer the reader to [L, Introduction and Chap. 1] for the notion of Jordan pairs. Since this notion is perhaps less familiar to the reader than the notion of Jordan algebras, let us give some motivation and a brief introduction.

A Freuthendal-Jordan algebra or a cubic norm structure  $J$  comes equipped with a norm form, which is a cubic form  $N_J : J \rightarrow F$ . The similitude group of this cubic form

$$\text{sim}(J, N_J) = \{(g, t) \in \text{GL}(J) \times \mathbb{G}_m : N_J \circ g = t \cdot N_J\}$$

is called the structure group of  $J$ , and we will call the isometry group  $\text{iso}(J, N_J)$  of  $N_J$  the reduced or special structure group (this consists of those pairs  $(g, t)$  with  $t = 1$ ). If  $J$  is the 27-dimensional Jordan algebra, for example,  $\text{sim}(J, N_J)$  is the group  $GE_6$  and  $\text{iso}(J, N_J)$  is the simply-connected  $E_6$ .

Now the group  $\text{sim}(J, N_J)$  acts irreducibly on  $J$  and its linear dual  $J^*$ , but these two representations are not isomorphic (as their central characters are different) and there is no reason to favour one of these representation over the other (a more familiar example is: a group isomorphic to  $\text{GL}(V)$  has two standard representations). Indeed, the trace bilinear form on  $J$  allows us to identify  $J$  with  $J^*$ , and this defines an outer automorphism of  $\text{sim}(J, N_J)$  which is the inverse map on the center of  $\text{sim}(J, N_J)$  and which interchanges the two representations. In addition, in the context of  $E_6$ , when one considers a quasi-split  $E_6$  associated to a quadratic field extension  $K/F$ , the two 27-dimensional representations are fused together to give a single rational representation over  $F$ . Hence, such a quasi-split  $E_6$  cannot be realized as the isometry group of a cubic form over  $F$ , unlike the split form.

The theory of Jordan pairs, introduced by Loos [L], treats both these representations  $J$  and  $J^*$  on equal footing and realizes  $\text{sim}(J, N_J)$  as the automorphism group of an algebraic structure on the pair  $\{J, J^*\}$ . More formally, a Jordan pair over  $F$  consists of the data:

- a pair  $(J^+, J^-)$  of  $F$ -vector spaces;
- a pair of quadratic maps defined over  $F$ :

$$Q^+ : J^+ \rightarrow \text{Hom}_F(J^-, J^+) \quad \text{and} \quad Q^- : J^- \rightarrow \text{Hom}_F(J^+, J^-),$$

satisfying the following axioms [L, Pg. 1, Def. 1.2] for  $\epsilon = \pm$  and any  $F$ -algebra  $K$ :

(JP1)

$$\{x, y, Q^\epsilon(x)(z)\}^\epsilon = Q^\epsilon(x) (\{y, x, z\}^{-\epsilon}) \quad \text{for any } x \in J^\epsilon(K) \text{ and } y, z \in J^{-\epsilon}(K);$$

(JP2)

$$\{Q^\epsilon(x)(y), y, z\}^\epsilon = \{x, Q^{-\epsilon}(y)(x), z\}^\epsilon \quad \text{for any } x, z \in J^\epsilon(K) \text{ and } y \in J^{-\epsilon}(K);$$

(JP3)

$$Q^\epsilon(Q^\epsilon(x)(y)) = Q^\epsilon(x) \circ Q^{-\epsilon}(y) \circ Q^\epsilon(x) \text{ for any } x \in J^\epsilon(K) \text{ and } y \in J^{-\epsilon}(K),$$

where we have set

$$\{x, y, z\}^\epsilon = Q^\epsilon(x+z)(y) - Q^\epsilon(x)(y) - Q^\epsilon(z)(y) \in J^\epsilon$$

for the linearization of the map  $(x, y) \mapsto Q^\epsilon(x)(y)$ . These axioms may look a bit unwieldy for the uninitiated (including ourselves), but we will not seriously make use of them in this paper.

A homomorphism from one Jordan pair  $(J^+, J^-)$  to another  $(V^+, V^-)$  is a pair of linear maps

$$\phi^\epsilon : J^\epsilon \longrightarrow V^\epsilon \quad (\text{with } \epsilon = \pm)$$

such that

$$\phi^\epsilon(Q_J^\epsilon(x)(y)) = Q_V^\epsilon(\phi^\epsilon(x))(\phi^{-\epsilon}(y)) \quad \text{for } x \in J^\epsilon \text{ and } y \in J^{-\epsilon}.$$

Hence, one has a notion of the automorphism group of a Jordan pair  $(J^+, J^-)$ , which is a subgroup of  $\text{GL}(J^+) \times \text{GL}(J^-)$ . Via the projection onto each of the factors,  $\text{Aut}(J^+, J^-)$  has two natural representations.

**6.6. Examples.** We give two pertinent examples here:

- For a vector space  $V$ , set

$$(V^+, V^-) = (V, V^*),$$

and define:

$$Q^+(v)(v^*) = \langle v, v^* \rangle \cdot v, \quad Q^-(v^*)(v) = \langle v^*, v \rangle \cdot v^*.$$

This defines a Jordan pair whose automorphism group is the subgroup

$$\text{GL}(V) \hookrightarrow \text{GL}(V^+) \times \text{GL}(V^-).$$

Thus, this gives a description of the general linear group without favouring one of its two standard representations.

- For a cubic norm structure  $J$ , the pair

$$(J^+, J^-) = (J, J^*)$$

inherits the structure of a Jordan pair from its Jordan algebra structure and the automorphism group of  $(J^+, J^-)$  is precisely the structure group of  $J$ . More precisely, recall that in addition to the cubic norm form  $N_J$  and an identity element  $1_J$ , a cubic norm structure  $J$  comes equipped with

- a nondegenerate symmetric bilinear trace form  $T : J \times J \rightarrow F$ ;
- a quadratic map  $x \mapsto x^\#$  from  $J$  to itself, with linearization

$$x \times y = (x + y)^\# - x^\# - y^\#.$$

Given these, one sets

$$U_x(y) = T(x, y)x - x^\# \times y \quad \text{for } x, y \in J$$

and observes that  $U$  is quadratic in  $x$  and linear in  $y$ . The reader familiar with the notion of quadratic Jordan algebras will recognize that this is the  $U$ -operator in that theory. One then has a Jordan pair defined by setting:

$$(J^+, J^-) = (J, J) \quad \text{and} \quad (Q^+, Q^-) = (U, U).$$

Since  $T$  gives an identification of  $J$  with  $J^*$ , we may also describe this Jordan pair as  $(J, J^*)$ .

**6.7. Dual pairs.** After this brief sidetrack, we can now introduce a family of dual pairs in exceptional group of the form [MS, Ru]

$$i : \mathrm{SL}_3 \times \mathrm{iso}(J, N) \longrightarrow \mathcal{E}$$

where  $J$  is a Freudenthal Jordan algebra. For split groups, this dual pair is obtained by removing from the extended Dynkin diagram of  $\mathcal{E}$  (of type  $F_4$  or  $E_n$ ) the simple vertex joined to the unique vertex attached to the extra vertex. One sees that the extended Dynkin diagram breaks into two pieces, with one of them of type  $A_2$ .

By the notion of Jordan pairs introduced above, we recognize that this dual pair is of the form

$$\mathrm{Aut}(V^+, V^-)^{der} \times \mathrm{Aut}(J^+, J^-)^{der} = \mathrm{SL}(V) \times \mathrm{iso}(J, N_J),$$

where  $\dim V = 3$  and the superscript *der* signifies the derived group.

The following summarizes the algebras and groups which occur, where  $J(C)$  the space of 3 by 3 hermitian symmetric matrices with coefficients in a composition algebra  $C$  ( $F$ , quadratic  $K$ , quaternion  $B$ , or octonion  $\mathbb{O}$ ).

$\mathcal{E}$	$F_4$	$E_6 \rtimes \mathrm{Gal}(K/F)$	$E_7$	$E_8$
$C$	$F$	$K$	$B$	$\mathbb{O}$
$\dim J(C)$	6	9	15	27
$\mathrm{iso}(J, \det)$	$\mathrm{SL}_3(F)$	$\mathrm{SL}_3(K)/\mathrm{Res}_{K/F}^1(\mu_3) \rtimes \mathrm{Gal}(K/F)$	$\mathrm{SL}_3(B)/\mu_2$	$E_6^{sc}$

Observe that the centers of  $\mathrm{SL}(V)$  and  $\mathrm{iso}(J, \det)$  are isomorphic to  $\mu_3$  in all cases and

$$\mathrm{Ker}(i) = \mu_3^\nabla.$$

Hence, we take  $T = \mathbb{G}_m$ , so that

$$\tilde{J}^{\mathrm{sim}} = (\mathrm{GL}(V) \times \mathrm{sim}(J, N_J))^{\mathrm{sim}}.$$

PROPOSITION 6.1. *If  $J = J(C)$  is the space of 3 by 3 hermitian symmetric matrices with coefficients in a composition algebra  $C$ , and  $N_J = \det$ , then the similitude character  $\text{sim} : \text{sim}(J, N_J)(F) \rightarrow F^\times$  is surjective.*

PROOF. Let  $t \in F^\times$ . Let  $g : J \rightarrow J$  be the linear transformation such that for any  $x \in J$ ,  $y = g(x)$  is obtained from  $x$  by rescaling the entries in the following way. The entries in the upper-left  $2 \times 2$  block are multiplied by  $t$ , the remaining diagonal entry is divided by  $t$  and the other entries are left unchanged. Then  $\det(y) = t \det(x)$ . □

Thus in this case  $\tilde{G}(F) = \tilde{G}(F)^+$  and  $\tilde{H}(F) = \tilde{H}(F)^+$  and the resulting similitude dual pair is

$$\text{Aut}(V^+, V^-) \times \text{Aut}(J^+, J^-) = \text{GL}(V) \times \text{sim}(J, N_J).$$

Moreover, the group  $\text{sim}(J, N_J)(F)$  is given as follows:

$$\begin{aligned} \text{sim}(J, N_J)(F) = & \\ \left\{ \begin{array}{ll} \text{GL}_3(F), & \text{if } C = F; \\ (\text{GL}_3(K) \times F^\times) / \{(t, N_{K/F}(t)^{-1}) : t \in K^\times\} \rtimes \text{Gal}(K/F), & \text{if } C = K; \\ (\text{GL}_3(B) \times F^\times) / \{(t, t^{-2}) : t \in F^\times\}, & \text{if } C = B; \\ \text{GE}_6(F), & \text{if } C = \mathbb{O}. \end{array} \right. \end{aligned}$$

6.8. **Twisted Jordan pairs.** It is in fact better to slightly repackage the definition of Jordan pairs introduced above, by setting

$$J^\square = J^+ \times J^- \quad \text{regarded as an } F \times F\text{-module}$$

and

$$Q^\square = Q^+ \times Q^- : J^\square \longrightarrow \text{Hom}_{F \times F}((J^\square)^\sigma, J^\square)$$

where  $\sigma$  is the nontrivial automorphism of the  $F$ -algebra  $F \times F$  given by switching the two components. The axioms (JP1-3) for a Jordan pair can be accordingly reformulated in terms of  $(J^\square, Q^\square)$ . A homomorphism  $\phi : (J^\square, Q^\square_J) \longrightarrow (V^\square, Q^\square_V)$  is then a  $F \times F$ -module homomorphism  $J^\square \longrightarrow V^\square$  such that

$$\phi(Q^\square_J(x)(y)) = Q^\square_V(\phi(x))(\phi(y)) \quad \text{for } x, y \in J^\square.$$

The advantage of such a repackaging is that it allows one to introduce a twisted version (or  $F$ -rational forms) of Jordan pairs, where one replaces  $F \times F$  by a separable quadratic field extension  $K/F$ .

Moreover, in the definition of an automorphism of  $(J^\square, Q^\square)$ , we may consider those  $\phi$ 's which are  $(F \times F, \sigma)$ -linear instead of  $(F \times F)$ -linear. This gives rise to a larger  $F$ -automorphism group  $\text{Aut}_F(J^\square, Q^\square)$  containing the subgroup  $\text{Aut}_{F \times F}(J^\square, Q^\square)$  as a normal subgroup of index 2; these extra  $F$ -automorphisms are outer automorphisms of  $\text{Aut}_{F \times F}(J^\square, Q^\square)$  and their action on  $J^\square$  exchanges the two factors  $J^+$  and  $J^-$ .

Now let  $K$  be a separable étale  $F$ -algebra with nontrivial automorphism  $\sigma \in \text{Aut}(K/F)$ . We shall define a notion of Jordan pair  $(J, Q)$  with respect to  $K/F$ , following a paper of de Medts [dM] where it was introduced under the guise of *Hermitian cubic norm structure* [dM, §4]. Such a Jordan pair consists of:

- a  $K$ -vector space  $J$ ;
- a  $K$ -quadratic map  $Q : J \longrightarrow \text{Hom}_K(J^\sigma, J)$

satisfying the reformulated analog of (JP1-3) (we will not elaborate further here). Let us give two examples:

- Suppose  $(V, h)$  is a  $K$ -vector space equipped with a Hermitian form  $h : V \times V^\sigma \rightarrow K$ . Then defining  $Q$  by

$$Q(v)(w) = h(v, w) \cdot v \quad \text{for } v, w \in V$$

gives a Jordan pair relative to  $K/F$ . The automorphism group of  $(V, Q)$  is precisely the unitary group  $U(V, h)$ .

- In [dM, Thm. 4.6], it was explained how a cubic norm structure  $J$  over  $K$  and a “ $\sigma$ -linear self adjoint autotopy” of  $J$  give rise to a Hermitian cubic norm structure on  $J$ , from which one can deduce a Jordan pair  $(J, Q)$  over  $K/F$  with  $Q$  given by the  $U$ -operator for the Hermitian cubic norm structure. If  $J$  is the exceptional Jordan algebra, then  $\text{Aut}_K(J, Q)^{\text{der}}$  is the quasi-split simply connected  $E_6^K$  associated to  $K/F$  and

$$\text{Aut}_K(J, Q)/\text{Aut}_K(J, Q)^{\text{der}} \simeq \text{Res}_{K/F}^1(\mathbb{G}_m)$$

is the anisotropic 1-dimensional torus associated to  $K$ .

Associated to these two examples and as a twisted version of the dual pair  $\text{SL}_3 \times \text{iso}(J, N_J)$  introduced in the previous subsection, one has the quasi-split but non-split dual pair

$$\text{SU}(V, h) \times \text{Aut}_K(J, Q)^{\text{der}} \longrightarrow \mathcal{E},$$

where we assume for simplicity that  $\mathcal{E}$  is split. The centers of these groups are isomorphic to

$$Z = \text{Res}_{K/F}^1(\mu_3).$$

Following Example (3) in §2, we choose  $T = \text{Res}_{K/F}(\mathbb{G}_m)$  and note that there is a short exact sequence

$$1 \longrightarrow Z \longrightarrow T \longrightarrow T \longrightarrow 1$$

defined by the map

$$x \mapsto x^3/N_{K/F}(x) \quad \text{on } T.$$

The associated similitude dual pair is:

$$\text{GU}(V, h) \times \text{GAut}_K(J, Q)$$

where

$$\mathrm{GAut}_K(V, Q) = (\mathrm{Res}_{K/F}(\mathbb{G}_m) \times \mathrm{Aut}_K(J, Q)^{\mathrm{der}}) / Z^\nabla = (\mathbb{G}_m \times \mathrm{Aut}_K(J, Q)) / \mu_2^\Delta.$$

We note here that the similitude character on  $\mathrm{GU}(V, h)$  is not the usual similitude character  $\mathrm{sim} : \mathrm{GU}(V, h) \rightarrow \mathbb{G}_m$ . Rather, by construction, it is the homomorphism

$$\mathrm{GU}(V, h) \rightarrow \mathrm{Res}_{K/F}(\mathbb{G}_m)$$

given by

$$g \mapsto \det(g) / \mathrm{sim}(g).$$

This is surjective onto  $K^\times$ , as one can see by restricting it to a maximal torus in a Borel subgroup (or by observing that  $\mathrm{SU}(V, h)$  is simply-connected when  $F$  is nonarchimedean). Hence  $\mathrm{GAut}_K(J, Q)(F)^+ = \mathrm{GAut}_K(J, Q)(F)$ . We will leave it to the reader to work out what  $\mathrm{GU}(V, h)(F)^+$  is in the various cases.

### 7. Seesaw Duality

In the theory of theta correspondence, seesaw dual pairs and the associated seesaw dualities serve as useful tools. In this section, we examine how this seesaw duality is impacted by the extension to similitude dual pairs. Hence, suppose one has a seesaw diagram of dual pairs in  $\mathcal{E}$ :

$$\begin{array}{ccc} G' & & H \\ & \searrow & \swarrow \\ & & H' \\ & \swarrow & \searrow \\ G & & H' \end{array}$$

For  $\pi \in \mathrm{Irr}(G(F))$  and  $\sigma \in \mathrm{Irr}(H'(F))$ , the associated seesaw identity reads:

$$\mathrm{Hom}_{H'}(\Theta(\pi), \sigma) \simeq \mathrm{Hom}_{G \times H'}(\Omega, \pi \otimes \sigma) \simeq \mathrm{Hom}_G(\Theta(\sigma), \pi).$$

Now let us make the following hypotheses:

- The pair  $(G, H)$  gives rise to a corresponding similitude dual pair  $(\tilde{G}, \tilde{H})$ , as we explained earlier in this paper;
- for the other dual pair  $(G', H')$ ,  $G' \times H'$  maps injectively into  $\mathcal{E}$  (so that there is no associated similitude pair).

Observe also that

$$G \times H' \subset \tilde{J}^{\mathrm{sim}} = (\tilde{G} \times \tilde{H})^{\mathrm{sim}}(F).$$

Then we have:

**PROPOSITION 7.1.** *Assume the above hypotheses. Then for  $\tilde{\pi} \in \mathrm{Irr}(\tilde{G}(F))$  and  $\sigma \in \mathrm{Irr}(H'(F))$ , one has the similitude seesaw identity:*

$$\mathrm{Hom}_G(\Theta(\sigma), \tilde{\pi}) \simeq \mathrm{Hom}_{H'}(\Theta(\tilde{\pi}), \sigma).$$

PROOF. For  $\tilde{\pi} \in \text{Irr}(\tilde{G}(F))$  and  $\sigma \in \text{Irr}(H'(F))$ , we consider

$$\text{Hom}_{G \times H'}(\Omega, \tilde{\pi} \otimes \sigma).$$

On the one hand, this is isomorphic to

$$\text{Hom}_{G \times H'}(\Omega \otimes \sigma^\vee, \tilde{\pi}) = \text{Hom}_G((\Omega \otimes \sigma^\vee)_{H'}, \tilde{\pi}) \simeq \text{Hom}_G(\Theta(\sigma), \tilde{\pi}).$$

On the other hand, by Lemma 5.2(i), we also get

$$\text{Hom}_{G \times H'}(\Omega \otimes \tilde{\pi}^\vee, \sigma) = \text{Hom}_{H'}((\Omega \otimes \tilde{\pi}^\vee)_G, \sigma) = \text{Hom}_{H'}(\Theta(\tilde{\pi}), \sigma).$$

Hence, one deduces that

$$\text{Hom}_G(\Theta(\sigma), \tilde{\pi}) \simeq \text{Hom}_{H'}(\Theta(\tilde{\pi}), \sigma),$$

as desired. □

In other words, there is no significant problem in extending the seesaw identity to the similitude setting, at least under the hypotheses in the proposition.

Let us give some examples of the above situation, using the exceptional dual pairs we discussed above. A first example is the seesaw diagram:

$$\begin{array}{ccc} \text{Aut}(\mathbb{O}) = G_2 & & \text{iso}(J, N) \\ | & \diagdown & | \\ \text{Aut}(V^\square, Q^\square)^{\text{der}} = \text{SL}_3 & & \text{Aut}(J) \end{array}$$

where

- $J$  is a cubic norm structure over  $F$  with norm form  $N$ ;
- $V$  is a 3-dimensional vector space over  $F$ , giving rise to a Jordan pair  $(V^\square, Q^\square)$ ;
- $\mathbb{O}$  is the octonion  $F$ -algebra constructed as  $\mathbb{O} = F^2 \times V^\square$  (as a structurable algebra [dM, §4], a notion we did not introduce).

We have explained that the dual pair  $\text{Aut}(J) \times \text{Aut}(\mathbb{O})$  does not have a similitude version, whereas the other pair  $\text{SL}(V) \times \text{iso}(J, N)$  does.

As another example, we have:

$$\begin{array}{ccc} \text{Aut}(E \oplus C^b) & & \text{Aut}_E(\mathbb{O}^b) = \text{Spin}_8^E \\ | & \diagdown & | \\ \text{Aut}_E(C^b) & & \text{Aut}(\mathbb{O}) = G_2 \end{array}$$

where

- $\mathbb{O}$  is the octonion  $F$ -algebra giving rise to the twisted octonion algebra  $\mathbb{O}^b = \mathbb{O} \otimes_F E$  over  $E/F$ ;

- $C^b$  is a twisted composition algebra relative to  $E/F$  giving rise to a cubic norm structure (or Freudenthal-Jordan algebra)  $J = E \oplus C^b$  via the Springer decomposition [KMRT, §38A, Pg. 522].

As above, the dual pair  $\text{Aut}(\mathbb{O}) \times \text{Aut}(J)$  does not have a similitude version whereas the pair  $\text{Aut}(\mathbb{O}^b) \times \text{Aut}_E(C^b)$  does.

Observe that in forming both these seesaw diagrams, the initial data consists of giving two algebraic structures on the bottom row of the diagram, which then induces the algebraic structures in the top row. Moreover, observe that the two seesaw diagrams can be combined into a single one involving 3 dual pairs.

### 8. Global Theta Correspondence

In this section, we consider the global theta correspondence. Hence, let  $k$  be a number field with adèle ring  $\mathbb{A}$ . Suppose that we have a dual pair

$$i : G \times H \longrightarrow \mathcal{E}$$

as in the introduction, with  $\text{Ker}(i) = Z^\nabla$ .

Let  $\Omega$  be the global minimal representation of  $\mathcal{E}$  and suppose one has an automorphic realization

$$\theta : \Omega \longrightarrow \mathcal{A}(\mathcal{E}),$$

where  $\mathcal{A}(\mathcal{E})$  denotes the space of smooth (but not necessarily  $K_\infty$ -finite) automorphic forms (see [W] or [GZ]). For an irreducible cuspidal representation  $\pi \subset \mathcal{A}_{cusp}(G)$  of  $G$ , one has the usual notion of global theta lifting. More precisely, for  $\phi \in \Omega$  and  $f \in \pi$ , one sets

$$\theta(\phi, f)(h) = \int_{[G]} \theta(\phi)(gh) \cdot \overline{f(g)} dg$$

for  $h \in H(\mathbb{A})$ . Then the global theta lift of  $\pi$  is the  $H(\mathbb{A})$ -submodule

$$\Theta(\pi) = \langle \theta(\phi, f) : \phi \in \Omega, f \in \pi \rangle \subset \mathcal{A}(H).$$

Moreover the map  $(\phi, f) \mapsto \theta(\phi, f)$  is a  $G(\mathbb{A})$ -invariant and  $H(\mathbb{A})$ -equivariant map

$$\theta : \Omega \otimes \bar{\pi} \longrightarrow \Theta(\pi).$$

Now suppose we have chosen data (a) and (b) as in §2 and hence have the similitude groups  $\tilde{G}$  and  $\tilde{H}$ . On taking points, we have the groups

$$\tilde{G}(k)^+ \subset \tilde{G}(\mathbb{A})^+ = \prod_v \tilde{G}(k_v)^+$$

and likewise for  $\tilde{H}(k)^+ \subset \tilde{H}(\mathbb{A})^+$ . Moreover, by construction, one has a group homomorphism

$$\iota : \tilde{J}^{\text{sim}} = (\tilde{G} \times \tilde{H})^{\text{sim}} \longrightarrow \mathcal{E}.$$



Let  $\tilde{\pi}$  be an irreducible cuspidal representation of  $\tilde{G}(\mathbb{A})^+ = \prod_v \tilde{G}(F_v)^+$  with a realization

$$\tilde{\pi} \subset \mathcal{A}_{cusp}(\tilde{G}(k)^+ \backslash \tilde{G}(\mathbb{A})^+).$$

We would like to define its similitude global theta lift to the space  $\mathcal{A}(\tilde{H}^+)$  of automorphic forms on  $\tilde{H}(\mathbb{A})^+$ . For  $f \in \tilde{\pi}$ ,  $\phi \in \Omega$  and  $y \in \tilde{J}^{\text{sim}}(\mathbb{A})$ , we set

$$\theta(\phi, f)(y) = \int_{[G]} \theta(\phi)(g \cdot \iota(y)) \cdot \overline{f(g)} dg.$$

Here (and elsewhere),  $[G]$  is used to denote the quotient  $G(F) \backslash G(\mathbb{A})$ . This function of  $\tilde{J}^{\text{sim}}(\mathbb{A})$  descends to a function on  $\tilde{H}(\mathbb{A})^+$  which is left invariant under  $\tilde{H}(k)^+$ , thus giving an element of  $\mathcal{A}(\tilde{H}^+)$ .

A more concrete way of defining the function  $\theta(\phi, f)$  as a function on  $\tilde{H}(\mathbb{A})^+$  is as follows. Given  $h \in \tilde{H}(\mathbb{A})^+$ , one can find an element  $t(h) \in \tilde{G}(\mathbb{A})^+$  such that

$$\text{sim}_{\tilde{G}}(t(h)) = \text{sim}_{\tilde{H}}(h).$$

Then one has an element

$$(t(h), h) \in \tilde{J}(\mathbb{A})^{\text{sim}} = (\tilde{G} \times \tilde{H})^{\text{sim}}(\mathbb{A}).$$

Then

$$\theta(\phi, f)(h) = \int_{[G]} \theta(\phi)(\iota(gt(h), h)) \cdot \overline{f(g)} dg.$$

Since this is merely an explication of the more natural definition of  $\theta(\phi, f)$  given initially, the choice of the element  $t(h)$  is immaterial.

The span of the functions  $\theta(\phi, f)$ , as  $\phi \in \Omega$  and  $f \in \tilde{\pi}$  vary, is the global similitude theta lift  $\Theta(\tilde{\pi})$  of  $\tilde{\pi}$ . We note:

**PROPOSITION 8.1.** *Let  $\tilde{\pi} \subset \mathcal{A}(\tilde{G}^+)$  be an irreducible cuspidal representation of  $\tilde{G}(\mathbb{A})^+$ . Assume that*

- $\Theta(\tilde{\pi}) \subset \mathcal{A}_2(\tilde{H}^+)$  is nonzero and contained in the space of square-integrable automorphic forms (with a fixed central character);
- the local Howe duality property holds at all places  $v$  of  $k$  for  $G(k_v) \times H(k_v)$  or equivalently  $\tilde{G}(k_v)^+ \times \tilde{H}(k_v)^+$ ;

Then

$$\Theta(\tilde{\pi}) \simeq \bigotimes_v \theta(\tilde{\pi}_v).$$

In particular,  $\Theta(\tilde{\pi})$  is irreducible and one has the compatibility of the local and global similitude theta correspondences.

### 9. Appendix: Clifford theory for $p$ -adic groups

Let  $G$  be a locally compact group and  $N$  be a normal open subgroup of  $G$  of finite index. Let  $A = G/N$ , and  $m = |A|$ . We assume that the topology on  $N$  (and hence on  $G$ ) is defined by a sequence of open compact groups. We assume that  $G/K$  is countable for some open compact subgroup  $K$ , and hence for all open compact groups of  $G$ . This condition assures that Schur's lemma holds for smooth irreducible representations of  $G$  and  $N$ . The following elementary result is based on [BH, §2.7].

**PROPOSITION 9.1.** *Let  $(\pi, V)$  be a smooth finite length representation of  $G$ . Then*

1.  $V$  is a finite length  $N$ -module.
2.  $V$  is semisimple  $G$ -module if and only if it is a semisimple  $N$ -module.
3. The  $N$ -socle of  $V$  is equal to the  $G$ -socle of  $V$ .

**PROOF.** (1) We can assume that  $V$  is irreducible. Since  $V$  is finitely generated over  $N$ , it admits an irreducible  $N$ -quotient  $U$ . Then, by Frobenius reciprocity,  $V$  is a submodule of  $\text{Ind}_N^G U$ . The latter is a semisimple  $N$ -module of finite length. Thus  $V$  is not only finite length but also a semisimple  $N$ -module. This completes (1) and gives one direction of (2). Now assume that  $V$  is  $N$ -semisimple. Let  $W \subset V$  be a  $G$ -submodule. Since  $V$  is  $N$ -semisimple, there exists an  $N$ -invariant projection  $P$  of  $V$  onto  $W$ . Then

$$\frac{1}{m} \sum_{g \in G/N} \pi(g)P\pi(g)^{-1}$$

is a  $G$ -invariant projection of  $V$  onto  $W$ . Let  $U \subset V$  be the kernel of this projection. Then  $V = W \oplus U$  and  $V$  is  $G$ -semisimple. (3) The  $G$ -socle of  $V$  is  $H$ -semisimple by (2) hence it is contained in the  $H$ -socle. On the other hand, the  $H$ -socle is  $G$ -stable, and  $G$ -semisimple by (2). Hence it is contained in the  $G$ -socle.  $\square$

Next we want to describe smooth irreducible representations of  $G$  in terms of those of  $N$ . Let  $V$  be an irreducible representation of  $G$  and  $U$  an irreducible  $N$ -quotient. By Frobenius reciprocity,  $V$  is an irreducible submodule of  $\text{Ind}_N^G(U)$ . Thus we need to analyze the induction from  $N$  to  $G$ . For every  $g \in G$ , let  $U^g$  be the  $g$ -conjugate of  $(\pi, U)$ , that is, the representation of  $N$  on  $U$  where every  $n \in N$  acts by  $\pi(gng^{-1})$ . Let  $G_U$  be the stabilizer of  $U$  in  $G$ , that is, the subgroup of  $G$  consisting of all  $g \in G$  such that  $U$  is isomorphic to  $U^g$ , as  $N$ -modules.

**LEMMA 9.2.** *Let  $V$  be an irreducible representation of  $G_U$  containing  $U$ . Then*

$$\text{Ind}_{G_U}^G V$$

*is an irreducible representation of  $G$ .*

PROOF. Observe that the restriction of  $V$  to  $N$  is a multiple of  $U$ . Let  $U_1, \dots, U_l$  be all non-isomorphic  $G$ -conjugates of  $U$ . We have a decomposition, as an  $N$ -module,

$$\mathrm{Ind}_{G_U}^G V = \bigoplus_{i=1}^l V_i$$

where  $V_i$  is a multiple of  $U_i$ . Observe that  $G$  permutes these summands, and that each  $V_i$  is an irreducible  $G_{U_i}$ -module. Furthermore, for each  $i$ ,

$$V_i' = \bigoplus_{j \neq i} V_j$$

is a  $G_{U_i}$ -module. Now let  $W$  be a  $G$ -submodule of  $\mathrm{Ind}_{G_U}^G V$ . Fix an  $i$ . It is easy to see that any  $G_i$ -submodule of  $\mathrm{Ind}_{G_U}^G V$ , in particular  $W$ , either contains  $V_i$  or is contained in  $V_i'$ . Since  $W$  is a  $G$ -module, it follows that  $W = \mathrm{Ind}_{G_U}^G V$  or 0.  $\square$

Next we need to understand irreducible representation of  $G_U$  containing  $U$ . To simplify notation assume that  $G = G_U$ . Then, for every  $g \in G$ , there exists  $A_g : U \rightarrow U$  that intertwines  $U$  and  $U^g$ . By Schur's lemma  $A_g$  is unique up to an element in  $\mathbb{C}^\times$ . Thus  $U$  is naturally a module for a central extension

$$1 \rightarrow \mathbb{C}^\times \rightarrow \tilde{G} \rightarrow G \rightarrow 1.$$

This extension splits over  $N$ , indeed, we have a canonical choice  $A_n = \pi(n)$  for all  $n \in N$ . After taking the quotient by  $N$  we get a central extension

$$1 \rightarrow \mathbb{C}^\times \rightarrow \tilde{A} \rightarrow A \rightarrow 1.$$

Next, we have

$$U \otimes \mathrm{Ind}_{\mathbb{C}^\times}^{\tilde{A}}(\epsilon) \simeq \mathrm{Ind}_N^G(U)$$

where  $U$  is viewed as  $\tilde{G}$ -module and  $\epsilon : \mathbb{C} \rightarrow \mathbb{C}$  is the inverse character. This isomorphism is realized by

$$u \otimes f \mapsto (g \mapsto f(g)\pi(g)(u)).$$

Thus in order to decompose  $\mathrm{Ind}_N^G(U)$  it suffices to decompose  $\mathrm{Ind}_{\mathbb{C}^\times}^{\tilde{A}}(\epsilon)$ .

Claim: The extension  $\tilde{A}$  is defined by a co-cycle with values in  $\mu_m$ . To see this, define a section  $s : A \rightarrow \tilde{A}$  such that  $s(a)$ , for every  $a \in A$ , acts on  $\mathrm{Ind}_{\mathbb{C}^\times}^{\tilde{A}}(\epsilon)$  as a linear transformation of determinant 1. Then the extension  $\tilde{A}$  is determined by the co-cycle  $c(a, b)$  defined by

$$s(a)s(b) = c(a, b)s(ab).$$

After taking determinant of both sides, we get  $c(a, b)^m = 1$ , as claimed.

Using well known facts from representations of finite groups, we can now write

$$\mathrm{Ind}_{\mathbb{C}^\times}^{\tilde{A}}(\epsilon) \simeq \bigoplus_E \dim(E) \cdot E$$

where the sum runs over all irreducible representations  $E$  of  $\tilde{A}$  such that  $\mathbb{C}^\times \subset \tilde{A}$  acts by  $\epsilon$ . Call such representations genuine.

LEMMA 9.3. *Let  $U$  be an irreducible representation of  $N$  such that  $G_U = G$ . With notation as above*

$$\mathrm{Ind}_N^G U = \bigoplus_E \dim(E) \cdot U \otimes E$$

where the sum is over all irreducible genuine representations  $E$  of  $\tilde{A}$ . Representations  $U \otimes E$  are irreducible and mutually non-isomorphic.

PROOF. It remains to prove the last sentence. Let  $E_1$  and  $E_2$  be any two genuine  $\tilde{A}$ -modules. Since  $U$  is irreducible  $N$ -module,

$$\mathrm{Hom}_N(U \otimes E_1, U \otimes E_2) \simeq \mathrm{Hom}_{\mathbb{C}}(E_1, E_2).$$

Thus

$$\mathrm{Hom}_G(U \otimes E_1, U \otimes E_2) \simeq \mathrm{Hom}_{\tilde{A}}(E_1, E_2).$$

The lemma follows.  $\square$

We now briefly discuss the lemma when  $A$  is abelian. Then  $\tilde{A}$  is a two-step nilpotent group. The commutator of elements in  $\tilde{A}$  defines a skew-linear form on  $A \times A$

$$\langle a, b \rangle = [s(a), s(b)].$$

Since the order of  $A$  is  $m$ , the skew form, and therefore the commutator, takes values in  $\mu_m$ . The dimension of any genuine irreducible representation  $E$  is equal to  $\sqrt{|\bar{A}|}$  where  $\bar{A}$  is the quotient of  $A$  by the kernel of the skew form  $\langle a, b \rangle$ . Furthermore, any two irreducible representations of  $G$  containing  $U$  are  $A$ -character twists of one another. Moreover, two character twists are isomorphic if and only if the characters coincide on the kernel of the skew form.

Finally we combine the two lemmas to obtain the general case.

PROPOSITION 9.4. *Let  $U$  be an irreducible  $N$ -module, and  $G_U$  its stabilizer in  $G$ . Let  $A_U = G_U/N$ . Let  $\tilde{A}_U$  be the central extension of  $A_U$  arising from the projective action of  $G_U$  on  $U$ . Then*

$$\mathrm{Ind}_N^G U = \bigoplus_E \dim(E) \cdot \mathrm{Ind}_{G_U}^G (U \otimes E)$$

where the sum is over all irreducible genuine representations  $E$  of  $\tilde{A}_U$ . This is a decomposition of  $\mathrm{Ind}_N^G U$  into isotypic summands.

We remark that a study of the type of problems considered in this appendix can be found in the paper [T] of M. Tadić, where it was shown that the restriction of an irreducible representation of  $\mathrm{GL}_n(F)$  to  $\mathrm{SL}_n(F)$  is multiplicity-free.

## ACKNOWLEDGEMENTS.

This paper is dedicated to Marko Tadić on the occasion of his 70th birthday. W.T.G. is partially supported by a Singapore government MOE Tier 1 grant R-146-000-320-114 and the Tan Chin Tuan Centennial Professorship at NUS. G. Savin is partially supported by a National Science Foundation grant DMS-1901745 and by a Simons Foundation gift 946504. The authors would like to thank Skip Garibaldi for pointing them to the notion of Jordan pairs developed in Loos' book [L].

## REFERENCES

- [BH] C. J. Bushnell and G. Henniart, *Local Langlands conjecture for  $GL(2)$* , Springer-Verlag, Berlin-Heidelberg, 2006.
- [dM] T. De Medts, *Structurable algebras of skew-dimension one and Hermitian cubic norm structures*, *Comm. Algebra* **47** (2019), 154–172.
- [GS] W. T. Gan and G. Savin, *On minimal representations definitions and properties*, *Represent. Theory* **9** (2005), 46–93.
- [GS2] W. T. Gan and G. Savin, *Twisted composition algebras and Arthur packets of triality  $Spin(8)$* , *Pure and Applied Math. Q.* **18** (2022), 1951–2130.
- [GZ] H. Grobner and S. Žunar, *On the notion of the parabolic and the cuspidal support of smooth-automorphic forms and smooth-automorphic representations*, Preprint, [arXiv:2108.06369](https://arxiv.org/abs/2108.06369).
- [Kn] M. Kneser, *Galois-Kohomologie halbeinfacher algebraischer Gruppen über  $p$ -adischen Körpern II*, *Math. Z.* **89** (1965), 250–272.
- [KMRT] M. Knus, A. Merkurjev, M. Rost and J.-P. Tignol, *The Book of Involutions*, AMS Colloquium Publications, Vol. **44**, American Mathematical Society, Providence, RI, 1998.
- [Ku] S. Kudla, *On the local theta-correspondence*, *Invent. Math.* **83** (1986), 229–255.
- [L] O. Loos, *Jordan pairs*, *Lecture Notes in Math.*, Vol. **460**, Springer-Verlag, Berlin-New York, 1975.
- [MS] K. Magaard and G. Savin, *Exceptional  $\Theta$ -correspondences I*, *Compositio Math.* **107** (1997), 89–123.
- [R] B. Roberts, *The theta correspondence for similitudes*, *Israel J. Math.* **94** (1996), 285–317.
- [Ru] K. Rummelhart, *Minimal representations of exceptional  $p$ -adic groups*, *Represent. Theory* **1** (1997), 133–181.
- [T] M. Tadić, *Notes on representations of non-archimedean  $SL(n)$* , *Pacific J. Math.* **152** (1992), 375–396.
- [W] N. Wallach,  *$C^\infty$  vectors*, in: *Representations of Lie groups and quantum groups* (Trento, 1993), *Pitman Res. Notes Math. Ser.*, **311**, Longman Sci. Tech., Harlow, 1994, pp. 205–270.

## Iznimne theta korespondencije za grupe sličnosti

*Petar Bakić, Wee Teck Gan i Gordan Savin*

SAŽETAK. U ovom radu opisujemo sistematičan način konstrukcije dualnih parova grupa sličnosti. Proučavamo theta korespondencije do kojih dolazi na ovaj način, te dokazujemo da Howeova dualnost vrijedi za dualni par grupa sličnosti ako i samo ako vrijedi za reduktivni dualni par koji koristimo u našoj konstrukciji. Posebnu pažnju posvećujemo primjerima iznimnih theta korespondencija.

Petar Bakić  
Department of Mathematics  
University of Utah  
Salt Lake City, UT, USA  
*E-mail:* bakic@math.utah.edu

Wee Teck Gan  
Department of Mathematics  
National University of Singapore  
10 Lower Kent Ridge Road, Singapore 119076  
*E-mail:* matgwt@nus.edu.sg

Gordan Savin  
Department of Mathematics  
University of Utah  
Salt Lake City, UT, USA  
*E-mail:* savin@math.utah.edu

*Received:* 22.6.2023.

*Revised:* 17.10.2023.

*Accepted:* 14.11.2023.