

Changes in grassland vegetation on the island of Plavnik (Croatia) over 100 years

Massimo Terzi^{1*}, Nenad Jasprica²

¹ Institute of Bioscience and Bioresources (IBBR), National Council of Research (CNR), via Amendola 165/A, IT-70126, Bari, Italy

² University of Dubrovnik, Institute for Marine and Coastal Research, Kneza Damjana Jude 12, HR-20000 Dubrovnik, Croatia

Abstract – The changes in the grassland vegetation that have occurred over the last almost 100 years on the northeastern Adriatic island of Plavnik (Croatia) were studied. For this purpose, 29 phytosociological relevés of standard plot size were collected using the Braun-Blanquet phytosociological approach and compared with those of 100 years ago. With the aim of defining the differences among the communities and identifying the main environmental gradients, cluster analysis (Ward's method on a Chord distance matrix) and non-metric multidimensional scaling ordination were applied. Taxa score-weighted means of the Ellenberg-type indicator values of main environmental variables were used to assess ecological differences among communities, and disturbance-related indicator values were calculated to assess the disturbance regime in shaping plant communities. According to the syntaxonomic framework for the grassland associations, three associations and two subassociations were found. Among them the *Festuco valesiacae-Poetum bulbosae* (*Poetalia bulbosae*, *Poetea bulbosae*), the *Helichrysetum italicum salvietosum officinalis* (*Scorzoneretalia villosae*, *Festuco-Brometea*), and the *Hedypnoido rhagadioloidis-Asphodeletum ramosi* were proposed as new. The *Festuco valesiacae-Poetum bulbosae* documents for the first time, based on phytosociological data, the occurrence of the *Poetea bulbosae* in Croatia. Three associations were considered extinct, possibly as a result of changes in grazing pressure and type.

Keywords: eastern Adriatic, grazing pressure, *Helichrysum italicum*, Mediterranean Islands, *Poetea bulbosae*, *Scorzoneretalia villosae*, syntaxonomy, vegetation changes

Introduction

The Mediterranean basin is considered a biodiversity hotspot on account of its exceptional concentration of endemic species and the loss of habitats suitable for them (Myers et al. 2000). A significant component of this biodiversity is preserved in the Mediterranean islands due to the presence of steno-endemic taxa and distinctive plant assemblages (Médail 2017, Vargas 2020). The islands of the eastern Adriatic, with its multitude of archipelagos, are particularly important from this point of view, and they are characterised by a very high plant richness, comprising almost 1800 vascular taxa (Nikolić et al. 2008).

The phytosociological investigation of the north-eastern Adriatic using the approach of Braun-Blanquet (Westhoff and van der Maarel 1980) began in the 1930s, with the pioneering studies of Croatian botanist Stjepan Horvatić on some islands, such as Pag and Rab (Trinajstić 2001 and

references therein). On those islands, one of the most widespread aspects of vegetation was represented by open Mediterranean-sub-Mediterranean grasslands the particular floristic composition of which justified their attribution to an endemic alliance (*Chrysopogono grylli-Saturejion subspicatae* Horvat et Horvatić 1934 *nom. inval.*). Subsequently, that alliance was classified in an order (*Scorzonero villosae-Chrysopogonetalia grylli* Horvatić et Horvat in Horvatić 1963) and class also endemic to the Western Balkans (*Brachypodio-Chrysopogonetea* Horvatić 1963) (see Terzi 2011).

According to the current European syntaxonomic framework, the so-called EuroVegChecklist (EVC, Mucina et al. 2016), those grasslands mostly belong to the western Balkan alliances *Chrysopogono grylli-Koelerion splendidis* and *Scorzonerion villosae*, and to the trans-Adriatic order

* Corresponding author e-mail: massimo.terzi@cnr.it

Scorzoneretalia villosae, included in turn in the class *Festuco-Brometea* (but see also Terzi et al. 2023).

The first survey – the subject of Horvatić's doctoral thesis (Trinajstić 2001) – concerned the vegetation of the island of Plavnik, in the Kvarner area (Croatia), where Horvatić (1927) described five grassland syntaxa, namely *Festucetum valesiaca*, *Asphodeletum microcarpi*, *Helichrysetum italici*, “*Helichrysetum salviosum*” and *Andropogonetum grylli*. For each of them, a diagnosis was provided consisting of a variable number (from 5 to 10) of quadrats of 1 square metre each, with a complete list of taxa accompanied by quantitative scores. Although each vegetation type was sampled on only a small area, Horvatić (1927) gave a fairly precise idea of the main dominant and codominant taxa present at that time.

Quite surprisingly, neither Horvatić's later papers (e.g. Horvatić 1963, 1973) nor the subsequent syntaxonomic revisions of Croatian grasslands of the order *Scorzoneretalia villosae* (e.g. Horvat et al. 1974, Trinajstić 2008, Terzi 2015) mentioned these five syntaxa. Moreover, the herbaceous vegetation of the island of Plavnik has never been studied subsequently.

This island presents a rather interesting situation from a vegetation point of view because it has remained uninhabited for a long period of time, being used as a hunting reserve and only occasionally visited by hunters and tourists (along the coastal areas). The conspicuous presence of animals, sheep and fallow deer, left in the wild, has presumably significantly increased the grazing pressure on the grassland ecosystems.

The present work therefore aims to (i) provide new relevés for the herbaceous vegetation of the island of Plavnik, using a more reliable plot size than that used by Horvatić (1927), (ii) assess the changes in vegetation that have occurred over almost 100 years, (iii) provide a syntaxonomic framework for the grassland associations occurring on the island, and (iv) identify the main ecological drivers of grassland diversity.

Materials and methods

Study area

The island of Plavnik (surface area 8.63 km², coastline length 18.47 km) belongs to the north-eastern Adriatic group of islands in the Kvarner Bay, Croatia (Fig. 1). It is located between the islands of Cres and Krk and it stretches in a northwest-southeast direction for a length of 6.3 km. The coastline of the island is not indented: the northern coast is very steep and poorly accessible, while the southern is flat, gentle, accessible and with several coves.

The island has high terrain ruggedness. The northern and central part is a karstic plateau with an elevation of approximately 90–180 m a.s.l., bordered by steep rocky slopes exceeding 30°. The highest peak reaches an elevation of 193.6 m a.s.l. To the south-east, the karstic plateau gradually passes into lowlands with altitudes lower than 30 m a.s.l. (Fig. 2).

The bedrock consists mainly of carbonate rocks (Fuček et al. 2015). The Upper Cretaceous carbonate deposits can be recognized in different types of vertically alternating lithotypes. Multi-layered and massive recrystallized limestones with radiolithid-chondrodotic lithosomes are developed on both western and eastern coasts, while in the central part there are medium-thick layered shallow-sea mudstones and granular limestones with benthic foraminifera, radiolithid rudists and chondrodonts. The brown soils on limestone are predominant on the island (Vukadinović 2021).

The area is influenced by both Mediterranean and continental climates and characterized by intrusions of cold air during the winter and convective precipitation in the summer. These specific conditions result in the lack of a distinct dry period typical of the Mediterranean climate (Šegota and Filipčić 2003). The average annual air temperature is 15.3 °C and precipitation averages 1253.7 mm yr⁻¹ (data from the town of Krk station for 1998-2022, Croatian Meteorological and Hydrological Service). The highest monthly average



Fig. 1. The island of Plavnik is located between the islands of Cres and Krk in the Kvarner Bay (Croatia). The square on the map in the lower right corner shows the study area in the NE Adriatic context.




Fig. 2. Physiognomy of the plant associations on the island of Plavnik in May 2023: A – *Helichrysetum italici* Horvatić 1927 on the island's eastern coast, B – *Helichrysetum italici salvietosum officinalis subass. nov.* covers the open areas on the island's western slopes, C – *Hedynoides rhagadioloidis-Asphodeletum ramosi ass. nov.*, in the southernmost part of the island, D – *Festuco valesiacae-Poetum bulbosae ass. nov.* occurs on the karstic plateau in the north-central part of the island.

temperature is 30.5 °C in July, and the lowest 2.5 °C in January. The absolute maximum of 38.8 °C was recorded on 21 July 2015, and the absolute minimum (-9.0 °C) on 24 January 2006. The greatest rainfall comes in September, October and November with averages of 155.6, 162.0 and 178.0 mm, respectively. In the period from June to August the total rainfall is 184.2 mm. Northern winds (NW, ENE, NE) prevail throughout the year. This area has 2521 hours of sunshine per year (data for the island of Rab station for 1998-2022). On average the relative air humidity is 70%.

Phytoclimatic indices were calculated according to Rivas-Martínez et al. (2011): annual positive temperature (Tp) is 1,834; continentality index (Ic) is 18.9; thermicity index (It) is 238; ombrothermic index (Io) is 0.68. According to these indices, the neighbouring Krk station lies within the temperate macrobioclimate, meso-temperate thermotype and semiarid ombrotype.

The observed global and regional warming has been unequivocal since the middle of the 20th century (IPCC 2014) and is present along the eastern Adriatic coast (Branković et al. 2013). Climate change shows a tendency to manifest in drier conditions for the Adriatic (Patarčić et al. 2014, WBG 2021), which is a common feature for the broader European part of the Mediterranean (EEA 2019). However, for example, the precipitation change is regionally highly variable and further analyses are needed to define the assessment of climate change impacts at the regional and local scale (Gajić-Čapka et al. 2018).

From the phytogeographic point of view, the eastern and southwestern part of the island is covered by stands of the *Fraxino orni-Ostryion* alliance i.e. *Quercus pubescenti-Carpinetum orientalis* Horvatić 1939 association. The meso-Mediterranean evergreen vegetation of the *Fraxino orni-Quercion ilicis* alliance is developed only in the north-

western part of the island. However, woodlands have been significantly changed due to the long-term anthropogenic influences and nowadays only degraded vegetation types, i.e. maquis, garrigue and grasslands, may be found. Large areas of trees were felled between the two world wars to obtain open pastures, intended for sheep rearing. The island is classified among Croatia's Important Plant Areas (IPAs) and is part of the Natura 2000 ecological network (Topić and Vuković 2010, Official Gazette 2019).

The island has had no permanent residents or settlements during the last centuries. Between the two world wars, only a few herding groups lived on the island to provide living conditions for livestock, mainly sheep, while goats made up only 10-20% of the total number of head (Dumančić 1988). After the Second World War, such a simple nomadic way of life gradually faded away. The sheep population has drastically decreased, while goats, in general, have practically disappeared on the eastern coast of the Adriatic. There are no available data on the number of sheep in Plavnik over last century. In addition, the non-native European fallow deer (*Dama dama*) was introduced in 1995. According to the data of the local hunting association Farma Plavnik d.o.o. (personal communication), a number of the fallow deer, with regular annual shooting of ca. 50 nap, are stable and amounts to about 160 nap. The maximum nutritional requirements for females are during lactation, and this period coincides with the maximum development of vegetation.

Vegetation sampling and data analysis

The map of the distribution areas of the five types of plant communities on the island of Plavnik provided by Horvatić (1927: p. 44) was compared with the most recent satellite images available on Google Earth Engine, to

identify areas still covered by herbaceous vegetation. Those areas, and in particular the locations mentioned in that paper, were surveyed. The vegetation types were identified in the field based on their dominant and codominant taxa. Only for the “*Andropogonetum grylli*”, whose dominant taxon should have been *Chrysopogon gryllus*, we were unable to find any phytocoenosis clearly dominated by this plant, which is nowadays rather rare on the island.

The remaining four vegetation types were sampled through 29 relevés, made in May 2023 according to the phytosociological approach (see Westhoff and van der Maarel 1980). The relevés were sampled in quadrats of 25 m² (5 m × 5 m). All vascular plants in the plots were recorded and their cover-abundance values were visually estimated using the Braun-Blanquet cover-abundance scale with the subdivisions 2m, 2a and 2b as proposed by Barkman, Doing and Segal (in: van der Maarel 1979). This extended Braun-Blanquet scale (i.e. r, +, 1, 2m, 2a, 2b, 3, 4, 5) was replaced by the 9-point ordinal scale (i.e. 1-9, see Westhoff and van der Maarel 1980), before the statistical analyses. The main physical features of the plots (e.g., geographic coordinates, slope, exposure, vegetation cover, rockiness, and stoniness) were also recorded. Stoniness is defined here as the percentage of plot coverage by stones (incoherent material), regardless of their size. Rockiness, on the other hand, indicates the percentage of coverage of the plot determined by the outcropping rocky substrate.

The data matrix (29 relevés × 115 taxa) was subjected to hierarchical clustering using Ward's method on a Chord distance matrix (i.e., Euclidean distance on pre-transformed data through normalization by sites). Four interpretable clusters were subjectively defined by visual examination of the dendrogram.

To visualize the floristic and ecological relationships among the four clusters of relevés, nonmetric multidimensional scale (NMS) ordination was performed by using the Chord distance. Both cluster analysis and ordination were performed through the PcOrd 6.22 software package (McCune and Mefford 2011), the NMS with the "slow and thorough" option of the autopilot mode.

To assess ecological differences among the four plant communities, taxa score-weighted means of the Ellenberg-type Indicator Values (IVs) of temperature (T), soil reaction (R), moisture, and nutrient content were calculated for each relevé, based on the EIVs provided for Italy – which is the closest country geographically and ecologically to the study area – in Tichý et al. (2023).

In addition, given the presence of high numbers of sheep and fallow deer on the island of Plavnik and the possible importance of the disturbance regime in shaping the different plant communities, three additional disturbance-related IVs, taken from Midolo et al. (2023), were considered. In more detail, the following IVs, were calculated on taxa score-weighted means for each relevé: Disturbance severity (Ds), Disturbance frequency (Df), and Grazing pressure

(Gr). Joint plots were used to represent linear correlations between IVs and ordination scores, with an r^2 cut-off of 0.3.

Indicator species analysis (ISA) (Dufrêne and Legendre 1997) was used to identify the indicator taxa of the four main clusters of relevés obtained by cluster analysis, and their combinations in larger groups (see De Cáceres et al. 2010). The indicator values were calculated for all taxa present in at least three relevés. The statistical significance ($P < 0.01$) of the indicator values was tested by a Monte Carlo test with 10000 permutations. The ISA was run in R software (R Core Team 2023) through the package *indicpecies* 1.7.12 (De Cáceres and Legendre 2009).

In order to verify whether the relevés from the island of Plavnik in which *Asphodelus ramosus* was dominant (i.e. cluster 'He' in Figs. 3 and 4) could represent an association independent of the others described so far in the eastern Adriatic, we compared these relevés with the associations *Bromo erecti-Chrysopogonetum grylli* Horvatić 1934 and *Narcisso tazettae-Asphodeletum microcarpi* Šegulja 1969. In more detail, a second data matrix (55 relevés × 283 taxa) was obtained by adding the seven relevés reported in Tab. 3 to the other 48 relevés carried out by i) Horvatić (1934, rels. 11-23 of table XXVIII) on the island of Pag, Croatia; ii) Stanišić-Vujačić et al. (2022, rels. 1-17 of table 1) at Čemovsko polje, Montenegro, iii) Šegulja (1970, rels. 1-8 of table II) in Labin, Istria, Croatia, and iv) Hećimović (1984, *Narcisso tazettae-Asphodeletum microcarpi sisymbrietosum officinalis* Hećimović 1984, rels. 1-10 of table 5) on the islets of Bobara and Mrkan, in southern Croatia.

Using the same methods as described above, an NMS ordination was performed on that matrix to visualise the floristic relationships between the relevés, and then an indicator species analysis was run to identify the taxa that distinguish the *Asphodelus ramosus* community on Plavnik Island from those elsewhere.

The taxonomic nomenclature follows Euro+Med (2006-2023). *Bromopsis erecta* and *B. condensata* were treated under the same tag, the latter being more abundant and frequent on the island of Plavnik.

Syntaxonomic nomenclature of alliances, orders and classes follows the EuroVegChecklist (<https://floraveg.eu/vegetation/>, retrieved on 10 October 2023, as modified by Terzi et al. 2024a), whose correct names are given without author citations. Syntaxon names not included in the EVC, but are instead provided with author citations. Names reported exactly as mentioned in the original papers are reported in quotation marks.

Results

The dendrogram (Fig. 3) and further NMS (Fig. 4) showed four main clusters of relevés corresponding to the plant communities dominated by (Fe) *Festuca valesiaca* and *Poa bulbosa*, (As) *Asphodelus ramosus*, (He) *Helichrysum italicum*, and (Sa) *Salvia officinalis* and *Helichrysum italicum*.




Fig. 3. Hierarchical clustering (Ward's method, Chord distance) shows four main groups of relevés corresponding to the plant communities: As – *Hedypnoido rhagadioloidis-Asphodeletum ramosi*, He – *Helichrysetum italicum*, Sa – *Helichrysetum italicum salvietosum officinalis*, “sal” indicates a group of relevés clearly dominated by *Salvia officinalis*, Fe – *Festuco valesiaca-Poetum bulbosae*.

The cluster Fe groups together relevés from the summit plateau of the island, at more than 150 m a.s.l., where Horvatić (1927) had previously described the association *Festucetum valesiaca*. The mean number of taxa is 33 whereas in all the relevés of this cluster 53 taxa were found. In addition to *F. valesiaca* and *P. bulbosa* many species were recorded with high frequency (see Tab. 1).




Fig. 4. Non-metric multidimensional scaling of relevés in the dataset. Abbreviations: As – *Hedypnoido rhagadioloidis-Asphodeletum ramosi*, He – *Helichrysetum italicum*, Sa – *Helichrysetum italicum salvietosum officinalis*, “sal” indicates a group of relevés clearly dominated by *Salvia officinalis*, Fe – *Festuco valesiaca-Poetum bulbosae*. Ds – disturbance severity, Df – disturbance frequency, Gr – grazing pressure, T – temperature, R – soil reaction, st – stoniness, sl – slope.

Tab. 1. *Festuco valesiaca-Poetum bulbosae* ass. nov. on the island of Plavnik. Life-forms (LF): Ch – chamaephytes, G – geophytes, H – hemicryptophytes, P – phanerophytes, T – therophytes. Asterisk (*) indicates holotypus (rel. 1).

Rel. number	LF	1*	2	3	4	5	6
Exposition (°)		0	0	0	0	0	0
Slope (%)		0	0	0	0	0	0
Rockiness (%)		30	40	30	60	40	20
Stoniness (%)		10	30	10	10	20	30
Vegetation cover (%)		70	60	65	50	60	70
<i>Festuco valesiaca-Poetum bulbosae</i>							
<i>Poa bulbosa</i>	H	6	6	6	5	6	4
<i>Festuca valesiaca</i>	H	5	6	6	5	6	6
<i>Poetalia bulbosae, Poetea bulbosae</i>							
<i>Trifolium suffocatum</i>	T	6	4	4	4	3	3
<i>Plantago lanceolata</i>	H	3	2	2	2	2	3
<i>Erodium cicutarium</i>	T	2	2	2	2	2	2
<i>Trifolium subterraneum</i>	T	3	2	.	2	3	2
<i>Trifolium scabrum</i>	T	2	2	2	2	.	2
<i>Stipo-Trachynietea distachyae</i>							
<i>Trifolium campestre</i>	T	4	4	5	5	2	5
<i>Filago germanica</i>	T	4	4	3	3	4	2
<i>Galium murale</i>	T	3	3	3	2	3	2
<i>Sherardia arvensis</i>	T	3	2	3	2	2	3
<i>Gastridium ventricosum</i>	T	2	2	3	2	2	3
<i>Sideritis romana</i>	T	2	2	2	3	3	2
<i>Catapodium rigidum</i>	T	2	2	2	2	2	2
<i>Ornithogalum collinum</i>	G	2	2	.	2	.	.
<i>Ononis reclinata</i>	T	2	.	.	2	.	.

Tab. 1. continued

Rel. number	LF	1*	2	3	4	5	6
Tuberarietea guttatae							
<i>Crepis neglecta</i> subsp. <i>neglecta</i>	T	2	3	2	2	2	2
<i>Vulpia ciliata</i>	T	2	2	3	3	2	2
<i>Medicago minima</i>	T	3	2	2	2	2	2
<i>Galium divaricatum</i>	T	.	2	3	2	2	2
<i>Aira elegantissima</i>	T	2	2	2	2	.	2
<i>Asterolinon linum-stellatum</i>	T	.	2	.	2	2	2
<i>Plantago bellardii</i>	T	.	.	3	3	2	.
<i>Minuartia hybrida</i>	T	4	2	.	2	.	.
<i>Lotus hispidus</i>	T	.	2	.	.	2	.
Festuco-Brometea							
<i>Koeleria splendens</i>	H	3	4	2	2	3	3
<i>Thymus longicaulis</i>	Ch	2	2	3	.	2	3
<i>Teucrium capitatum</i> subsp. <i>capitatum</i>	Ch	2	2	2	.	2	2
<i>Carduus nutans</i> subsp. <i>micropterus</i>	H	.	.	.	2	.	2
<i>Eryngium amethystinum</i>	H	.	.	.	2	.	2
<i>Salvia pratensis</i>	H	2	2
Other taxa							
<i>Petrorhagia saxifraga</i>	H	3	2	2	2	2	2
<i>Anagallis arvensis</i>	T	2	.	2	2	2	2
<i>Scorzonerooides autumnalis</i>	H	2	2	2	.	.	2
<i>Cerastium pumilum</i> subsp. <i>glutinosum</i>	T	2	2	2	2	.	.
<i>Torilis nodosa</i>	T	2	2	.	.	2	.
<i>Geranium molle</i>	T	2	.	2	.	2	.
<i>Marrubium incanum</i>	H	.	.	2	2	2	.
<i>Medicago monspeliaca</i>	T	.	.	2	.	2	.
<i>Sedum acre</i>	Ch	2	.	2	.	.	.
<i>Euphorbia helioscopia</i>	T	2	.	.	.	2	.

Date, coordinates and sporadic species (with 2) of the relevés: Rel. 1 (holotypus): 28.V.2023, lat. 44° 58' 41", long. 14° 31' 27", *Arenaria serpyllifolia*, *Linum tenuifolium*; Rel. 2: 28.V.2023, lat. 44° 58' 42.5", long. 14° 31' 23", *Asparagus acutifolius*, *Brachypodium sylvaticum*, *Euphorbia exigua*, *Veronica arvensis*; Rel. 3: 28.V.2023, lat. 44° 58' 41", long. 14° 31' 21", *Euphorbia cyparissias*, *Helichrysum italicum*; Rel. 4: 28.V.2023, lat. 44° 58' 39", long. 14° 31' 23", *Valantia muralis*, *Achnatherum bromoides*, Rel. 5: 28.V.2023, lat. 44° 58' 38", long. 14° 31' 27.5", *Ficaria verna*; Rel. 6: 28.V.2023, lat. 44° 58' 35", long. 14° 31' 32", *Carex halleriana*.

The indicator species associated to this cluster are *Trifolium suffocatum*, *Erodium cicutarium*, *Crepis neglecta* subsp. *neglecta*, *Petrorhagia saxifraga*, and *Marrubium incanum*.

The second partitioning level separates the cluster As. This cluster represents the vegetation clearly dominated by *Asphodelus ramosus*, the latter being always recorded with a cover greater than 75%, except in one relevé. This vegetation type is well represented in the southern part of the island. The indicator taxa of this cluster are *Asphodelus ramosus*, *Hedypnois rhagadioloides*, *Crepis zacintha*, and *Centaureum maritimum*. The mean number of taxa per relevé is 32.

The third cluster He represents the most widespread type on the island: open vegetation dominated by *Helichrysum italicum* and *Koeleria splendens*, and with other taxa typical of the *Festuco-Brometea* class occurring with high frequency (e.g. *Bromopsis erecta*, *Euphorbia cyparissias*, Tab. 2).

The indicator species of this cluster are *Reichardia picroides*, *Bupleurum veronense*, *Plantago holostium*, and *Hippocrepis comosa*. This vegetation type usually develops on rocky and stony ground whereas on steep slopes on the eastern part of the island, in some places, it gives way to a *Salvia officinalis*-*Helichrysum italicum* community, represented by the cluster Sa. The latter corresponds to the "*Helichrysetum salviosum*" described by Horvatić (1927) and includes seven relevés, with a mean number of taxa of 21. This cluster is actually heterogeneous with some relevés which represent a variant of the *Helichrysetum italicum* due to the presence and co-dominance of both *Helichrysum italicum* and *Salvia officinalis*, whereas two relevés (indicated as "sal" in Figs. 3 and 4) are clearly dominated by the latter. *Salvia officinalis* is the only indicator species of this cluster.

The NMS (Fig. 4) resulted in a two-axis solution, with a final stress 8.96, indicating a fairly good ordination.

Tab. 2. *Helichrysetum italici typicum* (rels. 1-10) and *Helichrysetum italici salvietosum officinalis* (rels. 11-14); transition to *Stipo-Salvietum officinalis* (rels. 15-16) on the island of Plavnik. For abbreviations of Life-forms [LF], see caption of Tab. 1. Asterisk (*) indicates neotypus (rel. 3), and holotypus (rel. 13), respectively.

Rel. number	LF	1	2	3*	4	5	6	7	8	9	10	11	12	13*	14	15	16
Exposition (°)		20	50	0	130	230	50	140	0	160	70	160	150	270	50	270	60
Slope (%)		5	10	10	2	3	10	2	0	2	2	15	20	10	20	30	25
Rockiness (%)		30	40	40	10	70	30	20	10	40	40	40	40	30	30	30	30
Stoniness (%)		40	50	30	40	20	70	20	40	40	40	70	40	40	50	70	70
Vegetation cover (%)		60	60	70	60	60	70	60	70	70	75	50	55	60	60	70	60
<i>Helichrysetum italici</i>																	
<i>Helichrysum italicum</i>	Ch	7	7	8	6	6	7	7	8	7	7	6	6	6	5	2	2
<i>Salvia officinalis</i>	Ch	6	5	4	7	7	8	8
<i>Chrysopogono grylli-Koelerion splendidis, Scorzoneretalia villosae</i>																	
<i>Bromopsis condensata</i> (+ <i>Br. erecta</i>)	H	4	4	3	2	4	3	2	4	3	5	2	2	5	3	4	3
<i>Koeleria splendens</i>	H	5	5	5	4	5	4	3	4	5	4	.	.	2	2	.	.
<i>Festuca valesiaca</i>	H	.	.	2	2	.	.	2	2	4	3	.	.	3	2	.	.
<i>Bupleurum veronense</i>	T	2	2	2	2	2	2
<i>Carduus nutans</i> subsp. <i>micropterus</i>	H	2	2	2	2	.
<i>Centaurea tommasinii</i>	H	2	2	2	2
<i>Teucrium montanum</i>	Ch	2	2	2	.	2	.	.	.
<i>Aethionema saxatile</i>	Ch	2	2	2	.	.	.
<i>Medicago prostrata</i>	H	.	.	2	2	2
<i>Chrysopogon gryllus</i>	H	2	2
<i>Eryngium amethystinum</i>	H	2	.	.	.	2	.	.
<i>Genista sylvestris</i> subsp. <i>dalmatica</i>	Ch	2
<i>Stipa eriocalis</i>	H	2
<i>Festuco-Brometea</i>																	
<i>Euphorbia cyparissias</i>	H	4	3	2	2	3	5	2	.	2	2
<i>Linum tenuifolium</i>	Ch	2	2	.	2	2	2	.	.	2	2	2	.	2	.	.	.
<i>Melica ciliata</i>	H	.	.	2	.	.	.	2	2	2	2	2	2	.	.	2	2
<i>Salvia pratensis</i>	H	2	.	3	2	3	2	2	2	.	2
<i>Sanguisorba minor</i>	H	2	2	2	.	2	3	2	2	.	2
<i>Sedum acre</i>	Ch	2	.	2	2	2	2
<i>Thymus longicaulis</i>	Ch	2	.	2	2	2	.	.	2
<i>Convolvulus cantabrica</i>	H	.	.	.	2	.	2	2	2
<i>Hippocrepis comosa</i>	H	2	2	.	2	.	2
<i>Plantago holosteum</i>	H	4	4	4	.	.	4
<i>Achnatherum bromoides</i>	H	2	2
<i>Carex halleriana</i>	H	2	4	.	.	.
<i>Centaureum erythraea</i>	H	.	.	2	2
<i>Dorycnium pentaphyllum</i> subsp. <i>herbaceum</i>	H	2
<i>Galium corrudifolium</i>	H	.	2
<i>Knautia purpurea</i>	H	2
<i>Pilosella piloselloides</i>	H	2	.	.	.
<i>Teucrium chamaedrys</i>	Ch	2
<i>Stipo-Trachynietea distachyae</i>																	
<i>Catapodium rigidum</i>	T	3	2	2	2	2	3	2	3	3	2	2	2	2	3	.	.
<i>Trifolium campestre</i>	T	2	2	3	3	3	2	4	4	5	3	2	3	2	2	.	.
<i>Trachynia distachya</i>	T	3	2	3	3	3	2	2	2	3	2	4	3	2	.	.	.

Tab. 2. continued

Rel. number	LF	1	2	3*	4	5	6	7	8	9	10	11	12	13*	14	15	16
<i>Euphorbia exigua</i>	T	2	2	2	2	2	.	2	2	2	2	3	3	2	.	.	.
<i>Gastridium ventricosum</i>	T	2	2	2	2	2	2	3	2	2	2	.	2	2	.	.	.
<i>Sideritis romana</i>	T	2	.	2	5	2	2	2	2	2	2	2	2	.	2	.	.
<i>Medicago minima</i>	T	2	2	2	3	2	.	3	3	3	2	2	3
<i>Valantia muralis</i>	T	3	3	2	3	2	4	2	.	2	.	2	2	.	2	.	.
<i>Trifolium scabrum</i>	T	2	2	2	3	.	.	3	3	3	2	.	2	2	.	.	.
<i>Filago germanica</i>	T	.	.	2	2	2	.	2	2	2	.
<i>Sherardia arvensis</i>	T	.	.	2	2	.	.	2	3	2	2	.	.
<i>Asterolinon linum-stellatum</i>	T	.	2	2	2	2	.	.	2
<i>Ononis reclinata</i>	T	.	.	2	2	2	.	.	.	2	2
<i>Plantago bellardii</i>	T	.	.	2	3	3	.	.	2
<i>Crepis neglecta</i> subsp. <i>neglecta</i>	T	.	.	2	2	2	.
<i>Galium murale</i>	T	2	2	.	2
<i>Ornithogalum collinum</i>	G	2	2	2
<i>Valerianella eriocarpa</i>	T	.	.	2	2	.	.	2
Other taxa																	
<i>Dactylis glomerata</i> subsp. <i>hispanica</i>	H	2	2	2	2	.	3	2	3	2	3	.	3	2	2	3	.
<i>Linum trigynum</i>	T	3	2	2	2	2	2	2	2	2	3	2	.	3	2	.	.
<i>Anagallis arvensis</i>	T	3	2	2	2	2	2	2	2	2	2	2	2
<i>Carlina corymbosa</i>	H	2	.	2	2	2	2	.	2	2	2	.	.	2	.	.	.
<i>Reichardia picroides</i>	H	2	2	2	2	2	2	2	.	.	2	.	.	2	.	.	.
<i>Juniperus oxycedrus</i>	P	2	2	2	2	2	2	.	.	.	4	2	.
<i>Plantago lanceolata</i>	H	3	2	2	2	2	2	2	.	2
<i>Sonchus bulbosus</i> subsp. <i>bulbosus</i>	G	3	4	2	2	.	2	2	2	2	.
<i>Teucrium capitatum</i> subsp. <i>capitatum</i>	Ch	2	.	2	.	2	.	2	2	2	2	.	2
<i>Geranium purpureum</i>	T	.	.	2	.	.	.	2	2	2	2	.	.	2	2	.	.
<i>Orobanche minor</i>	T	.	.	.	2	2	.	2	2	2	.	.	.	2	.	.	.
<i>Torilis nodosa</i>	T	2	2	2	2	.	2	.	2	.	.
<i>Petrorhagia saxifraga</i>	H	.	.	.	2	.	.	2	2	2	.	.	2
<i>Poa bulbosa</i>	H	.	.	2	2	.	.	3	3	2
<i>Scorpiurus muricatus</i>	T	.	2	2	.	3	3	2
<i>Aira elegantissima</i>	T	.	.	2	.	.	.	2	2
<i>Allium</i> sp.	G	2	2	.	.	2	.	.
<i>Asparagus acutifolius</i>	P	2	.	2	.	.	.	2	.	.	.
<i>Cerastium pumilum</i> subsp. <i>glutinatum</i>	T	2	.	.	.	2	.	2	.	.
<i>Galium divaricatum</i>	T	.	.	.	2	.	.	.	2	2
<i>Lotus corniculatus</i>	H	2	2	2

Dates, coordinates and sporadic species (with 2): Rel. 1: 25.V.2023, lat. 44° 58' 28", long. 14° 32' 25", *Euphorbia peplus*, *Thesium divaricatum*; Rel. 2: 25.V.2023, lat. 44° 58' 13", long. 14° 32' 36", *Thesium divaricatum*, *Anthyllis vulneraria* subsp. *rubriflora*; Rel. 3 (neotypus): 25.V.2023, lat. 44° 58' 00", long. 14° 32' 42", *Asphodelus ramosus*, *Hedypnois rhagadioloides*, *Leontodon crispus*, *Pilosella officinarum*, *Convolvulus althaeoides* subsp. *tenuissimus*; Rel. 4: 25.V.2023, lat. 44° 57' 54", long. 14° 32' 47", *Asphodelus ramosus*, *Crepis zacintha*, *Hedypnois rhagadioloides*, *Leontodon tuberosus*, *Crepis foetida*; Rel. 5: 25.V.2023, lat. 44° 57' 50", long. 14° 32' 56", *Pilosella officinarum*; Rel. 6: 25.V.2023, lat. 44° 57' 55", long. 14° 32' 54", *Lotus hispidus*, *Rubus ulmifolius*; Rel. 7: 27.V.2023, lat. 44° 58' 04", long. 14° 31' 30", *Clematis vitalba*, *Vulpia ciliata*, *Oxalis corniculata*, *Quercus ilex*, *Stachys officinalis*; Rel. 8: 27.V.2023, lat. 44° 58' 09.5", long. 14° 31' 15", *Clematis vitalba*, *Cynosurus echinatus*, *Euphorbia peplus*, *Geranium molle*, *Minuartia hybrida*, *Vulpia ciliata*, *Trifolium suffocatum*, *Ficaria verna*; Rel. 9: 27.V.2023, lat. 44° 58' 13.5", long. 14° 31' 02", *Paliurus spina-christi*, *Scorzoneroideis autumnalis*; Rel. 10: 28.V.2023, lat. 44° 58' 49", long. 14° 31' 42", *Crepis zacintha*, *Minuartia hybrida*, *Paliurus spina-christi*, *Sonchus asper*, *Carthamus lanatus*; Rel. 11: 26.V.2023, lat. 44° 57' 27.5", long. 14° 31' 59", *Urospermum picroides*; Rel. 12: 26.V.2023, lat. 44° 57' 28", long. 14° 31' 55", *Cynosurus echinatus*; Rel. 13 (holotypus): 27.V.2023, lat. 44° 58' 11", long. 14° 31' 13", Rel. 14: 28.V.2023, lat. 44° 58' 42", long. 14° 31' 38", *Geranium molle*; Rel. 15: 27.V.2023, lat. 44° 58' 06.5", long. 14° 31' 07", *Sesleria autumnalis*, *Sonchus asper*, *Myosotis arvensis*, *Phillyrea latifolia*; Rel. 16: 28.V.2023, lat. 44° 58' 39", long. 14° 31' 39", *Clematis flammula*.

The two axes accounted for 79.5 of the total variation (axis 1 = 65.3%, axis 2 = 14.2%). The four clusters occupy different parts of the ordination space without overlapping each other. Axis 1 separates the clusters He and Sa among them and from Fe and As. This axis is positively correlated with Df, Ds, Gr and temperature whereas it is negatively correlated with slope, stoniness and reaction. The correlation strengths of the EIVs of moisture and nutrient content with the ordination axes were lower than the selected r^2 cut-off and we will not comment on those variables further. Therefore, the Sa cluster groups relevés taken on steeper slopes than the other clusters and with high percentages of stones. The Fe and As clusters have the highest values of disturbance, in terms of both severity and frequency. The lowest reaction values for these two clusters are probably related to disturbance. Axis 2 clearly separates Fe from As, the former with greater values of grazing, the second with greater values of temperature, being in the southern part of the island at the lowest altitude and mostly on the south-facing side.

The NMS ordination of the relevés of the As cluster together with those of the associations *Bromo erecti-Chrysopogonietum grylli* and *Narcisso tazettae-Asphodeletum microcarpi* resulted in a three-axis solution with a final stress of 7.5 (Fig. 5).

Most of the variance is represented by axis 1 (55.1 %) whereas axes 2 and 3 account for 22.6 % and 2.5 %, respectively. The relevés of the two associations *Bromo erecti-Chrysopogonietum grylli* (BC1 and BC2 in Fig. 5) and *Narcisso tazettae-Asphodeletum microcarpi* (NA1 and NA2) are separated on axis 1, whereas the relevés of the cluster As are in the middle. The latter are clearly separated from the others on axis 3 (Fig. 5). Interestingly, the relevés of the two phytosociological tables of the *Bromo erecti-Chrysopogonietum*

grylli (i.e. BC1 and BC2) and those of the two tables of the *Narcisso tazettae-Asphodeletum microcarpi* (i.e. NA1 and NA2) form four well separated groups of relevés.

The cluster As was found to be well distinguished in the species-space from the other associations with which it has no overlap. The indicator species analysis indicated the following 11 taxa as diagnostic for the As cluster (indicator values in brackets): *Hedypnois rhagadioloides* (74), *Linum trigynum* (72.3), *Galium murale* (70.6), *Euphorbia cyparissias* (68.9), *Geranium purpureum* (67.2), *Sideritis romana* (62.4), *Vulpia ciliata* (60.8), *Centaurium maritimum* (56.3), *Plantago bellardii* (56.3), *Scorpiurus muricatus* (56.3), and *Lotus hispidus* (42.3).

Discussion

A direct quantitative comparison of Horvatić's data with the new data is not completely possible because the plot sizes used in the two surveys are too different (see also Dengler et al. 2009). For example, the five plots used by Horvatić (1927) to represent the *Festucetum valesiacae* sample a total area of 5 m² and include 17 taxa. The new dataset comprises six relevés of the same area – each with a plot size of 25 m² and containing an average of 33 taxa – for a total number of taxa of 53. Furthermore, the quantitative values of the taxa in Horvatić's article are estimated on a scale comprising at least 10 (undefined) values (-, -1, 1, 1-2, 2, 2-3, 3, 3-4, 4, 5), which are not directly comparable with the 9-points scale used in this paper. However, on the basis of qualitative considerations, it can be deduced whether the plant communities remained fairly similar or not during the 100 years.

As written above, although Horvatić (1927) described five grassland syntaxa on the island of Plavnik, we were on-




Fig. 5. Non-metric multidimensional scaling (with axes 1, 2 and 3) of eastern Adriatic associations with *Asphodelus ramosus*. Abbreviations: HA – *Hedypnois rhagadioloides-Asphodeletum ramosi*, BC – *Bromo erecti-Chrysopogonietum grylli* (BC1 from island of Pag, Croatia, and BC2 from Montenegro), NA – *Narcisso tazettae-Asphodeletum microcarpi* (NA1 from Istria and NA2 from islands of Bobara and Mrkan, Croatia).

ly able to find four grassland community types. In fact, the area originally covered by *Andropogonetum grylli* is completely occupied by a plant community dominated by *Asphodelus ramosus*, whereas *Chrysopogon gryllus* – i.e. the dominant taxon of the *Andropogonetum grylli* – is quite rare on the island.

Festuco valesiaca-*Poetum bulbosae*

The *Festucetum valesiaca* described by Horvatić (1927) for the continuous herbaceous expanses on the summit of the island is no longer present. This association was dominated by *Festuca valesiaca*, followed as cover by *Bromopsis erecta* and *Plantago holostium*, and included several other species of the class *Festuco-Brometea* class (e.g. *Thymus longicaulis*, *Chrysopogon gryllus*, *Koeleria splendens*), to which it clearly belonged. In that area, the physiognomic situation appears quite different nowadays, with herbaceous phytocoenoses interspersed with shrubs and trees of *Juniperus oxycedrus*, generally on flat surfaces, where sheep and fallow deer find shelter (Fig. 2D). The herbaceous vegetation is dominated by *Poa bulbosa* and *F. valesiaca*, followed by some *Trifolium* species (*T. campestre*, *T. suffocatum*, *T. subterraneum*, *T. scabrum*), and *Koeleria splendens*. The results show that this plant community is clearly distinct from the others in terms of floristic composition and ecology. Indeed, it is the most disturbed in terms of intensity and frequency of disturbance (Ds and Df in Fig. 3) together with *Asphodeletum ramosi* but, unlike the latter, it has a higher grazing pressure (Gr). Therefore, the plant community dominated by *Poa bulbosa* and *Festuca valesiaca* (Fe in Fig. 3) is here considered to be a new association, the *Festuco valesiaca*-*Poetum bulbosae* *ass. nov. hoc loco* (holotypus rel. 1 in Tab. 1), possibly derived from *Festucetum valesiaca* – of which *Festuca valesiaca*, *Koeleria splendens*, and other taxa are still present – as a consequence of the prolonged stay of sheep and deer and intensive grazing.

The presence with high cover and/or frequency of *Poa bulbosa*, *Trifolium suffocatum*, *T. subterraneum*, *Erodium cicutarium*, *Plantago lanceolata*, and other species, indicates that the *Festuco valesiaca*-*Poetum bulbosae* belongs to the *Poetea bulbosae*, i.e. a class representing heavily grazed pastures, which are mainly present in the western Mediterranean basin (e.g. Rivas Goday and Rivas-Martínez 1963). This class comprises a single order, *Poetalia bulbosae*, and several alliances, none of which occur in Croatia (Preislerová et al. 2022), with the exception of the *Romuleion* (Škvorc et al. 2017). This alliance was originally invalidly described by Oberdorfer (1954, see Terzi et al. 2024b) for the south and southeast of the Balkan Peninsula, and was identified by 20 character species, none of which are present in our relevés (at least in the phenological stage of vegetation at the time of sampling). The alliance was originally classified in the class “*Thero-Brachypodietea*” (Oberdorfer 1954), and then moved to the “*Brachypodio-Chrysopogonetea*” (Horvat et al. 1974), “*Saginetea maritimae*” (Rodwell et al. 2002), and *Tuberarietea guttatae* (Čarni et al. 2014). The EVC has instead classified the *Romuleion* in the *Poetea*

bulbosae (EVC) to represent “Macedonian seasonal perennial pastures on acidic substrates”. The *Poetea bulbosae*, indeed, are most frequently found on acid substrates (Oberdorfer 1954, Čarni et al. 2014).

The numerous taxa of the *Stipo-Trachynietea distachyae* found in the *Festuco valesiaca*-*Poetum bulbosae* are linked to the basic substratum of the island of Plavnik, while those of the *Tuberarietea guttatae* highlight the dynamic relationships with the oligotrophic grasslands of this class, where the *Romuleion* was framed by some authors (e.g. Čarni et al. 2014). Two associations of the *Poetalia bulbosae* were recently described near Podgorica, Montenegro, on fluvio-glacial deposits and calcareous rocky substrate and framed in the *Romuleion* (Stanišić-Vujačić et al. 2023). The floristic composition of those associations, however, includes only three character species of this alliance (*Romulea linaresii* subsp. *graeca*, and the very rare *Hypochaeris cretensis* and *Hedypnois rhagadioloides*). The associations from Montenegro were found in an area where rocky grasslands of the alliance *Chrysopogono grylli-Koelerion splendentis* are widespread and indeed at least one of those associations (*Romuleo bulbocodii-Poetum bulbosae* Stanišić-Vujačić et al. 2023) is characterized by the presence of numerous *Festuco-Brometea* species (Stanišić-Vujačić et al. 2022, 2023). The *Festuco valesiaca*-*Poetum bulbosae*, derived from the *Festucetum valesiaca*, also has dynamic relationships with that alliance and in fact shares several species (e.g., *Koeleria splendens*, *Thymus longicaulis*) with the Montenegrin associations. However, the natural potential vegetation of the Plavnik area is different from that of the surrounding of Podgorica, in Montenegro.

The floristic relationships of the *Festuco valesiaca*-*Poetum bulbosae* with the *Romuleion* are, however, even weaker, given that none of the character species of this alliance were detected in our relevés. For this reason, the *Festuco valesiaca*-*Poetum bulbosae* is only provisionally classified in the *Romuleion*, pending the expansion of knowledge about the variability of the *Poetalia bulbosae* in the Balkans by new phytosociological studies. It is indeed plausible to admit that a second alliance of the *Poetalia bulbosae*, in addition to the *Romuleion*, is present in the Western Balkans and that it is dynamically linked to a degradation of the grasslands of the *Scorzoneretalia villosae* (not present in Greece, where the *Romuleion* was originally described), but a large-scale comparison is needed to confirm this hypothesis, which is beyond the scope of this paper. In any case, the *Festuco valesiaca*-*Poetum bulbosae* documents for the first time, based on phytosociological data, the presence of *Poetalia bulbosae* and *Poetea bulbosae* in Croatia.

Helichrysetum italicum

The most frequent vegetation type on the island is represented by the *Helichrysum italicum* dominated plant community (“He” in Figs. 3 and 4), which grows on rocky and stony substrates throughout the island. *Helichrysum italicum* is widespread in southern Europe with several subspecies and the nominal one, *Helichrysum italicum* subsp. *italicum*,

which is the only one found on the island of Plavnik, has a range in Europe that includes southern France, Italy and the Balkans (Euro+Med 2023). This subspecies grows in different types of vegetation, such as thermophilic scree communities of the *Scrophulario-Helichrysetalia* (Brullo and Spampinato 1991), and coastal sub-aerohaline dwarf scrub vegetation of the *Helichrysetalia italici* (Foucault 2020). In the Balkans, it has been considered a character taxon of the “*Brachypodio-Chrysopogonetea*” (Horvatić 1963, Horvat et al 1974). *Helichrysum italicum* is a suffruticose chamaephyte and, where it is dominant, the vegetation physiognomy takes on the appearance of a chamaephytic garrigue (Horvatić 1927). In the Dinaric Arc mountain range, this type of vegetation is often rich in many species typical of grasslands, so that it is not always possible to distinguish chamaephytic garrigues from ‘pure’ grasslands from a syntaxonomic point of view. In fact, the alliance *Chrysopogono grylli-Koelerion splendidis*, of the order *Scorzoneretalia villosae*, includes vegetation types dominated by both hemicyptophytes and dwarf chamaephytes, such as *Helichrysum italicum* and *Salvia officinalis*.

The same situation can be observed on the island of Plavnik where the *Helichrysetum italici* includes *Helichrysum italicum*, as the dominant, together with many grassland taxa, such as *Bromopsis condensata*, *Trifolium campestre*, *Dactylis glomerata* subsp. *hispanica*, *Salvia pratensis*, *Euphorbia cyparissias*, *Koeleria splendens*, *Hippocrepis comosa*. This plant community grows on incoherent substrata with a floristic composition that resembles that sampled by Horvatić (1927) a hundred years ago, except for the absence of *Chrysopogon gryllus*, which has evidently become rather rare on the island (Fig. 2A). Although the total number of species found in the new relevés is far higher than that recorded by Horvatić (1927), this difference is almost certainly due to the different sizes of the plots used in the two sampling surveys. The species that shape the physiognomy of the *Helichrysetum italici* are *Helichrysum italicum*, *Bromopsis condensata*, *Koeleria splendens*, and *Trifolium campestre*. The association clearly belongs to the *Chrysopogono grylli-Koelerion splendidis* due to the presence of characteristic species of the class *Festuco-Brometea* (e.g. *Bromopsis condensata*, *Euphorbia cyparissias*, *Linum tenuifolium*, *Festuca valesiaca*, *Hippocrepis comosa*), of the order *Scorzoneretalia villosae* and of the alliance (e.g. *Thymus longicaulis*, *Centaurea tommasinii*, *Plantago holosteum*, *Bupleurum veronense*) (see Terzi 2015). Although *Helichrysum italicum* is quite common in other associations of this alliance, the specific floristic composition of the *Helichrysetum italici* is different from that of all the other associations already described. In any case, even if this association was considered as a syntaxonomic synonym of another association belonging to the alliance, the name *Helichrysetum italici* would retain priority over names described later.

The *Helichrysetum italici* occurs in its typical aspect (subass. *typicum*, represented by rels. 1-9, in Tab. 2) on most of the island, and only on some spots does it associate with *Salvia officinalis* to give rise to the subassociation described

in the next section. The typical aspect corresponds to that described by Horvatić (1927) with 10 plots of 1 square meter each. The size of these individual plots thus appears to be much smaller than that normally used to sample the plant communities of the *Scorzoneretalia villosae*, which varies in 90% of cases between 10 and 150 square meters (Terzi 2015). Because of the small area of the plots sampled by Horvatić (1927), and perhaps also because of the omission of some species, each of those plots counts only a very limited number of taxa. Moreover, the quantitative values associated with these taxa are on a scale hardly comparable with those routinely used in modern phytosociological studies.

Although none of the 10 relevés originally published by Horvatić (1927), if taken alone, can be considered a representative relevé of the *Helichrysetum italici* for the reasons expressed above, the association is nevertheless validly described because at least the synoptic column of the association, i.e. the last column of table IV (“*Skrižaljka IV*”, on p. 33, in Horvatić 1927), constitutes a sufficient original diagnosis according to Article 7 of the International Code of Phytosociological Nomenclature (ICPN, Theurillat et al. 2021).

Therefore, we do not designate a lectotype, which would be incomplete and unrepresentative of the typical aspect of the association. Thus, not having a suitable relevé of the original diagnosis to be selected as lectotype of the association (see also Def. VIII of the ICPN), we find it more correct and useful to designate the relevé 3 in Tab. 2 as neotypus hoc loco of the *Helichrysetum italici* and of its subassociation *typicum*.

Helichrysetum italici salvietosum officinalis

Along some steep, mainly north-facing slopes on incoherent substrates or on limestone outcrops, the typical aspect of the *Helichrysetum italici* gives way to another type of vegetation where *Salvia officinalis* becomes dominant or co-dominant with *Helichrysum italicum* (Fig. 2B). This aspect was erroneously described by Horvatić (1927) as “*Helichrysetum salviosum*” (Art. 3e of the ICPN, Theurillat et al. 2021), to indicate a subtype of the *Helichrysetum italici*. Its floristic composition indeed retains many species of the *Helichrysetum italici* with the addition of a few others, including the dominant *Salvia officinalis*, thus forming the subassociation *Helichrysetum italici salvietosum officinalis subass. nov. hoc loco* (holotypus rel. 13 of Tab. 2). This subassociation has fewer species than the typical one.

The grassland vegetation of the Western Balkans dominated by *Salvia officinalis* belongs to different vegetation types, mainly framed in the alliances *Chrysopogono grylli-Koelerion splendidis* and *Saturejion subspicatae* (Terzi et al. 2023). In our case, the *Helichrysetum italici salvietosum officinalis* definitely belongs to the first alliance and is perhaps transitional to the *Stipo-Salvietum officinalis*, whose nomenclatural type was recorded on the island of Unije, also located in Kvarner, about 50 km (as the crow flies) from Plavnik. In particular, two relevés could belong to this association (subcluster ‘sal’ in Fig. 3), which show the lowest

number of taxa, the clear dominance of *Salvia officinalis*, and the presence of *Helichrysum italicum* with very low coverage values.

Hedypnoido rhagadioloidis-Asphodeletum ramosi

The vegetation with *Asphodelus ramosus* develops mainly in the southern part of the island, with large and surprising expanses absolutely dominated by this species, with a cover usually greater than 90% (Fig. 2C). Two relevés recorded in the vicinity of the *Helichrysetum italicum* distribution area represent a transition towards this association and indeed there *Helichrysum italicum* is the second dominant species with a cover of 25-50%. A very different situation is described by Horvatić (1927) who reports local patches of *Asphodeletum microcarpi* interspersed in a matrix of other associations (*Andropogonetum grylli* and *Helichrysetum italicum*) and sharing several species with them. He notes that this sharing of species between the *Asphodeletum microcarpi* and the other associations is even more evident during the winter, when *Asphodelus ramosus*, the main dominant species, survives belowground. In the plots sampled by Horvatić (1927), *Asphodelus ramosus* was recorded with a score of “2” (or “3” in only one case) together with other species with

similar or greater cover, so that some of its plots look like local variants of the *Andropogonetum grylli*.

On the other hand, as written above, during our surveys on the island we found few plants of *Chrysopogon gryllus* for which it was not possible to identify the *Andropogonetum grylli* nor transitions between this association and the *Asphodeletum microcarpi*. In the most typical situation, *Asphodelus ramosus* is by far the dominant species, and with the exception of the cases representing a transition with the *Helichrysetum italicum* (rels. 5 and 6 in Tab. 3), in all other relevés the second dominant species never scored higher than 2a (i.e. up to 12% cover).

This plant community comprises numerous annual taxa of the classes *Stipo-Trachynietea distachyae* and *Tuberarietea guttatae*. The presence of so many annual plants is not a novelty for this kind of vegetation and has been observed in many other areas of the Mediterranean (see Terzi 2023 and references therein). The plant community dominated by *Asphodelus ramosus* was found to be the most disturbed grassland type on the island, together with the *Festuco valesiacae-Poetum bulbosae*. The thriving of *Asphodelus ramosus* is usually associated with intense grazing pressure and frequent fires. These factors might also be the ecologi-

Tab. 3. *Hedypnoido rhagadioloidis-Asphodeletum ramosi* ass. nov. on the island of Plavnik. For abbreviations of Life-forms [LF], see caption of Tab. 1. Asterisk (*) indicates holotypus (rel. 2).

Rel. number	LF	1	2*	3	4	5	6	7
Exposition (°)		90	0	0	40	0	0	0
Slope (%)		5	0	0	2	0	0	0
Rockiness (%)		3	10	0	0	20	20	10
Stoniness (%)		3	20	5	0	40	30	10
Vegetation cover (%)		90	100	100	95	98	85	90
<i>Hedypnoido rhagadioloidis-Asphodeletum ramosi</i>								
<i>Asphodelus ramosus</i>	G	9	9	9	9	9	7	9
<i>Helichrysum italicum</i>	Ch	2	3	2	3	7	7	5
<i>Crepis zacintha</i>	T	.	4	3	3	2	.	2
<i>Hedypnois rhagadioloides</i>	T	2	2	2	2	2	2	.
<i>Centaurium maritimum</i>	T	.	.	2	3	.	2	2
<i>Stipo-Trachynietea distachyae</i>								
<i>Gastridium ventricosum</i>	T	2	3	4	3	4	2	3
<i>Trifolium campestre</i>	T	2	2	3	3	3	2	3
<i>Trifolium scabrum</i>	T	3	3	2	2	2	.	3
<i>Sherardia arvensis</i>	T	2	2	2	2	2	2	2
<i>Catapodium rigidum</i>	T	3	2	2	.	2	3	2
<i>Sideritis romana</i>	T	2	.	2	.	2	5	2
<i>Filago germanica</i>	T	2	2	2	3	.	.	2
<i>Galium murale</i>	T	2	2	3	2	.	.	2
<i>Trachynia distachya</i>	T	.	2	.	.	3	3	3
<i>Medicago minima</i>	T	2	.	.	.	2	3	2
<i>Euphorbia exigua</i>	T	.	.	2	2	2	.	2
<i>Trifolium subterraneum</i>	T	2	.	2	2	.	.	.
<i>Valantia muralis</i>	T	.	2	.	.	.	3	.

Tab. 3. continued

Rel. number	LF	1	2*	3	4	5	6	7
Tuberarietea guttatae								
<i>Linum trigynum</i>	T	2	3	3	2	2	2	2
<i>Galium divaricatum</i>	T	2	2	3	3	2	.	2
<i>Plantago bellardii</i>	T	2	2	2	2	2	.	2
<i>Vulpia ciliata</i>	T	5	2	4	3	.	.	2
<i>Scorpiurus muricatus</i>	T	2	2	2	.	2	2	.
<i>Aira elegantissima</i>	T	.	3	3	3	.	.	.
<i>Lotus hispidus</i>	T	4	2	3
Festuco-Brometea								
<i>Euphorbia cyparissias</i>	H	5	4	4	2	2	.	3
<i>Sanguisorba minor</i>	H	2	2	2	2	2	2	2
<i>Festuca valesiaca</i>	H	3	2	2	.	2	2	2
<i>Linum tenuifolium</i>	Ch	2	2	.	2	2	2	2
<i>Bromopsis condensata</i> (+ <i>Br. erecta</i>)	H	.	.	.	2	2	4	2
<i>Convolvulus cantabrica</i>	H	.	2	2	.	2	2	.
<i>Koeleria splendens</i>	H	.	2	.	.	3	2	.
<i>Salvia pratensis</i>	H	.	2	.	3	.	2	.
<i>Thymus longicaulis</i>	Ch	2	2	2
<i>Galium corrudifolium</i>	H	2	4	.
<i>Bupleurum veronense</i>	T	2	.
<i>Medicago prostrata</i>	H	2
<i>Chrysopogon gryllus</i>	H	2	.	.
<i>Melica ciliata</i>	H	2
Other taxa								
<i>Dactylis glomerata</i> subsp. <i>hispanica</i>	H	4	.	2	2	2	3	4
<i>Anagallis arvensis</i>	T	2	3	2	2	2	2	2
<i>Poa bulbosa</i>	H	3	.	2	2	2	2	2
<i>Geranium purpureum</i>	T	.	2	2	2	2	2	2
<i>Teucrium capitatum</i> subsp. <i>capitatum</i>	Ch	2	2	.	.	2	2	2
<i>Plantago lanceolata</i>	H	.	.	.	2	.	2	.
<i>Orobancha minor</i>	T	2	2

Date, coordinates and sporadic species (with 2, except where indicated) of the relevés: Rel. 1: 25.V.2023, lat. 44° 58' 02", long. 14° 32' 41", *Daucus carota*; Rel. 2 (holotypus): 26.V.2023, lat. 44° 57' 26", long. 14° 33' 01", *Carlina corymbosa*; Rel. 3: 26.V.2023, lat. 44° 57' 28.5", long. 14° 32' 59"; Rel. 4: 26.V.23, lat. 44° 57' 36", long. 14° 32' 58", *Scorzoneroides autumnalis* (3); Rel. 5: 26.V.2023, lat. 44° 57' 31", long. 14° 32' 51"; Rel. 6: 25.V.2023, lat. 44° 57' 51", long. 14° 32' 52", *Euphorbia peplus*, *Leontodon tuberosus*, *Ononis reclinata*, *Anisantha madritensis*, *Asterolinon linum-stellatum*, *Sedum acre*; Rel. 7: 26.V.2023; lat. 44° 57' 33", long. 14° 32' 42", *Urospermum dalechampii*.

cal drivers on the island of Plavnik, where we hypothesise that at least one of them (grazing or fires) has presumably occurred on the island with greater intensity and/or frequency than was the case 100 years ago. Increased grazing pressure is plausible due to the high number of sheep easily observed on that side of the island, while it is more difficult to assess the role of fires as data on their intensity and frequency are not available.

Thus, the present plant community dominated by *Asphodelus ramosus* on Plavnik Island differs both floristically and ecologically from the *Asphodeletum microcarpi* described by Horvatić (1927). It is also clearly different from the *Bromo erecti-Chrysopogonetum grylli* (i.e. *Asphodelo-Chrysopogonetum grylli* Horvatić 1963, cf. Terzi 2015) and *Narcisso tazettae-Asphodeletum microcarpi*. In this case, the Plavnik plant community differs in physiognomy, given the

clear dominance of *Asphodelus ramosus*, and floristic composition (Fig. 5). The indicator species that distinguish the plant community of Plavnik Island from those of other locations in the eastern Adriatic are mainly annual species, some of which indicate a slightly acidic to almost neutral substrate (e.g. *Centaureum maritimum*, *Linum trigynum*), as is to be expected in heavily grazed karst areas.

In view of the above, the plant community dominated by *Asphodelus ramosus* on the island of Plavnik is considered here a new association: *Hedypnoido rhagadioloidis-Asphodeletum ramosi* ass. nov. hoc loco (holotypus rel. 2 of Tab. 3). However, its classification at a higher syntaxonomic rank is rather difficult (see Terzi 2023 and references therein). Biondi et al. (2016) proposed a new class, order and alliance to classify the vegetation dominated by *Asphodelus ramosus* (and other geophytes) but they have not been ac-

cepted because a comprehensive syntaxonomic review on the subject is still lacking (Biurrun and Willner 2020, Terzi 2023). In Montenegro, *Asphodelus ramosus* is included within communities of the *Artemisietea vulgaris*, although it is most abundant in the *Bromo erecti-Chrysopogonetum grylli* (Stanišić-Vujačić et al. 2022). As a result of this as yet undefined syntaxonomic framework, we refrain from classifying the new association into one of the accepted classes.

With the exception of *Festucetum valesiacaе*, *Andropogonetum grylli* and *Asphodeletum microcarpi*, which can be considered extinct, the other plant communities on Plavnik Island can be classified as in the following syntaxonomic scheme:

Festuco-Brometea Br.-Bl. et Tx. ex Soó 1947

Scorzoneretalia villosae Kovačević 1959

Chrysopogono grylli-Koelerion splendidis

Horvatić 1973

Helichrysetum italici Horvatić 1927 *typicum*

Helichrysetum italici salvietosum officinalis
subass. nov. hoc loco

Poetea bulbosae Rivas Goday et Rivas-Mart. in
Rivas-Mart. 1978

Poetalia bulbosae Rivas Goday et Rivas-Mart. in
Rivas Goday et Ladero 1970

Romuleion Oberd. 1954 *nom. inval. (?)*

Festuco valesiacaе-Poetum bulbosae ass. nov.
hoc loco

Class: ?

Hedypnoido rhagadioloidis-Asphodeletum ramosi
ass. nov. hoc loco

Acknowledgments

The authors thank Steve Latham (United Kingdom) for improving the English. We thank the Croatian Geological Survey for providing the Geological Map of the Republic of Croatia (sheet Cres and Lošinj). We also thank the Croatian Meteorological and Hydrological Service for providing climatological data from the meteorological station in the town of Krk, island of Krk, for the period 1998-2022. We would also like to thank two anonymous reviewers for their helpful suggestions.

The study was financed with funds from the Multi-purpose Institutional Financing Program supported by the Ministry of Science and Education of the Republic of Croatia.

Author contribution statement

MT and NJ conceived the study, sampled the vegetation, made the statistical analysis, and share first authorship.

References

- Biondi, E., Pesaresi, S., Galdenzi, D., Gasparri, R., Biscotti, N., Del Viscio, G., Casavecchia, S., 2016: Post-abandonment dynamic on Mediterranean and sub-Mediterranean perennial grasslands: the edge vegetation of the new class *Charyb-dido pancratii-Asphodeletea ramosi*. *Plant Sociology* 53(2), 3–18. <https://doi.org/10.7338/pls2016532/01>
- Biurrun, I., Willner, W., 2020: First report of the European Vegetation Classification Committee (EVCC). *Vegetation Classification and Survey* 1, 145–147. <https://doi.org/10.3897/VCS/2020/60352>
- Branković, Č., Güttler, I., Gajić-Čapka, M., 2013: Evaluating climate change at the Croatian Adriatic from observations and regional climate models' simulations. *Climate Dynamics* 41(9), 2353–2373. <https://doi.org/10.1007/s00382-012-1646-z>
- Brullo, S., Spampinato, G., 1991: La vegetazione dei corsi d'acqua della Sicilia. *Bollettino dell'Accademia Gioenia di Scienze Naturali* 23, 119–252.
- Čarni, A., Matevski, V., Šilc, U., Čušterevska, R., 2014: Early spring ephemeral therophytic non-nitrophilous grasslands as a habitat of various species of *Romulea* in the southern Balkans. *Acta Botanica Croatica* 73(1), 155–177. <https://doi.org/10.2478/botcro-2013-0017>
- De Cáceres, M., Legendre, P., 2009: Associations between species and groups of sites: indices and statistical inference. *Ecology* 90(12), 3566–3574. <https://doi.org/10.1890/08-1823.1>
- De Cáceres, M., Legendre, P., Moretti, M., 2010: Improving indicator species analysis by combining groups of sites. *Oikos* 119(10), 1674–1684. <https://doi.org/10.1111/j.1600-0706.2010.18334.x>
- Dengler, J., Löbel, S., Dolnik, C., 2009: Species constancy depends on plot size – a problem for vegetation classification and how it can be solved. *Journal of Vegetation Science* 20(4), 754–766. <https://doi.org/10.1111/j.1654-1103.2009.01073.x>
- Dufrêne, M., Legendre, P., 1997: Species assemblages and indicator species: the need for a flexible asymmetrical approach. *Ecological Monographs* 67(3), 345–366. [https://doi.org/10.1890/0012-9615\(1997\)067\[0345:SAAI\]2.0.CO;2](https://doi.org/10.1890/0012-9615(1997)067[0345:SAAI]2.0.CO;2)
- Dumančić, D., 1988: Unapređenje proizvodnosti travnjaka mediteranskog područja kao preduvjet razvoja ovčarstva i kozarstva [Improving the productivity of grasslands in the Mediterranean area as a prerequisite for the development of sheep and goat farming]. *Agronomski glasnik* 1, 105–111.
- EEA (European Environment Agency), 2019: The European environment - state and outlook 2020. Knowledge for transition to a sustainable Europe. Retrieved October 8, 2023 from <https://www.eea.europa.eu/soer/publications/soer-2020>
- Euro+Med. 2006-2023: Euro+Med PlantBase - the information resource for Euro-Mediterranean plant diversity. Retrieved September 1, 2023 from <http://www.europlusmed.org/>
- Foucault, B. (de), 2020: Sur la synsystème des *Rosmarineta officinalis* ouest-méditerranéens. *Carnets Botaniques* 4, 1–27. <https://doi.org/10.34971/s9d8-qj43>
- Fuček, L., Matičec, D., Vlahović, I., Oštrić, N., Prtoljan, B., Kroljija, B., Korbar, T., Husinec, A., Palenik, D., 2015: Osnovna geološka karta Republike Hrvatske M 1:50 000: list Cres i Lošinj [Basic geological map of the Republic of Croatia, scale 1:50 000: sheet Cres and Lošinj]. Hrvatski geološki institut, Zagreb.
- Gajić-Čapka, M., Güttler, I., Cindrić, K., Branković, Č., 2018: Observed and simulated climate and climate change in the lower Neretva river basin. *Journal of Water and Climate Change* 9(1), 124–136. <https://doi.org/10.2166/wcc.2017.034>
- Hećimović, S., 1984: Vegetation der Inseln Bobara und Mrkan. *Acta Botanica Croatica* 43(1), 109–118.
- Horvat, I., Horvatić, S., 1934: *Chrysopogoneto-Satureion subspicatae*: ein neuer Verband der *Brometalia erecti* Braun-Blanquet. *Acta Botanica Instituti Botanici Universitatis Zagrebensis* 9(1), 8–12.

- Horvat, I., Glavaš, V., Ellenberg, H., 1974: Vegetation Südosteuropas. Geobotanica selecta 4. Gustav Fischer Verlag, Stuttgart.
- Horvatić, S., 1927: Flora i vegetacija otoka Plavnika [Flora and vegetation of the island of Plavnik]. Acta Botanica Instituti Botanici Universitatis Zagrebensis 2(1), 1–56.
- Horvatić, S., 1963: Vegetacijska karta otoka Paga s općim pregledom vegetacijskih jedinica Hrvatskog primorja [Vegetation map of the island of Pag with survey of the plant communities of Hrvatsko Primorje]. Prirodoslovna Istraživanja 33. Acta Biologica IV, 3–187.
- Horvatić, S., 1973: Syntaxonomic analysis of the vegetation of dry grassland and stony meadows in Eastern Adriatic coastal Karts district based on the latest phytocoenological research. Fragmenta Herbológica Jugoslavica 32, 1–15.
- IPCC (Intergovernmental Panel on Climate Change), 2014: Climate Change 2014: Synthesis Report. In: Core Writing Team, Pachauri, R. K., Meyer, L. A., (eds.), Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. IPCC, Geneva, Switzerland.
- McCune, B., Mefford, M. J., 2011: PC-ORD. Multivariate analysis of ecological data. Version 6.0 MjM Software, Gleneden Beach, Oregon, USA.
- Médail, F., 2017: The specific vulnerability of plant biodiversity and vegetation on Mediterranean islands in the face of global change. Regional Environmental Change 17(6), 1775–1790. <https://doi.org/10.1007/s10113-017-1123-7>
- Midolo, G., Herben, T., Axmanová, I., Marcenò, C., Pätsch, R., Bruelheide, H., Karger, D. N., Ačić, S., Bergamini, A., Bergmeier, E., Biurrun, I., Bonari, G., Čarni, A., Chiarucci, A., De Sanctis, M., Demina, O., Dengler, J., Dziuba, T., Fanelli, G., Garbolino, E., Giusso del Galdo, G., Goral, F., Güler, B., Hinojos-Mendoza, G., Jansen, F., Jiménez-Alfaro, B., Lengyel, A., Lenoir, J., Pérez-Haase, A., Pielech, R., Prokhorov, V., Rašomavičius, V., Ruprecht, E., Růšņa, S., Šilc, U., Škvorc, Ž., Stančić, Z., Tatarenko, I., Chytrý, M., 2023: Disturbance indicator values for European plants. Global Ecology and Biogeography 32(1), 24–34. <https://doi.org/10.1111/geb.13603>
- Mucina, L., Bültman, H., Dierssen, K., Theurillat, J.-P., Dengler, J., Čarni, A., Šumberová, K., Raus, T., Di Pietro, R., Gavilán García, R., Chytrý, M., Iakushenko, D., Schaminée, J. H. J., Bergmeier, E., Santos Guerra, A., Daniëls, F. J. A., Ermakov, N., Valachovič, M., Pigantti, S., Rodwell, J. S., Pallas, J., Capelo, J., Weber, H. E., Lysenko, T., Solomeshch, A., Dimopoulos, P., Aguiar, C., Freitag, H., Hennekens, S. M., Tichý, L., 2016: Vegetation of Europe: Hierarchical floristic classification system of plant, lichen, and algal communities. Applied Vegetation Science 19(S1), 3–264. <https://doi.org/10.1111/avsc.12257>
- Myers, N., Mittermeier, R. A., Mittermeier, C. G., da Fonseca, G. A., Kent, J., 2000: Biodiversity hotspots for conservation priorities. Nature 403, 853–858. <https://doi.org/10.1038/35002501>
- Nikolić, T., Antonić, O., Alegro, A., Dobrović, I., Bogdanović, S., Liber, Z., Rešetnik, I., 2008: Plant species diversity of Adriatic islands: an introductory survey. Plant Biosystems 142(3), 435–445. <https://doi.org/10.1080/11263500802410769>
- Oberdorfer, E., 1954: Nordägyptische kraut- und zwergrsträucher im vergleich mit den entsprechenden Vegetationseinheiten des westlichen Mittelmeergebietes. Vegetatio 5(1), 88–96. <https://doi.org/10.1007/BF00299558>
- Official Gazette, 2019: Uredba o ekološkoj mreži i nadležnostima javnih ustanova za upravljanje područjima ekološke mreže [Regulation on the ecological network and the competences of public institutions for the management of ecological network areas]. 80. Croatian Government, Zagreb.
- Patarčić, M., Gajić-Čapka, M., Cindrić, K., Branković, Č., 2014: Recent and near-future changes in precipitation-extremes indices over the Croatian Adriatic coast. Climate Research 61(2), 157–176. <https://doi.org/10.3354/cr01250>
- Preislerová, Z., Jiménez-Alfaro, B., Mucina, L., Berg, C., Bonari, G., Kuzemko, A., Landucci, F., Marcenò, C., Monteiro-Henriques, T., Novák, P., Vynokurov, D., Bergmeier, E., Dengler, J., Apostolova, I., Bioret, F., Biurrun, I., Campos, J. A., Capelo, J., Čarni, A., Čoban, S., Csiky, J., Čuk, M., Čušterevska, R., Daniëls, F. J. A., De Sanctis, M., Didukh, Ya., Dítě, D., Fanelli, F., Golovanov, Y., Golub, V., Guarino, R., Hájek, M., Iakushenko, D., Indreica, A., Jansen, F., Jašková, A., Jiroušek, M., Kalníková, V., Kavğacı, A., Kucherov, I., Kůzmič, F., Lebedeva, M., Loidi, J., Lososová, Z., Lysenko, T., Milanović, Đ., Onyshchenko, V., Perrin, G., Peterka, T., Rašomavičius, V., Rodríguez-Rojo, M. P., Rodwell, J. S., Růšņa, S., Sánchez Mata, D., Schaminée, J. H. J., Semenishchenkov, Y., Shevchenko, N., Šibík, J., Škvorc, Ž., Smagin, V., Stešević, D., Stupar, V., Šumberová, K., Theurillat, J.-P., Tikhonova, E., Tzonev, R., Valachovič, M., Vassilev, K., Willner, W., Yamalov, S., Večeřa, M., Chytrý, M., 2022: Distribution maps of vegetation alliances in Europe. Applied Vegetation Science 25(1), e12642. <https://doi.org/10.1111/avsc.12642>
- R Core Team, 2023: R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
- Rivas Goday, S., Rivas-Martínez, S., 1963: Estudio y clasificación de los pastizales españoles. Publicaciones del Ministerio de Agricultura, Madrid.
- Rivas-Martínez, S., Sáenz, S. R., Penas, A., 2011: Worldwide bioclimatic classification system. Global Geobotany 1, 1–634.
- Rodwell, J. S., Schaminée, J. H. J., Mucina, L., Pignatti, S., Dring, J., Moss, D., 2002: The diversity of European vegetation. an overview of phytosociological alliances and their relationships to EUNIS habitats. National reference centre for agriculture, nature and fisheries, Wageningen.
- Stanišić-Vujačić, M., Stešević, D., Hadžiablahović, S., Caković, D., Šilc, U., 2022: An *Asphodelus ramosus* dominated plant community in Montenegro: fringe or grassland? Acta Botanica Croatica 81(1), 12–22. <https://doi.org/10.37427/botcro-2022-027>
- Stanišić-Vujačić, M., Stešević, D., Hadžiablahović, S., Šilc, U., 2023: Ecological and syntaxonomical characteristics of early spring therophytic ephemeral grasslands (alliance *Romuleion*) in the northeastern Mediterranean. Plant Biosystems 157(3), 540–563. <https://doi.org/10.1080/11263504.2023.2165570>
- Šegota, T., Filipčić, A., 2003: Köppenova podjela klime i hrvatsko nazivlje [Köppen climate division and Croatian nomenclature]. Geoadria 8(1), 17–37. <https://doi.org/10.15291/geoadria.93>
- Šegulja, N. 1970: Vegetacija sjeveroistočnog dijela Labinštine u Istri [Vegetation of the north-eastern part of Labin in Istria]. Acta Botanica Croatica 29, 157–172.
- Škvorc, Ž., Jasprica, N., Alegro, A., Kovačić, S., Franjić, J., Krstonošić, D., Vraneša, A., Čarni, A., 2017: Vegetation of Croatia: phytosociological classification of the high-rank syntaxa. Acta Botanica Croatica 76(2), 200–224. <https://doi.org/10.1515/botcro-2017-0014>
- Terzi, M., 2011: Nomenclatural revision for the order *Scorzonero-Chrysopogonetalia*. Folia Geobotanica 46(4), 411–444. <https://doi.org/10.1007/s12224-011-9100-2>

- Terzi, M., 2015: Numerical analysis of the order *Scorzoneretalia villosae*. *Phytocoenologia* 45(1–2), 11–32. <https://doi.org/10.1127/phyto/2015/0009>
- Terzi, M., 2023: A new *Asphodelus ramosus*-dominated association from the Murge Plateau (SE Italy). *Hacquetia* 22(2), 179–195. <https://doi.org/10.2478/hacq-2022-0020>
- Terzi, M., Jasprica, N., Pandža, M., Milović, M., Caković, D., 2023: Diversity and ecology of *Salvia officinalis* communities in the Western Balkans. *Plant Biosystems* 157(1), 175–187. <https://doi.org/10.1080/11263504.2022.2098868>
- Terzi, M., Fernández-González, F., Di Pietro, R., Theurillat, J.-P., 2024a: Phytosociological nomenclature of the class names *Helianthemetea guttati*, *Poetea bulbosae* and *Stipo giganteae-Agrostietea castellanae*. *Plant Biosystems* 158(1), 70–83. <https://doi.org/10.1080/11263504.2023.2287539>
- Terzi, M., Jasprica, N., Čarni, A., Matevski, V., Bergmeier, E., Theurillat, J.-P., 2024b: Nomenclature of the Balkan alliance *Romuleion graecae* (*Poetea bulbosae*). *Acta Botanica Croatica* 83(2), 87–91. <https://doi.org/110.37427/botcro-2024-020>
- Theurillat, J. P., Willner, W., Fernández-González, F., Bültmann, H., Čarni, A., Gigante, D., Mucina, L., Weber, H., 2021: International code of phytosociological nomenclature. 4th edition. *Applied Vegetation Science* 24(1), e1249. <https://doi.org/10.1111/avsc.12491>
- Tichý, L., Axmanová, I., Dengler, J., Guarino, R., Jansen, F., Mí-dolo, G., Nobis, M. P., Van Meerbeek, K., Ačić, S., Attorre, F., Bergmeier, E., Biurrun, I., Bonari, G., Bruelheide, H., Campos, J. A., Čarni, A., Chiarucci, A., Čuk, M., Čušterevska, R., Didukh, Y., Dítě, D., Dítě, Z., Dziuba, T., Fanelli, G., Fernández-Pascual, E., Garbolino, E., Gavilán, R.G., Gégout, J.-C., Graf, U., Güler, B., Hájek, M., Hennekens, S. M., Jandt, U., Jašková, A., Jiménez-Alfaro, B., Julve, P., Kambach, S., Karger, D. N., Karrer, G., Kavgaci, A., Knollová, I., Kuzemko, A., Kůzmič, F., Landucci, F., Lengyel, A., Lenoir, J., Marcenò, C., Moeslund, J.E., Novák, P., Pérez-Haase, A., Peterka, T., Pielech, R., Pignatti, A., Rašomavičius, V., Rüşa, S., Saatkamp, A., Šilc, U., Škvorc, Ž., Theurillat, J.-P., Wohlgemuth, T., Chytrý, M., 2023: Ellenberg-type indicator values for European vascular plant species. *Journal of Vegetation Science* 34(1), e13168. <https://doi.org/10.1111/jvs.13168>
- Topić, J., Vuković, N., 2010: Krk i Plavnik [Krk and Plavnik]. In: Nikolić, T., Topić, J., Vuković N. (eds.), *Botanički važna područja Hrvatske*, 217–224. Prirodoslovno-matematički fakultet Sveučilišta u Zagrebu, Školska knjiga d.o.o., Zagreb.
- Trinajstić, I., 2001: Stjepan Horvatić i Gabrijel Tomažič - pioniri fitocenoloških istraživanja travnjaka u Sloveniji [Stjepan Horvatić and Gabrijel Tomažič - the pioneers of phytosociological investigations of grasslands in Slovenia]. *Hladnikia* 12–13, 23–29.
- Trinajstić, I., 2008: Biljne zajednice republike Hrvatske [Plant communities of Croatia]. Akademija Šumarskih Znanosti, Zagreb.
- van der Maarel, E., 1979: Transformation of cover-abundance values in phytosociology and its effects on community similarity. *Vegetatio* 39(2), 97–114. <https://doi.org/10.1007/BF00052021>
- Vargas, P., 2020: The Mediterranean floristic region: High diversity of plants and vegetation types. In: Goldstein, M. I., DellaSala, D. A. (eds.), *Encyclopedia of the world's biomes*, vol. 3, 602–616. Elsevier, Oxford. <https://doi.org/10.1016/B978-0-12-409548-9.12097-4>
- Vukadinović, V., 2021: Pedološka karta Hrvatske [Pedological map of Croatia]. Retrieved October 8, 2023 from http://pedologija.com.hr/iBaza/DPKH_QGIS_VV.webp. Josip Juraj Strossmayer University, Osijek.
- WBG (World Bank Group), 2021: Climate Risk Country Profile - Croatia. Retrieved October 8, 2023 from <https://reliefweb.int/report/croatia/climate-riskcountry-profile-croatia>
- Westhoff, V., van der Maarel, E., 1980: The Braun-Blanquet approach. In: Whittaker, R. H. (ed.), *Classification of plant communities*, 287–399. 2nd edition. Dr. W. Junk Publishers, The Hague, Boston, London. https://doi.org/10.1007/978-94-009-9183-5_9