

## Antibiotic sensitivity patterns and molecular detection of Enterotoxin genes in *Staphylococcus aureus* isolated from frozen muscle foods

Olu<sup>1</sup>aseyi Akinware<sup>1</sup>, Clement Olusola Ogidi<sup>2\*</sup>, Bamidele Juliet Akinyele<sup>1</sup>

<sup>1</sup>Department of Microbiology, The Federal University of Technology, PMB 704 Akure, Nigeria

<sup>2</sup>Department of Food Science and Technology, School of Agriculture, Food and Natural Resources, Olusegun Agagu University of Science and Technology, PMB 353, Okitipupa, Nigeria

### ARTICLE INFO

#### Article history:

Received: August 26, 2022

Accepted: January 9, 2023

#### Keywords:

Enterotoxin  
polymerase chain reaction  
staphylococcal  
fish  
meat

### ABSTRACT

*Staphylococcus aureus* is considered to be the most common pathogen in commonly consumed and improperly stored frozen foods. Hence, this study investigated the antibiotic sensitivity patterns and the occurrence of enterotoxin genes (*sea* and *seb*) in *S. aureus* isolated from frozen poultry meat of chicken, turkey and fish, namely *Trachurus trachurus*, *Scomber scombrus* and *Merluccius merluccius*. A total of 500 samples of raw meat from chicken ( $n = 130$ ), turkey ( $n = 130$ ), and different fish ( $n = 240$ ) were randomly purchased from various markets and shopping malls in Akure between September 2020 and December 2021. The antibiotic sensitivity of isolated *S. aureus* was determined by disk diffusion method. *S. aureus* was analyzed for enterotoxin genes by multiplex polymerase chain reaction (PCR). The highest staphylococcal count ( $7.70 \times 10^5$  CFU/g) was observed in *T. trachurus*. One hundred and twenty five *S. aureus* were isolated from meat and fish with the highest occurrence (40.80%) in turkey meat. *S. aureus* is highly resistant (33.3 to 100%) to the tested antibiotics. The PCR assay, respectively, showed *sea* and *seb* coded for staphylococcal enterotoxin A and B. The presence of staphylococcal enterotoxin A and B in frozen muscle foods could be attributed to unhygienic conditions during processing and storage. The occurrence of antibiotic resistant and enterotoxigenic *S. aureus* in muscle foods could pose a serious threat to people when consumed. Hence, urgent attention is required to avert the incidences of staphylococcal food poisoning after consuming frozen meat and fish.

### Introduction

Frozen meat, fish, poultry and sea foods are important source of proteins in diets and mostly consumed by people of all social strata (Laskowski et al., 2018). Tissues from healthy animals are sterile, but many factors can influence their microbial contamination during slaughtering. Microbial contamination is possible through the water, air, soil, from the workers and the equipment involved and it often leads to spoilage if not properly handled, processed and preserved (Alegbeleye et al., 2018). The most common method to preserve muscle foods is by

chilling and freezing, while super-chilling is also an attractive technique of preserving muscle foods to maintain their freshness, enhancing the shelf-life during storage, transportation and at retail (Banerjee and Maheswarappa, 2019). Freezing preserves muscle foods for extended periods by suppressing the growth and multiplication of microorganisms that could cause food spoilage and foodborne illnesses (Archer, 2004; Zhan et al., 2018). In frozen products, some microorganisms are killed, while others might only be sub-lethally damaged and can recover upon thawing if storage is above  $-10$  °C. Gram negative bacteria are more susceptible to freezing injury than Gram positive

\*Corresponding author E-mail: [clementogidi@yahoo.com](mailto:clementogidi@yahoo.com)

bacteria. Hence, it supports the occurrence of *Staphylococcus* species in frozen meats and fish (Ogofure and Igbinsosa, 2021).

*S. aureus* is a Gram-positive bacteria and facultative anaerobe, which can live as a commensal organism on the skin as well as mucosal membranes. *S. aureus* recovers and survives sub-lethal injuries from freezing with cell membrane synthesis-related proteins, oxidative stress resistance-related proteins, and metabolism-related proteins with their virulence factors exhibit distinct expression patterns during resuscitation (Suo et al., 2018). The resuscitation mechanisms adopted by *S. aureus* together with their multiple antibiotic resistance pose a threat to food safety and are now of public health concern.

The multiple resistance of *S. aureus* to all  $\beta$ -lactam antibiotics has been linked to the presence of an enzyme;  $\beta$ -lactamase and the *mecA* gene that encode penicillin-binding protein (PBP2a), which allows the synthesis of the cell wall even at lethal concentrations of  $\beta$ -lactam antibiotics (Atanassova et al., 2017). *S. aureus* produces a variety of enzymes, toxins (cytotoxins, exotoxins, and exfoliative) with multiple virulence factors encoded by phages, plasmids, and pathogenicity, which has the ability to bind the major histocompatibility complex proteins of their host's immune system (Tam and Torres, 2019). Staphylococcal food poisoning is an intoxication that occurs after consuming improperly prepared foods contaminated with *S. aureus* enterotoxins as a result of unhygienic practices including improper handling of cooked or processed muscle foods, followed by their storage conditions (Argudín et al., 2010). *S. aureus* is a major cause of food poisoning induced by heat resistant enterotoxin and is one of the leading causes of foodborne illnesses (Varshney et al., 2009; Pinchuk et al., 2010). *S. aureus* strains produce a large variety of enterotoxins A, B, C, and D that have been frequently detected in poultry meat products (Pepe et al., 2006; Akkaya et al., 2014). Enterotoxin *S. aureus* is considered to be the most common pathogen causing outbreaks of food poisoning, characterized by symptoms associated with gastroenteritis, including vomiting, nausea, abdominal pain, cramps, diarrhea and causing significant morbidity (Ortega et al., 2010). The contamination of foods with pathogenic microorganisms with reoccurrence of foodborne diseases face significant challenges in modern healthcare services and have substantial negative impacts on economy. Enterotoxins are primarily responsible for foodborne illnesses and morbidity. Hence, frequent screening of staphylococcal enterotoxins (SE) in commonly consumed foods like animal muscle foods, poultry foods, and dairy products is paramount since they are foods of proteins source. This study, therefore,

investigated the antibiotics sensitivity patterns and occurrence of enterotoxins *S. aureus* in selected frozen poultry meat (frozen chicken and turkey) and fish like *Scomber scombrus*, *Trachurus trachurus* and *Merluccius merluccius* vended in Akure metropolis.

## Materials and methods

### Collection of meat and fish

Samples of frozen meat of chicken (*Gallus gallus domesticus*) = 130, turkey (*Meleagris gallopavo domesticus*) = 130, and different fish species, namely, horse mackerel (*Trachurus trachurus*) = 80, mackerel (*Scomber scombrus*) = 80, and hake (*Merluccius merluccius*) = 80, were randomly purchased at different vendors in Akure. Samples were collected in ice packs and transported to Postgraduate Research Laboratory, Department of Microbiology, The Federal University of Technology, Akure, for microbiological analysis.

### Isolation and identification of *Staphylococcus aureus*

One gram of tissue from each sample was crushed in sterile crucible and serially diluted to ten-fold dilutions, each dilution ( $10^4$ ) was plated on mannitol salt agar using pour plate method and incubated at 37 °C for 24 hours. The colonies were counted using colony counter (TT-20, Techmel and Techmel, USA). The colony was sub-cultured to obtain pure isolate. The morphological, Gram's reaction and biochemical tests like catalase, coagulase, oxidase, methyl red, Voges Proskauer, and sugar fermentation were carried out according to methods of Olutiola et al. (2000) and Cheesbrough (2006). The identity of isolates was determined based on the biochemical reactions using Krieg et al. (2010).

### Antibiotic susceptibility of *Staphylococcus*

The modified Kirby-Bauer method was used to determine the susceptibility of obtained isolates to some antibiotics as described by The Clinical and Laboratory Standards Institute (CLSI, 2017). The inoculum size in nutrient broth (18 h old culture) was adjusted to 0.5 McFarland turbidity standards to  $1.3 \times 10^5$  CFU/ml. Thereafter, 0.1 ml of the suspension was transferred to prepared Mueller-Hinton agar and spread with a sterilized glass spreader. The surface of the agar was allowed to dry, antibiotic discs were placed at the equidistance of the plate by sterile forceps and aseptically placed on top of agar plate. The plates were incubated at 37 °C for 18 hours. After incubation, a clear zone of no growth in the immediate

vicinity of antibiotic disk was measured and recorded as diameter of zone of inhibition in millimeter (mm), interpreted using CLSI interpretative chart as resistance, intermediate and susceptible (CLSI, 2017).

*Genomic DNA extraction and PCR test for staphylococcal enterotoxin genes*

*S. aureus* isolates were further examined for the presence of enterotoxin genes using specific primer in a multiplex PCR assay. DNA of *S. aureus* was extracted with PureLink Genomic DNA Mini Kit (Thermo Fisher Scientific, USA) according to manufacturer’s instructions. The primers used for the detection of SE genes are listed in Table 1 (Johnson et al., 1991). For polymerase chain reaction amplification, the reaction mixture contained: 2.5 µl 10 X PCR buffer (Invitrogen), 1 µl DNA, 1 µl of primer F (10 pmol), 1 µl of primer R (10 pmol), 0.5 µl dNTP (10 mM, Invitrogen), 1.5 µl MgCl<sub>2</sub> (Invitrogen), 0.5 U Taq DNA polymerase (Invitrogen) and final volume was adjusted to 25 µl by adding sterile ultrapure water. DNA amplification was performed in a thermal cycler with initial denaturation at 94 °C for 5 min followed by 30 cycles of denaturation at 94 °C for 1 min, annealing at 58 °C for 1 min and extension at 72 °C for 1 min, followed by a final extension at 72 °C for 5 min. The amplified PCR products were electrophoresed in a 1.5 % agarose gel (Sigma–Aldrich) containing 0.5 mg/ml ethidium bromide, TBE buffer (0.09 M Tris–HCl, 0.09 M boric acid, 2 mM EDTA, pH 8.3) for 30 min at 100 V and visualized under UV trans-illumination.

*Data analysis*

Data was statistically analyzed using SPSS version 20, was separated using Duncan’s New Multiple Range test and significant difference will be value at  $p \leq 0.05$ .

**Results and discussion**

Table 2 shows total Staphylococcal count ( $\times 10^5$  CFU/g) of frozen muscle foods.

The staphylococcal count of  $0.70$  to  $7.50 \times 10^5$  CFU/g were obtained for meat from chicken,  $1.50$  to  $7.40 \times 10^5$  CFU/g,  $2.10$  to  $7.70 \times 10^5$  CFU/g,  $3.30$  to  $5.10 \times 10^5$  CFU/g and  $1.00$  to  $6.50 \times 10^5$  CFU/g were recorded for turkey meat, *T. trachurus*, *S. scombrus* and *M. merluccius*, respectively. Findings of Bodunde et al. (2019) reported Staphylococcus count of  $1.70 \times 10^4$  to  $6.0 \times 10^5$  CFU/g for different muscle foods like beef (*Bos taurus*), chicken (*Gallus gallus domesticus*), turkey (*Meleagris gallopavo*), pork (*Sus scrofa domesticus*), chevon (*Capra aegagrus hircus*), mackerel (*Scomber scombrus*), horse mackerel (*Trachurus trachurus*), herrings (*Clupea pallasii*), blue whiting (*Merluccius merluccius*), and croaker (*Micropogonias undulatus*). The higher *S. aureus* count in meat and fish is a result of improper hygienic practices at the point of handling by slaughter personnel during meat production. Findings of Thwala et al. (2021) reported that a total of 2853 samples containing beef, pork, goat meat, camel meat, sheep/lamb from different African countries were highly contaminated with higher load of *S. aureus*. Species of *Staphylococcus* like *S. aureus*, *S. epidermidis*, *S. xylosus*, *S. sciuri*, *S. warneri*, *S. saprophyticus*, *S. schleiferi* and *S. auricularis* were identified from ready-to-eat fish products (Sergelidis et al. 2014). Humans and animals such as cattle, pigs, chickens, turkey, horses, and sheep can be colonised by *S. aureus* on their skin and in their nares (Lozano et al., 2016). The frequency of *Staphylococci* in muscle foods (meat and fish) is revealing how commensal bacteria from human skin and mucosal surfaces contributed to the contamination during different stages of processing; slaughtering, transportation, chopping, storage and persons involved in the marketing (Wu et al., 2018). Table 3 reveals the occurrence of *S. aureus* in selected frozen muscle foods, meat from chicken, turkey and fish. Frozen meat of turkey has got the highest occurrence of *S. aureus* 40.80%, followed by meat of chicken with 30.40%, but *S. scombrus* has got the least occurrence of 2.40%.

**Table 1.** Primers used for the detection of *Staphylococcus aureus* enterotoxin genes

| Gene       | Primers | Sequences 5’-3’       | Gene location | Size (bp) of PCR product |
|------------|---------|-----------------------|---------------|--------------------------|
| <i>sea</i> | SEA-F   | ttgaaacggttaaaccgaa   | 490-509       |                          |
|            | SEA-R   | gaaccttccatcaaaaaca   | 591-610       | 120                      |
| <i>seb</i> | SEB-F   | tcgcatcaaaactgacaaacg | 634-653       |                          |
|            | SEB-R   | gcaggtactctataagtgcc  | 1091-1110     | 478                      |

Source: (Johnson et al., 1991)

Findings of Ogofure and Igbinsosa (2021) showed that beef had the highest frequency of *S. aureus* contamination (46.7%), followed by chicken (40.0%) and fish (30.0%). Wu *et al.* (2018) also reported 35% of *S. aureus* from 1,850 samples of frozen meat and meat products sold in 39 cities and provincial capitals of China. Savariraj *et al.* (2021) isolated 66.67% of *S. aureus* from 120 chicken meat marketed in retail outlets of Chennai, India. Higher prevalence of *S. aureus* (64%) has been reported from frozen meat and fresh meat from Karbala province in Iraq (Namir *et al.*, 2017). Likewise, Oranusi *et al.* (2014); Ogidi *et al.* (2016); and Bodunde *et al.* (2019) reported the prevalence of *S. aureus* in muscle foods and ready-to-eat foods examined in Nigeria. The prevalence of *S. aureus* could be attributed to poor hygienic practice of personnel, equipment in slaughterhouses, salesmen and women. Table 4 shows the resistance percentage of *S. aureus* isolates to rifampin, gentamycin, ampicillin, nitrofurantoin, amoxicillin, oxacillin, fluoroquinolones, streptomycin, vancomycin, erythromycin and trimethoprim/sulfamethoxazole. The resistance percent ranged from 33.3 to 100%. Findings of Parvin *et al.* (2021) revealed the highest resistance percentage of 73.9% to 100% for *S. aureus* isolated from frozen chicken meat against ceftiofur, nalidixic acid, ampicillin and oxacillin, colistin, amoxicillin-clavulanic acid and amoxicillin, penicillin-G and cloxacillin, oxytetracycline, and cefixime. Findings of Wang *et al.* (2017) revealed a total of 1,150 *S. aureus* isolated from 27,000 retail foods with 97.6% of *S. aureus* displayed resistant to at least one antimicrobial compound like penicillin, oxacillin, ceftiofur, vancomycin, daptomycin, erythromycin, gentamicin, tetracycline, ciprofloxacin, clindamycin, trimethoprim-sulfamethoxazole, chloramphenicol, linezolid and 57.5% of these were multi drug resistant. *S. aureus* isolated from frozen meat and fish were highly resistant to some of the commercially available antibiotics (Ogofure and Igbinsosa, 2021). The researchers reported the antibiotic resistance profile of *S. aureus* with high resistance to erythromycin (94%), amoxicillin/clavulanic acid (87.5%) and trimethoprim-sulfamethoxazole (81%). Findings of Wu *et al.* (2018) revealed that *S. aureus* isolated from 1,850 retail meat and meat products displayed resistance of 11.0% to 85.4% to ampicillin, penicillin, erythromycin, tetracycline, kanamycin, telithromycin, clindamycin, streptomycin, norfloxacin, gentamicin, fusidic acid, ciprofloxacin, chloramphenicol, amoxicillin/clavulanic acid. *S. aureus* isolated from Oklahoma retail chicken and turkey meats displayed multiple resistance (5.4 to 94.6%) to ampicillin, tetracycline, penicillin, doxycycline, oxacillin, azithromycin, erythromycin, vancomycin, ciprofloxacin, ceftiofur, gentamicin, kanamycin, clindamycin, rifampin, trimethoprim/sulfamethoxazole, chloramphenicol with 12

Methicillin-Resistant *S. aureus* (MRSA) showed 100% resistance to ampicillin, penicillin, ceftiofur with 2% NaCl, oxacillin with 2% NaCl, azithromycin, and erythromycin (Abdalrahman *et al.*, 2015). The use of medically important antibiotics in livestock production is tremendously contributing to multiple antibiotic resistance and pathogenicity of *S. aureus* (Park and Ronholm, 2021). The ability of various pathogenic bacteria to withstand antibiotic residues in muscle foods contributed to reoccurrence emergency of foodborne diseases, which have become a matter of food security worldwide (Kumar *et al.*, 2020). Figure 1 shows agarose gel electrophoresis of the PCR product of *sea* gene in selected bacteria isolates (120 bp). Gel image indicates a positive amplification with the presence of *sea* gene in all 5 isolates loading arrangement (1-5 represent different strains of *S. aureus*). Figure 2 shows agarose gel electrophoresis of the PCR product of *seb* gene in two *S. aureus* (Band size approximately 478 bp). Table 5 revealed the occurrence of enterotoxin producing *S. aureus*. The use of DNA hybridization and PCR approaches are reliable biological methods to detect staphylococcal enterotoxins using gene-specific nucleotide sequences as probes (Wu *et al.*, 2016). In this study there was a low prevalence of enterotoxin producing *S. aureus* isolated from meats from chicken, turkey and fish. Findings of Ali (2014) and Arslan and Özdemir (2017) revealed the presence of enterotoxin genes in *S. aureus* isolated from fish. *S. aureus* 18/31 (58.1%) isolated from raw lamb and beef meat were positive for the presence of at least one or more SE genes, while no SE genes were found in strains isolated from cooked meat (Haghi *et al.*, 2021). Findings of Şanlıbaba (2022) indicated *sea* and *seb* in *S. aureus* isolated from retail raw beef, sheep, and lamb meat in Turkey, which were directly connected to the contamination arose from human and animal origins. Findings of Rodríguez-Lázaro *et al.* (2017) examined 868 animal-derived food items; diverse meat from antelope, duck, guinea pig, pork, rodents, turkey, dairy products and eggs that were confiscated from the non-European Union passengers or illegally sold in an open market near an European Union border. The researchers revealed the presence of MRSA isolates with 73% tested positive for one or more enterotoxin genes A, B, C, D, G, H, I, and J. Kitai *et al.* (2005) revealed that 78 enterotoxigenic *S. aureus* isolated from raw chicken meat; thighs, breasts, wings, livers, gizzards, hearts and ovaries belong to human biotype and poultry. SEs producing *S. aureus* are widely distributed in animals and humans and are frequently contaminating muscle foods leading to make Staphylococcal food poison, thereby causing a serious threat to consumers when bacteria multiply and release toxins in uncooked or inadequately cooked foods (Grispoldi *et al.*, 2021).

**Table 2.** Total Staphylococcal count ( $\times 10^5$  CFU/g) of frozen muscle foods vended in Akure

| Locations       | Points of collection | Chicken (130)          | Turkey (130)           | <i>T. trachurus</i> (80) | <i>S. scombrus</i> (80) | <i>M. merluccius</i> (80) |
|-----------------|----------------------|------------------------|------------------------|--------------------------|-------------------------|---------------------------|
| Kings market    | A                    | 5.40±0.40 <sup>b</sup> | 0.00                   | 2.20±0.10 <sup>d</sup>   | 4.20±0.31 <sup>b</sup>  | 6.40±0.29 <sup>a</sup>    |
|                 | B                    | 3.20±0.16 <sup>d</sup> | 7.50±0.11 <sup>a</sup> | 0.00                     | 0.00                    | 0.00                      |
|                 | C                    | 1.40±0.02 <sup>f</sup> | 4.70±0.93 <sup>b</sup> | 0.00                     | 0.00                    | 0.00                      |
|                 | D                    | 3.00±0.01 <sup>d</sup> | 3.30±0.03 <sup>c</sup> | 7.70±0.55 <sup>a</sup>   | 0.00                    | 0.00                      |
|                 | E                    | 0.00                   | 1.50±0.11 <sup>e</sup> | 2.50±0.71 <sup>d</sup>   | 0.00                    | 0.00                      |
| FUTA South Gate | A                    | 4.30±0.01 <sup>c</sup> | 3.10±0.06 <sup>c</sup> | 2.20±0.10 <sup>d</sup>   | 0.00                    | 2.40±0.02 <sup>c</sup>    |
|                 | B                    | 3.80±0.11 <sup>c</sup> | 0.00                   | 7.70±0.12 <sup>a</sup>   | 0.00                    | 1.30±0.00 <sup>d</sup>    |
|                 | C                    | 7.50±0.93 <sup>a</sup> | 1.70±0.86 <sup>e</sup> | 4.35±0.01 <sup>c</sup>   | 3.30±0.01 <sup>c</sup>  | 0.00                      |
|                 | D                    | 3.30±0.05 <sup>d</sup> | 2.40±0.02 <sup>d</sup> | 5.40 ±0.32 <sup>b</sup>  | 0.00                    | 0.00                      |
| Araromi Market  | A                    | 0.00                   | 7.40±0.07 <sup>a</sup> | 4.40±0.18 <sup>c</sup>   | 0.00                    | 1.00±0.00 <sup>d</sup>    |
|                 | B                    | 4.90±0.01 <sup>c</sup> | 1.80±0.01 <sup>e</sup> | 2.10±0.08 <sup>d</sup>   | 0.00                    | 3.20±0.44 <sup>b</sup>    |
|                 | C                    | 5.70±0.36 <sup>b</sup> | 3.30±0.11 <sup>c</sup> | 0.00                     | 0.00                    | 0.00                      |
| Isinkan Market  | A                    | 3.90±0.27 <sup>c</sup> | 0.00                   | 0.00                     | 0.00                    | 0.00                      |
|                 | B                    | 0.70±0.00 <sup>i</sup> | 5.50±0.50 <sup>b</sup> | 7.50±0.06 <sup>a</sup>   | 5.10 ±0.02 <sup>a</sup> | 0.00                      |
|                 | C                    | 2.10±0.00 <sup>e</sup> | 3.10±0.30 <sup>c</sup> | 5.10±0.13 <sup>b</sup>   | 0.00                    | 6.50±0.32 <sup>a</sup>    |
| Shopping Malls  |                      | 0.00                   | 2.00±0.07 <sup>e</sup> | 0.00                     | 0.00                    | 0.00                      |

Values are mean of replicates (n=3). Values carrying the same superscript at the column are not significantly different at  $p < 0.05$ .

**Table 3.** Occurrence (%) of *Staphylococcus aureus* in examined frozen muscle foods


| Locations       | Chicken (130) | Turkey (130) | <i>T. trachurus</i> (80) | <i>S. scombrus</i> (80) | <i>M. merluccius</i> (80) |
|-----------------|---------------|--------------|--------------------------|-------------------------|---------------------------|
| King's market   | 8             | 13           | 5                        | 1                       | 6                         |
| FUTA South Gate | 15            | 8            | 3                        | 1                       | 0                         |
| Araromi Market  | 8             | 15           | 4                        | 0                       | 3                         |
| Isinkan Market  | 7             | 11           | 8                        | 1                       | 4                         |
| Shopping malls  | 0             | 4            | 0                        | 0                       | 0                         |
| n               | 38            | 51           | 20                       | 3                       | 13                        |
| % occurrence    | 30.40         | 40.80        | 16.00                    | 2.40                    | 10.40                     |

n= number of *Staphylococcus aureus* isolates from meat and fish


**Table 4.** Resistance profile (%) of *S. aureus* isolated from selected frozen muscle Foods in Akure

| Antibiotics                   | Chicken n=38 | Turkey n=51 | <i>T. trachurus</i> n=20 | <i>S. scombrus</i> n=3 | <i>M. merluccius</i> n=13 |
|-------------------------------|--------------|-------------|--------------------------|------------------------|---------------------------|
| Rifampin                      | 20(52.60)    | 33(64.70)   | 13(65.00)                | 1(33.30)               | 7(53.80)                  |
| Gentamycin                    | 28(73.70)    | 40(78.40)   | 10(50.00)                | 0                      | 6(46.20)                  |
| Ampicillin                    | 16(42.10)    | 20(39.20)   | 8(40.00)                 | 0                      | 5(38.50)                  |
| Nitrofurantoin                | 25(65.80)    | 31(60.80)   | 11(55.00)                | 0                      | 8(61.50)                  |
| Amoxicillin                   | 16(40.10)    | 42(82.40)   | 17(85.00)                | 2(66.70)               | 7(53.80)                  |
| Oxacillin                     | 30(78.90)    | 37(72.50)   | 10(50.00)                | 3(100)                 | 4(30.70)                  |
| Fluoroquinolones              | 18(47.40)    | 32(62.75)   | 8(40.00)                 | 3(100)                 | 4(30.70)                  |
| Streptomycin                  | 20(52.60)    | 29(56.90)   | 15(75.00)                | 0                      | 8(61.50)                  |
| Vancomycin                    | 14(36.80)    | 18(35.30)   | 5(25.00)                 | 1(33.30)               | 4(30.70)                  |
| Erythromycin                  | 21(55.30)    | 30(58.80)   | 14(70.00)                | 2(66.70)               | 6(46.20)                  |
| Trimethoprim/sulfamethoxazole | 18(47.40)    | 28(54.90)   | 8(40.00)                 | 3(100)                 | 6(46.20)                  |

n= number of *S. aureus* isolates. Values in bracket represent % resistance to antibiotics.



**Figure 1.** PCR amplification for the detection of *sea* gene in *S. aureus*. Lane M: 100 bp marker. Lanes 1-5 indicate *sea* gene (120 bp).



**Figure 2.** PCR amplification for the detection of *seb* gene in *S. aureus*. Lane M: 100 bp marker. Lanes 3 and 5 indicate *seb* gene (478 bp).

**Table 5.** Occurrence and antibiotic profile of enterotoxin producing *S. aureus* isolated from frozen meat and fish

| Sample location | Muscle foods        | Antibiotic resistance profile                    | Staphylococcal genes |
|-----------------|---------------------|--|----------------------|
| Kings' market   | Turkey              | GEN, AMP, STR, VAN, AMX, OXA, FLU, ERY           | <i>sea</i>           |
| FUTA South Gate | Turkey              | RFM, GEN, AMP, VAN, AMX, OXA, FLU                | <i>sea</i>           |
| Kings' market   | Chicken             | RFM, AMP, VAN, AMX, VAN, FLU, ERY, TMS           | <i>sea</i>           |
| FUTA South Gate | Chicken             | GEN, AMP, VAN, AMX, NIT, OXA, FLU, ERY           | <i>sea</i>           |
| Isinkan Market  | Chicken             | RFM, VAN, AMX, STR, VAN, FLU, ERY, TMS           | <i>sea</i>           |
| Araromi Market  | Chicken             | RFM, AMP, VAN, AMX, NIT, OXA, VAN, FLU, ERY, TMS | <i>seb</i>           |
| FUTA South Gate | <i>T. trachurus</i> | GEN, AMP, STR, VAN, NIT, AMX, OXA, FLU, ERY      | <i>seb</i>           |

RFM: Rifampin, GEN: Gentamycin, AMP: Ampicillin, NIT: Nitrofurantoin, AMX: Amoxicillin OXA: Oxacillin, FLU: Fluoroquinolones, STR: Streptomycin, VAN: Vancomycin, ERY: Erythromycin, TMS: Trimethoprim/sulfamethoxazole

## Conclusions

The isolation and identification of *S. aureus* from selected muscle foods could be due to various types of contamination during processing steps and poor personnel hygiene of the food handlers. The low prevalence of enterotoxin producing *S. aureus* from frozen meat and fish can still be considered as a potential source of foodborne diseases. In order to safeguard public health, public enlightenment should be sustained to raise awareness on the proper cooking, packaging and storage of poultry products. The illegal importation of poultry products should be curbed by the creation of right policies and implementation of existing laws. The elimination of plethora pathogenic bacteria from frozen foods should be emphasized. With the increase in the prevalence of antibiotic resistant bacteria in frozen foods, public health awareness is essential as a preventative measure to control the use of antibiotics in livestock production.

**Author Contributions:** COO and BJA conceived, designed the experiments, and supervised the research work. COO and OA conducted the experiments, analyzed and interpreted the data, and wrote the paper. All the authors read the manuscript and approved.

**Funding:** This research received no external funding.

**Conflict of Interest:** The authors declare no conflict of interest.

## References

Abdallahman, L. S., Stanley, A., Wells, H., Fakhr, M. K. (2015): Isolation, Virulence, and Antimicrobial

Resistance of Methicillin-Resistant *Staphylococcus aureus* (MRSA) and Methicillin Sensitive *Staphylococcus*

*aureus* (MSSA) Strains from Oklahoma Retail Poultry Meats, *Intern. J. Environ. Res. Public Health* 12(6), 6148–6161. <https://doi.org/10.3390/ijerph120606148>

Akkaya, L., Kara, V. R., Yaman, H. (2014): Enterotoxin production by *Staphylococcus aureus* (A, B, C, D) during the ripening of sucuk (Turkish dry fermented sausage), *CyTA – J. Food* 12(2), 127-133.

Alegbeleye, O. O., Singleton, I., Sant'Ana, A. S. (2018): Sources and contamination routes of microbial pathogens to fresh produce during field cultivation: A review, *Food Microbiol.* 73, 177–208. <https://doi.org/10.1016/j.fm.2018.01.003>

Ali H. H. (2014): Isolation and identification of *Staphylococcus* bacteria from fish of fresh water and its antibiotics sensitivity in Mosul city, *Basrah J. Vet. Res.* 13(1), 33–42. <https://doi.org/10.33762/bvetr.2014.88123>

Archer D. L. (2004): Freezing: an underutilized food safety technology?, *Intern J Food Microbiol.* 90(2), 127–138. [https://doi.org/10.1016/s0168-1605\(03\)00215-0](https://doi.org/10.1016/s0168-1605(03)00215-0)

Argudín, M. Á., Mendoza, M. C., Rodicio, M. R. (2010): Food poisoning and *Staphylococcus aureus* enterotoxins, *Toxins* 2(7), 1751–1773. <https://doi.org/10.3390/toxins2071751>

Arslan S., Özdemir F. (2017): Molecular characterization and detection of enterotoxins, methicillin resistance genes and antimicrobial resistance of *Staphylococcus aureus* from fish and ground beef, *Polish J. Veter. Sci.* 20(1), 85–94. <https://doi.org/10.1515/pjvs-2017-0012>.

Atanassova, V., Meindl, A., Ring, C. (2017): Prevalence of *Staphylococcus aureus* and staphylococcal enterotoxins in raw pork and uncooked smoked ham—a comparison of classical culturing detection and RFLP-PCR, *Intern. J. Food Microbiol.* 68, 105–113.

Banerjee, R., Maheswarappa, N. B. (2019): Superchilling of muscle foods: Potential alternative for chilling and freezing, *Cri. Rev Food Sci. Nutr.* 59(8), 1256–1263. <https://doi.org/10.1080/10408398.2017.1401975>

Bodunde, R.S., Ogidi, C.O., Akinyele, B.J. (2019): Load and antibiotic susceptibility pattern of microorganisms

- in muscle foods sold in Akure, Southwest Nigeria, *J. Food Qual. Hazards Control.* 6, 30-36.
- Cheesbrough, M. (2006). *District Laboratory Practice in Tropical Countries.* 2nd Edition., Cambridge University Press, Cambridge, UK.
- CLSI-Clinical Laboratory Standards Institute (2017): *Performance Standards for Antimicrobial Susceptibility Testing; Twenty-Fourth Informational Supplement.* CLSI document M100-S24 Wayne, Pennsylvania, 19087, USA.
- Fawole, M. O., Oso, B. A. (2004): Characterization of bacteria: *Laboratory Manual of Microbiology.* 4th Edition, Spectrum Book Ltd., Ibadan, Nigeria, 24-33.
- Grispoldi, L., Karama, M., Armani, A., Hadjicharalambous, C., Cenci-Goga, B. T. (2021): *Staphylococcus aureus* enterotoxin in food of animal origin and staphylococcal food poisoning risk assessment from farm to table, *Italian J Ani. Sci.* 20(1), 677-690.
- Haghi, F., Zeighami, H., Hajiloo, Z., Torabi, N., Derakhshan, S. (2021): High frequency of enterotoxin encoding genes of *Staphylococcus aureus* isolated from food and clinical samples, *Journal of health, population, and nutrition* 40(1), 27. <https://doi.org/10.1186/s41043-021-00246-x>
- Johnson, W. M., Tyler, S. D., Ewan, E. P., Ashton, F. E., Pollard, D. R., Rozee, K. R. (1991): Detection of genes for enterotoxins, exfoliative toxins, and toxic shock syndrome toxin 1 in *Staphylococcus aureus* by the polymerase chain reaction, *J. Clinical Microbiol.* 29(3), 426-430. <https://doi.org/10.1128/jcm.29.3.426-430.1991>
- Kitai, S., Shimizu, A., Kawano, J., Sato, E., Nakano, C., Kitagawa, H., Fujio, K., Matsumura, K., Yasuda, R. Inamoto, T. (2005): Prevalence and characterization of *Staphylococcus aureus* and enterotoxigenic *Staphylococcus aureus* in retail raw chicken meat throughout Japan, *J. Veter Med. Sci.* 67 (3), 269-274. <https://doi.org/10.1292/jvms.67.269>
- Krieg, N.R., Staley, J.T., Brown, D.R., Hedlund, B.P., Paster, B.J., Ward, N.L., Ludwig, W., Whitman, W.B. (2010): *The Bacteroidetes, Spirochaetes, Tenericutes (Mollicutes), Acidobacteria, Fibrobacteres, Fusobacteria, Dictyoglomi, Gemmatimonadaetes, Lentisphaerae, Verrucomicrobia, Chlamydiae, and Planctomycetes,* 2nd ed., vol. 4. New York: Springer. <https://doi.org/10.1007/978-0387-68572-4>.
- Kumar, H., Bhardwaj, K., Kaur, T., Nepovimova, E., Kuča, K., Kumar, V., Bhatia, S. K., Dhanjal, D. S., Chopra, C., Singh, R., Guleria, S., Bhalla, T. C., Verma, R., Kumar, D. (2020): Detection of bacterial pathogens and antibiotic residues in chicken meat: a review, *Foods* 9(10), 1504. <https://doi.org/10.3390/foods9101504>
- Laskowski, W., Górska-Warsewicz, H., Kulykovets, O. (2018): Meat, meat products and seafood as sources of energy and nutrients in the average Polish diet, *Nutrients* 10(10), 1412. <https://doi.org/10.3390/nu10101412>
- Lozano, C., Gharsa, H., Ben Slama, K., Zarazaga, M., Torres, C. (2016): *Staphylococcus aureus* in animals and food: methicillin resistance, prevalence and population structure. A review in the African continent, *Microorganisms* 4(1), 12. <https://doi.org/10.3390/microorganisms4010012>
- Namir, I. M., Mohammad, J. A. (2017): Isolation and Identification of *Staphylococcus aureus* strains from fresh and frozen meat in Karbala Province, *Int J Sci Nat.* 8 (3), 704-709.
- Ogidi, C. O., Oyetayo, V. O., Akinyele, B. J. (2016): Microbial quality and antibiotic sensitivity profile of microorganisms isolated from street foods sold in Akure metropolis, Nigeria, *Jordan J Biol. Sci.* 9(4), 227 – 234.
- Ogofure, A. G., Igbiosa, E. O. (2021). Effects of rinsing on *Staphylococcus aureus* load in frozen meats and fish obtained from open markets in Benin City, Nigeria, *Afr. J. Clin. Exper. Microbiol.* 22 (2), 294-299.
- Olutiola, P. O., Famurewa, O., Sonntag, H. G. (2000): *An introduction to general microbiology. A practical approach,* 1st ed. Heidelbergger Verlagsanstalt, Heidelberg, Germany.
- Oranusi, S., Obioha, T. U., and Adekeye, B. T. (2014): Investigation on the microbial profile of frozen foods: Fish and Meat, *Int. J Adv Res Biol Sci.* 1 (2), 71-78.
- Ortega, E., Abriouel, H., Lucas, R., Gálvez, A. (2010): Multiple roles of *Staphylococcus aureus* enterotoxins: pathogenicity, superantigenic activity, and correlation to antibiotic resistance, *Toxins* 2(8), 2117–2131. <https://doi.org/10.3390/toxins2082117>
- Park, S., Ronholm, J. (2021): *Staphylococcus aureus* in agriculture: lessons in evolution from a multispecies pathogen, *Clinical Microbiol. Rev.* 34(2), e00182-20. <https://doi.org/10.1128/CMR.00182-20>
- Parvin, Mst. S., Ali, Md. Y., Talukder, S., Nahar, A., Chowdhury, E. H., Rahman, Md. T., Islam, Md. T. (2021): Prevalence and multidrug resistance pattern of methicillin resistant *S. aureus* isolated from frozen chicken meat in Bangladesh, *Microorganisms* 9(3), 636. <https://doi.org/10.3390/microorganisms9030636>
- Pepe, O., Blaiotta, G., Bucci, F., Anastasio, M., Aponte, M., Villani, F. (2006): *Staphylococcus aureus* and staphylococcal enterotoxin A in breaded chicken products: detection and behavior during the cooking process, *Appl. Environ. Microbiol.* 72(11), 7057–7062. <https://doi.org/10.1128/AEM.00198-06>.
- Pinchuk, I. V., Beswick, E. J., Reyes, V. E. (2010): Staphylococcal enterotoxins, *Toxins* 2, 2177-2197.
- Şanlıbaba, P. (2022): Prevalence, antibiotic resistance, and enterotoxin production of *Staphylococcus aureus* isolated from retail raw beef, sheep, and lamb meat in Turkey, *Intern. J. Food Microbiol.* 361, 109461. <https://doi.org/10.1016/j.ijfoodmicro.2021.109461>
- Savariraj, W. R., Ravindran, N. B., Kannan, P., Rao, V. A. (2021): Occurrence and enterotoxin gene profiles of *Staphylococcus aureus* isolated from retail chicken meat, *Food Sci. Technol. Intern.* 27(7), 619–625. <https://doi.org/10.1177/1082013220980204>
- Sergelidis, D., Abraham, A., Papadopoulos, T., Soultos, N., Martziou, E., Koulourida, V., Govaris, A., Pexara, A., Zdragas, A., Papa, A. (2014). Isolation of methicillin-

- resistant *Staphylococcus* spp. from ready-to-eat fish products, *Letters Appl Microbiol.* 59(5), 500–506. <https://doi.org/10.1111/lam.12304>
- Suo, B., Yang, H., Wang, Y., Lv, H., Li, Z., Xu, C., Ai, Z. (2018): Comparative proteomic and morphological change analyses of *Staphylococcus aureus* during resuscitation from prolonged freezing, *Front. Microbiol.* 9, 866. <https://doi.org/10.3389/fmicb.2018.00866>
- Tam, K., Torres, V. J. (2019). *Staphylococcus aureus* secreted toxins and extracellular enzymes, *Microbiol Spectrum* 7(2), <https://doi.org/10.1128/microbiolspec.GPP3-0039-2018>
- Thwala, T., Madoroba, E., Basson, A., Butaye, P. (2021): Prevalence and characteristics of staphylococcus aureus associated with meat and meat products in African Countries: A review, *Antibiotics* 10(9), 1108. <https://doi.org/10.3390/antibiotics10091108>
- Varshney, A. K., Mediavilla, J. R., Robiou, N. Guh, A., Wang, X., Gialanella, P., Levi, M. H., Kreiswirth, B. N., Fries, B. C. (2009): Diverse enterotoxin gene profiles among clonal complexes of *Staphylococcus aureus* isolates from the Bronx, New York, *Appl Environ. Microbiol.* 75, 6839–6849.
- Wang, W., Baloch, Z., Jiang, T., Zhang, C., Peng, Z., Li, F., Fanning, S., Ma, A., Xu, J. (2017): Enterotoxigenicity and Antimicrobial Resistance of *Staphylococcus aureus* Isolated from Retail Food in China, *Front. Microbiol.* 8, 2256. <https://doi.org/10.3389/fmicb.2017.02256>
- Wu, S., Duan, N., Gu, H., Hao, L., Ye, H., Gong, W., Wang, Z. (2016): A Review of the methods for detection of *Staphylococcus aureus* enterotoxins, *Toxins* 8(7), 176. <https://doi.org/10.3390/toxins8070176>
- Wu, S., Huang, J., Wu, Q., Zhang, J., Zhang, F., Yang, X., Wu, H., Zeng, H., Chen, M., Ding, Y., Wang, J., Lei, T., Zhang, S., Xue, L. (2018): *Staphylococcus aureus* isolated from retail meat and meat products in China: incidence, antibiotic resistance and genetic diversity, *Front. Microbiol.* 9, 2767. <https://doi.org/10.3389/fmicb.2018.02767>
- Zhan, X., Sun, D. W., Zhu, Z., Wang, Q. J. (2018): Improving the quality and safety of frozen muscle foods by emerging freezing technologies: A review, *Critical Rev. Food Sci. Nutr.* 58(17), 2925–2938. <https://doi.org/10.1080/10408398.2017.1345854>