

ON THE BOUNDEDNESS OF EULER-STIELTJES CONSTANTS FOR THE RANKIN-SELBERG L -FUNCTION

MEDINA ZUBAČA

University of Sarajevo, Bosnia and Herzegovina

ABSTRACT. Let E be a Galois extension of \mathbb{Q} of finite degree and let π and π' be two irreducible automorphic unitary cuspidal representations of $GL_m(\mathbb{A}_E)$ and $GL_{m'}(\mathbb{A}_E)$, respectively. Let $\Lambda(s, \pi \times \tilde{\pi}')$ be a Rankin-Selberg L -function attached to the product $\pi \times \tilde{\pi}'$, where $\tilde{\pi}'$ denotes the contragredient representation of π' , and let its finite part (excluding Archimedean factors) be $L(s, \pi \times \tilde{\pi}')$. The Euler-Stieltjes constants of the Rankin-Selberg L -function are the coefficients in the Laurent (Taylor) series expansion around $s = 1 + it_0$ of the function $L(s, \pi \times \tilde{\pi}')$. In this paper, we derive an upper bound for these constants.

1. INTRODUCTION

The classical Euler constant

$$\gamma = \gamma_0 = \lim_{x \rightarrow \infty} \left(\sum_{n < x} \frac{1}{n} - \log x \right) = 0.57721 \dots,$$

discovered and computed correctly up to five decimal places by L. Euler [12] in 1731. is the constant term in the Laurent series expansion of the Riemann zeta function at $s = 1$,

$$\zeta(s) = \frac{1}{s-1} + \gamma + \sum_{k=1}^{\infty} \gamma_k (s-1)^k = \frac{1}{s-1} + \sum_{k=0}^{\infty} \gamma_k (s-1)^k.$$

In 1885, T. J. Stieltjes [16] pointed out that each γ_n can be obtained as

$$(1.1) \quad \gamma_k = \frac{(-1)^k}{k!} \lim_{x \rightarrow \infty} \left(\sum_{n < x} \frac{\log^k n}{n} - \frac{\log^{k+1} x}{k+1} \right).$$

2020 *Mathematics Subject Classification.* 11M26, 11S40.

Key words and phrases. Euler-Stieltjes constants, Rankin-Selberg L -function.

The proof of equation (1.1) can be found in [3, 6]. Therefore, the constants γ_k ($k \geq 0$) are named the Stieltjes constants, the generalized Euler constants or the Euler-Stieltjes constants.

The Euler-Stieltjes constants γ_k are closely related (see e.g. [4]) to coefficients η_k of the Laurent series expansion of the logarithmic derivative of the Riemann zeta function at $s = 1$

$$\frac{\zeta'}{\zeta}(s) = -\frac{1}{s-1} + \sum_{k=0}^{\infty} \eta_k (s-1)^k, \quad |s-1| < 3.$$

Constants η_k can be evaluated as (see e.g. [9])

$$\eta_k = \frac{(-1)^{k-1}}{k!} \lim_{x \rightarrow \infty} \left(\sum_{n < x} \frac{\Lambda(n) \log^k n}{n} - \frac{\log^{k+1} x}{k+1} \right),$$

where $\Lambda(n)$ is the von Mangoldt function [25, 40]. Usually, constants γ_k are called the Euler-Stieltjes constants of the first kind, while constants η_k are called the Euler-Stieltjes constants of the second kind.

The Euler-Stieltjes constants of the first and the second kinds are important in both theoretical and computational analytic number theory since they appear in various estimations and as a result of asymptotic analysis. For example, the Euler-Stieltjes constants of the first kind can be used to determine a zero-free region of the Riemann zeta function near the real axis in the critical strip $0 < \text{Res} < 1$ [1]. The Euler-Stieltjes constants of the second kind are related to the Li positivity criterion for the Riemann hypothesis [4] since they appear in the arithmetic formula for the non-archimedean part of the Li coefficient. Numerical evaluation and estimations are given in [23].

The Euler-Stieltjes constants of the first and the second kinds and their relation to the Li criterion for the Riemann hypothesis were further investigated by M. Coffey in [8, 10] and by C. Knessl and M. Coffey in [20]. Some interesting formulas and bounds are recently derived in [30].

This concept is generalized in many different settings. Coefficients appearing in the Laurent (Taylor) series representation of a zeta or L -function or its logarithmic derivative are called generalized Euler-Stieltjes constants of the first and the second kinds. Different kinds of formulas, properties or bounds are derived.

Results related to the Hurwitz zeta function are given in [3], those for the Dedekind zeta function in [15, 33], for the general setting of a non-co-compact Fuchsian group with unitary representation in [2], for a class of functions possessing an Euler product representation in [14], for a subclass \mathcal{S}^b of the Selberg class in [39], for the extended Selberg class in [17] and for the Rankin-Selberg L -functions in [27, 28]. Also, some investigations are done in the case of zeta functions with multiple variables, introducing multiple Stieltjes constants, for example, see [22, 34]. q -analogues of these coefficients are investigated in [7].

In this paper, we investigate generalized Euler-Stieltjes constants attached to the Rankin-Selberg L -functions associated with two representations. We precisely define coefficients under consideration in the sequel. Let E be a Galois extension of \mathbb{Q} of finite degree and let π and π' be two irreducible automorphic unitary cuspidal representations (see e.g. [11]) of $GL_m(\mathbb{A}_E)$ and $GL_{m'}(\mathbb{A}_E)$, respectively. The generalized Euler-Stieltjes constants of the first kind $\gamma_{\pi, \pi'}(k)$ attached to the finite part of Rankin-Selberg L -function $L(s, \pi \times \tilde{\pi}')$ (an analogue of classical ζ function) are defined as coefficients in the Laurent (Taylor) series representation of $L(s, \pi \times \tilde{\pi}')$ at $s = 1 + it_0$:

$$(1.2) \quad L(s, \pi \times \tilde{\pi}') = \sum_{k=-\delta(t_0)}^{\infty} \gamma_{\pi, \pi'}(k)(s-1-it_0)^k,$$

where $\delta(t_0) = 1$ if and only if $m = m'$ and $\pi' \cong \pi \otimes |\det|^{it_0}$, for some $t_0 \in \mathbb{R}$, where \cong denotes isomorphic representations. Otherwise, $\delta(t_0) = 0$.

In this paper, the finite part of Rankin-Selberg L -function we denote by $L(s, \pi \times \tilde{\pi}')$ and call the Rankin-Selberg L -function, and its completed function (including Archimedean factors) we denote by $\Lambda(s, \pi \times \tilde{\pi}')$.

The purpose of this paper is to derive an upper bound for coefficients $\gamma_{\pi, \pi'}(k)$ appearing in (1.2). The Rankin-Selberg L -functions attached to a convolution of two irreducible, unitary cuspidal representations of $GL_m(\mathbb{A}_E)$ and $GL_{m'}(\mathbb{A}_E)$ over number field E do not always belong to the extended Selberg class \mathcal{S}^\sharp , which is introduced in [19] (nor to the class of functions considered in [14]). In the case when $m = m'$ and $\pi' \cong \pi \otimes |\det|^{it_0}$, for some $t_0 \in \mathbb{R} \setminus \{0\}$ the Rankin-Selberg L -function possesses pole at $s = 1 + it_0 \neq 1$. Hence, they do not satisfy axiom (ii) of the class \mathcal{S}^\sharp . Furthermore, coefficients μ_j appearing in the functional equation for the Rankin-Selberg L -functions unconditionally satisfy the bound $\operatorname{Re} \mu_j > -1$, different from the bound $\operatorname{Re} \mu_j \geq 0$, posed in axiom (iii) of the class \mathcal{S}^\sharp .

The rest of the paper is organized as follows. In section 2 we give a complete overview of the setting we are dealing with, introduce necessary notation and recall some known results that will be used for the proofs. Section 3 contains some preliminary results about functions under consideration, while the main results are stated and proved in sections 4 and 5. In section 4 integral representation of coefficients under consideration is derived, while their bounds are proved in 5.

2. PRELIMINARIES AND NOTATIONS

Let E be a Galois extension of \mathbb{Q} of degree d , and let \mathbb{A}_E denote the ring of adèles over E . For every place v , let E_v be the completion of a number field E at v , and let f_p denote the modular degree of E_v over the field of p -adic numbers \mathbb{Q}_p for $v|p$, where p is a prime. Let S_∞ denotes a set of infinite places v of the number field E . The Rankin-Selberg L -function attached to

the product $\pi \times \tilde{\pi}'$ of irreducible cuspidal representations of $GL_m(\mathbb{A}_E)$ and $GL_{m'}(\mathbb{A}_E)$ with a unitary central character (see e.g. [11]), respectively, is given by absolutely convergent Euler product of local factors

$$L(s, \pi \times \tilde{\pi}') = \prod_{v < \infty} L_v(s, \pi_v \times \tilde{\pi}'_v),$$

for $\text{Res} > 1$, see e.g. [18, Th. 5.3.], where $\tilde{\pi}$ denotes the contragredient representation of π . For finite place v at which π_v and π'_v are unramified, the local factors of $L(s, \pi \times \pi')$ are given by

$$(2.1) \quad L_v(s, \pi \times \tilde{\pi}') = \prod_{j=1}^m \prod_{k=1}^{m'} \left(1 - \alpha_\pi(v, j) \overline{\alpha_{\pi'}(v, k)} p^{-fs}\right)^{-1},$$

where $\{\alpha_\pi(v, j)\}_{j=1}^m$ and $\{\alpha_{\pi'}(v, k)\}_{k=1}^{m'}$ are corresponding sets of Satake parameters associated to π and π' , respectively. If π_v or $\pi_{v'}$ ramified, we can also write the local factors at ramified places v in the same form (2.1) with the convention that some of $\alpha_\pi(v, j)$ and $\alpha_{\pi'}(v, k)$ may be zero (see e.g. [27]).

The function $L(s, \pi \times \tilde{\pi}')$ has a Dirichlet series expansion of the form

$$(2.2) \quad L(s, \pi \times \tilde{\pi}') = \sum_{n=1}^{\infty} \frac{a_{\pi \times \tilde{\pi}'}(n)}{n^s},$$

that is valid for $\text{Res} > 1$.

Similarly, at the infinite place $v \in S_\infty$, the archimedean local factor $L_v(s, \pi_v \times \tilde{\pi}'_v)$ can be written as a product

$$L_v(s, \pi_v \times \tilde{\pi}'_v) = \prod_{j=1}^m \prod_{k=1}^{m'} \Gamma_v(s + \mu_{\pi \times \tilde{\pi}'}(v, j, k)),$$

where $\mu_{\pi \times \tilde{\pi}'}(v, j, k) = \mu_\pi(v, j) + \overline{\mu_{\pi'}(v, k)}$, at the infinite places v unramified for both π and π' , $\{\mu_\pi(v, j)\}_{j=1}^m$ and $\{\mu_{\pi'}(v, j)\}_{j=1}^{m'}$ are the Langlands parameters associated to π_v and π'_v respectively and $\Gamma_v(s) = \pi^{-s/2} \Gamma(s/2)$, if v is real and $\Gamma_v(s) = 2(2\pi)^{-s} \Gamma(s)$, if v is complex. In the case when infinite place v is ramified for π or π' , parameters $\mu_{\pi \times \tilde{\pi}'}(v, j, k)$ are described in [31, Appendix], where it is also proved that $\mu_{\pi \times \tilde{\pi}'}(v, j, k)$, for all $j = 1, \dots, m$ and $k = 1, \dots, m'$ satisfy the trivial bound $\text{Re} \mu_{\pi \times \tilde{\pi}'}(v, j, k) > -1$.

As proved in [11, Th. 9.1. and Th. 9.2.], the completed Rankin-Selberg L -function

$$\Lambda(s, \pi \times \tilde{\pi}') = L(s, \pi \times \tilde{\pi}') \prod_{v \in S_\infty} L_v(s, \pi_v \times \tilde{\pi}'_v)$$

extends to a meromorphic function of order one on the whole complex plane, bounded (away from its possible poles) in the vertical strip. The functional equation, which is due to F. Shahidi ([36–38]),

$$(2.3) \quad \Lambda(s, \pi \times \tilde{\pi}') = \varepsilon(\pi \times \tilde{\pi}') Q_{\pi \times \tilde{\pi}'}^{\frac{1}{2}-s} \Lambda(1-s, \tilde{\pi} \times \pi')$$

is valid for all s , where $Q_{\pi \times \tilde{\pi}'} > 0$ is the arithmetic conductor and $\varepsilon(\pi \times \tilde{\pi}')$ is a complex number of modulus 1. The function $\Lambda(s, \pi \times \tilde{\pi}')$ has simple poles at $s = 1 + it_0$ and $s = it_0$, arising from $L(s, \pi \times \tilde{\pi}')$ if and only if $m = m'$ and $\pi' \cong \pi \otimes |\det|^{it_0}$, for some $t_0 \in \mathbb{R}$. Otherwise, it is an entire function.

Following [13] let us define

$$(2.4) \quad \delta(t_0) = \begin{cases} 1, & m = m' \text{ and } \pi' \cong \pi \otimes |\det|^{it_0}, \text{ for some } t \in \mathbb{R}; \\ 0, & \text{otherwise,} \end{cases}$$

then the functional equation (2.3) can be written as

$$(2.5) \quad L(s, \pi \times \tilde{\pi}') \Psi_{\pi, \pi'}(s) = \bar{L}(1 - s, \pi \times \tilde{\pi}'),$$

where $\bar{L}(s, \pi \times \tilde{\pi}') = \overline{L(\bar{s}, \pi \times \tilde{\pi}')}$ and the factor $\Psi_{\pi, \pi'}(s)$ is given by

$$(2.6) \quad \Psi_{\pi, \pi'}(s) = \frac{Q_{\pi \times \tilde{\pi}'}^{s - \frac{1}{2}}}{\varepsilon(\pi \times \tilde{\pi}')} \prod_{v \in S_\infty} \prod_{j=1}^m \prod_{k=1}^{m'} \frac{\Gamma_v(s + \mu_{\pi \times \tilde{\pi}'}(v, j, k))}{\Gamma_v\left(1 - s + \overline{\mu_{\pi \times \tilde{\pi}'}(v, j, k)}\right)}.$$

As in [26], it follows that (2.6) can be written in more convenient form, as

$$(2.7) \quad \Psi_{\pi, \pi'}(s) = \frac{\left(Q_{\pi \times \tilde{\pi}'} \pi^{-dmm'}\right)^{s - \frac{1}{2}}}{\varepsilon(\pi \times \tilde{\pi}')} \prod_{l=1}^{dmm'} \frac{\Gamma\left(\frac{1}{2}(s + \mu_{\pi \times \tilde{\pi}'}(l))\right)}{\Gamma\left(\frac{1}{2}\left(1 - s + \overline{\mu_{\pi \times \tilde{\pi}'}(l)}\right)\right)},$$

where $|\varepsilon(\pi \times \tilde{\pi}')| = 1$ and $\mu_{\pi \times \tilde{\pi}'}(l) = \mu_{\pi \times \tilde{\pi}'}(v, j, k)$, for $r_1 + r_2$ places $v \in S_\infty$ and $\mu_{\pi \times \tilde{\pi}'}(l) = \mu_{\pi \times \tilde{\pi}'}(v, j, k) + 1$, for the rest of r_2 places $v \in S_\infty$ ($j = 1, \dots, m, k = 1, \dots, m'$) and r_1 denotes number of real places $v \in S_\infty$ and r_2 denotes number of complex places $v \in S_\infty$.

The zeros of $\Lambda(s, \pi \times \tilde{\pi}')$ are called non-trivial zeros of $L(s, \pi \times \tilde{\pi}')$. They lie in the strip $0 < \text{Res} < 1$, see [35]. The function $L(s, \pi \times \tilde{\pi}')$ may also have trivial zeros, which arise from the poles of the local L -factors at infinite places. There are finitely many of them inside the critical strip $0 \leq \text{Res} \leq 1$ at points $s = -\mu_{\pi \times \tilde{\pi}'}(v, j, k)$, for those $v \in S_\infty, j \in \{1, \dots, m\}$ and $k \in \{1, \dots, m'\}$ such that $\text{Re} \mu_{\pi \times \tilde{\pi}'}(v, j, k) \leq 0$.

3. SOME PROPERTIES OF THE RANKIN-SELBERG L -FUNCTIONS

In the following proposition, we give some asymptotic bounds for the Rankin-Selberg L -functions and the factor $\Psi_{\pi, \pi'}(s)$ of the functional equation. These results are used in proof of the main result of the paper.

PROPOSITION 3.1. *Let E be a Galois extension of \mathbb{Q} of finite degree d and let π and π' be two irreducible automorphic unitary cuspidal representations of $GL_m(\mathbb{A}_E)$ and $GL_{m'}(\mathbb{A}_E)$. The function $\Psi_{\pi, \pi'}(s)$ satisfies relation*

$$(3.1) \quad |\Psi_{\pi, \pi'}(\sigma + it)| \sim_\sigma \left(\frac{Q_{\pi \times \tilde{\pi}'}}{(2\pi)^{dmm'}} \right)^{\sigma - \frac{1}{2}} |t|^{(\sigma - \frac{1}{2})dmm'},$$

as $|t| \rightarrow +\infty$. Further, for an arbitrary $\varepsilon > 0$ the function $L(s, \pi \times \tilde{\pi}')$ satisfies

$$(3.2) \quad L(\sigma + it, \pi \times \tilde{\pi}') = \begin{cases} O_\varepsilon(1) & \text{if } \sigma \geq 1 + \varepsilon, \\ O_\varepsilon \left(|t|^{\frac{dmm'}{2}(1-\sigma+\varepsilon)} \right) & \text{if } -\varepsilon \leq \sigma \leq 1 + \varepsilon, \\ O_{\varepsilon, \sigma} \left(|t|^{\frac{dmm'}{2}(1-2\sigma)} \right) & \text{if } \sigma \leq -\varepsilon. \end{cases}$$

PROOF. The function $\Psi_{\pi, \pi'}(s)$ can be written as

$$\begin{aligned} \Psi_{\pi, \pi'}(s) &= \frac{1}{\varepsilon(\pi \times \tilde{\pi}')} \left(Q_{\pi \times \tilde{\pi}'} \pi^{-dmm'} \right)^{s-\frac{1}{2}} \\ &\times \exp \left[\sum_{l=1}^{dmm'} \left(\log \left[\Gamma \left(\frac{s + \mu_{\pi \times \tilde{\pi}'}(l)}{2} \right) \right] - \log \left[\Gamma \left(\frac{1-s + \overline{\mu_{\pi \times \tilde{\pi}'}(l)}}{2} \right) \right] \right) \right]. \end{aligned}$$

By applying the asymptotic series expansion of function $\log \Gamma(z+a)$ (see [21, Section 2.11, relation (4)]) on the functions $\log \left[\Gamma \left(\frac{s + \mu_{\pi \times \tilde{\pi}'}(l)}{2} \right) \right]$ and $\log \left[\Gamma \left(\frac{1-s + \overline{\mu_{\pi \times \tilde{\pi}'}(l)}}{2} \right) \right]$, with $z = \frac{it}{2}$ and $z = \frac{-it}{2}$ respectively, we obtain relation (3.1).

For $\text{Res} = \sigma \geq 1 + \varepsilon > 1$ the Rankin-Selberg L -function $L(s, \pi \times \tilde{\pi}')$ is given by an absolutely convergent Euler product for $\text{Res} > 1$, so

$$L(\sigma + it, \pi \times \tilde{\pi}') = O_\varepsilon(1), \quad \text{for } \sigma \geq 1 + \varepsilon,$$

where O_ε denotes that a constant appearing in O notation depends on ε . For $\text{Res} = \sigma \leq -\varepsilon < 0$, the functional equation for the Rankin-Selberg L -function given by (2.5) and relation (3.1) imply

$$L(\sigma + it, \pi \times \tilde{\pi}') = O_{\varepsilon, \sigma} \left(|t|^{\frac{dmm'}{2}(1-2\sigma)} \right),$$

as $|t| \rightarrow +\infty$, where $O_{\varepsilon, \sigma}$ denotes that a constant appearing in O notation depends on σ and ε . In special case, if σ lies in a closed and bounded subset of \mathbb{R} , a constant in O notation is uniform in σ and depends on ε .

For σ such that $-\varepsilon \leq \sigma \leq 1 + \varepsilon$, Phragmén-Lindelöf theorem for strip can be used to derive the desired result. Basically, since the function

$$(s - it_0)^{\delta(t_0)} (s - 1 - it_0)^{\delta(t_0)} L(s, \pi \times \tilde{\pi}'),$$

where $\delta(t_0)$ is defined by (2.4), is an entire of finite order, the bound

$$|L(s, \pi \times \tilde{\pi}')| = O(\exp(\exp(\delta|t|))),$$

holds true for sufficiently large $|t|$ and any $\delta > 0$. Application of the result [29, Proposition 8.15] to the Rankin-Selberg L -function in the strip $-\varepsilon \leq \sigma \leq 1 + \varepsilon$ implies

$$L(\sigma + it, \pi \times \tilde{\pi}') = O_\varepsilon \left(|t|^{\frac{dmm'}{2}(1-\sigma+\varepsilon)} \right),$$

as $|t| \rightarrow +\infty$. The proof is complete. \square

4. INTEGRAL REPRESENTATION OF THE GENERALIZED EULER-STIELTJES CONSTANTS ASSOCIATED TO THE RANKIN-SELBERG L -FUNCTION

In this section, we derive an integral representation for coefficients in the Laurent (Taylor) series expansion of the Rankin-Selberg L -function given by (1.2) using a classical method in the analytic number theory based on contour integrals (see e.g. [40, Section 4.14], [17]). A key idea in the method is to apply the Cauchy integral formula to obtain an integral expression for coefficients, and then deform the contour appearing in the integral expression to a line from $a - i\infty$ to $a + i\infty$. Cauchy integral formula implies

$$(4.1) \quad \gamma_{\pi, \pi'}(k) = \frac{1}{2\pi i} \int_C \frac{L(s, \pi \times \tilde{\pi}')}{(s-1-it_0)^{k+1}} ds,$$

where contour C is a positively oriented circle with centre $s = 1 + it_0$ and radius r such that it contains $s = 1 + it_0$ as the only singularity of the integrand¹. If $\delta(t_0) = 0$, for all $t_0 \in \mathbb{R}$, then (1.2) gives Taylor series expansions of function $L(s, \pi \times \tilde{\pi}')$ and in that case, let $t_0 = 0$.

PROPOSITION 4.1. *Let E be a Galois extension of \mathbb{Q} of finite degree d and let $L(s, \pi \times \tilde{\pi}')$ be Rankin-Selberg L -function attached to the product $\pi \times \tilde{\pi}'$ be two irreducible automorphic unitary cuspidal representations of $GL_m(\mathbb{A}_E)$ and $GL_{m'}(\mathbb{A}_E)$. Let k be a positive integer and a be a real number such that $1 < 1 + \varepsilon < a < \frac{k+1}{dmm'} + \frac{1}{2}$ and $\frac{1}{2}(1 - a + \operatorname{Re}\mu_{\pi \times \tilde{\pi}'}(l)) \notin \mathbb{Z}$ for all $l = 1, \dots, dmm'$. Then,*

$$(4.2) \quad \gamma_{\pi, \pi'}(k) = \frac{(-1)^k}{2\pi i} \int_{a-i\infty}^{a+i\infty} \frac{\overline{L(\bar{s}, \pi \times \tilde{\pi}')} G_L(s)}{(s+it_0)^{k+1}} ds + \delta(t_0)(-1)^{k+1} \operatorname{Res}_{s=it_0} L(s, \pi \times \tilde{\pi}'),$$

where

$$(4.3) \quad G_L(s) = \frac{\epsilon(\pi \times \tilde{\pi}') Q_{\pi \times \tilde{\pi}'}^{s-\frac{1}{2}}}{(\pi^{dmm'})^{s+\frac{1}{2}}} \prod_{l=1}^{dmm'} \left[\Gamma\left(\frac{s + \overline{\mu_{\pi \times \tilde{\pi}'}(l)}}{2}\right) \times \Gamma\left(\frac{1+s - \mu_{\pi \times \tilde{\pi}'}(l)}{2}\right) \sin \frac{\pi}{2} (1-s + \mu_{\pi \times \tilde{\pi}'}(l)) \right].$$

PROOF. The proof is based on integral representation (4.1). The contour \mathcal{C} is deformed to a suitable rectangular $\mathcal{R}_{a,A,T}$ and the integral is decomposed into integrals over its sides.

¹Since the function $L(s, \pi \times \tilde{\pi}')$ might have two poles, $s = it_0$ and $s = 1 + it_0$, we can choose for radius r any positive number less than $\frac{1}{2}$.

Let A and T be sufficiently large positive numbers. Let $\mathcal{R}_{a,A,T}$ be a positively oriented rectangle determined by vertices $-a+1-iT$, $A-iT$, $A+iT$ and $-a+1+iT$. Compared to the integral over C , the additional contribution can be from a simple pole $s = it_0$ of the function $L(s, \pi \times \tilde{\pi}')$ if it exists. By the Cauchy's formula, we can write

$$\frac{1}{2\pi i} \int_{\mathcal{R}_{a,A,T}} \frac{L(s, \pi \times \tilde{\pi}')}{(s-1-it_0)^{k+1}} ds = \gamma_{\pi, \pi'}(k) + \delta(t_0) \operatorname{Res}_{s=it_0} \frac{L(s, \pi \times \tilde{\pi}')}{(s-1-it_0)^{k+1}}.$$

Therefore,

$$(4.4) \quad \gamma_{\pi, \pi'}(k) = \frac{1}{2\pi i} \int_{\mathcal{R}_{a,A,T}} \frac{L(s, \pi \times \tilde{\pi}')}{(s-1-it_0)^{k+1}} ds + \delta(t_0) (-1)^k \operatorname{Res}_{s=it_0} L(s, \pi \times \tilde{\pi}').$$

Now, integral over $\mathcal{R}_{a,A,T}$ can be written as a sum of integrals over line segments \mathcal{S}_1 , \mathcal{S}_2 , \mathcal{S}_3 and \mathcal{S}_4 joining $-a+1+iT$, $-a+1-iT$, $A-iT$, $A+iT$ and $-a+1+iT$, respectively.

For integral over \mathcal{S}_2 , we have

$$\begin{aligned} \int_{\mathcal{S}_2} \frac{L(s, \pi \times \tilde{\pi}')}{(s-1-it_0)^{k+1}} ds &= \int_{-a+1-iT}^{A-iT} \frac{L(s, \pi \times \tilde{\pi}')}{(s-1-it_0)^{k+1}} ds \\ &= \left(\int_{-a+1-iT}^{-\varepsilon-iT} + \int_{-\varepsilon-iT}^{1+\varepsilon-iT} + \int_{1+\varepsilon-iT}^{A-iT} \right) \frac{L(s, \pi \times \tilde{\pi}')}{(s-1-it_0)^{k+1}} ds. \end{aligned}$$

Using Proposition 3.1 we obtain following asymptotic bounds

$$\begin{aligned} \left| \int_{-a+1-iT}^{-\varepsilon-iT} \frac{L(s, \pi \times \tilde{\pi}')}{(s-1-it_0)^{k+1}} ds \right| &= O_\varepsilon \left(\left| \frac{T}{T+t_0} \right|^{k+1} |T|^{(a-\frac{1}{2})dmm'-k-1} \right), \\ \left| \int_{-\varepsilon-iT}^{1+\varepsilon-iT} \frac{L(s, \pi \times \tilde{\pi}')}{(s-1-it_0)^{k+1}} ds \right| &= O_\varepsilon \left(\left| \frac{T}{T+t_0} \right|^{k+1} |T|^{\frac{dmm'}{2}(1+2\varepsilon)-k-1} \right), \end{aligned}$$

and

$$\left| \int_{1+\varepsilon-iT}^{A-iT} \frac{L(s, \pi \times \tilde{\pi}')}{(s-1-it_0)^{k+1}} ds \right| = O_\varepsilon \left(\frac{1}{|T+t_0|^{k+1}} \right),$$

where O_ε denotes that constants appearing in O notation are uniform in $\operatorname{Res} = \sigma$, for $s \in \mathcal{S}_2$, and might depend on ε .

Hence, for $1+\varepsilon < a < \frac{k+1}{dmm'} + \frac{1}{2}$ and $k > -1$, we obtain

$$(4.5) \quad \int_{\mathcal{S}_2} \frac{L(s, \pi \times \tilde{\pi}')}{(s-1-it_0)^{k+1}} ds \rightarrow 0, \quad \text{as } |T| \rightarrow \infty.$$

Integral over \mathcal{S}_4 can be bounded completely analogously, i.e. we get

$$(4.6) \quad \int_{\mathcal{S}_4} \frac{L(s, \pi \times \tilde{\pi}')}{(s-1-it_0)^{k+1}} ds \rightarrow 0, \quad \text{as } |T| \rightarrow \infty.$$

Next, we consider the integral over \mathcal{S}_3 . Here $s = A + it$, and by choice of A we are in the region of absolute convergence of the Rankin-Selberg L -function, thus from Proposition 3.1 and by substitution $u = t - t_0$ follows

$$\left| \int_{\mathcal{S}_3} \frac{L(s, \pi \times \tilde{\pi}')}{(s-1-it_0)^{k+1}} ds \right| \leq 2K \int_0^{+\infty} \frac{du}{((A-1)^2 + u^2)^{\frac{k+1}{2}}},$$

where K is a positive constant such that $|L(A + it, \pi \times \tilde{\pi}')| \leq K$. From Lebesgue's convergence theorem, when $A \rightarrow \infty$, it follows that the contribution of the integral over \mathcal{S}_3 tends to zero, as $|T| \rightarrow \infty$. Namely, for the integrand

$$f_A(t) = \frac{1}{((A-1)^2 + t^2)^{\frac{k+1}{2}}},$$

and function

$$g(t) = \begin{cases} 1, & t \in [0, 1]; \\ \frac{1}{t^{k+1}}, & t > 1, \end{cases}$$

holds $f_A(t) \leq g(t)$ on $[0, +\infty)$, for $k > 0$ and $g(t)$ is integrable. Then, since $\lim_{A \rightarrow +\infty} f_A(t) = 0$, we have

$$\lim_{A \rightarrow +\infty} \lim_{T \rightarrow +\infty} \int_{\mathcal{S}_3} \frac{L(s, \pi \times \tilde{\pi}')}{(s-1-it_0)^{k+1}} ds = 0.$$

Thus, the only contribution to the integral in (4.4), when $|T| \rightarrow \infty$, is from the integral over \mathcal{S}_1 . So, for $k > \max\{0, (\frac{1}{2} + \varepsilon) dmm' - 1\}$, we have

$$\begin{aligned} \gamma_{\pi, \pi'}(k) &= \frac{1}{2\pi i} \int_{-a+1-i\infty}^{-a+1+i\infty} \frac{L(s, \pi \times \tilde{\pi}')}{(s-1-it_0)^{k+1}} ds + \delta(t_0)(-1)^k \operatorname{Res}_{s=it_0} L(s, \pi \times \tilde{\pi}') \\ &= \frac{(-1)^k}{2\pi i} \int_{a-i\infty}^{a+i\infty} \frac{L(1-s, \pi \times \tilde{\pi}')}{(s+it_0)^{k+1}} ds + \delta(t_0)(-1)^k \operatorname{Res}_{s=it_0} L(s, \pi \times \tilde{\pi}'). \end{aligned}$$

Functional equation (2.5) for the Rankin-Selberg L -function and definition (4.3) of the function $G_L(s)$, combined with formula $\Gamma(s)\Gamma(1-s) = \frac{\pi}{\sin \pi s}$, which is valid for all $s \notin \mathbb{Z}$, applied to the gamma functions appearing in gamma factor of the functional equation imply

$$L(1-s, \pi \times \tilde{\pi}') = \overline{L(\bar{s}, \pi \times \tilde{\pi}')} G_L(s),$$

for $\frac{1}{2}(1-s + \mu_{\pi \times \tilde{\pi}'}(l)) \notin \mathbb{Z}$.

Hence, relation (4.2) holds true for all $k > \max\{0, (\frac{1}{2} + \varepsilon) dmm' - 1\}$, where $a \in (1 + \varepsilon, \frac{k+1}{dmm'} + \frac{1}{2})$ is chosen such that $\frac{1}{2}(1 - a + \operatorname{Re}\mu_{\pi \times \tilde{\pi}'}(l)) \notin \mathbb{Z}$ for all $l = 1, 2, \dots, dmm'$. This completes the proof of Proposition 4.1. \square

5. BOUNDS FOR THE GENERALIZED EULER-STIELTJES CONSTANTS ASSOCIATED TO THE RANKIN-SELBERG L -FUNCTION

In this section, we prove the main result of the paper, the theorem that gives an upper bound for the Euler-Stieltjes coefficients $\gamma_{\pi, \tilde{\pi}'}(k)$ defined by (1.2). The proof is based on integral representation (4.2) derived in the previous section. Firstly, in the following lemma, we prove a bound for the function $G_L(s)$ appearing in the integrand in (4.2).

LEMMA 5.1. *Let E be a Galois extension of \mathbb{Q} of finite degree d and let $L(s, \pi \times \tilde{\pi}')$ be Rankin-Selberg L -function attached to the product $\pi \times \tilde{\pi}'$ two irreducible automorphic unitary cuspidal representations of $GL_m(\mathbb{A}_E)$ and $GL_{m'}(\mathbb{A}_E)$. Let $\mu_R = \max_{l=1, \dots, dmm'} |\operatorname{Re}\mu_{\pi \times \tilde{\pi}'}(l)|$, $\mu_I = \max_{l=1, \dots, dmm'} |\operatorname{Im}\mu_{\pi \times \tilde{\pi}'}(l)|$. For $a > \max\{1 + \varepsilon, \mu_R\}$, where $\varepsilon > 0$, we have*

$$(5.1) \quad |G_L(a + it)| \leq Q_{\pi \times \tilde{\pi}'}^{a - \frac{1}{2}} C_L(a) \times \left[\left(\frac{1 + a + \mu_R}{2} \right)^2 + \left(\frac{|t| + \mu_I}{2} \right)^2 \right]^{dmm' \frac{2a-1}{4}},$$

where constant $C_L(a)$ is given by

$$C_L(a) = \left(\frac{2}{\pi^{a-\frac{1}{2}}} \right)^{dmm'} \exp \left(\sum_{l=1}^{dmm'} \frac{2a+1}{6(a + \operatorname{Re}\mu_{\pi \times \tilde{\pi}'}(l))(1 + a - \operatorname{Re}\mu_{\pi \times \tilde{\pi}'}(l))} \right).$$

PROOF. From definition (4.3) of function G_L for $s = a + it$, and having in mind that $\epsilon(\pi \times \tilde{\pi}')$ is a complex number of modulus 1, one obtains

$$(5.2) \quad |G_L(a + it)| = \frac{Q_{\pi \times \tilde{\pi}'}^{a - \frac{1}{2}}}{(\pi^{dmm'})^{a + \frac{1}{2}}} \prod_{l=1}^{dmm'} \left[\left| \sin \frac{\pi}{2} (1 - a - it + \mu_{\pi \times \tilde{\pi}'}(l)) \right| \times \left| \Gamma \left(\frac{1 + a + it - \mu_{\pi \times \tilde{\pi}'}(l)}{2} \right) \Gamma \left(\frac{a + it + \overline{\mu_{\pi \times \tilde{\pi}'}(l)}}{2} \right) \right| \right].$$

Factors containing sine function, we bound using a simple representation in terms of exponential functions, precisely for $z \in \mathbb{C}$,

$$(5.3) \quad |\sin z| \leq e^{|\operatorname{Im}z|}.$$

While bounds for the factors containing gamma functions will be based on Binet formula [41, p. 258]

$$(5.4) \quad \begin{aligned} \log |\Gamma(z)| &= \left(\operatorname{Re} z - \frac{1}{2} \right) \log |z| - \operatorname{Im} z \arctan \frac{\operatorname{Im} z}{\operatorname{Re} z} - \operatorname{Re} z + \frac{1}{2} \log(2\pi) \\ &+ \operatorname{Re} \left[\int_0^{+\infty} \left(\frac{1}{2} - \frac{1}{t} + \frac{1}{e^t - 1} \right) \frac{e^{-tz}}{t} dt \right], \end{aligned}$$

valid for $\operatorname{Re} z > 0$. A simple calculation implies that the second term can be additionally simplified, i.e.

$$-\operatorname{Im} z \arctan \frac{\operatorname{Im} z}{\operatorname{Re} z} - \operatorname{Re} z \leq -\frac{\pi}{2} |\operatorname{Im} z|.$$

The properties of the function $g(t) = \left(\frac{1}{2} - \frac{1}{t} + \frac{1}{e^t - 1} \right) \frac{1}{t}$, specially, the fact that it attains its maximum $1/12$, at $t = 0$, gives us a bound

$$\operatorname{Re} \left[\int_0^{+\infty} \left(\frac{1}{2} - \frac{1}{t} + \frac{1}{e^t - 1} \right) \frac{e^{-tz}}{t} dt \right] \leq \frac{1}{12 \operatorname{Re} z}.$$

So, for $\operatorname{Re} z > 0$, relation (5.4) implies

$$(5.5) \quad \log |\Gamma(z)| \leq \left(\operatorname{Re} z - \frac{1}{2} \right) \log |z| - |\operatorname{Im} z| \frac{\pi}{2} + \frac{1}{2} \log(2\pi) + \frac{1}{12 \operatorname{Re} z}.$$

For the arguments appearing in (5.2), bound (5.3) implies

$$(5.6) \quad \left| \sin \frac{\pi}{2} (1 - a - it + \mu_{\pi \times \tilde{\pi}'(l)}) \right| \leq \exp \left(\frac{\pi}{2} |t - \operatorname{Im} \mu_{\pi \times \tilde{\pi}'(l)}| \right),$$

for all $l = 1, \dots, dmm'$. Since, by the assumption, $a > \max\{1 + \varepsilon, \mu_R\}$, and coefficients $\mu_{\pi \times \tilde{\pi}'(l)}$ for the Rankin-Selberg L -function satisfy bound $\operatorname{Re} \mu_{\pi \times \tilde{\pi}'} > -1$, we have

$$\operatorname{Re} \left(\frac{a + it + \overline{\mu_{\pi \times \tilde{\pi}'(l)}}}{2} \right) > 0 \quad \text{and} \quad \operatorname{Re} \left(\frac{1 + a + it - \mu_{\pi \times \tilde{\pi}'(l)}}{2} \right) > 0,$$

for all $l = 1, \dots, dmm'$, thus inequality (5.5) may be applied for the gamma factors in (5.2).

In addition, definition of numbers μ_R and μ_I implies the following inequalities

$$\begin{aligned} (t - \operatorname{Im} \mu_{\pi \times \tilde{\pi}'(l)})^2 &\leq (|t| + \mu_I)^2, \\ (a + \operatorname{Re} \mu_{\pi \times \tilde{\pi}'(l)})^2 &\leq (1 + a + \mu_R)^2, \\ (1 + a - \operatorname{Re} \mu_{\pi \times \tilde{\pi}'(l)})^2 &\leq (1 + a + \mu_R)^2, \end{aligned}$$

and from (5.5) we obtain

$$\begin{aligned} & \log \left| \Gamma \left(\frac{a + it + \overline{\mu_{\pi \times \tilde{\pi}'(l)}}}{2} \right) \right| + \log \left| \Gamma \left(\frac{1 + a + it - \mu_{\pi \times \tilde{\pi}'(l)}}{2} \right) \right| \\ & \leq \frac{2a-1}{4} \log \left(\left(\frac{1+a+\mu_R}{2} \right)^2 + \left(\frac{|t|+\mu_I}{2} \right)^2 \right) - \frac{\pi}{2} |t - \text{Im} \mu_{\pi \times \tilde{\pi}'(l)}| \\ & \quad + \frac{1}{6} \frac{2a+1}{(a + \text{Re} \mu_{\pi \times \tilde{\pi}'(l)})(1+a - \text{Re} \mu_{\pi \times \tilde{\pi}'(l)})} + \log 2\pi, \end{aligned}$$

for all $l = 1, \dots, dmm'$. This bound combined with (5.6) implies

$$\begin{aligned} & \left| \Gamma \left(\frac{a + it + \overline{\mu_{\pi \times \tilde{\pi}'(l)}}}{2} \right) \Gamma \left(\frac{1 + a + it - \mu_{\pi \times \tilde{\pi}'(l)}}{2} \right) \right| \\ & \quad \times \left| \sin \frac{\pi(1-a-it + \mu_{\pi \times \tilde{\pi}'(l)})}{2} \right| \\ & \leq \exp \left[\frac{2a-1}{4} \log \left(\left(\frac{1+a+\mu_R}{2} \right)^2 + \left(\frac{|t|+\mu_I}{2} \right)^2 \right) \right. \\ & \quad \left. + \frac{2a+1}{6(a + \text{Re} \mu_{\pi \times \tilde{\pi}'(l)})(1+a - \text{Re} \mu_{\pi \times \tilde{\pi}'(l)})} + \log 2\pi \right]. \end{aligned}$$

Substituting it into (5.2), we obtain (5.1), and the proof is complete. \square

The first explicit upper bound for coefficients in the Laurent series expansion of the Riemann zeta function about $s = 1$ has been given by Briggs [5]. Then, Matsuoka studied the asymptotic behaviour of these coefficients and he gave an excellent upper bound for its in [24]. Results related to upper bound for Stieltjes constants for the Dirichlet L-function when χ is a primitive character modulo q is given in [32], those for the Hurwitz zeta function in [3]. The investigation of Stieltjes constants for functions from the extended Selberg class \mathcal{S}^{\sharp} is done and an upper bound for these coefficients is obtained in [17].

The following theorem is the main result of the paper, it gives a bound for the coefficients under consideration.

THEOREM 5.2. *Let E be a Galois extension of \mathbb{Q} of finite degree d and let $L(s, \pi \times \tilde{\pi}')$ be Rankin-Selberg L-function attached to the product $\pi \times \tilde{\pi}'$ two irreducible automorphic unitary cuspidal representations of $GL_m(\mathbb{A}_E)$ and $GL_{m'}(\mathbb{A}_E)$ with pole at $s = 1 + it_0$ if $m = m'$ and $\pi' \cong \pi \otimes |\det|^{it_0}$, otherwise $t_0 = 0$. Let $\mu_R = \max_{l=1, \dots, dmm'} |\text{Re} \mu_{\pi \times \tilde{\pi}'(l)}|$, $\mu_I = \max_{l=1, \dots, dmm'} |\text{Im} \mu_{\pi \times \tilde{\pi}'(l)}|$ and $\mu_{R,I} = \max\{\mu_R, \mu_I + t_0 - 1\}$. Let $a > \max\{1 + \varepsilon, \mu_{R,I}, |t_0| + \mu_I - \mu_{R,I}\}$ and $\frac{1}{2}(1 - a + \text{Re} \mu_{\pi \times \tilde{\pi}'(l)}) \notin \mathbb{Z}$ for all $l = 1, \dots, dmm'$. For positive integer k such*

that $k > dmm'(a - \frac{1}{2})$ we have

$$(5.7) \quad |\gamma_{\pi, \pi'}(k)| \leq D_L(a) a^{-k} \left(2 + \mu_{R,I} + \mu_I + \frac{4}{k - dmm' \frac{2a-1}{2}} \right) + \delta(t_0) \left| \operatorname{Res}_{s=it_0} L(s, \pi \times \tilde{\pi}') \right|,$$

where constant $D_L(a)$ is defined by

$$D_L(a) = \exp \left(\frac{2a+1}{6} \sum_{l=1}^{dmm'} \frac{1}{(a + \operatorname{Re} \mu_{\pi \times \tilde{\pi}'}(l))(1 + a - \operatorname{Re} \mu_{\pi \times \tilde{\pi}'}(l))} \right) \times 2^{\frac{dmm'}{2}} (3a + \frac{1}{2}) \frac{Q_{\pi \times \tilde{\pi}'}^{a-\frac{1}{2}}}{\pi} \left(\frac{a}{\pi} \right)^{dmm'(a-\frac{1}{2})} \left(\sum_{n=1}^{+\infty} \frac{|a_{\pi \times \tilde{\pi}'}(n)|}{n^a} \right).$$

PROOF. From the integral representation of generalized Euler-Stieltjes coefficients given in Proposition 4.1, and using the bound obtained in Lemma 5.1, we have

$$|\gamma_{\pi, \pi'}(k)| \leq C_L(a) \frac{Q_{\pi \times \tilde{\pi}'}^{a-\frac{1}{2}}}{2\pi} \int_{-\infty}^{+\infty} \left[\left(\frac{1+a+\mu_R}{2} \right)^2 + \left(\frac{|t|+\mu_I}{2} \right)^2 \right]^{dmm' \frac{2a-1}{4}} \times \frac{|L(a-it, \pi \times \tilde{\pi}')|}{(a^2 + (t+t_0)^2)^{\frac{k+1}{2}}} dt + \delta(t_0) \left| \operatorname{Res}_{s=it_0} L(s, \pi \times \tilde{\pi}') \right|,$$

where $C_L(a)$ is defined in Lemma 5.1.

Since the Rankin-Selberg L -function possesses a Dirichlet series representation (2.2) that converges absolutely for $\operatorname{Re} s > 1$, for $a > 1 + \varepsilon > 1$, one yields

$$\left| \overline{L(a-it, \pi \times \tilde{\pi}')} \right| \leq \sum_{n=1}^{+\infty} \frac{|a_{\pi \times \tilde{\pi}'}(n)|}{n^a} < +\infty,$$

hence

$$(5.8) \quad |\gamma_{\pi, \pi'}(k)| \leq C_L(a) \frac{Q_{\pi \times \tilde{\pi}'}^{a-\frac{1}{2}}}{2\pi} \sum_{n=1}^{+\infty} \frac{|a_{\pi \times \tilde{\pi}'}(n)|}{n^a} I + \delta(t_0) \left| \operatorname{Res}_{s=it_0} L(s, \pi \times \tilde{\pi}') \right|,$$

where

$$I = \int_{-\infty}^{+\infty} \left[\left(\frac{1+a+\mu_R}{2} \right)^2 + \left(\frac{|t|+\mu_I}{2} \right)^2 \right]^{dmm' \frac{2a-1}{4}} \frac{dt}{(a^2 + (t+t_0)^2)^{\frac{k+1}{2}}}.$$

Thus, it is left to derive a bound for the integral I . Depending on the value of t_0 , we examine two cases.

(i) Let $t_0 \geq 0$. Then

$$(5.9) \quad I = \int_0^{+\infty} \left(\frac{1}{(a^2 + (t + t_0)^2)^{\frac{k+1}{2}}} + \frac{1}{(a^2 + (t - t_0)^2)^{\frac{k+1}{2}}} \right) \\ \times \left[\left(\frac{1 + a + \mu_R}{2} \right)^2 + \left(\frac{t + \mu_I}{2} \right)^2 \right]^{dmm' \frac{2a-1}{4}} dt.$$

The interval of integration we derive into two parts. Denote by I_1 and I_2 integrals that correspond to intervals $(0, B)$ and $(B, +\infty)$, respectively, where $B = 1 + a + \mu_{R,I} - \mu_I > t_0 + 1$.

For I_1 we have

$$(5.10) \quad I_1 \leq 2(2 + \mu_{R,I} + \mu_I) 8^{dmm' \frac{2a-1}{4}} a^{-k + \frac{2a-1}{2} dmm'},$$

since $1 + a + \mu_R \leq 1 + a + \mu_{R,I} < 4a$ and $\frac{B}{a} \leq 2 + \mu_{R,I} + \mu_I$, by assumptions of the theorem.

For integral I_2 , we have $t \geq B$,

$$\left(\frac{1 + a + \mu_R}{2} \right)^2 + \left(\frac{t + \mu_I}{2} \right)^2 \leq 2 \left(\frac{t + \mu_I}{2} \right)^2,$$

and $(t + t_0)^2 \geq (t - t_0)^2$, so

$$I_2 \leq \int_B^{+\infty} \frac{2}{(a^2 + (t - t_0)^2)^{\frac{k+1}{2}}} \left[2 \left(\frac{t + \mu_I}{2} \right)^2 \right]^{dmm' \frac{2a-1}{4}} dt \\ \leq \int_{B-t_0}^{+\infty} \left(\frac{t + t_0 + \mu_I}{t} \right)^{k+1} \frac{2^{1-dmm' \frac{2a-1}{4}}}{(t + t_0 + \mu_I)^{k+1}} (t + t_0 + \mu_I)^{dmm' \frac{2a-1}{2}} dt.$$

Furthermore, since the function $g(t) = \frac{t+t_0+\mu_I}{t}$ is monotonically decreasing for $t \geq B - t_0$, $g(t) > 1$ and $\lim_{t \rightarrow +\infty} g(t) = 1$, it follows that maximal value of $g(t)$ is at point $t = B - t_0$ and it is equal to $\frac{B+\mu_I}{B-t_0}$. Hence,

$$I_2 \leq \left(\frac{B + \mu_I}{B - t_0} \right)^{k+1} 2^{1-dmm' \frac{2a-1}{4}} \\ \times \int_{B-t_0}^{+\infty} (t + t_0 + \mu_I)^{-(k+1) + dmm' \frac{2a-1}{2}} dt.$$

For constant a under consideration, we have $a < \frac{1}{2} + \frac{k}{dmm'}$, thus the above integral converges and yields

$$I_2 \leq \frac{2^{1-dmm' \frac{2a-1}{4}}}{k - dmm' \frac{2a-1}{2}} \frac{(1 + a + \mu_{R,I})^{1+dmm' \frac{2a-1}{2}}}{(1 + a + \mu_{R,I} - \mu_I - t_0)^{k+1}}.$$

Additionally, since $\mu_{R,I} = \max\{\mu_R, \mu_I + t_0 - 1\}$ inequalities $1 + a + \mu_{R,I} - \mu_I - t_0 > a > 1 + \varepsilon > 1$ hold true. Also, $1 + a + \mu_{R,I} < 4a$. Thus

$$(5.11) \quad I_2 \leq \frac{8^{1+dm} m'^{\frac{2a-1}{4}}}{k - dmm'^{\frac{2a-1}{2}}} a^{-k+dm} m'^{\frac{2a-1}{2}}.$$

Substituting (5.10) and (5.11) into (5.9), combined with (5.8) implies (5.7).

(ii) The result for the case $t_0 < 0$ can be derived completely analogously as in (i) using simple substitution $-t_0 = t_1 > 0$.

The proof is complete. \square

ACKNOWLEDGEMENTS.

I would like to thank the anonymous referee for his/her comments and suggestions that helped me to improve the paper.

REFERENCES

- [1] J. A. Adell, *Asymptotic estimates for Stieltjes constants: a probabilistic approach*, Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci. **467** (2011), 954–963.
- [2] M. Avdispahić and L. Smajlović, *Euler constants for a Fuchsian group of the first kind*, Acta Arith. **131** (2008), 125–143.
- [3] B. C. Berndt, *On the Hurwitz zeta-function*, Rocky Mountain J. Math. **2** (1972), 151–157.
- [4] E. Bombieri and J. C. Lagarias, *Complements to Li's criterion for the Riemann hypothesis*, J. Number Theory **77** (1999), 274–287.
- [5] W. E. Briggs, *Some constants associated with the Riemann zeta-function*, Michigan Math. J. **3** (1955/56), 117–121.
- [6] W. E. Briggs and S. Chowla, *The power series coefficients of $\zeta(s)$* , Amer. Math. Monthly **62** (1955), 323–325.
- [7] T. Chatterjee and S. Garg, *On q -analogue of Euler-Stieltjes constants*, Proc. Amer. Math. Soc. **151** (2023), 2011–2022.
- [8] M. W. Coffey, *New results on the Stieltjes constants: asymptotic and exact evaluation*, J. Math. Anal. Appl. **317** (2006), 603–612.
- [9] M. W. Coffey, *The Stieltjes constants, their relation to the η_j coefficients, and representation of the Hurwitz zeta function*, Analysis (Munich) **30** (2010), 383–409.
- [10] M. W. Coffey, *Series representations for the Stieltjes constants*, Rocky Mountain J. Math. **44** (2014), 443–477.
- [11] J. W. Cogdell, *L-functions and converse theorems for GL_n* , in: Automorphic forms and applications, Amer. Math. Soc., Providence, RI, 2007, 97–177.
- [12] L. Euler, *De progressionibus harmonicis observationes*, Comment. acad. sci. Petrop. **7** (1740), 150–161. (Opera Omnia, Series 1, Vol. 14, 87–100.)
- [13] T. Gillespie and G. Ji, *Prime number theorems for Rankin-Selberg L-functions over number fields*, Sci. China Math. **54** (2011), 35–46.
- [14] Y. Hashimoto, *The Euler-Selberg constants for non-uniform lattices of rank one symmetric spaces*, Kyushu J. Math. **57** (2003), 347–370.
- [15] Y. Hashimoto, Y. Iijima, N. Kurokawa and M. Wakayama, *Euler's constants for the Selberg and the Dedekind zeta functions*, Bull. Belg. Math. Soc. Simon Stevin **11** (2004), 493–516.
- [16] C. Hermite and T. J. Stieltjes, *Correspondance d'Hermite et de Stieltjes*, edited by B. Baillaud and H. Bourget, Gauthier-Villars, Paris, (1905).

- [17] S. Inoue, S. Saad Eddin and A. I. Suriajaya, *Stieltjes constants of L -functions in the extended Selberg class*, Ramanujan J. **55** (2021), 609–621.
- [18] H. Jacquet and J. A. Shalika, *On Euler products and the classification of automorphic representations. I*, Amer. J. Math. **103** (1981), 499–558.
- [19] J. Kaczorowski and A. Perelli, *On the structure of the Selberg class. I. $0 \leq d \leq 1$* , Acta Math. **182** (1999), 207–241.
- [20] C. Knessl and M. W. Coffey, *An asymptotic form for the Stieltjes constants $\gamma_k(a)$ and for a sum $S_\gamma(n)$ appearing under the Li criterion*, Math. Comp. **80** (2011), 2197–2217.
- [21] Y. L. Luke, *The special functions and their approximations*. Academic Press, New York-London, 1969.
- [22] K. Matsumoto, T. Onozuka and I. Wakabayashi, *Laurent series expansions of multiple zeta-functions of Euler-Zagier type at integer points*, Math. Z. **295** (2020), 623–642.
- [23] K. Maślanka, *Li's criterion for the Riemann hypothesis—numerical approach*, Opuscula Math. **24** (2004), 103–114.
- [24] Y. Matsuoka, *Generalized Euler constants associated with the Riemann zeta function*, in: Number theory and combinatorics. Japan 1984 (Tokyo, Okayama and Kyoto, 1984), World Sci. Publishing, Singapore, 1985, 279–295.
- [25] M. R. Murty, *Problems in analytic number theory*, Springer-Verlag, New York, 2001.
- [26] A. Odžak and L. Smajlović, *On Li's coefficients for the Rankin-Selberg L -functions*, Ramanujan J. **21** (2010), 303–334.
- [27] A. Odžak and L. Smajlović, *Euler-Stieltjes constants for the Rankin-Selberg L -function and weighted Selberg orthogonality*, Glas. Mat. Ser. III **51(71)** (2016), 23–44.
- [28] A. Odžak and L. Smajlović, *On the generalized Euler-Stieltjes constants for the Rankin-Selberg L -function*, Int. J. Number Theory **13** (2017), 1363–1379.
- [29] M. Overholt, *A course in analytic number theory*, American Mathematical Society, Providence, RI, 2014.
- [30] M. Prévost and T. Rivoal, *New convergent sequences of approximations to Stieltjes' constants*, J. Math. Anal. Appl. **524** (2023), Paper No. 127091, 20.
- [31] Z. Rudnick and P. Sarnak, *Zeros of principal L -functions and random matrix theory*, 1996, 269–322.
- [32] S. Saad Eddin, *Explicit upper bounds for the Stieltjes constants*, J. Number Theory **133** (2013), 1027–1044.
- [33] S. Saad Eddin, *The signs of the Stieltjes constants associated with the Dedekind zeta function*, Proc. Japan Acad. Ser. A Math. Sci. **94** (2018), 93–96.
- [34] B. Saha, *Multiple Stieltjes constants and Laurent type expansion of the multiple zeta functions at integer points*, Selecta Math. (N.S.) **28** (2022), Paper No. 6, 41.
- [35] F. Shahidi, *On nonvanishing of L -functions*, Bull. Amer. Math. Soc. (N.S.) **2** (1980), 462–464.
- [36] F. Shahidi, *Fourier transforms of intertwining operators and Plancherel measures for $GL(n)$* , Amer. J. Math. **106** (1984), 67–111.
- [37] F. Shahidi, *Local coefficients as Artin factors for real groups*, Duke Math. J. **52** (1985), 973–1007.
- [38] F. Shahidi, *A proof of Langlands' conjecture on Plancherel measures; complementary series for p -adic groups*, Ann. of Math. (2) **132** (1990), 273–330.
- [39] L. Smajlović, *On Li's criterion for the Riemann hypothesis for the Selberg class*, J. Number Theory **130** (2010), 828–851.
- [40] E. C. Titchmarsh, *The theory of functions*, 2nd ed., Oxford University Press, Oxford, 1958.
- [41] E. T. Whittaker, G. N. Watson, *A Course of Modern Analysis*, Cambridge University Press, Cambridge, 1902.

M. Zubača

Department of Mathematical and Computer Sciences, University of Sarajevo

Zmaja od Bosne 35, 71 000 Sarajevo

Bosnia and Herzegovina

E-mail : medina.zubaca@pmf.unsa.ba

Received: 13.7.2023.

Revised: 8.12.2023.

O OGRANIČENOSTI EULER-STIELTJESOVIH KONSTANTI ZA RANKIN-SELBERGOVU L -FUNKCIJU

MEDINA ZUBAČA

SAŽETAK. Neka je E Galoisovo proširenje od \mathbb{Q} konačnog stupnja i neka su π i π' dvije ireducibilne automorfne unitarne kupidalne reprezentacije od $GL_m(\mathbb{A}_E)$ i $GL_{m'}(\mathbb{A}_E)$, redom. Neka je $\Lambda(s, \pi \times \tilde{\pi}')$ Rankin-Selbergova L -funkcija pridružena produktu $\pi \times \tilde{\pi}'$, gdje $\tilde{\pi}'$ označava kontragradijentu reprezentaciju od π' , a neka njegov konačni dio (bez Arhimedovih faktora) bude $L(s, \pi \times \tilde{\pi}')$. Euler-Stieltjesove konstante Rankin-Selbergove L -funkcije su koeficijenti u razvoju u Laurentov (Taylorov) red oko $s = 1 + it_0$ funkcije $L(s, \pi \times \tilde{\pi}')$. U ovom radu izvodimo gornju među ovih konstanti.