

# BENTHIC MACROINVERTEBRATE COMMUNITIES CHANGE ALONG A KARSTIC BARRAGE-LAKE SYSTEM

**SANJA SVIBEN<sup>1</sup>, RENATA MATONIČKIN KEPČIJA<sup>1</sup>, MARINA VILENICA<sup>2,\*</sup>,  
IVANČICA KRULIK<sup>1</sup>, VLATKA MIČETIĆ STANKOVIĆ<sup>3</sup>, PETAR KRUŽIĆ<sup>1</sup>  
& MLADEN KUČINIĆ<sup>1</sup>**

<sup>1</sup>University of Zagreb, Faculty of Science, Department of Biology, Horvatovac 102a, 10 000 Zagreb, Croatia (e-mails: sanjasviben@gmail.com; renata.matonickin.kepcija@biol.pmf.hr; ivancica.krulik@biol.pmf.hr; petar.kruzic@biol.pmf.hr; mladen.kucinic@biol.pmf.hr)

<sup>2</sup>University of Zagreb, Faculty of Teacher Education, Trg Matice Hrvatske 12, 44250 Petrinja, Croatia (e-mail: marina.vilenica@ufzg.hr)

<sup>3</sup>Croatian Natural History Museum, Demetrova 1, 10000 Zagreb, Croatia (e-mail: vlatka.micetic-stankovic@hpm.hr)

**Sviben, S., Matoničkin Kepčija, R., Vilenica, M., Krulik, I., Mičetić Stanković, V., Kružić, P. & Kučinić, M.: Benthic macroinvertebrate communities change along a karstic barrage-lake system. Nat. Croat., Vol. 33, No. 1., 105-121, 2024, Zagreb.**

Barrage-lake systems are a rare natural phenomenon with specific ecological characteristics that are greatly affected by tufa formation. Our main goal was to investigate benthic macroinvertebrate communities and their longitudinal variations within the lakes arranged in cascades in the Dinaric karst hydrosystem. Each lake within the Plitvice Lakes system studied presented with a distinct community, showing the effects of the barrage-lake system on the macroinvertebrates in these lakes. We observed a downstream decrease in taxa richness, abundance and Shannon diversity index along the longitudinal profile of the studied barrage-lake system. This pattern is most probably a result of decreased organic matter and increased tufa deposition. Overall, our results suggest that the hydrosystem studied represents a sequence of sediment sinks in which lakes retain and accumulate organic matter and fauna. Therefore, each lake acts as a filtering screen and consequently appears to be a small ecosystem. Each lake has its own community characteristics, regardless of the fact that water flows from one lake to the next, connecting them and creating very similar abiotic conditions in each lake along the system.

**Key words:** Dinaric karst, tufa deposition, barrage lakes, retention effect, macrozoobenthos

**Sviben, S., Matoničkin Kepčija, R., Vilenica, M., Krulik, I., Mičetić Stanković, V., Kružić, P. & Kučinić, M.: Zajednice bentičkih makrobekralješnjaka u kaskadnom jezerskom sustavu. Nat. Croat., Vol. 33, No. 1., 105-121, 2024, Zagreb.**

Kaskadni jezerski sustavi rijedak su prirodni fenomen obilježen specifičnim ekološkim značajkama na koje uvelike utječe stvaranje sedre. Glavni cilj ovog rada bio je istražiti zajednice bentičkih makrobekralješnjaka i njihove razlike duž kaskadnog jezerskog kompleksa u hidrološkom sustavu Dinarskog krša. Svako jezero unutar istraživanog dijela kompleksa Plitvičkih jezera ima specifičnu zajednicu bentičkih makrobekralješnjaka, što je posljedica učinka kaskadnog sustava jezera na makrozoobentos. Uzduž longitudinalnog profila jezerskog kompleksa zabilježeno je smanjenje bogatstva svojiti, brojnosti jedinki i Shannonovog indeksa raznolikosti, što je vjerojatno posljedica smanjenja količine organske tvari i povećanja taloženja sedre. U ovakvim kaskadnim jezerskim sustavima jezera zadržavaju i akumuliraju organsku tvar i faunu zbog čega je svako jezero naseljeno specifičnim zajed-

\*Corresponding author

nicama makrobekralješnjaka, bez obzira na to što voda teče iz jednog jezera u drugo, povezuje ih i osigurava vrlo slične abiotičke uvjete u svakom jezeru duž istraživanog jezerskog sustava.

**ključne riječi:** dinarski krš, taloženje sedre, kaskadna jezera, retencijski učinak, makrozoobentos

## INTRODUCTION

Despite the considerable amount of research on the process of freshwater carbonate precipitation and tufa formation (e.g. SRDOČ *et al.*, 1985; EMEIS *et al.*, 1987; CHAFETZ *et al.*, 1991; FORD & PEDLEY, 1996; BELLINZONI *et al.*, 2003), the components involved in these processes are still the subject of research. Generally, tufa is a porous sedimentary rock typical of carbonate rich areas where the spring waters appear supersaturated with calcium carbonate. Restoring the chemical equilibrium, some of the calcium carbonate is precipitated from the solution (GOLUBIĆ, 1969; RIDING, 1991). This results in tufa deposition, and the initiation of this process appears to be result of three factors – the biological, geological and chemical (micro) environments (EMEIS *et al.*, 1987; FORD & PEDLEY, 1996). Organisms have an inevitable role (CHAFETZ *et al.*, 1994) since they provide a substrate for calcite nucleation which accelerates tufa deposition (KEMPE & EMEIS, 1985; MERZ-PREISS & RIDDING, 1999; MATONIČKIN KEPČIJA *et al.*, 2006). A wide range of organisms is associated with tufa, as it is a suitable substratum that provides shelter and food in the form of encrusted algae and detritus for invertebrates (PENTECOST, 2005). Nevertheless, intensive carbonate deposition and encrustation create unfavorable conditions for these organisms (GOLUBIĆ, 1969; EMEIS *et al.*, 1987), which could have negative effects on macroinvertebrate abundance and community structure (RUNDIO, 2009). Many invertebrates, especially insects, have evolved peculiar adaptations which enable them to exist in these extreme conditions (PENTECOST, 2005).

Tufa deposits are quite common, yet the development of tufa dams which lead to the differentiation of a river profile into a series of lakes and waterfalls is a rare phenomenon (GOLUBIĆ, 1969). As a result of this process, barrage-lake systems are formed, such as the Ruidera Pools Natural Park (Spain), Plitvice Lakes National Park (Croatia), d'Immouzer and Ida du Tanane (Marocco), Turner Falls (USA) and tufa springs in Queensland (Australia) (FORD & PEDLEY, 1996). In a stream containing tufa dams, organic matter tends to accumulate, resulting in tight spirals, delaying nutrient loss (PENTECOST, 2005). So far, the effects of the dams have not been investigated in terms of nutrient retention (PENTECOST, 2005), however, a significant decrease in the organic matter downstream the Plitvice Lakes barrage-system in Croatia has already been observed (OBELIĆ *et al.*, 2005; HORVATINČIĆ *et al.*, 2006; MILIŠA *et al.*, 2006). Also, according to the developmental scheme of a barrage system proposed by GOLUBIĆ (1969), tufa deposition should be increasing downstream. For now, Plitvice Lakes are the only barrage system confirming both these hypotheses (SRDOČ *et al.*, 1985) which makes them very interesting in the context of ecological research.

The Plitvice Lakes are a karstic hydrosystem with well-defined hydrobiological (GOLUBIĆ, 1969; EMEIS *et al.*, 1987), hydrological (Petrik, 1958), mineralogical (STOFFERS, 1975), chemical (KEMPE & EMEIS, 1985; FRANČIŠKOVIĆ-BILINSKI *et al.*, 2004) and ecological (ŠEGULJA & HRŠAK, 1994; HABDIJA *et al.*, 2004; MILIŠA *et al.*, 2014) characteristics. In this study, we aimed to determine whether the organisms within the tufa barrage system would show distinctive quantitative or qualitative downstream patterns. Macrozoobenthos provides an excellent opportunity to address this type of question in


community ecology due to its interactions with tufa. Despite the increasing amount of research conducted in the area of Plitvice Lakes National Park (NP) during the last two decades (e.g. POPIJAČ & SIVEC, 2009; IVKOVIĆ *et al.*, 2012, 2014; ŠEMNIČKI *et al.*, 2012; KUČINIĆ *et al.*, 2008; IVKOVIĆ & PONT, 2016; VILENICA *et al.*, 2014; 2017; MIČETIĆ STANKOVIĆ *et al.*, 2019), macroinvertebrates within the lakes have not yet been investigated in detail. Previous studies were mainly focused on the organisms living on the barriers where calcium precipitation is very pronounced due to water splashing and aeration (e.g. ŠPOLJAR *et al.*, 1994; PRIMC-HABDIJA *et al.*, 2001; MILIŠA *et al.*, 2006; 2014). Thus, we sampled the littoral zone, which in comparison to other lake zones supports larger and more diverse populations of benthic invertebrates (WIEDERHOLM, 1984) due to substrate heterogeneity (STRAYER, 1985). Having in mind the complexity of barrage systems, the main goals of this study were to determine: a) benthic macroinvertebrate community composition and its seasonal variations within the lakes, b) influence of different substrate types on the abundance and distribution of benthic macroinvertebrates and c) quantitative and qualitative patterns of benthic macroinvertebrates downstream the barrage-lake system of the Plitvice Lakes.

## METHODS

### Study area

The Plitvice Lakes are located in the karstic region of the NW Dinaric Mountains in Croatia (Fig. 1). This natural system is composed of 16 barrage-lakes where water flows from one lake to the next over tufa barriers, which are the main factor in creating waterfalls and cascades (FORD & PEDLEY, 1996). Water splashes in the waterfalls, stimulating the extraction of carbon dioxide and sedimentation of calcium carbonate from the water. The overall result is the growth of tufa barriers, causing a rise in the water level (ZWICKER & RUBINIĆ, 2005). The lakes receive water from the two main suppliers, Bijela rijeka and Crna rijeka rivers, which join into the Matica River, and two tributaries, the Rječica and Plitvica streams. The altitude of the area ranges between 636 and 503 m a.s.l. The lake system is 9.05 km long and because of its geomorphological features, it is divided into 12 upper and four lower lakes (RIĐANOVIĆ & BOŽIČEVIĆ, 1996). Upper lakes are located in a dolomite valley while lower lakes flow through a rudist limestone canyon, where the last lake drains into the karst lowland Korana River (RIĐANOVIĆ & BOŽIČEVIĆ, 1996).

In our study, we investigated four lakes: Prošće, Ciginovac, Kozjak and Kaluđerovac. Their characteristics are given in Tab. 1. The first three lakes, Prošće, Ciginovac and Kozjak, are part of the upper section while the fourth, Kaluđerovac Lake, is a part of the lower section. Prošće is the first lake in the barrage system. Progressive filling with sediment and aquatic vegetation was recorded there 40 years ago (IVEKOVIĆ, 1971). At the outlet, water from the Prošće Lake flows partially into Ciginovac Lake and partially into Okrugljak Lake. Some of the water that flows into Ciginovac Lake is probably drained under the ground through the very porous tufa barrier (BERAKOVIĆ, 2005). Kozjak Lake is the last in the string of upper lakes. It is the largest lake, and its deepest point is 46 m, which makes it also the deepest lake in the hydrosystem. Except for a few perennial streams and temporary flows, the watercourse that mostly affects the water regime of Kozjak Lake is the Rječica Stream (IVEKOVIĆ, 1971). Kaluđerovac Lake is specific because of its pronounced tufa deposition, greater than that of other lakes.



**Fig. 1.** a) Study area and location of the sampling sites, b) and c) Lakes divided by tufa barriers in Plićvice Lakes barrage-system. (Photo: P. Kružić and R. Matonićkin Kepčija).

Biologically induced calcite precipitation is probably enhanced due to the increase in primary production (HORVATINČIĆ *et al.*, 2008).

### Sampling protocol

At each lake, one littoral station was selected near forested banks at depth not more than one metre. To determine macroinvertebrate abundance and taxa richness, samples were collected monthly from April 2007 to March 2008. Sampling points were characterized with respect to the most dominant substrates (Tab. 1) in the littoral zone of each lake. In Prošće Lake, we sampled *Chara* sp. and mosaic substrate sand/leaf litter, and in Ciginovac Lake cobbles and leaf litter were sampled. In the Kozjak Lake, macroinvertebrate communities were sampled on *Chara* sp. and leaf litter. The algae were sampled over a period of only eight months, as it vanished during October, November, December and January, when it was replaced with sandy substrate. Thus, during those four months, sand was sampled instead. In Kaluđerovac Lake, we sampled only the

**Tab. 1.** Main characteristics of the four studied lakes in the Plitvice Lakes barrage-system (P = Prošće, C = Ciginovac, Ko = Kozjak, Ka = Kaluđerovac).

Lake characteristics	P	C	Ko	Ka
Elevation (m a.s.l.)	637	620	534	505
Mean depth (m)	37	11	46	13
Length (m)	2100	470	2350	225
Width (m)	180-400	120	135-670	70-100
Surface area (km <sup>2</sup> )	0.68	0.07	0.83	0.02
Dominant riparian vegetation:				
<i>Fagus sylvatica</i> L.	+	+	+	
<i>Carpinus betulus</i> L.				+
Substrate type:				
<i>Chara</i> sp.	+		+	
Cobbles		+		
Leaf litter		+	+	
Sand			+	
Cobbles/Leaf litter				+
Sand/leaf litter	+			

mosaic combination of two substrates, leaf litter and cobbles. To equalize the area sampled in each lake, we collected duplicates of microhabitat with leaf litter and cobbles in Kaluđerovac Lake during each sampling event.

Macroinvertebrates were collected using a D-frame net sampler with a surface area of 0.1 m<sup>2</sup> and mesh size 500 µm. Sampled substrates were carefully washed in the field. Material was placed in plastic containers and preserved in 96 % ethanol. In the laboratory, 24 hours after sampling, material was transferred to 80 % ethanol. Macroinvertebrates were sorted, counted, identified under a stereomicroscope and stored in 70 % ethanol. According to the identification keys (e.g. ENGBLOM, 1996; MEINANDER, 1996; NILSSON *et al.*, 1996; GERKEN & STERNBERG, 1999; WARINGER & GRAF, 1997; 2011; BAUERNEFIND & HUMPESECH, 2001; HÖLZEL, 2002; JÄCH & BROJER, 2006) benthic macroinvertebrates were identified to the lowest possible taxonomic level with a focus on insect larvae.

Analysis of physico-chemical water properties was conducted according to the procedures in APHA (1998). The following physico-chemical water properties were measured at each sampling event: water temperature, dissolved oxygen concentration (using an oximeter (WTW Oxi 330/SET), pH (using a pH-meter (WTW ph 330), conductivity (using a conductometer (WTW LF 330)), nutrients (nitrate levels (by chromotropic acid method), nitrites (by sulfanilic acid and 1-Naphthylamine in the colorimetric method), ortho-phosphates (by the molybdophosphoric acid method) and ammonia (by salicylic acid)). Although the ortho-phosphate and nitrite levels were measured during the entire study period, the sensitivity of the method was too low for their detection.

## Data analysis

The benthic macroinvertebrate abundance, taxa richness and Shannon diversity index were calculated for each investigated lake.

To examine similarities in benthic macroinvertebrate communities between different lakes and different substrates, NMDS analyses were conducted. The analyses were based on the monthly samples and were conducted using a Bray-Curtis resemblance coefficient. Prior to analysis, all data were  $\log(X+1)$ -transformed. Samples without macroinvertebrate records were excluded from the analyses. Shannon diversity and NMDS analyses were performed in the software package PRIMER 6 (CLARKE & GORLEY, 2006).

## RESULTS

### Physico-chemical water properties

Mean water temperature ranged between 11.6°C in Prošće Lake and 14.2°C in Kozjak Lake. Mean concentration of dissolved oxygen in water ranged between 9.8 mg L<sup>-1</sup> in Kozjak Lake and 11.6 mg L<sup>-1</sup> in Prošće Lake. Mean value of pH was 8.2 in Prošće and Ciginovac lakes, and 8.4 in Kozjak and Kaluđerovac lakes. Conductivity ranged between 366  $\mu\text{S cm}^{-1}$  in Kaluđerovac Lake and 405  $\mu\text{S cm}^{-1}$  in Prošće Lake. Mean concentration of nitrates in water ranged between 0.5 mg L<sup>-1</sup> in Kaluđerovac Lake and 0.8 mg L<sup>-1</sup> in Prošće Lake. Mean concentration of ammonia was higher in Kaluđerovac Lake (0.06 mg L<sup>-1</sup>) than in the other three lakes (Tab. 2).

**Tab. 2.** Physico-chemical water properties measured from April 2007 to March 2008 in the four lakes of the Plitvice Lakes barrage-system presented as mean values  $\pm$  SD.

Abiotic parameter	Prošće	Ciginovac	Kozjak	Kaluđerovac
Water temperature (°C)	11.6 $\pm$ 7.0	12.9 $\pm$ 7.7	14.2 $\pm$ 7.9	13.0 $\pm$ 7.9
Dissolved oxygen (mg O <sub>2</sub> L <sup>-1</sup> )	11.6 $\pm$ 1.7	10.9 $\pm$ 1.3	9.8 $\pm$ 1.6	10.5 $\pm$ 1.7
pH	8.2 $\pm$ 0.5	8.2 $\pm$ 0.5	8.4 $\pm$ 0.2	8.4 $\pm$ 0.1
Conductivity ( $\mu\text{S cm}^{-1}$ )	405 $\pm$ 28	383 $\pm$ 256	382 $\pm$ 39	366 $\pm$ 22
Nitrates (mg N L <sup>-1</sup> )	0.8 $\pm$ 0.1	0.6 $\pm$ 0.1	0.6 $\pm$ 0.2	0.5 $\pm$ 0.3
Ammonia (mg N L <sup>-1</sup> )	0.02 $\pm$ 0.01	0.03 $\pm$ 0.02	0.02 $\pm$ 0.01	0.06 $\pm$ 0.10

### Macroinvertebrate community composition

In all, 16 benthic macroinvertebrate taxa groups were recorded in the studied four lakes (Appendix 1). Altogether 26082 individuals of benthic macroinvertebrates were collected and analysed. Overall, the most abundant macroinvertebrates were Diptera larvae, with Chironomidae as the numerically dominant group. High frequency of Ephemeroptera, especially *Caenis horaria* (Linnaeus, 1758), was recorded only in Prošće Lake. Mollusca, with the most abundant representative *Pisidium* sp., were the second most abundant taxon in Ciginovac Lake, where a high abundance of Isopod *Asellus aquaticus* (Linnaeus, 1758) was also recorded. Macroinvertebrate community compositions of the Kozjak and Kaluđerovac lakes were similar, with Diptera and Oligochaeta as dominant groups.

In the studied lakes, Plecoptera were the least abundant insect group. Ten Odonata taxa and four species of Megaloptera were recorded. Trichoptera larvae were infrequent, but they showed the highest richness with 13 recorded genera. Heteroptera

were recorded in a high abundance only in the first lake in the hydrosystem, Prošće Lake. They were also present sporadically in Ciginovac Lake, while they were completely absent from the other two lakes. Abundance of Hirudinea was also decreasing downstream the barrage-lake system, being the most abundant in Prošće Lake and completely absent from Kaluđerovac Lake.

### Seasonal variations in the community composition

Generally, macroinvertebrate abundance was the highest in winter in Prošće Lake, while the lowest value was also recorded in winter, but in Kaluđerovac Lake. During all seasons, Dipteran larvae were a dominant taxon (Fig. 2).

In the upper lakes section (i.e. Prošće, Ciginovac and Kozjak lakes), the abundance of benthic macroinvertebrates was reduced downstream the barrage system during spring and autumn. During summer and winter, a decrease in the abundance was observed along the whole longitudinal profile (Fig. 2).




Fig. 2. Seasonal variations of the benthic macroinvertebrate communities at investigated sites in the Plitvice Lakes barrage-system. For the abbreviations of the study sites see Tab. 1. Abundance is expressed as number of individuals per m<sup>2</sup>.

### Changes in communities of benthic macroinvertebrates along a barrage-lake hydrosystem

Taxon richness and Shannon diversity index decreased downstream the hydro-system (Tab. 3). The abundance showed a similar pattern in the downstream decrease, except for Kaluđerovac Lake, where we recorded higher values than in Kozjak Lake.


Among the investigated substrates, the highest taxa richness was recorded on *Chara* sp. and mosaic substrate sand/leaf litter in Prošće Lake, and the lowest on sand in Kozjak Lake (Tab. 3).

According to the NMDS analyses (Fig. 3), macroinvertebrate communities of each lake grouped together. Macroinvertebrate communities on different substrates in each lake were more similar to each other than macroinvertebrate communities on the same substrates in different lakes.



**Tab. 3.** Benthic macroinvertebrate taxa richness (S), abundance (N) and Shannon diversity index (H') in the four lakes and taxa richness (S) at various substrates in the Plitvice Lakes barrage-system. Abundance is expressed as number of collected individuals during the study.

Lake	S	N	H'	Substrate	S
Prošće	51	15762	1.81	<i>Chara</i> sp.	46
				Sand/Leaf litter	40
Ciginovac	38	4783	1.59	Cobbles	34
				Leaf litter	33
Kozjak	33	2306	1.40	<i>Chara</i> sp.	24
				Leaf litter	24
				Sand	9
Kaluđerovac	32	3231	1.36	Cobbles/Leaf litter	32



**Fig. 3.** NMDS analysis of the monthly macroinvertebrate samples at different substrate types in four lakes in the Plitvice Lakes barrage-system based on: a) each lake and b) substrate type.



## DISCUSSION

The chemistry of carbonate deposition in freshwater habitats is relatively well understood, yet the biological factors included in this process are still intriguing and a subject of research (e.g. BAYARI & KURTTAS, 1997; DRYSDALE, 1999; PENTECOST & ZHANG, 2000; BELLINZONI *et al.*, 2003; GOLUBIĆ *et al.*, 2008). Our study confirmed the results of previous studies, which reported a wide range of organisms associated with harsh environment in tufa-depositing habitats (PENTECOST, 2005). Of all freshwater invertebrates, the insects are generally the most diverse, and tufa-depositing systems like the Plitvice Lakes are no exception (PENTECOST, 2005). As expected, Diptera dominated in all four investigated lakes (MINCKLEY, 1963; HUMPHREYS *et al.*, 1995; CARTHEW *et al.*, 2003) due to their wide ecological preferences in terms of feeding behaviour, habitat and microhabitat selection as well as their tolerance to the environmental factors (MALMQVIST *et al.*, 1999; EDWARDS & USHER, 1985; SHIMADA *et al.*, 1991; IVKOVIĆ & PONT, 2016). High taxa richness of Megaloptera recorded in the Plitvice Lakes was previously reported from carbonate-depositing ecosystems (LEONARD, 1939), where they contribute to the maintenance of the populations structure as predators (DIEHL, 1995). On the other hand, rather low taxa richness of predatory Odonata could be a result of a limitations of a sampling methodology used, as the sampling net collects their relatively large nymphs from a rather small surface area (e.g. HORNING & POLLARD, 1978). Trichoptera larvae represent a popular subject in studies focused on tufa diagenesis (DURRENFELDT, 1978; DRYSDALE, 1999; PAPROCKI *et al.*, 2003; HABIJA *et al.*, 2004) as their biological structures are often found fossilized within tufa (PENTECOST, 2005). In the Plitvice Lakes hydrosystem, Trichoptera were abundant only in the area of tufa barriers and water suppliers (PREVIŠIĆ *et al.*, 2007; ŠEMNIČKI *et al.*, 2012), characterized by high water flow, compared to the lentic conditions in lakes. Although infrequent in lakes, their rather high taxa richness was not surprising, since the Balkan Peninsula was already recognized as an important evolutionary centre for a number of trichopteran genera (e.g. KUMANSKI & MALICKY, 1999; KUČINIĆ *et al.*, 2015; PREVIŠIĆ *et al.*, 2014; VITECEK *et al.*, 2015). As the majority of mayfly species prefer rhithral sections of lotic habitats (BAUERNEFEIND & SOLDÁN, 2012), relatively low diversity was recorded in the studied lakes. The presence of species such as *Caenis horaria* is generally believed to be an important environmental indicator of oligotrophic to mesotrophic conditions (BAUERNEFEIND & MOOG, 2000). Although the lakes' trophic state is important feature for Ephemeroptera assemblages, presence of rare species like *Leptophlebia vespertina* (Linnaeus, 1758) and *Paraleptophlebia werneri* Ulmer, 1920 (VILENICA *et al.*, 2015) seem to depend more on special conditions in the context of carbonate depositing area than on trophic state (e.g. MENETREY *et al.*, 2008). Very low abundance of Plecoptera is not surprising as majority of species prefer headwaters of lotic habitats (GRAF *et al.*, 2002; 2009; 2023). A substantial decrease of organic matter downstream the Plitvice Lakes barrage system (OBELIĆ *et al.*, 2005; HORVATINČIĆ *et al.*, 2006; MILIŠA *et al.*, 2006; IVEKOVIĆ, 1971) is most probably the reason for restricted distribution of Heteroptera along this hydrosystem due to the food availability. As majority of species are gatherers/collectors or predators (ZETTEL, 1995; SCHMEDITJE & COLLING, 1996), the highest amount of sedimented fine particulate organic matter in the Prošće Lake, and consequently highest abundance of some other macroinvertebrates, such as mayflies, could have provided optimal food resources for these insects. Regarding the other recorded macroinvertebrates, it is important to mention that Mollusca and Oligochaeta were present and abundant in all four lakes due to their wide ecological tolerance and/or preference for lentic habitats

(HÖRNER *et al.*, 1995; NESEMANN & REISCHÜTZ, 1995). Surprisingly, Isopoda, which are usually common in a typical benthic-lake macrofauna and have already been recorded in tufa-depositing systems (e.g. MINCKLEY, 1963), were collected only in the Ciginovac Lake. Absence of the Isopoda species in the studied lakes is probably consequence of geological and climatological history of this area which was, and still is, because of the tufa deposition, transforming from the stream-river to the lake system (KOSTIĆ-BRNEK & BRNEK-KOSTIĆ, 1974). Finally, a decrease in abundance of other macroinvertebrates most probably caused decrease in abundance of predatory and parasitic Hirudinea downstream the hydrosystem (NESEMANN & MOOG, 1995; SCHMEDITJE & COLLING, 1996).

A substantial decrease of organic matter downstream the Plitvice Lakes barrage system (OBELIĆ *et al.*, 2005; HORVATINČIĆ *et al.*, 2006; MILIŠA *et al.*, 2006; IVEKOVIĆ, 1971) is most probably the reason for the restricted distribution of Heteroptera along this hydro-system due to food availability. As the majority of species are gatherers/collectors or predators (ZETTEL, 1995; SCHMEDITJE & COLLING, 1996), the highest amount of sedimented fine particulate organic matter in the Prošće Lake, and consequently highest abundance of some other macroinvertebrates, such as mayflies, could have provided optimal food resources for these insects. Similarly, a decrease in the abundance of other macroinvertebrates most probably caused a decline in the abundance of predatory and parasitic Hirudinea downstream the hydrosystem (NESEMANN & MOOG, 1995; SCHMEDITJE & COLLING, 1996). The very low abundance of Plecoptera is not surprising as the majority of species prefer the headwaters of lotic habitats (GRAF *et al.*, 2002; 2009; 2023).

Development of tufa dams along a river profile creates a series of lakes and waterfalls. When the pooled waterbodies in the barrage system are formed, they become effective traps of eroded material (GOLUBIĆ, 1969). Water behind the tufa dams accumulates in the form of lakes containing their own plankton, benthos and littoral communities (GOLUBIĆ *et al.*, 2008). Accordingly, NMDS analyses showed that macroinvertebrate composition in the Plitvice Lakes hydrosystem was specific for each lake, regardless of the substrate type. Thus, differences in abundance and composition of benthic macroinvertebrate communities formed differences between the lakes. Dynamics of the physico-chemical environment, which in barrage-lake systems could be a consequence of the damming process, surely controls the biodiversity at relevant scales, and should also affect the community composition (HUSTON, 1994). In the case of the Plitvice Lakes, physico-chemical fluctuations did not explain the biological patterns that were observed. On the contrary, although the physical stream continuum has been disturbed by the development of tufa barriers (HABDIJA *et al.*, 1994), lakes in the barrage system can be taken as a continuum of physical and chemical conditions. These results corroborate results of previous a study where effects of the dams were not evident from examination of water quality variables (NICHOLS *et al.*, 2006). It seems that position of the lakes within the system plays a crucial role in the explanation of the observed patterns in benthic macroinvertebrates. Therefore, our results suggest that observed biodiversity patterns are, together with biology of individual taxa, also a consequence of the geomorphology of the Plitvice Lakes. Prošće Lake is the first lake in the barrage system, acting like a filter for organic matter, sediments and even fauna (IVEKOVIĆ, 1971), which explains a higher diversity observed there than in the other three lakes. All food and sediment particles coming with the inflowing water from the two main springs are first being retained in the Prošće Lake. Due to that process, the progressive filling with sediment and aquatic vegetation, like *Chara* sp., has been recorded there (IVEKOVIĆ, 1971). Therefore, we assume

that lakes in barrage systems are not just effective traps of eroded material but also “small filtering screens”. Similar to lakes that are inserted into river systems, these lakes in the cascade probably also function as sediment sinks (e.g. ARP *et al.*, 2007). They retain organic matter and faunal components like benthic macroinvertebrates (MILIŠA *et al.*, 2014), the distribution of which in the Plitvice Lakes does not correspond to the river continuum concept (RCC) hypothesis (HABDIJA *et al.*, 1994; ŠEMNIČKI *et al.*, 2012; VILENICA *et al.*, 2017). Furthermore, as a result of the retention effect in barrage systems, organisms should show specific quantitative and qualitative patterns. Indeed, taxa richness and diversity showed a decreasing trend downstream the Plitvice Lakes. This trend was partially related to the abundance, which decreased in the upper section, where Prošće, Ciginovac and Kozjak lakes are placed. It is possible that Rječica Stream, a tributary that flows into the Kozjak Lake, represents a disturbance (e.g. KATANO *et al.*, 2009), which caused a reduction in total macroinvertebrate abundance in the Kozjak Lake (e.g. OBELIĆ *et al.*, 2005). This observation could additionally be explained by habitat characteristics. The substrates sampled in Kaluđerovac Lake were stable microhabitats, while there was dynamics of appearance and disappearance of sampled substrates in Kozjak Lake. Substrate heterogeneity is also an interesting consideration in barrage systems. If we consider a barrage system as a set of spatial discontinuities then, according to the serial discontinuity concept (SDC), environmental heterogeneity exhibits a pattern along the stream profile that is similar to biotic diversity (WARD & STANFORD, 1983). Such a conclusion must, however, remain highly speculative at this time, since further studies are necessary.

Generally, the observed decreasing trend of macroinvertebrate components should also be discussed and further studied in the context of tufa deposition, which has an increasing trend downstream the Plitvice Lakes (SRDOČ *et al.*, 1985). In this study, we also measured conductivity, which can be used as a surrogate measure of Ca<sup>2+</sup> concentration (DRYSDALE *et al.*, 2002; CHEN *et al.*, 2004; LIU *et al.*, 2006). Thus, according to the results, tufa formation was pronounced downstream and followed by a significantly decreasing diversity and abundance of organisms. Furthermore, several studies have suggested that deposition of carbonates has negative effects on macroinvertebrate communities (MINCKLEY, 1963; CARTER & MARKS, 2007). Specifically, RUNDIO (2009) showed that taxa richness was lower at sites with tufa in comparison to the non-tufa sites.

## CONCLUSION

The barrage-lake system is a model of spatial discontinuities, where retention effects and tufa deposition cause notable alterations in ecosystem patterns, such as changes in taxon richness and the composition and structure of communities. These spatial discontinuities constitute sediment sinks, where lakes retain and accumulate organic matter and fauna. Therefore, each lake acts as a filtering screen and consequently appears to be a small ecosystem. Moreover, each lake has its own community characteristics, regardless of the fact that water flows from one lake to the next, connecting them and resulting in similar abiotic conditions in each lake along the system.

## ACKNOWLEDGMENTS

We thank Igor Stanković for assistance with measurements of physico-chemical water properties. He and Miran Katar prepared the artwork. Many thanks to Ivan Vučković for his valuable comments and support. We would also like to thank the

following students who helped us with the fieldwork: Andreja Komljenović, Martina Krbavčić, Sanja Žalac, Bruno Polak and Matija Šimac. This research was supported by Croatian Ministry of Science, Education and Sports as a part of the Projects No. 119-1193080-1206 granted to Mladen Kučinić.

Received June 15, 2023

## REFERENCES

- APHA (American Public Health Association), 1998: Standard methods for the examination of water and wastewater. American Public Health Association, Washington, DC.
- ARP, C.D., SCHMIDT, J.C., BAKER, M.A. & MYERS, A.K., 2007: Stream geomorphology in a mountain lake district: Hydraulic geometry, sediment sources and sinks, and downstream lake effects. *Earth Surface Processes and Landforms* **32**, 525-543.
- BAUERNFEIND, E. & HUMPEŠCH, U.H., 2001: Die Eintagsfliegen Zentraleuropas: Bestimmung und Ökologie. Verlag des Naturhistorischen Museums Wien, Wien, pp. 239.
- BAUERNFEIND, E. & MOOG O., 2000: Mayflies (Insecta: Ephemeroptera) and the assessment of ecological integrity: a methodological approach. *Hydrobiologia* **422**, 71-83.
- BAUERNFEIND E., & SOLDÁN T., 2012: The mayflies of Europe (Ephemeroptera). Apollo Books, Ollerup, Denmark, pp. 781.
- BAYARI, C. S. & KURTAS T., 1997: Algae: an important agent in deposition of karstic travertines: observations on natural-bridge Yerköprü travertines, Aladaglar, Eastern Taurids, Turkey. In: GUNAY G. & JOHNSON A. I (eds.), *Karst Waters and Environmental Impacts*, Rotterdam, p. 269-280.
- BELLINZONI, A.M., CANEVA, G. & RICCI, S., 2003: Ecological trends in travertine colonisation by pioneer algae and plant communities. *International Biodeterioration and Biodegradation* **51**, 203-210.
- BERAKOVIĆ, M., 2005: Hydrological investigations - water balance of the Plitvice Lakes. *Croatian water management* **156**, 59-63.
- CARTER, C.D. & MARKS J.C., 2007: Influences of travertine dam formation on leaf litter decomposition and algal accrual. *Hydrobiologia* **575**, 329-341.
- CARTHEW, K.D., DRYSDALE R.N. & TAYLOR M.P., 2003: Tufa deposits and biological activity, Riversleigh, Northwestern Queensland. In: ROACH I.C. (ed.), *Advances in Regolith*, CRC LEME, p. 55-59.
- CHEN, J., ZHANG, D.D., WANG, S., XIAO, T. & HUANG, R., 2004: Factors controlling tufa deposition in natural waters at waterfall sites. *Sedimentary Geology* **166**, 353-366.
- CHAFETZ, H.S., RUSH, P.F. & UTECH, N.M., 1991: Microenvironmental controls on mineralogy and habit of CaCO<sub>3</sub> precipitates: an example from an active travertine system. *Sedimentology* **38**, 107-126.
- CHAFETZ, H.S., SRDOČ, D. & HORVATINČIĆ, N., 1994: Early diagenesis of Plitvice Lakes waterfall and barrier travertine deposits. *Géographie physique et Quaternaire* **48**, 247-255.
- CLARKE, K.R. & GORLEY, R.N., 2006: PRIMER v6: User manual/tutorial, PRIMER-E, Plymouth, UK.
- DIEHL S., 1995: Direct and indirect effects of omnivory in a littoral lake community. *Ecology* **76**, 1727-1740.
- DRYSDALE, R.N., 1999: The sedimentological significance of hydropsychid caddisfly larvae (Order: Trichoptera) in a travertine-depositing stream: Louie Creek, Northwest Queensland, Australia. *Journal of Sedimentary Research* **69**, 145-150.
- DRYSDALE, R.N., TAYLOR, M.P. & IHLENFELD, C., 2002: Factors controlling the chemical evolution of travertine-depositing rivers of the Barkly karst, northern Australia. *Hydrological Processes* **16**, 2941-2962.
- DURRENFELDT, A., 1978: Untersuchungen zur Besiedlungsbiologie von Kalktuff faunistische, ökologische und elektronenmikroskopische Befunde. *Archiv für Hydrobiologie* **54**, 1-79.
- EDWARDS, M. & USHER, M.B., 1985: The winged Antarctic midge *Parochlus steineii* (Gerke) (Diptera: Chironomidae) in the South Shetland Islands. *Biological Journal of the Linnean Society* **26**, 83-93.
- EMEIS, K.C., RICHNOW, H.H. & KEMPE, S., 1987: Travertine formation in Plitvice National Park, Yugoslavia: chemical versus biological control. *Sedimentology* **34**, 595-609.
- ENGBLOM, E., 1996: Ephemeroptera, Mayflies. In: NILSSON, A. (ed.), *Aquatic Insects of North Europe. A Taxonomic Handbook, Vol.1*, Apollo Books, Stenstrup, p. 13-53.
- FORD, T.D. & PEDLEY, H.M., 1996: A review of tufa and travertine deposits of the world. *Earth-Science Reviews* **41**, 117-175.
- FRANČIŠKOVIĆ-BILINSKI, S., BARIŠIĆ, D., VERTAČNIK, A., BILINSKI, H. & PROHIĆ, E., 2004: Characterization of tufa from the Dinaric Karst of Croatia: mineralogy, geochemistry and discussion of climate conditions. *Facies* **50**, 183-193.

- GERKEN, B. & STERNBERG, K., 1999: Die Exuvien Europäischer Libellen - The exuviae of European Dragonflies (Insecta, Odonata), Arnika & Eisvogel, Hökster & Jena, 354 pp.
- GOLUBIĆ, S., 1969: Cyclic and non-cyclic mechanisms in the formation of travertine. Verhandlungen des Internationalen Verein Limnologie 17, 956–961.
- GOLUBIĆ, S., VIOLANTE, C., PLENKOVIĆ-MORAJ, A. & GRGASOVIĆ, T., 2008: Travertines and calcareous tufa deposits: an insight into diagenesis. Geologia Croatica 61, 363–378.
- GRAF, W., GRASSER, U. & WEINZIERL, A., 2002: Plecoptera. In: MOOG, O. (ed.), Fauna Aquatica Austriaca, Lieferung 2002. Wasserwirtschaftskataster, Bundesministerium für Land- und Forstwirtschaft, Umwelt und Wasserwirtschaft, Wien.
- GRAF, W., LORENZ, A.W., TIerno DE FIGUEROA, J.M., LÜCKE, S., LÓPEZ-RODRÍGUEZ, M.J. & DAVIES, C., 2009: Volume 2 - Plecoptera. In: SCHMIDT-KLOIBER, A. & HERING, D. (eds.), Distribution and Ecological Preferences of European Freshwater Organisms, Pensoft Publishers (Sofia-Moscow), p. 1-262.
- GRAF, W., LORENZ, A.W., TIerno DE FIGUEROA, J.M., LÜCKE, S., LÓPEZ-RODRÍGUEZ, M.J., MURPHY, J. & SCHMIDT-KLOIBER, A., 2023: Dataset "Plecoptera". www.freshwaterecology.info - the taxa and autecology database for freshwater organisms, version 7.0 (accessed on 27.09.2023).
- HABDIJA, I., PRIMC-HABDIJA, B. & BELINIĆ, I., 1994: Functional community organization of macroinvertebrates in lotic habitats of the Plitvice Lakes. Acta Hydrochimica Et Hydrobiologica 22, 85-92.
- HABDIJA, I., PRIMC-HABDIJA, B., MATONIČKIN, R., KUČINIĆ, M., RADANOVIĆ, I., MILIŠA, M. & MIHALJEVIĆ, Z., 2004: Current velocity and food supply as factors affecting the composition of macroinvertebrates in bryophyte habitats in karst running water. Biologia 59, 577–593.
- HORNING, C.E. & POLLARD J.E., 1978: Macroinvertebrate sampling techniques for streams in semi-arid regions: comparison of the Surber method and unit-effort traveling Kick method, Report number EPA 600-4-78-040, US EPA, Washington, 37 p.
- HORVATINČIĆ, N., BRIANSÓ, J.L., OBELIĆ, B., BAREŠIĆ, J. & BRONIĆ, I.K., 2006: Study of pollution of the Plitvice Lakes by water and sediment analyses. Water, Air, & Soil Pollution 6, 475-485.
- HORVATINČIĆ, N., BAREŠIĆ, J., BABINKA, S., OBELIĆ, B., KRAJCAR BRONIĆ, I., VREČA, P. & SUCKOW, A., 2008: Towards a deeper understanding how carbonate isotopes ( $^{14}\text{C}$ ,  $^{13}\text{C}$ ,  $^{18}\text{O}$ ) reflect environmental changes: A study with recent  $^{210}\text{Pb}$ -dated sediments of the Plitvice Lakes, Croatia. Radiocarbon 50, 233-253.
- HÖLZEL, H., 2002: Insecta: Megaloptera, Neuroptera, Lepidoptera. In: FISHER, G. (ed.), Süßwasserfauna von Mitteleuropa, Spektrum Akademischer Verlag Heidelberg, Berlin, p. 3-26.
- HÖRNER, K., MOOG, O. & SPORKA, F., 1995: Oligochaeta. In: Moog, O. (ed.), Fauna Aquatica Austriaca, Lieferungen 1995, 2002. Wasserwirtschaftskataster, Bundesministerium für Land- und Forstwirtschaft, Umwelt und Wasserwirtschaft, Wien.
- HUMPHREYS, W.F., AWRAMIK, S.M. & JEBB, M.H.P., 1995: Freshwater biogenic tufa in Madang Province, Papua New Guinea. Journal of the Royal Society of Western Australia 78, 43–54.
- HUSTON, M.A., 1994: Biological Diversity, The Coexistence of Species on Changing Landscapes, Cambridge University Press, New York, 681 pp.
- IVKOVIĆ, H., 1971: Samouništanje i samozaštita Plitvičkih jezera. In: VARIČAK, T. (ed.), Simpozij o zaštiti prirode u našem kršu, Zagreb, 2-3 December 1970, Croatia, p. 81-97.
- IVKOVIĆ, M., KESIĆ, M. & STILOUKALOVA, V., 2012: Contribution to the knowledge of blackfly (Diptera, Simuliidae) fauna at the Plitvice National Park. Natura Croatica 21, 263-268.
- IVKOVIĆ, M., KESIĆ, M., MIHALJEVIĆ, Z. & KUDELA, M., 2014: Emergence patterns and ecological associations of some hematophagous blackfly species along an oligotrophic hydrosystem. Medical and Veterinary Entomology 28, 94-102.
- IVKOVIĆ, M. & PONT, A.C., 2016: Long-time emergence patterns of *Limnophora* species (Diptera, Muscidae) in specific karst habitats: tufa barriers. Limnologica 61, 29-35.
- JÄCH, M.A. & BROJER, M., 2006: Wasserkäfer. Skriptum zum Spezialpraktikum. Forschungsinstitut für Insektenkunde, Naturhistorisches Museum, Wien, 426 pp.
- KATANO, I., NEGISHI, J.N., MINAGAWA, T., DOI, H., KAWAGUCHI, Y. & KAYABA, Y., 2009: Longitudinal macroinvertebrate organization over contrasting discontinuities: effects of a dam and a tributary. Journal of the North American Benthological Society 28, 331-351.
- KEMPE, S. & EMEIS, K., 1985: Carbonate chemistry and the formation of Plitvice Lakes. Geologisch-Paläontologisches Institut der Universität Hamburg 58, 351-383.
- KOŠTIĆ-BRNEK, L.J. & BRNEK-KOŠTIĆ, A., 1974: Vode Nacionalnog parka Plitvička jezera i život u njima. In: Gušić, B. & MARKOVIĆ, M. (eds.), Plitvička jezera - čovjek i priroda. National Park Plitvice Lakes, Zagreb, p. 37-49.

- KUČINIĆ, M., PREVIŠIĆ, A., GOTTSTEIN, A., HRAŠOVEC, B., STANIĆ-KOŠTROMAN, S., PERNEK, M. & DELIĆ, A., 2008: Description of the larvae of *Drusus radovanovici septentrionis* Marinković-Gospodnetić, 1976 and *Drusus croaticus* Marinković-Gospodnetić, 1971 (Trichoptera: Limnephilidae) from Bosnia and Herzegovina and Croatia. *Zootaxa* **1783**, 1-17.
- KUČINIĆ, M., PREVIŠIĆ, A., GRAF, W., MIHOČI, I., ŠOUFEK, M., STANIĆ-KOŠTROMAN, S., LELO, S., VITECEK, S. & WARINGER, J., 2015: Larval description of *Drusus bosniacus* Klapálek 1899 (Trichoptera: Limnephilidae) with distributional molecular and ecological features. *Zootaxa* **3957**, 85-97.
- KUMANSKI, K. & MALICKY, H., 1999: A survey of the genus *Potamophyllax* Wallengren, 1981 in the Balkan Peninsula, with description of two new species (Trichoptera: Limnephilidae). *Braueria* **26**, 27-30.
- LEONARD, J. W., 1939: Comments on the adequacy of accepted stream bottom sampling techniques, Transactions of 4th North American Wildlife Conference, p. 288-295.
- LIU, Z., LI, Q., SUN, H., LIA, C., LI, H., WANG, J. & WU, K., 2006: Diurnal variations of hydrochemistry in a travertine-depositing stream at Baishuitai, Yunnan, SW China. *Aquatic Geochemistry* **12**, 103-121.
- MALMQUIST, B., ZHANG, Y. & ADLER, P.H., 1999: Diversity, distribution and larval habitats of North Swedish blackflies (Diptera: Simuliidae). *Freshwater Biology* **42**, 301-314.
- MATONIČKIN KEPČIJA, R., HABDIJA, I., PRIMC-HABDIJA, B. & MILIŠA, M., 2006: Simuliid silk pads enhance tufa deposition. *Fundamental and Applied Limnology* **166**, 387-409.
- MEINANDER, M., 1996: Megaloptera, Alder Flies. In: NILSSON, A. (ed.), *Aquatic Insects of North Europe, A Taxonomic Handbook, Vol 1*, Apollo Books, Stenstrup, p. 105-110.
- MENETREY, N., OERTLI, B., SARTORI, M., WAGNER, A. & LACHAVANNE, J.B., 2008: Eutrophication: are mayflies (Ephemeroptera) good bioindicators for ponds? *Hydrobiologia* **597**, 125-135.
- MERZ-PREISZ, M. & RIDING, R., 1999: Cyanobacterial tufa calcification in two freshwater streams: ambient environment, chemical thresholds and biological processes. *Sedimentary Geology* **22**, 103-124.
- MIČEVIĆ STANKOVIĆ, V., JÄCH, M.A., IVKOVIĆ, M., STANKOVIĆ, I., KRUŽIĆ, P., KUČINIĆ, M., 2019: Spatio-temporal distribution and species traits of water beetles along an oligotrophic hydrosystem: a case study. *International Journal of Limnology* **55**, 1-22.
- MILIŠA, M., HABDIJA, I., PRIMC-HABDIJA, B., RADANOVIĆ, I. & MATONIČKIN KEPČIJA, R., 2006: The role of flow velocity in the vertical distribution of particulate organic matter on moss-covered travertine barriers of the Plitvice Lakes (Croatia). *Hydrobiologia* **553**, 231-243.
- MILIŠA, M., IVKOVIĆ, M. & MATONIČKIN KEPČIJA, R., 2014: Energy resources and feeding guild structure of macroinvertebrate assemblages in the hyporheic zone of calcite depositing lake outlets. *Limnologia* **44**, 66-71.
- Minckley, W.L., 1963: The ecology of a spring stream: Doe Run, Meade County, Kentucky. *Wildlife Monographs* **11**, 1-124.
- NESEMANN, H. & MOOG, O., 1995: Hirudinea. In: MOOG, O. (ed.), *Fauna Aquatica Austriaca, Lieferungen 1995, 2002*. Wasserwirtschaftskataster, Bundesministerium für Land- und Forstwirtschaft, Umwelt und Wasserwirtschaft, Wien.
- NESEMANN, H. & REISCHÜTZ, P.L., 1995: Bivalvia. In: MOOG, O. (ed.), *Fauna Aquatica Austriaca, Lieferungen 1995, 2002*. Wasserwirtschaftskataster, Bundesministerium für Land- und Forstwirtschaft, Umwelt und Wasserwirtschaft, Wien.
- NILSSON, A.N., HANSEN, M., KLAUSNITZER, B. & PALM, E., 1996: Coleoptera. In: NILSSON, A. (ed.), *Aquatic Insects of North Europe, A Taxonomic Handbook, Vol.1*, Apollo Books, Stenstrup, p. 115-222.
- NICHOLS, S., NORRIS, R., MAHER, W. & THOMAS, M., 2006: Ecological effects of serial impoundment on the Cotter River, Australia. *Hydrobiologia* **572**, 255-273.
- OBELIĆ, B., HORVATINČIĆ, N., BAREŠIĆ, J., BRIANSÓ, J.L., BABINKA, S. & SUCKOW, A., 2005: Anthropogenic pollution in karst lake sediments (Croatia). In: ÖZKUL, M., YAGIZ, S. & JONES, B. (eds.), *Proceedings of 1st International Travertine Symposium and Technologies Exhibition*, Pamukkale University, 21-25 September 2005, Denizli, Turkey, p. 188-196.
- PAPROCKI, H., HOLZENTHAL, R.W. & CRESSA, C., 2003: A new species of Smicridea McLachan (Trichoptera: Hydropsychidae) from Venezuela and its role in travertine biogenesis. *Journal of the North American Benthological Society* **22**, 401-409.
- PENTECOST, A., 2005: *Travertine*. Berlin, Heidelberg. Springer-Verlag, New York, 445 pp.
- PENTECOST, A. & ZHANG, Z., 2000: The travertine flora of Juizhaigou and Munigou, China and its relationship with calcium carbonate deposition. *Cave Karst Science* **27**, 71-78.
- PETRIK, M., 1958: Prinosi hidrologiji Plitvica, Poljoprivredni nakladni zavod, Zagreb, Croatia.
- POPIJAČ, A. & SIVEC, I., 2012: Diversity and distribution of stoneflies in the area of Plitvice Lakes National Park and along the Mediterranean River Cetina (Croatia). *Aquatic Insects* **31** (1), 731-742.

- PREVIŠIĆ, A., KEROVEC, M. & KUČINIĆ, M., 2007: Emergence and composition of Trichoptera from karst habitats, Plitvice Lakes region, Croatia. *International Review of Hydrobiology* **92**, 61–83.
- PREVIŠIĆ, A., GRAF, W., VITECEK, S., KUČINIĆ, M., BÁLINT, M., KERESZTES, L., PAULS, S.U. & WARINGER, J., 2014: Cryptic diversity of caddisflies in the Balkans: the curious case of *Eclisopteryx* species (Trichoptera: Limnephilidae). *Arthropod Systematics & Phylogeny* **72**, 309–329.
- PRIMC-HABDIJA, B., HABDIJA, I. & PLENKOVIĆ-MORAJ, A., 2001: Tufa deposition and periphyton overgrowth as factors affecting the ciliate community on travertine barriers in different current velocity conditions. *Hydrobiologia* **457**, 87–96.
- RIDING, R., 1991: Classification of microbial carbonates. In: RIDING R. (ed.), *Calcareous Algae and Stromatolites*, Springer-Verlag, Berlin, p. 21–51.
- RIBANOVIĆ, J. & BOŽIČEVIĆ, S., 1996: Geographic-physical conditions of runoff and hydrogeological characteristics of the Plitvice Lakes. *Acta Geographica Croatica* **31**, 7–26.
- RUNDIO, D.E., 2009: Community-habitat relationships in coastal streams in Big Sur, California, USA: travertine influences macroinvertebrate abundance and community structure. *Hydrobiologia* **620**, 91–108.
- SHIMADA, K., OHYAMA, Y. & PAN, C.X., 1991: Cold-hardiness of the Antarctic winged midge *Parochlus steinii* during the active season at King George Island. *Polar Biology* **11**, 311–314.
- SCHMEDTJE, U. & COLLING, M., 1996: Ökologische Typisierung der aquatischen Makrofauna. *Bayerisches Landesamt für Wasserwirtschaft* **4/96**, 1–543.
- SRDOČ, D., HORVATINČIĆ, N., OBELIĆ, B., KRAJCAR, I. & SLIEPČEVIĆ, A., 1985: Calcite deposition processes in karst waters with special emphasis on the Plitvice Lakes, Yugoslavia. *Carsus Yugoslaviae* **11**, 101–204.
- STOFFERS, P., 1975: Recent carbonate sedimentation in the lakes of Plitvice (Yug.). *Neues Jahrbuch für Mineralogie* **9**, 412–418.
- STRAYER, D.L., 1985: The benthic micrometazoans of Mirror Lake, New Hampshire. *Archiv für Hydrobiologie* **72**, 287–426.
- ŠEGULJA, N. & HRŠAK, V., 1994: Some floristic and ecological characteristics of association Bromo-Plantaginatum mediae Ht. (31) 49 on permanent plot No. 82 (National Park Plitvice Lakes - Croatia). *Plant Ecology* **110**, 103–114.
- ŠEMNIČKI, P., PREVIŠIĆ, A., IVKOVIĆ, M., ČMRLEC, K. & MIHALJEVIĆ, Z., 2012: Tufa barriers from caddisfly's point of view: streams or lake outlets? *International Review of Hydrobiology* **97**, 465–484.
- ŠPOLJAR, M., PRIMC-HABDIJA, B. & HABDIJA, I., 1994: Transport of seston in the karstic hydrosystem of the Plitvice Lakes (Croatia). *Hydrobiologia* **579**, 199–209.
- VILENICA, M., GATTOLLIAT, J., IVKOVIĆ, M., KUČINIĆ, M., MIČETIĆ STANKOVIĆ, V., MIHALJEVIĆ, Z. & SARTORI, M., 2014: The mayfly fauna (Insecta, Ephemeroptera) of the Plitvice Lakes National Park, Croatia. *Natura Croatica* **23**, 349–363.
- VILENICA, M., GATTOLLIAT J.-L., MIHALJEVIĆ, Z. & SARTORI, M., 2015: Croatian mayflies (Insecta, Ephemeroptera): species diversity and distribution patterns. *ZooKeys* **523**, 99–127.
- VILENICA, M., MIČETIĆ STANKOVIĆ, V., SARTORI, M., KUČINIĆ, M. & MIHALJEVIĆ, Z., 2017: Environmental factors affecting mayfly assemblages in tufa-depositing habitats of the Dinaric Karst. *Knowledge and Management of Aquatic Ecosystems* **418** (14).
- VITECEK, S., GRAF, W., PREVIŠIĆ, A., KUČINIĆ, M., OLÁH, J., BÁLINT, M., KERESZTES, L., PAULS, S.U. & WARINGER, J., 2015: A hairy case: The evolution of filtering carnivorous Drusinae (Limnephilidae, Trichoptera). *Molecular Phylogenetics and Evolution* **93**, 249–260.
- WARD, J.W. & STANFORD, J.A., 1983: The serial discontinuity concept of lotic ecosystems. In: FONTANE T.D. & BARTTELL S.M. (eds.), *Dynamics of Lotic Ecosystems*. Ann Arbor Science Publ., Ann Arbor, Michigan, p. 29–42.
- WARINGER, J. & GRAF, W., 1997: *Atlas der Österreichischen Köcherfliegenlarven (unter Einschluss der angrenzenden gebiete)*, Facultas Universitätsverlag, Wien.
- WARINGER, J. & GRAF, W., 2011: *Atlas of Central European Trichoptera Larvae*. Erik Mauch Verlag, Dinkelscherben.
- WIEDERHOLM, T., 1984: Responses of aquatic insects to environmental pollution. In: RESH, V.H. & ROSENBERG, D.M. (eds.), *The ecology of aquatic insects*, Praeger Publishers, New York, p. 508–557.
- ZETTEL, H., 1995: Heteroptera. In: MOOG, O. (ed.), *Fauna Aquatica Austriaca, Lieferungen 1995, 2002*. Wasserwirtschaftskataster, Bundesministerium für Land- und Forstwirtschaft, Umwelt und Wasserwirtschaft, Wien.
- ZWICKER, G. & RUBINIĆ, J., 2005: Water level fluctuations as an indicator of tufa barrier growth dynamics in the Plitvice Lakes. *Materials and Geoenvironment* **52**, 161–163.





MEGALOPTERA								
<i>Sialis fuliginosa</i> Pictet, 1836	2	7	3			1		13
<i>Sialis lutaria</i> (Linnaeus, 1758)	17	5	1	4				3
<i>Sialis morio</i> Klingstedt, 1932	13	1	2	5				1
<i>Sialis sordida</i> Klingstedt, 1932	26	8	4	11	1	2		5
TRICHOPTERA								
<i>Agrypnia varia</i> (Fabricius, 1973)					1			9
<i>Athripsodes</i> sp.	39	3	29	17				
<i>Ceraclea</i> sp.	18		1	1				
<i>Cyrmus trimaculatus</i> (Curtis, 1834)	2		1	3				49
<i>Hydropsyche instabilis</i> (Curtis, 1834)					10			1
<i>Hydropsyche saxonica</i> McLachlan, 1884					5			1
<i>Hydropsyche</i> sp.					5			
<i>Hydroptila</i> sp.	5							
<i>Limnephilus</i> sp.	1	1	1	1				
<i>Mystacides</i> sp.	2		3	5				20
<i>Notidobia ciliaris</i> Linnaeus, 1761	1	3		17				
<i>Oecetis testacea</i> (Curtis, 1834)	1		5	5	4	4		7
Hydroptilidae non det.	1	1						
<i>Philopotamus</i> sp.	1	2						
<i>Rhyacophila</i> sp.			2			1		
<i>Timodes</i> sp.	1	1	4	2				
DIPTERA								
Ceratopogonidae	102	63	4	6	15	11	9	105
Chironomidae	5586	1713	1829	1078	269	589	305	1740
Empididae	2							4
Limoniidae					1			
<i>Tipula</i> sp.								3
Simuliidae			12		18	1		8
HETEROPTERA	137	501	2	3				
COLLEMBOLA	1	1	1	1		2		4
NEMATODA	11	12	3	5				7
Total number of collected individuals	11284	4264	2997	1986	535	1301	485	3230
Taxa richness	46	40	34	33	24	24	9	32

