

# Seroprevalence of *Mycobacterium avium* subsp. *paratuberculosis* Infections in Small Ruminants in Europe – A Systematic Review



H. Quintas\*, J. Benavides, J. Jacob-Ferreira, P. Afonso and A. C. Coelho

## Abstract

Paratuberculosis, also known as Johne's disease, is a granulomatous enteritis in both domestic and wild ruminants caused by the bacterium *Mycobacterium avium* subsp. *paratuberculosis*. Understanding the prevalence of this disease in small ruminants is essential for disease control and prevention strategies. A systematic review of the literature was conducted using the PubMed, ScienceDirect and Scopus databases to identify all articles reporting *Mycobacterium avium* subsp. *paratuberculosis* (MAP) seroprevalence in sheep and goats in Europe, published from January 2006 to December 2023. The initial search for existing publications reporting systematic reviews and primary studies was carried out by searching the available databases. For the final selection of studies, an initial screen for basic eligibility and a detailed appraisal of quality were performed. After the study selection, the relevant data was extracted. The detailed appraisal generated 21 publications that reported 55 studies, 22 (40.0%) from sheep (12 at the animal-level and 10 at the flock-level) and 28

(50.9%) from goats (17 at the animal-level and 11 at the flock-level), and 5 (9.1%) from mixed small ruminant species at the animal level. In total, 34 (61.8%) were animal-level studies and 21 (38.2%) were flock-level studies. Population and inclusion criteria were highly variable among studies. Sample sizes ranged from 291 to 15,585 animals. Most studies reported testing adult animals (over 24 months of age). Commercial ELISA kits were used in most studies. The highest prevalence was obtained in sheep (100%) in Türkiye, and the lowest was found also in sheep (0.7%) in Austria. Overall, the results suggest that MAP antibodies have been frequently detected among small ruminants in some countries and there is a need for standardisation of case definitions to improve the accuracy of prevalence estimates. Further research is needed to understand the risk factors associated with MAP infection in small ruminants and to develop effective control and prevention strategies.

**Key words:** Europe; *Mycobacterium avium* subsp. *paratuberculosis*; systematic review; serology

Hélder QUINTAS\*, (Corresponding author, e-mail: helder5tas@ipb.pt), DVM, MSc, PhD, Mountain Research Center (CIMO), Bragança Polytechnic University (IPB), Campus de Santa Apolónia, 5300-253 Bragança, Portugal; Júlio BENAVIDES, Instituto de Ganadería de Montaña, CSIC-University of León, Grulleros, 24346 León, Spain; João JACOB-FERREIRA, CIMO, IPB, Campus de Santa Apolónia, Bragança, 5300-253, Portugal; Paulo AFONSO, DVM, MSc, PhD, Animal and Veterinary Research Centre (CECAV), University of Trás-os-Montes e Alto Douro (UTAD), Quinta de Prados, 5000-801 Vila Real, Portugal; and CIMO, IPB, Campus de Santa Apolónia, 5300-253 Bragança, Portugal; Ana Cláudia COELHO, DVM, MSc, PhD, Department of Veterinary Sciences, School of Agrarian and Veterinary Sciences (ECAV), CECAV, UTAD, Quinta de Prados, 5000-801 Vila Real, Portugal

## Introduction

Paratuberculosis in small ruminants is caused by *Mycobacterium avium* subsp. *paratuberculosis* (MAP), a pathogen dependent on mycobactin, grows slowly and is acid-fast. It is a chronic inflammatory disease affecting the intestinal tract and lymphoid organs. This disease leads to a gradual decline in body weight and is responsible for significant economic losses on a global scale (Idris et al., 2022). Paratuberculosis in small ruminants can cause various clinical signs, including weight loss, decreased milk production, and lethargy. The disease can progress slowly and may not show clinical signs in the early stages (Bauman et al., 2016). In some cases, goats may develop a swollen abdomen due to the thickening of the intestinal wall. Paratuberculosis impacts a variety of wild and domesticated ruminant species, such as sheep, goats, and deer, but also other non-ruminant species such as swine, birds, otters, foxes, camelids, and primates (Miranda et al., 2009, 2011; Matos et al., 2013, 2014, 2015, 2017; Garvey, 2020). Clinical manifestation of paratuberculosis occurs in the final phase of the disease, when sheep and goats can present a variety of clinical signs, including weight loss, decreased milk production, and lethargy. Additional clinical signs in goats include fragile skin, poor coat, submandibular oedema, dehydration, anaemia, and depression (Djønne, 2010). The disease decreases the slaughter value of ruminants, leads to premature culling, reduces milk production and reproductive performance, and raises replacement costs, resulting in significant financial losses for the livestock industry. The potential for MAP-infected animals to contract other illnesses complicates quantifying the actual economic damage (Garcia and Shalloo, 2015; Fanelli et al., 2022). MAP may also

be involved in human Crohn's disease, a chronic inflammatory bowel condition, though this remains under debate (Agrawal et al., 2021; Espeschit et al., 2023).

During the initial stage of infection, MAP is ingested and taken up by macrophages in the intestinal lining. These bacteria can survive and multiply within macrophages, evading the host's immune system (Whittington et al., 2019). In a subclinical stage, the infection is managed by a cell-mediated immune response, primarily characterised by a Th1 response. This phase involves the activation of macrophages and the production of key cytokines, such as interferon-gamma (IFN- $\gamma$ ), which are critical in containing the bacteria (Sharma et al., 2024). The immune system relies heavily on cell-mediated immunity, with the generation of T-cells and the activation of macrophages is essential for controlling the infection. This response is vital for the formation of granulomas, thereby limiting the spread of MAP (Sweeney, 2011). As the disease progresses, there is a notable shift from a cell-mediated (Th1) immune response to a humoral (Th2) immune response. This transition is characterised by the production of antibodies against MAP; however, these antibodies are generally ineffective in controlling the infection (Lee et al., 2023). In this stage, the immune response increasingly favours antibody production, yet this shift is often accompanied by a decline in the overall effectiveness of the immune response. Consequently, this decline allows the bacteria to proliferate and leads to more severe clinical manifestations of the disease (Mallikarjunappa et al., 2021).

Detecting paratuberculosis in animals during its latent period before become infectious is the most crucial factor in eradicating this chronic disease. However, this is difficult with current ante-mortem diag-

nostic tests. As the treatment is impractical, and despite the availability of a commercial vaccine, paratuberculosis remains a problem in European flocks. Therefore, it is important to assess its spread (Kawaji et al., 2011; Idris et al., 2022).

Although there is no reliable test for detecting MAP infection, several methods have been reported in the literature for diagnosing paratuberculosis and estimating its prevalence, including Ziehl-Neelsen staining bacterial culture, histopathology analysis, PCR of faeces, tissues or blood samples, and diagnostic tools to measure immune response such as agar gel immunodiffusion (AGID), enzyme-linked immunosorbent assay (ELISA), complement fixation tests and Interferon-Gamma Release Assay (IGRA) (Pérez et al., 1996; Garrido et al., 2000; Whitlock et al., 2000; Coelho et al., 2008a, 2008b, 2009, 2010, 2018; Singh et al., 2013; Garg et al., 2015; Bucziniski et al., 2019; De Grossi et al., 2020).

ELISA is commonly employed to detect clinically infected animals when the animal may be shedding. However, commercially available ELISA methods have varying sensitivities (Nielsen and Toft, 2009; Blanco et al., 2020). IGRA measures the immune response, specifically the production of interferon-gamma (IFN- $\gamma$ ) by T-cells when exposed to specific MAP antigens. This test can identify the early stages of infection; however, it has several limitations. It is often considered costly and impractical for widespread use, and its reliability can vary. IGRA may not consistently differentiate between infected animals with controlled infections and those with subclinical disease, potentially leading to the culling of healthy animals (Bulum et al., 2024). ELISA tests are the most logical option for epidemiological studies as they are cost-effective, although they will miss a percentage of subclinically infected animals (Swenney, 2011).

Diagnosing paratuberculosis in small ruminants based solely on clinical signs is inadequate, but by combining these signs with epizootiological data, it is possible to suspect the presence of MAP at the animal level (Whittington et al., 2019).

The seroprevalence of MAP infections in small ruminants is a significant concern across Europe. When making decisions or policies regarding an infection, the prevalence among the herd and individual animals is frequently a crucial factor that decisions or policymakers consider when determining the significance of infection and the appropriate measures to be taken (Nielsen and Toft, 2009). Understanding the current seroprevalence of the disease at both the herd and animal levels is necessary to prioritise and guide future research and control programmes (Rita et al., 2011).

Conducting a systematic review of the seroprevalence of paratuberculosis in small ruminants in Europe is of great importance for understanding the extent of the problem and developing effective prevention and control strategies. Authorities can put particular management measures in place to stop the spread of MAP among livestock by determining the prevalence of the disease. This systematic review aimed to gather and critically evaluate the available scientific evidence on the seroprevalence of paratuberculosis in small ruminants in Europe in recent years to provide a comprehensive overview of the current state of the disease and identify research gaps.

## Material and Methods

### Study design

The present study consisted of a systematic literature review to answer the following research question: What is the seroprevalence *Mycobacterium avium*

subsp. *paratuberculosis* reported in sheep and goats in Europe in the last 17 years? This study was conducted based on the methodological recommendations of the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) (Page et al., 2021).

### Article eligibility

Articles published in indexed journals cited in PubMed, ScienceDirect and Scopus were considered eligible if they consisted of research papers, short communications, and cross-sectional studies describing serological features (including species, population, prevalence/occurrence, and country). There were other restrictions regarding the eligibility criteria: only studies published between 1 January 2006 and 31 December 2023 and written in languages of Western Europe (comprising English, French, German, Italian, Portuguese, and Spanish) were included.

Publications included original research papers, short communications and cross-sectional studies that addressed issues within the following criteria: seroprevalence, small ruminants, sheep, goats, and Europe. Publications, including studies reporting seroprevalence, frequency, occurrence, or proportion of paratuberculosis in individuals and/or flocks located in European countries performed by serological methods in blood, were included. Reviews of the literature, case reports, research notes, editorials, experimental essays, textbook chapters, posters, abstracts, articles with no primary data, dissertations, and unpublished studies and data were excluded.

### Information sources and search strategies

The process of identifying articles in indexed journals was developed using PubMed (available at <http://www.ncbi.nlm.nih.gov/pubmed>), Scopus ([\[scopus.com\]\(https://www.scopus.com\)\), and ScienceDirect \(<https://www.sciencedirect.com/>\) databases. Studies conducted on other farm animal species or wild ruminants were excluded.](http://www.</a></p>
</div>
<div data-bbox=)

The combination of search terms in English applied included: {[Europe] AND [*Mycobacterium avium* subsp. *paratuberculosis* OR paratuberculosis OR John's disease] AND [small ruminant OR sheep OR ovine OR goat OR caprine] AND [seroprevalence OR prevalence OR occurrence OR geoepidemiology OR ELISA OR Serology OR Serological diagnosis OR seropositivity]}. To prevent missing data, references of retrieved publications were also checked to identify additional papers. The searches were conducted between January 2023 and March 2024.

After a comprehensive systematic search, duplicate records were excluded. Then, two independent reviewers selected articles based on their titles and abstracts, followed by a complete reading of the text when the title or abstract met the inclusion criteria or could not be rejected with certainty. Any disagreements or divergences were resolved by discussion and consensus. Two researchers extracted the required data and added the information on an electronic spreadsheet, dividing them into five groups: (i) sheep, (ii) goat, (iii) sheep flock, (iv) goat flock, (v) small ruminant flock. The initial screen for basic eligibility was carried out using titles and abstracts of relevant studies found by searching databases and reference lists through the snowballing procedure for reports that match the general inclusion criteria on population, study factor, and outcome. Following the preliminary screening, a thorough quality assessment was carried out. Full-text versions of potentially eligible studies were evaluated for three principal issues: minimisation of selection bias of animals or flocks, an adequate assertion of out-

comes, and minimisation of measurement or misclassification bias.

### Data extraction

After the final selection of studies, quantitative data was extracted and included the reference (authors and year of publication), data on country and region, study period, population, inclusion criteria, selection, sensitivity, and specificity used (if available) for adjusting prevalence estimation, and results in terms of proportion were extracted. Studies from


which no crude numbers on frequency or proportion are included were excluded unless the information provided allowed its calculation.

## Results

### Selection of Papers

The search of the databases yielded 2122 results, once duplicates had been removed (Figure 1).

After screening titles and abstracts, 35



**Figure 1.** Flowchart of the search, selection, and inclusion process for studies in the systematic review based on PRISMA guidelines.



records potentially met the inclusion criteria. From the analysis of their full text, 14 articles were excluded, and 21 studies were included in this systematic review (Coelho et al., 2007; Ikonomopoulos et al., 2007; Falconi et al., 2009; Liapi et al., 2011; Rita et al., 2011; Corrias et al., 2012; Stau et al., 2012; Márquez et al., 2013; Buyuk et al., 2014; Mercier et al., 2015; Graham et al., 2016; Celik and Turutoglu, 2017; Galiero et al., 2017; Barrero-Domínguez et al., 2019; Cecchi et al., 2019; De Grossi et al., 2020; Iarussi et al., 2019; Quintas et al., 2021; Jiménez-Martín et al., 2022; Gaffuri et al., 2023; Schrott et al., 2023).

The study results were analysed according to animal species at the animal and flock levels. Articles that did not meet the inclusion criteria and duplicates were excluded.

### **Qualitative analysis of the prevalence of *Mycobacterium avium* subsp. paratuberculosis in small ruminants**

The final analysis included 21 of 35 eligible publications after quality appraisal. The 21 selected publications included 55 studies: 22 (40.0%) from sheep (12 at the animal-level and 10 at the flock-level), 28 (50.9%) from goats (17 at the animal-level and 11 at the flock-level), and 5 (9.1%) from mixed small ruminant species at the animal level. In total, 34 (61.8%) were animal-level studies, while 21 (38.2%) were flock-level studies (Tables 1, 2, 3, 4 and 5).

The studies were carried out in 10 countries: Austria ( $n=5$ ; 9.1%), France ( $n=2$ ; 3.6%), Germany ( $n=5$ ; 9.1%), Greece ( $n=5$ ; 9.1%), Italy ( $n=15$ ; 29.3%), Portugal ( $n=4$ ; 7.3%), Cyprus ( $n=4$ ; 7.3%), Scotland ( $n=1$ ; 1.8%), Spain ( $n=8$ ; 14.5%), and Türkiye ( $n=6$ ; 10.9%).

Forty-five studies reported information on the study period, and these studies were carried out between 2003 to 2021. However, they were required to have

been published between 2006 and 2023. The diagnostic tests used in the selected studies were ELISA in 50 studies and the complement fixation test in five studies. High variability was observed in the commercial ELISA diagnostic tests. Only 36 studies provided information on the sensitivity and specificity used for prevalence adjustment or true prevalence estimation. The remaining studies provided only apparent prevalence.

All studies reported information on the selection of animals or flocks included for MAP in small ruminant testing in the studies. Random selection of animals or flocks was reported in 54.5% (30/55) of the studies. Six studies were stratified by random sampling. Studies that report convenience sampling were kept in the study to summarise and synthesise their results if animals or flocks had no previous diagnosis of paratuberculosis.

### **Prevalence in sheep at the animal level**

Prevalence studies in sheep at the animal level ( $n = 12$ ) reported in nine countries are shown in Table 1. Of the studies included in the final analysis, three (25%) were from Italy, two (16.7%) from Türkiye, and one each (8.3%) from Austria, Germany, Greece, Portugal, Scotland, Spain, and Cyprus. Information on the study period was provided in 91.7% (11/12) of the studies.

Population and inclusion criteria were highly variable among studies. Sample sizes ranged from 291 (Ikonomopoulos et al., 2007) to 15,585 animals (Schrott et al., 2023).

Most studies tested adult animals (*i.e.*, over 24 months). A commercial ELISA kit was used in 91.7% of the cases (11/12).

The highest prevalence was obtained in the Republic of Türkiye (48.0%) (Celik and Turutoglu, 2017). The lowest level (0.7%) was detected in Austria (Schrott et al., 2023).

### **Prevalence in goats at the animal level**

**Table 1.** Studies in sheep at animal level

Study	Country/ Region	Study period	Population/ inclusion criteria	Selection	Diagnostic test	Results % (n/N)	Reference
1	Italy/ Ancona and Macerata	Between September 2008 and July 2010	2086 adult female sheep (over 2 years)	Stratified random sampling	ELISA (Pourquier®)	6.3% (129/2086)	Anna Rita et al. 2011
2	Republic of Türkiye /Kars region	Between January and August 2013	450 female sheep	Stratified random sampling	ELISA kit (IDEXX Laboratories®)	6.2% (28/450)	Buyuk et al. 2014
3	Portugal/ Trás-os- Montes and Alto Douro	Between September 2003 and May 2004	3900 sheep (over 2 years)	Stratified random sampling	ELISA (Pourquier®)	3.7% (144/3900)	Coelho et al., 2007
4	Republic of Türkiye /Burdur province	Between October 2014 and February 2015	150 female sheep (2 years and older)	Convenience sampling	ELISA (Parachek®)	48% (72/150)	Celik and Turutoglu, (2017)
5	Scotland/ St Kilda archipelago	Each August during the years 1997–2010	659 sheep	Convenience sampling	ELISA (not reported more information)	1.2% (8/659)	Graham et al., 2016
6	Italy/ Apulia	Between January 2015 to October 2016	13,714 sheep	Convenience sampling	ELISA (ID.vet®)	2.8% (382/13714)	Iarussi et al., 2019
7	Italy/ Apulia	Between January 2015 to October 2016	3189 (sheep in mixed flock)	Convenience sampling	ELISA (ID.vet®)	3.7% (117/3189)	Iarussi et al., 2019
8	Greece/ South- Western Greece	N.R.	291 sheep	Random sampling	Complement fixation test (CFT)	5.5% (16/291)	Ikono- mopoulos et al., 2007
9	Spain/ Andalusia	Between 2015 and 2017	2266 sheep	Random sampling	ELISA (Allied Monitor®)	8.1% (183/2266)	Jiménez- Martín et al., 2022
10	Republic of Cyprus	Between September 2005 and September 2006	3429 sheep	Random sampling	ELISA (Pourquier®)	9.9% (340/ 3429)	Liapi et al., 2011
11	Austria	2020	15,585 sheep	Random sampling	ELISA (ID.vet®)	0.7% (110/15,585)	Schrott et al., 2023
12	German/ Federal states except Berlin, Hamburg and Bremen	Between November 2009 and December 2010	1473 sheep	Convenience sampling	ELISA (CATTLETYPE® LDL)	14% (206/1473) <sup>a</sup>	Stau et al., 2012

N.R. Not reported

<sup>a</sup> Results of proportion were absent from the publication but were calculated using the data on sample size and apparent prevalence

Prevalence studies in goats at the animal level ( $n=17$ ) are shown in Table 2. The studies included in the final analysis were from Austria ( $n=1$ ; 5.9%), France ( $n=1$ ; 5.9%), Germany ( $n=1$ ; 5.9%), Greece ( $n=1$ ; 5.9%), Italy ( $n=6$ ; 35.3%), Portugal ( $n=1$ ; 5.9%), Cyprus ( $n=1$ ; 5.9%), Spain ( $n=4$ ; 23.5%), and Türkiye ( $n=1$ ; 5.9%). Most studies did not provide the age of the animals included in the sample. All studies reported information on the selection of animals tested for MAP. In these studies, random sampling was used in 41.2% ( $n=7$ ) and convenience sampling was used as the selection procedure in 58.8% ( $n=10$ ). The ELISA kit used to detect individual cases in the studies was diverse. Twelve (70.6%) studies reported information on the sensitivity and speci-

ficity of the test used.

Prevalence studies in goats at the animal level in European countries showed a prevalence that ranged between 2.0% and 26%. The highest paratuberculosis prevalence in goats at the animal level was found in Spain (26.0%) (Falconi et al., 2009). Low prevalence was obtained in Austria (2.0%) (Schrott et al., 2023) and France (2.9%) (Mercier et al., 2010).

### Flock level prevalence in sheep

Flock level prevalence studies in sheep ( $n=10$ ) are shown in Table 3. The studies were carried out from 2003 to 2020, though one study did not report the study period. Population and inclusion criteria differed between the studies. Sample sizes ranged from 19 (Ikonomopoulos, 2007) to

**Table 2.** Studies in goat at animal level

Study	Country/Region	Study period	Population/inclusion criteria	Selection	Diagnostic test	Results % (n/N)	Reference
1	Spain/Málaga	Between 2008 and 2009	670 from Malagueña goats, (3–7 years of age)	Random sampling	ELISA (IDEXX®)	7.6% (51/670) <sup>a</sup>	Astorga Márquez et al., 2013
2	Spain/Andalusia	N.R.	3312 (female goat)	Convenience sampling	ELISA (Ingenasa®)	15.4% (511/3312)	Barrero-Domínguez et al., 2019
3	Italy/Garfagnana district Tuscany	N.R.	269 females Age ranged from 2 to 9 years	Convenience sampling	ELISA (ID.vet®)	10% (27/269)	Cecchi et al., 2019
4	Republic of Türkiye /Burdur province	Between October 2014 and February 2015	150 goats, 2 years and older, female	Convenience sampling	ELISA kit (Parachek®)	24% (36/150)	Celik and Turutoglu, 2017
5	Italy/Tuscany	Between October 2008 and March 2009	320 goats (reproductive females (273), randomly chosen, and all reproductive males (47))	Convenience sampling	ELISA (Pourquier®)	5.3% (17/320) <sup>a</sup>	Corrias et al., 2012
6	Spain/north-western Spain	Between 2005 and 2008	219 goats	Convenience sampling	Housemade ELISA adsorbed ELISA	26.0% (57/219)	Falconi et al., 2009



7	Italy/ Northern: Piedmont, Trentino- Alto Adige, Emilia- Romagna, and Lombardy	Between 2019 and 2021	4431 goats	Convenience sampling	ELISA (ID.vet®)	3.7% (164/4431) <sup>a</sup>	Gaffuri et al., 2023
8	Italy/ Garfagnana district/Italy Tuscany	N.R.	294 goats	Convenience sampling	ELISA (ID.vet®)	6.8% (20/294)	Galiero et al., 2017
9	Italy/ Apulia	Between January 2015 and October 2016	7,377 goats (flocks with only goats)	Convenience sampling	ELISA (ID.vet®)	15.7% (1157/7,377)	Iarussi et al., 2019
10	Italy/ Apulia	Between January 2015 and October 2016	1992 goats (goats in mixed flocks)	Convenience sampling	ELISA (ID.vet®)	9.9% (197/1992)	Iarussi et al., 2019
11	Greece/ South- Western Greece	N.R.	117 adult goats	Random sampling	Complement fixation test (CFT)	10.2% (12/117)	Ikono- mopoulos et al., 2007
12	Spain/ Andalusia/ southern Spain	Between 2015 and 2017	1868 goats	Random sampling	ELISA (Allied Monitor)	20.0% (374/1868)	Jiménez- Martín et al., 2022
13	Republic of Cyprus	Between September 2005 and September 2006	4582 goats	Random sampling	ELISA (Pourquier®)	7.9% (362/4582)	Liapi et al., 2011
14	France	2007	11,847 goats	Random sampling	ELISA (Pourquier®)	2.9% (347/11847)	Mercier et al., 2015
15	Portugal/ Tras-os- Montes e Alto Douro	Between February 2014 and January 2015	936 goats	Random sampling	ELISA (ID.Vet®)	12.8% (120/936)	Quintas et al., 2021
16	Austria	2020	6434 goats	Random sampling	ELISA (ID.vet®)	2.0% (126/6434)	Schrott et al., 2023
17	German/ Federal states except Berlin, Hamburg and Bremen	Between November 2009 and December 2010	136 goats	Convenience sampling	ELISA (CATTLETYPE® LDL)	21% (28/136) <sup>a</sup>	Stau et al., 2012

N.R. Not reported

<sup>a</sup> Results of proportion were absent from the publication but were calculated using the data on sample size.

1032 (Schrott et al., 2023). In these studies, random sampling was used in 70% ( $n=7$ ) of studies (three studies used stratified random sampling), and convenience sampling was used as the selection procedure in 30% ( $n=3$ ). In 70% ( $n=7$ ) of studies, information on sensitivity and specificity was

reported. ELISA tests were used in 90% ( $n=9$ ) of studies.

Flock level prevalence studies in sheep in European countries revealed a prevalence between 8.9% (Schrott et al., 2023) and 100% (Celik and Turutoglu, 2017).

**Table 3.** Studies in a sheep flock

Study	Country/Region	Study period	Population/inclusion criteria	Selection	Diagnostic test	Results % (n/N)	Reference
1	Italy/ Ancona and Macerata	Between September 2008 and July 2010	38 sheep flock	Stratified random sampling	ELISA (Pourquier®)	73.7% (28/38)	Anna Rita et al., 2011
2	Republic of Türkiye / Kars Region	Between January and August 2013	26 sheep flock	Stratified random sampling	ELISA kit (IDEXX Laboratories®)	57.7% (13/26)	Buyuk et al., 2014
3	Republic of Türkiye /Burdur province	Between October 2014 and February 2015	15 sheep flock	Convenience sampling	ELISA (Parachek®)	100% (15/15)	Celik and Turutoglu, 2017
4	Portugal/ Trás-os- Montes and Alto Douro	Between September 2003 and May 2004	150 sheep flock	Stratified random sampling	ELISA (Pourquier®)	46.7% (70/150)	Coelho et al., 2007
5	Italy/ Apulia	Between January 2015 to October 2016	165 sheep flock	Convenience sampling	ELISA (ID.vet®)	44.8% (74/165)	Iarussi et al., 2019
	Greece/ South- Western Greece	N.R.	19 sheep flock	Random sampling	Complement fixation test (CFT)	36.8% <sup>a</sup> (7/19)	Ikono- mopoulos et al., 2007
6	Spain/ Andalusia	Between 2015 and 2017	83 sheep flock	Random sampling	ELISA (Allied Monitor®)	66.3% (55/83)	Jiménez- Martín et al., 2022
7	Republic of Cyprus	Between September 2005 until September 2006	49 sheep flock	Random sampling	ELISA (Pourquier®)	53.1% (26/49)	Liapi et al., 2011
8	Austria	2020	1032 sheep flock	Random sampling	ELISA (ID.vet®)	8.9% (92/1032)	Schrott et al., 2023
9	German/ Federal states except Berlin, Hamburg and Bremen	Between November 2009 to December 2010	150 sheep flock	Convenience sampling	ELISA (CATTLETYPE® LDL)	65% <sup>a</sup> (98/150)	Stau et al., 2012

N.R. Not reported

<sup>a</sup> Results of proportion were absent from the publication but were calculated using the data on sample size and apparent prevalence.

### Flock level prevalence in goats

Table 4 displays the results of flock level prevalence studies conducted in goats ( $n=11$ ). The studies were conducted between 2005 to 2021. Population and inclusion criteria differed between the studies. Sample sizes ranged

from 15 (Celik and Turutoglu, 2017) to 638 (Schrott et al., 2023) goat flocks. As the selection procedures, random sampling was used in 54.5% ( $n=6$ ) studies, and convenience sampling was used in 45.5% ( $n=5$ ). In 63.7% ( $n=7$ ) studies, information on sensitivity and specificity

**Table 4.** Studies in goat flock

Study	Country/Region	Study period	Population/inclusion criteria	Selection	Diagnostic test	Results % (n/N)	Reference
1	Spain/Andalusia	N.R.	48 dairy goat flock	Convenience sampling	ELISA (Ingenasa®)	87.5% (42/48)	Barrero-Domínguez et al., 2019
2	Republic of Türkiye /Burdur province	Between October 2014 and February 2015	15 goat flock	Convenience sampling	ELISA kit (Parachek®)	93.3% (14/15)	Celik and Turutoglu, 2017
3	Italy/Northern Piedmont, Trentino-Alto Adige, Emilia-Romagna, and Lombardy	Between 2019-2021	33 goat flock	Convenience sampling	ELISA (ID.vet®)	58% (19/33)	Gaffuri et al., 2023
4	Italy/Apulia	Between January 2015 to October 2016	163 goat flock	Convenience sampling	ELISA (ID.vet®)	82.2% (134/163)	Iarussi et al., 2019
5	Spain/Andalusia/southern Spain	Between 2015 and 2017	70 goat flock	Random sampling	ELISA (Allied Monitor®)	90.0% (63/70)	Jiménez-Martín et al., 2022
6	Greece/South-Western Greece	N.R.	70 goat flock	Random sampling	Complement fixation test (CFT)	54.5% (6/11)	Ikonomopoulos et al., 2007
7	Republic of Cyprus	Between September 2005 until September 2006	72 goat flock	Random sampling	ELISA kit (Pourquier®)	50% (36/72)	Liapi et al., 2011
8	France	2007	105 goat flock	Random sampling	ELISA kit (Pourquier®)	55.2% (58/105)	Mercier et al., 2015
9	Portugal/Trás-os-Montes and Alto Douro	Between February 2014 to January 2015	56 goat flock	Random sampling	ELISA (ID.vet®)	60.7% (34/56)	Quintas et al., 2021

10	Austria	2020	638 goat flock	Random sampling	ELISA (ID.vet®)	1.1% (71/638)	Schrott et al., 2023
11	German/ Federal states except Berlin, Hamburg and Bremen	Between November 2009 to December 2010	17 goat flock	Convenience sampling	ELISA (CATTLETYPE® LDL)	71% (12/17) <sup>a</sup>	Stau et al., 2012

N.R. Not reported

<sup>a</sup> Results of proportion were absent from the publication but were calculated using the data on sample size.

was reported. ELISA tests were used in 90.9% ( $n=10$ ) of studies.

Goat flock level prevalence studies in European countries have shown that the prevalence ranges between 1.1% (Schrott et al., 2023) and 93.3% (Celik and Turutoglu, 2017).

### Individual level prevalence in small ruminants

Table 5 shows prevalence studies for small ruminants at the animal level ( $n=5$ ). Of the studies included in the final analysis, one each was from Austria, Germany, and Greece, and two were from Italy.

**Table 5.** Studies in small ruminants individual

Study	Country/ Region	Study period	Population/ inclusion criteria	Selection	Diagnostic test	Results % (n/N)	Reference
1	Italy	N.R.	601 small ruminants	Convenience sampling	ELISA (ID.vet®)	14.3% <sup>a</sup> (86/601)	De Grossi et al., 2020
2	Italy/ Apulia	Between January 2015 to October 2016	26272 small ruminants	Random sampling	ELISA (ID.vet®)	7.0% (1853/26272)	Iarussi et al., 2019
3	Greece/ South-Western Greece	N.R.	408 small ruminants	Random sampling	Complement fixation test (CFT)	4.4% (28/408)	Ikonomopoulos et al., 2007
4	Austria	2020	22019 small ruminants	Random sampling	ELISA (ID.vet®)	1.1% (236/22019)	Schrott et al., 2023
5	German/ Federal states except Berlin, Hamburg and Bremen	Between November 2009 to December 2010	1609 small ruminants	Convenience sampling	ELISA (CATTLETYPE® LDL)	15% (240/1609)	Stau et al., 2012

N.R. Not reported

<sup>a</sup> Results of proportion were absent from the publication but were calculated using the data on sample size and apparent prevalence.

The study population and inclusion criteria were highly variable. Sample sizes ranged from 408 (Ikonomopoulos et al., 2007) to 26,272 (Iarussi et al., 2019) small ruminants.

A commercial ELISA kit was used in four of five studies. Random sampling was used as the selection procedure in three of five studies.

The lowest level (1.1%) was detected in Austria (Schrott et al., 2023) and the highest prevalence (15.0%) was obtained in Germany (Stau et al., 2012).

## Discussion

The seroprevalence of MAP in sheep and goats across Europe has been studied extensively due to its significant impact on animal health and agricultural productivity. This systematic review summarises individual and flock MAP seroprevalence throughout European countries and describes the diagnostic tests employed in recent decades. Detailed knowledge of the seroepidemiological status of paratuberculosis in sheep and goats is essential for assessing effective prevention measures against this disease in domestic animals (Whittington et al., 2019).

A systematic methodology was used to review studies on the prevalence of MAP in European countries (Coelho et al., 2007; Ikonomopoulos et al., 2007; Falconi et al., 2009; Liapi et al., 2011; Rita et al., 2011; Corrias et al., 2012; Stau et al., 2012; Márquez et al., 2013; Buyuk et al. 2014; Mercier et al., 2015; Graham et al., 2016; Celik and Turutoglu, 2017; Galiero et al., 2017; Barrero-Domínguez et al., 2019; Cecchi et al., 2019; De Grossi et al., 2020; Iarussi et al., 2019; Quintas et al., 2021; Jiménez-Martín et al., 2022; Gaffuri et al., 2023; Schrott et al., 2023). We aimed to gather all possible evidence on the frequency or seroprevalence of paratuberculosis in Europe. We

assumed that there are more similarities in production conditions, breeds, historical processes, and climates among European countries compared to countries in other regions of the world, such as America, Australia, or Asia. The number of publications on paratuberculosis in Europe was greater than expected, indicating an upward trend in paratuberculosis research and a growing interest in this disease and its negative effects. However, the number of publications and studies focused on estimating the seroprevalence or frequency of antibodies MAP was relatively low, although it was comparable to that of a previous review on the same topic in Europe (Nielsen and Toft, 2009).

The limited number of articles in this systematic review on paratuberculosis seroprevalence in these species suggests that research in this area is insufficient.

Over the past two decades, paratuberculosis has been classified as a notifiable disease in 65% of the 43 European countries examined. However, fewer than 50% of these countries have consistently implemented passive surveillance, and only 19% have engaged in active surveillance (Fanelli et al., 2022). This indicates that paratuberculosis may not be a priority for animal health in many European nations.

The absence of relevant information for study quality assessment was previously reported in Europe in a review of prevalence in farm animals (Nielsen and Toft, 2009). In the present review, case definitions, as well as variability in the kits of diagnostic tests and lack of random sampling for animal and flock selection, were the main study flaws included in the final analysis.

The present systematic review on the seroprevalence of sheep and goat paratuberculosis over the last 17 years revealed highly variable seroprevalence among studies, ranging from 0.7% found in sheep

in Austria (Schrott et al., 2023), up to 100% in sheep in Türkiye (Celik and Turutoglu, 2017).

The variability in the prevalence rate of MAP in small ruminants can be attributed to several factors, including geographic location and various unmeasured factors such as climate, production system regime, mechanisation, and professional assistance (Rangel et al., 2015). Improved diagnostic techniques and increased awareness may have led to better detection rates in some areas. Conversely, the implementation of control programmes in certain countries might have contributed to a reduction in prevalence.

The present review refrained from attempting the calculation of true prevalence due to the variability of tests used, and a lack of information on the sensitivity and specificity in some studies conducted in the populations under study. However, we were careful to exclude any studies that clearly exhibited selection bias toward clinical animals or animals with a history or diagnosis of paratuberculosis, to maintain the integrity and validity of our review. The diversity of diagnostic kits and other factors not studied in this work, such as the type of control programmes in those countries/regions, vaccination, and production system used, could explain the high heterogeneity of results from ELISA-based studies.

Our systematic review has several limitations. First, there is a chance that not all the publications related to sheep and goat paratuberculosis were included during document retrieval from the selected databases, although three search databases (PubMed, Scopus, Science Direct) were used, partially because of the keyword selection in the publication itself.

The quality of evidence regarding the seroprevalence of paratuberculosis in the region is restricted due to various flaws

in the study design. The main weakness of these studies was the absence of a clear definition of cases, variability in selected populations, and inclusion criteria for animals and flocks, which could affect the accuracy of prevalence estimates. Another limitation is the high variability in the commercial ELISA diagnostic tests used, variability or lack of information on sensitivity and specificity of diagnostic tests, and lack of random selection of animals and flocks, which also affect the accuracy of prevalence estimates. It is important to note that the apparent increase or decrease in seroprevalence may also be partly due to methodological differences between studies. This underscores the need for standardised approaches in future epidemiological studies to facilitate more accurate comparisons across time and regions.

## Conclusions

This systematic review has provided a comprehensive analysis of MAP seroprevalence in sheep and goats across Europe. The review highlights significant variability in seroprevalence that can be attributed to several factors, including geographic differences, climate, production systems, diagnostic methods, and the presence or absence of formal control programmes. Despite the upward trend in research and growing interest in paratuberculosis, our findings underscore a notable insufficiency in the number of high-quality studies focusing on MAP seroprevalence in small ruminants in Europe. The limited number of articles, coupled with common methodological flaws such as inconsistent case definitions, variability in diagnostic tests, and lack of random sampling, suggests a need for more robust and standardised research approaches in this field. While significant progress has been made in understanding the seroprevalence of MAP in



European small ruminants, there remains a critical need for more rigorous, standardised, and comprehensive research. Such efforts will be essential in developing effective prevention and control measures, ultimately enhancing European animal health and agricultural productivity.

## References

1. AGRAWAL, G., J. AITKEN, H. HAMBLIN, M. COLLINS and T. J. BORODY (2021): Putting Crohn's on the MAP: Five Common Questions on the Contribution of *Mycobacterium avium* subspecies *paratuberculosis* to the Pathophysiology of Crohn's Disease. *Dig. Dis. Sci.* 66, 348-358. 10.1007/s10620-020-06653-0
2. RITA, A. A., N. N. VICTOR, P. SILVIA, P. LUCIANA, D. ANASTASIA and C. VINCENZO (2011): Ovine paratuberculosis: a seroprevalence study in dairy flocks reared in the Marche region, Italy. *Vet. Med. Int.* 782875. 10.4061/2011/782875
3. BARRERO-DOMÍNGUEZ, B., I. LUQUE, B. HUERTA, J. GOMEZ-LAGUNA, Á. GALÁN-RELAÑO, L. GÓMEZ-GASCÓN, M. SÁNCHEZ and R. J. ASTORGA (2019): Paratuberculosis in dairy goat flocks from southern Spain: risk factors associated with seroprevalence. *Vet. Rec.* 185, 600. 10.1136/vr.105465
4. BAUMAN, C. A., A. JONES-BITTON, P. MENZIES, N. TOFT, J. JANSEN and D. KELTON (2016): Prevalence Of paratuberculosis in the Dairy Goat and Dairy Sheep Industries in Ontario. *Can. Vet. J.* 57, 169-175.
5. BUCZINSKI, S., J. ARSENAULT, P. KOSTOULAS, F. CORBIÈRE, G. FECTEAU and N. DENDUKURI (2019): Accuracy of paratuberculosis diagnostic tests in small ruminants: Protocol for a systematic review and meta-analysis. *Anim. Health Res. Rev.* 20, 98-102. 10.1017/S1466252319000082
6. BULUN, H., P. S. BRIDGER, S. SCHILLINGER, et al. (2024): Interferon-gamma producing CD4+ T cells quantified by flow cytometry as early markers for *Mycobacterium avium* ssp. *paratuberculosis* infection in cattle. *Vet. Res.* 55, 69. 10.1186/s13567-024-01324-8.
7. BUYUK, F., O. CELEBI, D. AKCA, S. OTLU, E. TAZEGUL, A. GULMEZ and M. SAHIN (2014): Estimated apparent and true prevalences of paratuberculosis in sheep herds of the Kars Region in Northeastern Turkey. *Vet. Med.* 59, 331-335. 10.17221/7619-VETMED
8. CECCHI, F., F. FRATINI, D. CERRI, P. BANDECHI, C. CANTILE and M. MAZZEI (2019): Small ruminant lentivirus and *Mycobacterium avium* subsp. *paratuberculosis*: co-infection prevalence and preliminary investigation on genetic resistance to both infections in a Garfagnina goat flock. *Large Anim. Rev.* 25, 89-92.
9. CELIK, A. and H. TURUTOGLU (2017): Seroprevalence of paratuberculosis in cattle, sheep and goats in Burdur, Southwestern Turkey. *Vet. World.* 10, 30-36.
10. COELHO, A. C., M. PINTO, S. SILVA, A. COELHO, J. RODRIGUES and R. JUSTE (2007): Seroprevalence of ovine paratuberculosis infection in the Northeast of Portugal. *Small Rumin. Res.* 71, 298-303. 10.1016/j.smallrumres.2006.07.009
11. COELHO, A. C., A. M. COELHO, J. GARCÍA-DIEZ, M. A. PIRES and M. L. PINTO (2018): Detection of *Mycobacterium avium* subsp. *paratuberculosis* by several diagnostics techniques in clinical suspected sheep. *J. Hell. Vet. Med. Soc.* 68, 167-174. 10.12681/jhvms.15601
12. COELHO, A. C., M. L. PINTO, A. M. COELHO and J. RODRIGUES (2009): Comparison of two techniques of isolation of *Mycobacterium avium* subsp. *paratuberculosis* in faecal samples of ovine with clinical suspicion of paratuberculosis. *Pesqui. Vet. Bras.* 2009, 29, 415-420. 10.1590/S0100-736X2009000500010
13. COELHO, A. C., M. L. PINTO, A. M. COELHO, J. RODRIGUES and R. JUSTE (2008): Estimation of the prevalence of *Mycobacterium avium* subsp. *paratuberculosis* by PCR in sheep blood. *Small Rumin. Res.* 76, 201-206. 10.1016/j.smallrumres.2007.12.003
14. COELHO, A. C., M. L. PINTO, A. MIRANDA, A. M. COELHO, M. A. PIRES and M. MATOS (2010): Comparative evaluation of PCR in Ziehl-Neelsen stained smears and PCR in tissues for diagnosis of *Mycobacterium avium* subsp. *paratuberculosis*. *Indian J. Exp. Biol.* 48, 948-950.
15. CORRIAS, F., G. BRAJON, F. SALARI, et al. (2012): Health evaluation in the native Garfagnina goat. *Small Ruminant Research.* 104, 191-194. 10.1016/j.smallrumres.2011.10.005
16. DE GROSSI, L., D. SANTORI, A. BARONE, S. ABBRUZZESE, M. RICCHI and G. A. MARCARIO (2019): Isolation of *Mycobacterium avium* subsp. *paratuberculosis* in the Feces and Tissue of Small Ruminants Using a Non-Automated Liquid Culture Method. *Animals* 10, 20. 10.3390/ani10010020
17. DJØNNE, B. (2010): Paratuberculosis in Goats. In: Behr, M. A. and D. M. Collins: Paratuberculosis: Organism, Disease, Control. CAB International, Oxfordshire, UK (169-178). 10.1079/9781845936136.0169
18. ESPESCHIT, I. F., D. S. S. BASTOS, A. FONSECA JUNIOR, S. A. CARDOSO, M. L. A. FERRARI and M. A. S. MOREIRA (2023): *Mycobacterium avium* subsp. *paratuberculosis* and Crohn's disease: characterization of the interaction with different aspects of the disease. *Braz. J. Microbiol.* 54, 1239-1249. 10.1007/s42770-023-00932-4.
19. FALCONI, C., Á. OLEAGA, J. R. LÓPEZ-OLVERA, R. CASAS, M. PRIETO and C. GORTÁZAR (2010): Prevalence of antibodies against selected agents shared between Cantabrian chamois (*Rupicapra pyrenaica parva*) and domestic goats. *Eur. J. Wildlife Res.* 56, 319-325. 10.1007/s10344-009-0322-z

20. FANELLI, A., M. GALGANO, A. SPOSATO and D. BUONAVOGLIA (2022): Assessment of Paratuberculosis international official reporting in Europe using the information supplied to the WOAHP by National Veterinary Services. *Vet. Ital.* 58, 2. 10.12834/VetIt.2625.16709.3
21. GAFFURI, A., F. BARSÌ, E. MAGNI, et al. (2023): Paratuberculosis, Animal Welfare and Biosecurity: A Survey in 33 Northern Italy Dairy Goat Farms. *Animals (Basel)*. 13, 2346. 10.3390/ani13142346
22. GALIERO, A., B. TURCHI, F. PEDONESE, R. NUVOLONI, C. CANTILE, G. COLOMBANI, M. FORZAN, D. CERRI, P. BANDECCHI and F. FRATINI (2017): Serological, culture and molecular survey of *Mycobacterium avium* subsp. *paratuberculosis* in a goat flock in Tuscany. *Folia Microbiol. (Praha)* 62, 471-477. 10.1007/s12223-017-0518-7
23. GARCIA, A. B. and L. SHALLOO (2015): Invited review: the economic impact and control of paratuberculosis in cattle. *J. Dairy Sci.* 98, 5019-5039. 10.3168/jds.2014-9241
24. GARG, R., P. K. PATIL, S. V. SINGH, et al. (2015): Comparative Evaluation of Different Test Combinations for Diagnosis of *Mycobacterium avium* subsp. *paratuberculosis* Infecting Dairy Herds in India. *Biomol. Res. Int.* 983978, 10.1155/2015/983978
25. GARRIDO, J. M., N. CORTABARRIA, J. A. OGUZZA, G. ADURIZ and R. A. JUSTE (2000): Use of a PCR method on faecal samples for diagnosis of sheep paratuberculosis. *Vet. Microbiol.* 77, 379-386. 10.1016/s0378-1135(00)00323-0
26. GARVEY, M. (2020): *Mycobacterium avium* paratuberculosis: A Disease Burden on the Dairy Industry. *Animals* 10, 1773. 10.3390/ani10101773
27. GRAHAM, A. L., D.H. NUSSEY, J. LLOYD-SMITH, et al. (2016): Exposure to viral and bacterial pathogens among Soay sheep (*Ovis aries*) of the St Kilda archipelago. *Epidemiol. Infect.* 144, 1879-1888. 10.1017/S0950268816000017
28. IARUSSI, F., P. PARADIES, R. SARDARO, G. RUBINO, D. SCALTRITO, E. PIERAGOSTINI and F. PETAZZI (2019): Epidemiology and risk factors of *Mycobacterium avium* subsp. *paratuberculosis* in semi-extensive dairy sheep and goat farms of Apulia, southern Italy. *Small Rumin. Res.* 177, 89-96. 10.1016/j.smallrumres.2019.06.016.
29. IDRIS, S. M., K. H. ELTOM, J. B. OKUNI, L. OJOK, W. A. ELMAGZOUB, A. A. EL WAHED, E. ELTAYEB and A. A. GAMEEL (2022): Paratuberculosis: The Hidden killer of small ruminants. *Animals* 12, 12. 10.3390/ani12010012
30. IKONOMOPOULOS, J., C. BALASKAS, B. KANTZOURA, E. FRAGIADAKI, I. PAVLIK, M. BARTOS, J. C. LUKAS and M.GAZOULI (2007): Comparative evaluation of positive tests to *Mycobacterium avium* subsp. *paratuberculosis* in clinically healthy sheep and goats in south-west Greece using molecular techniques, serology, and culture. *Vet. J.* 174, 337-343. 10.1016/j.tvjl.2006.09.004
31. JIMÉNEZ-MARTÍN, D., I. GARCÍA-BOCANEGRA, M. A. RISALDE, V. FERNÁNDEZ-MOLERA, S. JIMÉNEZ-RUIZ, J. ISLA and D. CANO-TERRIZA (2022): Epidemiology of paratuberculosis in sheep and goats in southern Spain. *Preventive Vet. Med.* 202, 105637. 10.1016/j.prevetmed.2022.105637
32. KAWAJI, S., D. J. BEGG, K. M. PLAIN and R. J. WHITTINGTON (2011): A longitudinal study to evaluate the diagnostic potential of a direct faecal quantitative PCR test for Johne's disease in sheep. *Vet. Microbiol.* 148, 35-44. 10.1016/j.vetmic.2010.07.022
33. LEE, J. H., H. T. PARK, S. SHIM, S. KIM, S. H. WOO, D. Y. KIM and H. S. YOO (2023): Immunopathological mechanisms in the early stage of *Mycobacterium avium* subsp. *paratuberculosis* infection via different administration routes in a murine model. *PLoS One*. 18, e0281880. 10.1371/journal.pone.0281880
34. LIAPI, M., L. LEONTIDES, P. KOSTOULAS, G. BOTSARIS, Y. IACOVOU, C. REES, K. GEORGIU and G. C. SMITH (2011): Bayesian estimation of the true prevalence of *Mycobacterium avium* subsp. *paratuberculosis* infection in Cypriot dairy sheep and goat flocks. *Small Rumin. Res.* 95, 174-178. 10.1016/j.smallrumres.2010.09.010.
35. MALLIKARJUNAPPA, S., L. F. BRITO, S. D. PANT, F. S. SCHENKEL, K. G. MEADE and N. A. KARROW (2021): Johne's Disease in Dairy Cattle: An Immunogenetic Perspective. *Front. Vet. Sci.* 8, 718987. 10.3389/fvets.2021.718987
36. MATOS, A. C., A. P. DIAS, M. MORAIS, L. FIGUEIRA, M. H. MARTINS, M. MATOS, M. L. PINTO and A. C. COELHO (2015): Granuloma coinfection with *Mycobacterium bovis*, *Mycobacterium avium* subsp. *paratuberculosis*, and *Corynebacterium pseudotuberculosis* in five hunted red deer (*Cervus elaphus*) in Portugal. *J. Wildl. Dis.* 51, 793-794. 10.7589/2014-09-240
37. MATOS, A. C., L. FIGUEIRA, M. H. MARTINS, F. LOUREIRO, M. L. PINTO, M. MATOS and A. C. COELHO (2014): Survey of *Mycobacterium avium* subspecies *paratuberculosis* in road-killed wild carnivores in Portugal. *J. Zoo Wildl. Med.* 45, 775-781. 10.1638/2014-0010.1
38. MATOS, A. C., L. FIGUEIRA, M. H. MARTINS, M. MATOS, S. ALVARES, M. L. PINTO and A. C. COELHO (2013): Disseminated *Mycobacterium avium* subsp. *paratuberculosis* infection in two wild Eurasian otters (*Lutra lutra* L.) from Portugal. *J. Zoo Wildl. Med.* 44, 193-195. 10.1638/1042-7260-44.1.193
39. MATOS, A. C., L. FIGUEIRA, M. H. MARTINS, M. MATOS, S. ÁLVARES, A. MENDES, M. L. PINTO and A. C. COELHO (2017): Detection of *Mycobacterium avium* subsp. *paratuberculosis* in kidney samples of red deer (*Cervus elaphus*) in Portugal: Evaluation of different methods. *J. Vet. Med. Sci.* 79, 692-698. 10.1292/jvms.16-0153
40. MERCIER, P., C. BAUDRY, F. BEAUDEAU, H. SEEGERS and X. MALHER (2010): Estimated prevalence of *Mycobacterium avium* subsp. *paratuberculosis* infection in herds of dairy goats in France. *Vet. Rec.* 167, 412-415. 10.1136/vr.c4454

41. QUINTAS, H., O. M. GONZÁLEZ, A. G. VILA, V. PÉREZ and A. C. COELHO (2021): A serosurvey of *Mycobacterium avium* subsp. *paratuberculosis* infection of goats in the North of Portugal. *Acta Vet. Hung.* 10.1556/004.2021.00051
42. MIRANDA, A., M. A. PIRES, M. L. PINTO, L. SOUSA, R. SARGO, J. RODRIGUES, A. C. COELHO, M. MATOS and A. M. COELHO (2009): *Mycobacterium avium* subsp. *paratuberculosis* in a diamond sparrow. *Vet. Rec.* 165, 184. 10.1136/vr.165.6.184-a
43. MIRANDA, C., M. MATOS, I. PIRES, P. RIBEIRO, S. ÁLVARES, M. VIEIRA-PINTO and A. C. COELHO (2011): *Mycobacterium avium* subsp. *paratuberculosis* infection in slaughtered domestic pigs for consumption detected by molecular methods. *Food Res. Int.* 44, 3276-3277. 10.1016/j.foodres.2011.09.010
44. NIELSEN, S. S. and N. TOFT (2009): A review of prevalences of paratuberculosis in farmed animals in Europe. *Prev. Vet. Med.* 88, 1-14. 10.1016/j.prevetmed.2008.07.003
45. PAGE, M. J. and J. E. MCKENZIE (2020): PRISMA 2020 explanation and elaboration: updated guidance and exemplars for reporting systematic reviews. *BMJ* 372. 10.1136/bmj.n160
46. PÉREZ, V., J. F. GARCÍA-MARÍM and J. J. BADIOLA (1996): Description and classification of different types of lesion associated with natural paratuberculosis infection in sheep. *J. Comp. Pathol.* 114, 107-122. 10.1016/s0021-9975(96)80001-6
47. RANGEL, S. J., J. PARÉ, E. DORÉ, J. C. ARANGO, G. CÔTÉ, S. BUCZINSKI, O. LABRECQUE, J. H. FAIRBROTHER, J. P. ROY, V. WELLEMANS and G. A. FECTEAU (2015): A systematic review of risk factors associated with the introduction of *Mycobacterium avium* spp. *paratuberculosis* (MAP) into dairy herds. *Can. Vet. J.* 56, 169-177.
48. SCHROTT, J., E. SODOMA, M. DÜNSER, A. TICHY and J. L. KHOL (2023): *Mycobacterium avium* subsp. *paratuberculosis* in Sheep and Goats in Austria: Seroprevalence, Risk Factors and Detection from Boot Swab Samples. *Animals* 13, 1517. 10.3390/ani13091517
49. SHARMA, S., K. SHARMA, R. KUMAR, D. DAYAL, S. DHANDA, N. KUMAR, K. K. CHAUBEY, S. V. SINGH, S. BANGER and V. SHARMA (2024): Evaluation of Immune Exhaustion and Co-Inhibitory Receptor Expression in *Mycobacterium avium* subsp. *paratuberculosis* (MAP) Seropositive Diarrhoeic Bovines. *Pathogens* 13, 473. 10.3390/pathogens13060473
50. SINGH, S. V., P. K. SINGH, S. GUPTA, K. K. CHAUBEY, B. SINGH, A. KUMAR, A. V. SINGH and N. KUMAR (2013): Comparison of microscopy and blood-PCR for the diagnosis of clinical Johne's disease in domestic ruminants. *Iranian J. Vet. Res.* 14, 345-349.
51. STAU, A., B. SEELIG, D. WALTER, C. SCHROEDER and M. GANTER (2012): Seroprevalence of *Mycobacterium avium* subsp. *paratuberculosis* in small ruminants in Germany. *Small Rumin. Res.* 105, 361-365. 10.1016/j.smallrumres.2012.03.008.
52. SWEENEY, R. W. (2011): Pathogenesis of paratuberculosis. *Vet. Clin. North Am.: Food Anim. Pract.* 27, 537-546. 10.1016/j.cvfa.2011.07.001
53. WHITLOCK, R. H., S. J. WELLS, R. W. SWEENEY and J. V. TIEM (2000): ELISA and faecal culture for paratuberculosis (Johne's disease): sensitivity and specificity of each method. *Vet. Microbiol.* 77, 387-398. 10.1016/s0378-1135(00)00324-2
54. WHITTINGTON, R., K. DONAT, M.F. WEBER, et al. (2019): Control of paratuberculosis: who, why and how. A review of 48 countries. *BMC Vet. Res.* 15, 198. 10.1186/s12917-019-1943-4

## Seroprevalencijska infekcija s *Mycobacterium avium* subsp. *paratuberculosis* u malih preživača u Europi – sistematski pregled

Hélder QUINTAS, DVM, MSc, PhD, Mountain Research Center (CIMO), Bragança Polytechnic University (IPB), Campus de Santa Apolónia, 5300-253 Bragança, Portugal; Júlio BENAVIDES, Instituto de Ganadería de Montaña, CSIC-University of León, Grulleros, 24346 León, Spain; João JACOB-FERREIRA, CIMO, IPB, Campus de Santa Apolónia, Bragança, 5300-253, Portugal; Paulo AFONSO, DVM, MSc, PhD, Animal and Veterinary Research Centre (CECAV), University of Trás-os-Montes e Alto Douro (UTAD), Quinta de Prados, 5000-801 Vila Real, Portugal; and CIMO, IPB, Campus de Santa Apolónia, 5300-253 Bragança, Portugal; Ana Cláudia COELHO, DVM, MSc, PhD, Department of Veterinary Sciences, School of Agrarian and Veterinary Sciences (ECAV), CECAV, UTAD, Quinta de Prados, 5000-801 Vila Real, Portugal

Paratuberkuloza, poznata i kao Johneova bolest, je granulomatozni enteritis u domaćih i divljih preživača koju prouzroči bakterija *Mycobacterium avium* subsp. *paratuberculosis*. Razumijevanje prevalencije

ove bolesti u malih preživača ključno je za strategije kontrole i prevencije bolesti. Sistematski pregled literature je proveden uporabom PubMed, ScienceDirect i Scopus baza podataka za pronalaženje

svih članaka u svezi seroprevalencije *Mycobacterium avium* subsp. *paratuberculosis* u ovaca i koza u Europi, objavljenih od siječnja 2006. do prosinca 2023. Početna pretraga postojećih publikacija koje su donosile sistematski pregled i izvjestile o primarnim studijama provedena je pretragom dostupnih baza podataka. Za konačni odabir studija, obavljen je početni probir osnovne podobnosti i detaljna procjena kvalitete. Nakon odabira studija, izvedeni su odgovarajući podatci. Detaljna procjena generala je 21 publikaciju s izvješćima o 55 studija, 22 (40,0 %) na ovcama (12 na razini životinja i 10 na razini stada) i 28 (50,9 %) na kozama (17 na razini životinja i 11 na razini stada) te 5 (9,1 %) na različitim malim preživcima na razini životinja. Sveukupno, 34 (61,8 %) bile su studije na razini životinja, dok je 21 studija (38,2 %) bila na razini stada. Populacija i kriteriji uključuje-

nja vrlo su se razlikovali među studijama. Veličina uzorka kretala se između 291 do 15.585 životinja. Većina studija izvještavala je o testiranju odraslih životinja (tj. više od 24 mjeseci starosti). Komercijalni ELISA kompleti su rabljeni u većini studija. Najveća prevalencija dobivena je u ovaca (100 %) u Republici Turskoj, a najmanja je otkrivena u ovaca (0,7 %) u Austriji. Općenito, rezultati su ukazali da se MAP antitijela često otkrivaju među malim preživcima u nekim zemljama i da postoji potreba za standardizacijom definicija slučajeva da bi se poboljšala točnost procjena prevalencije. Potrebno je dodatno istraživanje da bi se razumjeli faktori rizika povezani s MAP infekcijom u malih preživaca i za razvoj učinkovitih strategija kontrole i prevencije.

**Ključne riječi:** *Europa, Mycobacterium avium* subsp. *paratuberculosis*, sistematski pregled