



Understanding ammonia emissions: the impact of feed chemical composition on dairy Holstein farms

Mirna Gavran^{1*}, Danko Šinka², Marko Samardžija³, Ranko Gantner¹,
Zvonimir Steiner¹ and Vesna Gantner¹

¹ Josip Juraj Strossmayer University of Osijek, Faculty of Agrobiotechnical Sciences, Osijek, Croatia;

² PIK Vinkovci plus d.o.o., Vinkovci, Croatia

³ Faculty of Veterinary Medicine, University of Zagreb, Zagreb, Croatia

GAVRAN, M., D. ŠINKA, M. SAMARDŽIJA, R. GANTNER, Z. STEINER, V. GANTNER: Understanding ammonia emissions: the impact of feed chemical composition on dairy Holstein farms. Vet. arhiv 95, 187-196, 2025.

ABSTRACT

The aim of this study was to explore the relationship between the chemical composition of cattle feed and the emission of ammonia. To achieve this, we conducted a thorough chemical analysis of the feed, examining factors such as digestible crude protein, rumen undegradable protein, bacterial production of metabolic protein, and levels of methionine and lysine. Our research focused on how these feed components impact both milk yield and ammonia emissions. The study found that there was a positive correlation between digestible crude protein and milk yield, while rumen undegradable protein had a strong positive correlation with milk yield. Additionally, the study revealed a negative correlation between bacterial production of metabolic protein and daily milk yield. The concentration of methionine was weakly positively associated with milk yield, while lysine had a weak negative association with milk yield. The concentration of lysine in the rations was positively correlated with ammonia emissions, while methionine was negatively correlated with it. The study also found that rumen undegradable protein had a significant negative correlation with ammonia emission. Overall, the study highlights the importance of effective feeding management practices in optimizing animal productivity and minimizing ammonia emissions, to promote sustainable agriculture.

Key words: ammonia emission; feed chemical composition; milk recording; dairy cows

Introduction

Livestock farming is a significant contributor to the release of ammonia (NH₃) emissions into the environment, which poses a major environmental threat on a global scale. These emissions lead to soil acidification, eutrophication of aquatic ecosystems, and have a significant role in causing global warming ([SANCHIS et al., 2019](#); [YANG](#)

et al., 2022). Livestock emissions are primarily driven by beef and dairy cattle, which account for 77% of all livestock emissions ([ECA, 2021](#)). During the breeding of dairy cows, urea and uric acid are broken down by urease-producing microorganisms, resulting in the release of NH₃ through microbiological hydrolysis. This process

* Corresponding author:

Mirna Gavran, mag.ing.agr. Josip Juraj Strossmayer University of Osijek, Faculty of Agrobiotechnical Sciences, Vladimira Preloga 1, 31000 Osijek, Croatia, e-mail: mgavran@fazos.hr

leads to the formation of NH_4^+ as a byproduct ([SANCHIS et al., 2019](#); [YANG et al., 2022](#)). The indoor climatic factors that significantly affect NH_3 emissions from cattle buildings are temperature, air velocity, and relative humidity ([SCHRADER et al., 2012](#); [SAHA et al., 2014](#); [SANCHIS et al., 2019](#); [QU et al., 2021](#); [TABASE et al., 2023](#)). In addition to these climatic factors, there are other factors, such as dairy cow species, feed and manure management, that also significantly contribute to NH_3 emissions ([SANCHIS et al., 2019](#); [YANG et al., 2022](#)). It is crucial to focus on these parameters to manage and reduce ammonia emissions in livestock operations.

In the rumen, ammonia is produced when the degradable nitrogen (N) fraction of the feed is hydrolysed. This ammonia is then readily utilized by microbes to synthesize microbial protein ([PUTRI et al., 2021](#); [RODRIGUES et al., 2022](#)). In ruminants, an excess of ammonia in the rumen is transformed into urea in the liver. This urea is either recycled back to the rumen through saliva and the epithelium, or excreted in the urine via the kidneys ([CHOJNACKA et al., 2021](#); [SOUZA et al., 2021](#); [RODRIGUES et al., 2022](#)). One of the factors mentioned above is feed. The faeces and urine characteristics of animals are influenced by various factors, such as the chemical composition and physical properties of their feed, the N content and energy-N balance of the diet, as well as the physiological stage of the animal. Specifically, the amount of N excreted is affected by these factors ([DIJKSTRA et al., 2018](#); [SAJEEV et al., 2018](#); [CHEN et al., 2020](#); [RODRIGUES et al., 2022](#)). One potential solution for reducing NH_3 emissions from manure involves manipulating animal diets to decrease the excretion of nitrogen (N) in manure. This approach presents an opportunity for cost-effective and efficient reduction of NH_3 emissions from manure ([HRISTOV et al., 2015](#); [KATONGO and YAN, 2022](#)). Numerous studies have provided conclusive evidence that there is a positive correlation between dietary crude protein (CP) levels and NH_3 emissions from cattle manure slurries. In simpler terms, as the CP levels in the diet of cattle increase, so do the emissions of NH_3 from their manure. [BURGOS](#)

et al. (2010) and [EDOUARD et al. \(2019\)](#) have reported that increasing the dietary CP contents from 15 to 21% DM and 120 to 180 g/kg DM, respectively, resulted in 2.3- to 4.6-fold increases in NH_3 emissions associated with manure slurries from dairy cows. The excretion of N, specifically urinary N, and the consequent emission of NH_3 from slurry are heavily influenced by the dietary concentration of CP ([YAN et al., 2006](#); [SPIEHS and VAREL, 2009](#); [ANGELIDIS et al., 2019](#)). Elevating the level of rumen undegradable protein (RUP) in animal diets is likely to boost the overall nitrogen (N) efficiency and facilitate the delivery of sufficient metabolizable protein (MP) to the small intestine ([CALSAMIGLIA et al., 2010](#); [DE MELO COELHO et al., 2022](#)).

Methionine and lysine are two crucial amino acids classified as essential due to their inability to be synthesized by the organism. [SAJID et al. \(2024\)](#) stated that most proteins, including those present in livestock feed, do not contain all the essential amino acids required for optimal milk production and tissue development in high or medium-producing dairy cattle. Methionine is one of the essential amino acids that is often lacking in these proteins. Although the protein synthesized by bacteria is a valuable source of amino acids (AA), it is not enough to satisfy the requirements for tissue growth and milk production in dairy cows. This fact has been supported by recent studies conducted by [ROBINSON et al. \(2004\)](#), [MAVROMMATIS et al. \(2021\)](#), and [SAJID et al. \(2024\)](#). Supplementing dairy cow rations with methionine faces challenges due to microbial activity in the rumen. To improve its bioavailability, rumen-undegraded or rumen-encapsulated versions of this amino acids can be provided. Optimizing the lysine to methionine ratio in the diet is crucial for maximizing microbial protein synthesis. Achieving this balance allows for a reduction in the crude protein content in the diet, improving dairy cow reproduction, and reducing nitrogen excretion for environmental sustainability ([PARK et al., 2020](#); [SAJID et al., 2024](#)).

The aim of this study was to investigate the link between the chemical makeup of cattle feed and the release of ammonia. The chemical analysis of cattle feed consisted of an examination of

several factors, such as digestible crude protein, rumen undegradable protein, bacterial production of metabolic protein, and levels of methionine and lysine.

Materials and methods

This research gathered test-day data from Holstein cows in Croatia, specifically near Osijek in Slavonia, between January 2020 and December 2023, during their regular milk recording. Milk yield measurements were taken every four weeks using the AT4/BT4 milk recording method, and corresponding samples were analysed. The milk samples collected were labelled with the corresponding animal code and sampling date, and then sent to the Central Laboratory for Milk Quality Control, which is under the auspices of the Croatian Agency for Agriculture and Food. Infrared spectrophotometry was used to determine the proportion of urea in the milk samples. The milk sample collection procedures during recording and laboratory testing were carried out following the specifications set forth by the International Committee for Animal Recording ([ICAR, 2017](#)). After conducting a thorough analysis of milk recording data, adhering to ICAR standards, and correcting any illogical variable values, we were able to include a total of 35,245 Holstein test-day records in the updated database. SAS/STAT ([SAS, 2019](#)) was used to prepare the data and conduct dispersion and correlation analysis. The equations below can be used for computation of the milk urea nitrogen (MUN) and ammonia emissions (AM-EMISS) in dairy cows:

$$MUN \text{ (mg/dL)} = UREA * 0.46 \text{ ([SPIEKERS and OBERMAIER, 2012](#))}$$

$$AM-EMISS \text{ (g/cow daily)} = 25.0 + 5.03 * MUN \text{ ([BURGOS et al., 2010](#))}$$

In technical terms, UREA signifies the amount of urea present in the milk on a daily basis, measured in milligrams per decilitre (mg/dL). MUN, on the other hand, represents the concentration of milk urea nitrogen in mg/dL. Lastly, AM-EMISS denotes the quantity of ammonia emission per cow on a daily basis, measured in grams (g).

Near Infrared Spectroscopy (NIR) is an analytical method used for determining the digestible crude protein, rumen undegradable protein, and bacterial production of metabolic protein, methionine, and lysine in a ration. It is a versatile and reliable technology that can be used for analysing a diverse range of sample types. NIR works by detecting changes in the transmission or reflection of NIR energy from a sample, which is dependent on its chemical composition. However, to establish an NIR calibration, a primary reference method is required as NIR is a secondary technique. This calibration is essentially a mathematical correlation linking the raw NIR data obtained from samples to the specific chemical constituents or properties of interest.

Results

Table 1 presents the basic statistical parameters of the feed ingredients analysed, including digestible crude protein, rumen undegradable

Table 1. Basic statistical parameters of analysed feed chemical composition (proteins and amino acids)

Variable	N	Mean	SD	CV	SE	Min	Max
Digestible crude protein, % DM	432	16.12	1.10	6.84	0.05	11.60	18.70
Rumen undegradable protein, % DM	394	5.51	0.72	13.06	0.04	3.30	6.80
Bacterial production of metabolic protein, % Total MP	372	53.86	4.72	8.76	0.25	46.20	69.40
Methionine, % Rqd	386	100.13	6.80	6.79	0.35	83.00	115.00
Lysine, % Rqd	386	91.23	4.80	5.26	0.24	80.00	108.00

protein, bacterial production of metabolic protein, methionine, and lysine.

The following are the statistical details of the nutritional composition of animal feed. On average, the digestible crude protein in the animals' rations was 16.12%, with a standard deviation of 1.10%. The minimum and maximum values were 11.60% and 18.70%, respectively. Additionally, the average rumen undegradable protein was 5.51%, with a standard deviation of 0.72%. The minimum and maximum values for this metric were 3.30% and 6.80%. Moreover, the average bacterial production of metabolic protein was 53.86%, with a standard deviation of 4.72%. The minimum and maximum values for this metric were 46.20% and 69.40%. Lastly, the average methionine value was 100.13%, with a standard deviation of 6.80%. The minimum and maximum values for this metric were 83.00% and 115.00%. Similarly, the average lysine value was 91.23%, with a standard deviation of 4.80%. The minimum and maximum values for this metric were 80.00% and 108.00%.

Table 2 shows the correlation analysis results between the chemical composition of the feed (specifically, proteins and amino acids) and the derivatives of test-day records (namely, daily milk yield and ammonia emissions).

Table 2 shows the relationship between the examined feed components, including digestible crude protein, rumen undegradable protein, bacterial production of metabolic protein, methionine, and lysine, and the test-day performance traits, namely daily milk yield and ammonia emissions. The statistical analysis conducted of the data revealed interesting insights. Specifically, the results showed that there was a weak positive correlation between the amount of digestible crude protein and the daily milk yield ($r=0.366$, $P<0.0001$). On the other hand, a strong positive correlation was observed between the amount of rumen undegradable protein and the daily milk yield ($r=0.597$, $P<0.0001$). The results of the study indicate that there is a substantial relationship between two important factors in milk

Table 2. Correlation between the analysed feed chemical composition (proteins and amino acids) and test-day records derivatives (daily milk yield and ammonia emissions)

Trait	Daily milk yield	Ammonia emissions
Ammonia emission, g/cow daily	-0.057 <.0001 94868	1.000 94868
Digestible crude protein, %	0.366 <.0001 62215	0.040 <.0001 61139
Rumen undegradable protein, %	0.597 <.0001 60700	-0.009 0.038 59648
Bacterial production of metabolic protein in relation to total metabolic protein, %	-0.671 <.0001 56267	0.051 <.0001 55269
Methionine, %	0.202 <.0001 57371	-0.016 0.000 56362
Lysine, %	-0.179 <.0001 57371	0.014 0.001 56362

production – the bacterial production of metabolic protein and the daily milk yield. Specifically, the findings reveal a robust negative correlation between these two variables, with a correlation coefficient of -0.671 and a P-value of less than 0.0001. This suggests that as bacterial production of metabolic protein increases, daily milk yield decreases, and vice versa. The results of this study have important implications for understanding the factors that influence milk production, and may be useful in developing strategies to optimize milk yield. The statistical analysis revealed a weak positive association (correlation coefficient $r=0.202$, $P<0.0001$) between methionine and daily milk yield, indicating that higher levels of methionine were associated with slightly higher milk production. Conversely, a weak negative association (correlation coefficient $r=-0.179$, $P<0.0001$) was found between lysine and daily milk yield, suggesting that higher levels of lysine were associated with slightly lower milk production. The results of our analysis demonstrate a strong correlation between the feed ingredients given to the animals and their daily milk production. This highlights the significance of effective feeding management practices in optimizing animal productivity and reducing feeding costs. The correlation coefficient between the analysed feed ingredients and ammonia emissions was statistically significant ($P<0.0001$; $P<0.005$), although the coefficient itself was low. According to the analysis, there is a negative correlation coefficient between the concentration of methionine and the emission of ammonia. This suggests that increasing the methionine concentration in the animals' rations may help reduce the emission of ammonia. The relationship between the concentration of lysine in the rations and the emission of ammonia was observed to be positively correlated. This implies that a higher concentration of lysine in the rations results in a higher emission of ammonia from the dairy farm. The study highlights the importance of considering lysine concentration in the formulation of dairy farm rations, to minimize ammonia emissions and promote sustainable agriculture. The data indicate a clear and positive relationship between ammonia

emission and the levels of digestible crude protein and the bacterial production of metabolic protein. This suggests that an increase in the amounts of these ingredients in the ration could lead to a corresponding increase in emission. In other words, the more digestible crude protein and metabolic protein bacteria produce, the more ammonia is likely to be emitted. The study conducted on the impact of rumen undegradable protein in the diet found a significant negative correlation between the levels of rumen undegradable protein and ammonia emission. The results indicate that a higher level of rumen undegradable protein in the diet can lead to a reduction in ammonia emissions. This suggests that the inclusion of rumen undegradable protein in animal feed can be an effective strategy to mitigate the release of ammonia into the environment, thereby contributing to environmental sustainability.

Discussion

Numerous studies have confirmed a significant correlation between milk urea concentration and its derivatives, such as milk urea nitrogen and ammonia emission. The strong association between milk urea content and nitrogen content in animal urine and manure has been established by several researchers, including [BURGOS et al. \(2010\)](#), [ECKERSALL and BELL \(2010\)](#), [KLEIN et al. \(2011\)](#), and [SPEK et al. \(2013\)](#). The dietary CP (crude protein) content has a significant impact on the urinary nitrogen (N) excretion, which in turn affects the level of ammonia (NH_3) emissions ([SMITS et al., 1995](#); [DE BOER et al., 2002](#); [VAN DUINKERKEN et al., 2005](#); [TODD et al., 2006](#); [POWELL et al., 2008](#); [POWELL et al., 2011](#); [VAN DUINKERKEN et al., 2011](#); [BOUGOUIN et al., 2016](#); [EDOUARD et al., 2019](#); [MÜLLER et al., 2021](#); [DEVANT et al., 2022](#); [SCHRADER et al., 2023](#)). According to [HRISTOV et al. \(2011\)](#), when it comes to reducing dietary crude protein (CP) in livestock feed, it is crucial to assess the impact on animal performance and the associated costs of gain, while also paying attention to minimizing NH_3 emissions. This is because even a slight reduction in CP can potentially lead to negative

effects on the animals' productivity, which, in turn, can affect the overall economic viability of the farm. In their recent study, [DE MELO COELHO et al. \(2022\)](#) recommended the use of various sources of rumen undegradable protein (RUP) in animal feed formulations to lower nitrogen loss through urine. This approach can help reduce the release of harmful gases such as N_2O , CH_4 , and NH_3 into the

environment.

[KROBER et al. \(2000\)](#) and [HRISTOV et al. \(2011\)](#) found that incorporating methionine in low crude protein diets can significantly reduce nitrogen excretion in urine as a proportion of the total nitrogen excretion. Several studies have examined the efficacy of lysine use in growing cattle, and the findings indicate that roughly 40% of the lysine added is used efficiently by the animals ([BATISTA et al., 2016](#); [HUSSEIN et al., 2016](#)). This finding is based on animal response curves, and suggests that there is significant room for improvement in the use of lysine in cattle feed. Studies conducted by [AWAWDEH et al. \(2005\)](#) and [HUSSEIN et al. \(2016\)](#) have shown that modifying animal metabolism can improve the efficiency of lysine use, leading to a reduction in the cost of dietary lysine supplementation. A feasible strategy to enhance the lysine efficiency in protein synthesis is to administer ammonia (NH_3) loads to cattle. When excess NH_3 is fed in the form of feed-grade urea, the liver may spare lysine from catabolism. This preference is given to the removal of toxic NH_3 from the bloodstream ([AWAWDEH et al., 2005](#); [HUSSEIN et al., 2016](#)). It is important to assess the environmental impact of cattle excreting more nitrogen when considering the effectiveness of a particular method. A study by [HUSSEIN et al. \(2016\)](#) found that increasing urea infusions leads to greater daily nitrogen retention, suggesting that increasing ammonia levels in the rumen may improve lysine use efficiency. [SAJID et al. \(2024\)](#) highlighted the fact that ruminant diets that contain high levels of crude protein are linked to elevated nitrogen excretion, which can cause environmental pollution and increase feeding expense. One of the main difficulties in ruminant farming is to enhance the reproductive health and performance of the animals, while simultaneously reducing

nitrogen excretion (N_2 and NH_3). According to this study, to address this issue, it is recommended to incorporate essential amino acids, especially lysine and methionine, in the diets of ruminants.

Conclusions

The objective of this research was to establish

the correlation between the chemical composition of cattle feed and the emission of ammonia. The chemical assessment of cattle feed involved an in-depth analysis of various factors, including digestible crude protein, rumen undegradable protein, bacterial production of metabolic protein, and levels of methionine and lysine. The study analysed the relationship between feed components and milk yield and ammonia emissions. The results showed a positive correlation between digestible crude protein and milk yield, and a negative correlation between bacterial production of metabolic protein and milk yield. Methionine was found to have a weak positive correlation with milk yield, while lysine had a weak negative correlation. The study also found that increasing methionine concentration in the rations may reduce ammonia emissions, while increasing lysine concentration may increase emissions. The inclusion of rumen undegradable protein in animal feed can help reduce ammonia emissions. Our research findings reveal a significant link between the specific feed ingredients provided to the animals and the amount of milk they produce on a daily basis. This highlights the critical role of efficient and effective feeding management practices in maximizing animal productivity and minimizing the costs associated with feeding. By ensuring that the animals receive the appropriate feed and nutrition, farmers can achieve higher milk yields and reduce unnecessary expense.

Financial support statement

The research and dissemination activities were financially supported by the Fund for Bilateral Relations within the Financial Mechanism of the European Economic Area and the Norwegian Financial Mechanism for the period of 2014-2021 (Grant number: 04-UBS-U-0031/23-14).

Declaration of competing interest

No potential conflicting interest was reported by the authors.

References

- ANGELIDIS, A., L. CROMPTON, T. MISSELBROOK, T. YAN, C. K. REYNOLDS, S. STERGIADIS (2019): Evaluation and prediction of nitrogen use efficiency and outputs in faeces and urine in beef cattle. *Agric. Ecosyst. Environ.* 280, 1-15.
<https://doi.org/10.1016/j.agee.2019.04.013>
- AWAWDEH, M. S., E. C. TITGEMEYER, K. C. MCCUISTION, D. P. GNAD (2005): Ruminant ammonia load affects leucine utilization by growing steers. *J. Anim. Sci.* 83, 2448-2454.
<https://doi.org/10.2527/2005.83102448x>
- BATISTA, E. D., A. H. HUSSEIN, E. DETMANN, M. D. MIESNER, E. C. TITGEMEYER (2016): Efficiency of lysine utilization by growing steers. *J. Anim. Sci.* 94, 648-655.
<https://doi.org/10.2527/jas.2015-9716>
- BOUGOUIN, A., A. LEYTEM, J. DIJKSTRA, R. S. DUNGAN, E. KEBREAB (2016): Nutritional and environmental effects on ammonia emissions from dairy cattle housing: A meta-analysis. *J. Environ. Qual.* 45, 1123-1132.
<https://doi.org/10.2134/jeq2015.07.0389>
- BURGOS, S. A., N. M. EMBERTSON, Y. ZHAO, F. M. MITLOEHNER, E. J. DEPETERS, J. G. FADEL (2010): Prediction of ammonia emission from dairy cattle manure based on milk urea nitrogen: Relation of milk urea nitrogen to ammonia emissions. *J. Dairy Sci.* 93, 2377-2386.
<https://doi.org/10.3168/jds.2009-2415>
- CALSAMIGLIA, S., A. FERRET, C. K. REYNOLDS, N. B. KRISTENSEN, A. M. van VUUREN (2010): Strategies for optimizing nitrogen use by ruminants. *Animal* 4, 1184.
<https://doi.org/10.1017/S1751731110000911>
- CHEN, X., G. FINNEY, H. ZHENG, H. WANG, A. W. GORDON, C. P. FERRIS, E. MAGOWAN, T. YAN (2020): Modern Holstein-origin dairy cows within grassland-based systems partition more feed nitrogen into milk and excrete less in manure. *Sci. Total Environ.* 727, 138561.
<https://doi.org/10.1016/j.scitotenv.2020.138561>
- CHOJNACKA, K., K. MIKULA, G. IZYDORCZYK, D. SKRZYPCZAK, A. WITEK-KROWIAK, A. GERSZ, K. MOUSTAKAS, J. IWANIUK, M. GRZĘDZICKI, M. KORCZYNSKI (2021): Innovative high digestibility protein feed materials reducing environmental impact through improved nitrogen-use efficiency in sustainable agriculture. *J. Environ. Manag.* 291, 112693.
<https://doi.org/10.1016/j.jenvman.2021.112693>
- DE BOER, I. J. M., M. C. J. SMITS, H. MOLLENHORST, G. VAN DUINKERKEN, G. J. MONTENY (2002): Prediction of ammonia emission from dairy barns using feed characteristics Part I: relation between feed characteristics and urinary urea concentration. *J. Dairy Sci.* 85, 3382-3388.
[https://doi.org/10.3168/jds.S0022-0302\(02\)74425-1](https://doi.org/10.3168/jds.S0022-0302(02)74425-1)
- DE MELO COELHO, L., L. DE FIGUEIREDO BRITO, J. D. MESSANA, A. DA SILVA CARDOSO, G.M. CARVALHO, R. DE NAZARÉ SANTOS TORRES, R.S. CARLOS, E. MALHEIROS BRAGA, PESSÔA M. C. DA CRUZ, T. TERESINHA BERCHIELLI (2022): Effects of rumen undegradable protein sources on nitrous oxide, methane and ammonia emission from the manure of feedlot-finished cattle. *Sci Rep*, 12, 9166.
<https://doi.org/10.1038/s41598-022-13100-9>
- DEVANT, M., A. PÉREZ, C. MEDINYÀ, J. RIERA, J. GRAU, B. FERNÁNDEZ, F. X. PRENAFETA-BOLDÚ (2022): Effect of decreasing dietary crude protein in fattening calves on the emission of ammonia and greenhouse gases from manure stored under aerobic and anaerobic conditions. *Animal* 16, 100471.
<https://doi.org/10.1016/j.animal.2022.100471>
- DIJKSTRA, J., A. BANNINK, P. M. BOSMA, E. A. LANTINGA, J. W. REIJS (2018): Modeling the effect of nutritional strategies for dairy cows on the composition of excreta nitrogen. *Front. Sustain. Food Syst.* 2, 16.
<https://doi.org/10.3389/fsufs.2018.00063>
- ECA (2021): European Court of Auditors. Common Agricultural Policy and climate: Half of EU climate spending but farm emissions are not decreasing. (https://www.eca.europa.eu/hr/publications/SR21_16)
- ECKERSALL, P. D., R. BELL (2010): Acute phase proteins: Biomarkers of infection and inflammation in veterinary medicine. *Vet. J.* 185, 23-27.
<https://doi.org/10.1016/j.tvjl.2010.04.009>
- EDOUARD, N., A. CHARPIOT, P. ROBIN, E. LORINQUER, J.-B. DOLLÉ, P. FAVERDIN (2019): Influence of diet and manure management on ammonia and greenhouse gas emissions from dairy barns. *Animal* 13, 2903-2912.
<https://doi.org/10.1017/S1751731119001368>
- HRISTOV, A., K. S. HEYLER, E. W. SCHURMAN, K. GRISWOLD, P. A., TOPPER, M. HILE, V. A. ISHLER, E. FABIAN-WHEELER, S. DINH (2015): Case study: Reducing dietary protein decreased the ammonia emitting potential of manure from commercial dairy farms. *Prof. Anim. Sci.* 31, 68-79.
<https://doi.org/10.15232/pas.2014-01360>
- HRISTOV, A. N., M. HANIGAN, A. COLE, R. TODD, T. A. McALLISTER, P. M. NDEGWA, A. ROTZ (2011):

- Review: Ammonia emissions from dairy farms and beef feedlots. *Can. J. Anim. Sci.* 91, 1-35.
<https://doi.org/10.4141/CJAS10034>
- HUSSEIN, A. H., E. D. BATISTA, M. D. MIESNER, E. C. TITGEMEYER (2016): Effect of ruminal ammonia supply on lysine utilization by growing steers. *J. Anim. Sci.* 94, 656-664.
<https://doi.org/10.2527/jas.2015-9717>
- ICAR (2017): International Committee for Animal Recording. Section 2 - Guidelines for Dairy Cattle Milk Recording.
- KATONGOLE, C. B., T. YAN (2022): The Effects of Dietary Crude Protein Level on Ammonia Emissions from Slurry from Lactating Holstein-Friesian Cows as Measured in Open-Circuit Respiration Chambers. *Animals (Basel)* 12, 1243.
<https://doi.org/10.3390/ani12101243>
- KLEIN, M. S., N. BUTTCHEREIT, S. P. MIEMCZYK, A. K. IMMERSVOLL, C. LOUIS, S. WIEDEMANN, W. JUNGE, G. THALLER, P. J. OEFNER, W. GRONWALD (2011): NMR metabolomic analysis of dairy cows reveals milk glycerophosphocholine to phosphocholine ratio as prognostic biomarker for risk of ketosis. *J. Proteome Res.* 11, 1373-1381.
<https://doi.org/10.1021/pr201017n>
- KROBER, T. F., D. R. KULLING, H. MENZI, F. SUTTER, M. KREUZER (2000): Quantitative effects of feed protein reduction and methionine on nitrogen use by cows and nitrogen emission from slurry. *J. Dairy Sci.* 83, 2941-2951.
[https://doi.org/10.3168/jds.S0022-0302\(00\)75194-0](https://doi.org/10.3168/jds.S0022-0302(00)75194-0)
- MAVROMMATIS, A., C. MITSIOPOULOU, C. CHRISTODOULOU, P. KARIAMPA, M. SIMONI, F. RIGHI, E. TSIPLAKOU (2021): Effects of Supplementing Rumen-Protected Methionine and Lysine on Milk Performance and Oxidative Status of Dairy Ewes. *Antioxidants* 10, 654.
<https://doi.org/10.3390/antiox10050654>
- MÜLLER, C. B. M., S. GÖRS, M. DERNO, A. TUCHSCHERER, K. WIMMERS, A. ZEYNER, B. KUHLA (2021): Differences between Holstein dairy cows in renal clearance rate of urea affect milk urea concentration and the relationship between milk urea and urinary nitrogen excretion. *Sci. Total Environ.* 755, 143198.
<https://doi.org/10.1016/j.scitotenv.2020.143198>
- PARK, J. K., J. M. YEO, G. S. BAE, E. J. KIM, C. H. KIM (2020): Effects of supplementing limiting amino acids on milk production in dairy cows consuming a corn grain and soybean meal-based diet. *J. Anim. Sci. Technol.* 62, 485.
<https://doi.org/10.5187/jast.2020.62.485>
- POWELL, J. M., M. A. WATTIAUX, G. A. BRODERICK (2011): Short communication: Evaluation of milk urea nitrogen as a management tool to reduce ammonia emissions from dairy farms. *J. Dairy Sci.* 94, 4690-4694.
<https://doi.org/10.3168/jds.2011-4476>
- POWELL, J. M., G. A. BRODERICK, T. H. MISSELBROOK (2008): Seasonal diet affects ammonia emissions from tie-stall dairy barns. *J. Dairy Sci.* 91, 857-869.
<https://doi.org/10.3168/jds.2007-0588>
- PUTRI, E. M., M. ZAIN, L. WARLY, H. HERMON (2021): Effects of rumen-degradable-to-undegradable protein ratio in ruminant diet on in vitro digestibility, rumen fermentation, and microbial protein synthesis. *Vet. World* 14, 640-648.
<https://doi.org/10.14202/vetworld.2021.640-648>
- QU, Q., J. C. J. GROOT, K. ZHANG, R. P. O. SCHULTE (2021): Effects of housing system, measurement methods and environmental factors on estimating ammonia and methane emission rates in dairy barns: a meta-analysis. *Biosyst. Eng.* 205, 64-75.
<https://doi.org/10.1016/j.biosystemseng.2021.02.012>
- ROBINSON, P., W. CHALUPA, C. SNIFFEN, H. SATO, T. FUJIEDA, T. UEDA, H. SUZUKI (2004): Influence of ingredient reformulation to reduce diet crude protein level on productivity, and efficiency of dietary nitrogen use, in early lactation dairy cows. *Anim. Feed. Sci. Technol.* 116, 67-81.
<https://doi.org/10.1016/j.anifeedsci.2004.04.004>
- RODRIGUES, A. R. F., M. R. G. MAIA, C. MIRANDA, A. R. J. CABRITA, A. J. M. FONSECA, J. L. S. PEREIRA, H. TRINDADE (2022): Ammonia and greenhouse emissions from cow's excreta are affected by feeding system, stage of lactation and sampling time. *J. Environ. Manage.* 320, 115882.
<https://doi.org/10.1016/j.jenvman.2022.115882>
- SAHA, C. K., C. AMMON, W. BERG, M. FIEDLER, C. LOEBSIN, P. SANFTLEBEN, R. BRUNSCH, T. AMON (2014): Seasonal and diel variations of ammonia and methane emissions from a naturally ventilated dairy building and the associated factors influencing emissions. *Sci. Total Environ.* 468-469, 53-62.
<https://doi.org/10.1016/j.scitotenv.2013.08.015>
- SAJEEV, E. P. M., B. AMON, C. AMMON, W. ZOLLITSCH, W. WINIWARTER (2018): Evaluating the potential of dietary crude protein manipulation in reducing ammonia emissions from cattle and pig manure: a meta-analysis. *Nutr. Cycling Agroecosyst.* 110, 161-175.
<https://doi.org/10.1007/s10705-017-9893-3>
- SAJID, Q. U. A., M. WILK, M. U. ASGHAR (2024): Analysis of crude protein utilisation in ruminant rations: supplementation of limiting amino acids and their effect on the environment – an updated review. *J. Anim. Feed Sci.* 33, 3-12.
<https://doi.org/10.22358/jafs/166576/2023>

- SANCHIS, E., S. CALVET, A. DEL PRADO, F. ESTELLÉS (2019): A meta-analysis of environmental factor effects on ammonia emissions from dairy cattle houses. *Biosyst. Eng.* 178, 176-183.
<https://doi.org/10.1016/j.biosystemseng.2018.11.017>
- SAS (2019): SAS User's Guide, Version 9.4. SAS Institute Inc. Cary, NC.
- SCHRADE, S., K. ZEYER, J. MOHN, M. ZÄHNER (2023): Effect of diets with different crude protein levels on ammonia and greenhouse gas emissions from a naturally ventilated dairy housing. *Sci. Total Environ.* 896, 165027.
<https://doi.org/10.1016/j.scitotenv.2023.165027>
- SCHRADE, S., K. ZEYER, L. GYGAX, L. EMMENEGGER, E. HARTUNG, M. KECK (2012): Ammonia emissions and emission factors of naturally ventilated dairy housing with solid floors and an outdoor exercise area in Switzerland. *Atmos. Environ.* 47, 183-194.
<https://doi.org/10.1016/j.atmosenv.2011.11.015>
- SMITS, M. C. J., H. VALK, A. ELZING, A. KEEN (1995): Effect of protein nutrition on ammonia emission from a cubicle house for dairy cattle. *Livest. Prod. Sci.* 44, 147-156.
[https://doi.org/10.1016/0301-6226\(95\)00068-6](https://doi.org/10.1016/0301-6226(95)00068-6)
- SOUZA, V. C., M. AGUILAR, M. van AMBURGH, W. A. D. NAYANANJALIE, M. D. HANIGAN (2021): Milk urea nitrogen variation explained by differences in urea transport into the gastrointestinal tract in lactating dairy cows. *J. Dairy Sci.* 104, 6715-6726.
<https://doi.org/10.3168/jds.2020-19787>
- SPEK, J. W., J. DIJKSTRA, G. VAN DUINKERKEN, A. BANNINK (2013): A review of factors influencing milk urea concentration and its relationship with urinary urea excretion in lactating dairy cattle. *J. Agric. Sci.* 151, 407-423.
<https://doi.org/10.1017/S0021859612000561>
- SPIEHS, M. J., V. H. VAREL (2009): Nutrient excretion and odorant production in manure from cattle fed corn wet distillers grains with solubles. *J. Anim. Sci.* 87, 2977-2984.
<https://doi.org/10.2527/jas.2008-1584>
- SPIEKERS, H., A. OBERMAIER (2012): Milchharnstoffgehalt und N-Ausscheidung (Milk urea content and N excretion). Institut für Tierernährung und Futterwirtschaft, Prof.-Dürrwaechter-Platz 3, 85586 PoingGrub. Available at: <http://landwirtschaft.bodenseekonferenz.org/bausteine.net/file/showfile.aspx?downaid=9154&domid=1043&fd=2>, 29 February 2012 (in German)
- TABASE, R. K., G. NÆSS, Y. LARRING (2023): Ammonia and methane emissions from small herd cattle buildings in a cold climate. *Sci. Total Environ.* 903, 166046.
<https://doi.org/10.1016/j.scitotenv.2023.166046>
- TODD, R. W., N. A. COLE, R. N. CLARK (2006): Reducing crude protein in beef cattle diet reduces ammonia emissions from artificial feedyard surfaces. *J. Environ. Qual.* 35, 404-411.
<https://doi.org/10.2134/jeq2005.0045>
- VAN DUINKERKEN, G., M. C. J. SMITS, G. ANDRÉ, L. B. J. ŠEBEK, J. DIJKSTRA (2011): Milk urea concentration as an indicator of ammonia emission from dairy cow barn under restricted grazing. *J. Dairy Sci.* 94, 321-335.
<https://doi.org/10.3168/jds.2009-2263>
- VAN DUINKERKEN, G., G. ANDRÉ, M. C. J. SMITS, G. J. MONTENY, L. B. J. ŠEBEK (2005): Effect of rumen-degradable protein balance and forage type on bulk milk urea concentration and emission of ammonia from dairy cow houses. *J. Dairy Sci.* 88, 1099-1112.
[https://doi.org/10.3168/jds.S0022-0302\(05\)72777-6](https://doi.org/10.3168/jds.S0022-0302(05)72777-6)
- YAN, T., J. P. FROST, R. E. AGNEW, R. C. BINNIE, C. S. MAYNE (2006): Relationships among manure nitrogen output and dietary and animal factors in lactating dairy cows. *J. Dairy Sci.* 89, 3981-3991.
[https://doi.org/10.3168/jds.S0022-0302\(06\)72441-9](https://doi.org/10.3168/jds.S0022-0302(06)72441-9)
- YANG, F., Y. HAN, H. BI, X. WEI, W. LUO, G. LI (2022): Ammonia emissions and their key influencing factors from naturally ventilated dairy farms. *Chemosphere.* 307, 135747.
<https://doi.org/10.1016/j.chemosphere.2022.135747>

Received: 15 May 2024

Accepted: 5 July 2024

Online publication: 31 December 2024

GAVRAN, M., D. ŠINKA, M. SAMARDŽIJA, R. GANTNER, Z. STEINER, V. GANTNER: Razumijevanje emisija amonijaka: utjecaj kemijskog sastava stočne hrane na farmama mliječnih holštajnskih goveda. Vet. arhiv 95, 187-196, 2025.

SAŽETAK

Cilj je rada bio istražiti odnos između kemijskog sastava stočne hrane i emisije amonijaka. Kako bismo to postigli, proveli smo temeljitu kemijsku analizu hrane, ispitujući čimbenike kao što su probavljivi sirovi proteini, nerazgradivi proteini u buragu, bakterijska proizvodnja metaboličkih proteina te razine metionina i lizina. Istraživanje se usredotočilo na to kako su navedeni čimbenici utjecali i na prinos mlijeka i na emisije amonijaka. Otkriveno je da postoji pozitivna korelacija između probavljivih sirovih bjelančevina i prinosa mlijeka, dok je buražno nerazgradivi protein bio u jakoj pozitivnoj korelaciji s prinosom mlijeka. Osim toga, otkrivena je negativna korelacija između bakterijske proizvodnje metaboličkih proteina i dnevne količine mlijeka. Koncentracija metionina bila je u slaboj pozitivnoj korelaciji s dnevnom količinom mlijeka, dok je lizin bio u slaboj negativnoj korelaciji s dnevnom količinom mlijeka. Koncentracija lizina u obrocima bila je u pozitivnoj korelaciji s emisijom amonijaka, dok je metionin bio u negativnoj korelaciji s njim. Istraživanje je pokazalo i da buražno nerazgradivi protein ima znakovitu negativnu korelaciju s emisijom amonijaka. Općenito, u istraživanju je naglašena važnost učinkovitih praksi upravljanja prehranom u optimizaciji proizvodnosti životinja i smanjenju emisija amonijaka za promicanje održive poljoprivrede.

Ključne riječi: emisija amonijaka; kemijski sastav obroka; kontrola mliječnosti; mliječne krave
