

The association of bone mineral density Z-score with the early postoperative remission and characteristics of bone mineral loss in patients with Cushing's disease: a retrospective study

Emre Gezer¹, Zeynep Cantürk², Alev Selek², Berrin Çetinarslan², Mehmet Sözen², Özlem Elen³, İhsan Anık⁴, Savaş Ceylan⁴

¹Division of Endocrinology and Metabolism, Darıca Farabi Training and Research Hospital, Kocaeli, Turkey

²Department of Endocrinology, Kocaeli University, Kocaeli, Turkey

³Department of Internal Medicine, Kocaeli University, Kocaeli, Turkey

⁴Department of Neurosurgery, Kocaeli University, Kocaeli, Turkey

Aim To investigate the association of bone mineral density (BMD) Z-scores with early-postoperative remission rate and clinical parameters in patients with Cushing's disease (CD).

Methods We retrospectively evaluated the records of patients diagnosed with CD. After the exclusion of 230 patients, 87 CD patients were finally enrolled. BMD was determined by dual-energy x-ray absorptiometry (DXA) at the lumbar spine 1-4 (L₁₋₄) and left femur. Early-postoperative remission was defined as a morning cortisol concentration on the first day after surgery of less than 5 µg/dL. The diagnosis of BMD "below the expected range for age" was defined as a Z-score ≤ -2.00 standard deviations.

Results DXA results were not significantly associated with early postoperative remission. They also did not significantly differ between eugonadal and menopausal groups. Preoperative morning cortisol significantly negatively but weakly correlated with Z-score of the total femur, while preoperative adrenocorticotrophic hormone/cortisol ratio positively but weakly correlated with DXA results of L1-4.

Conclusion The severity of bone loss was not significantly related to the failure of transsphenoidal surgery for Cushing's disease.

Received: January 8, 2022

Accepted: December 12, 2022

Correspondence to:

Emre Gezer
Kocaeli Darıca Farabi Eğitim Ve
Araştırma Hastanesi
Fevziçakmak Mahallesi
Dr. Zeki Acar Cd. No:62
41700 Darıca/Kocaeli, Turkey
gezer_emre@hotmail.com

Cushing's disease (CD) is characterized by an overexpression of adrenocorticotrophic hormone (ACTH) as a result of pituitary corticotroph adenoma. It leads to endogenous hypercortisolism, one of the most important causes of bone mineral loss (1). The prevalence of osteoporosis due to CD ranges from 34% to 50% (2-4). Glucocorticoid (GC) excess impairs bone metabolism through various direct and indirect mechanisms: increased bone resorption, reduced bone formation, impaired renal and intestinal calcium reabsorption, inhibition of gonadal steroids and growth hormone, and reduced muscle strength (5,6). The severity of CD increases the loss in bone mineral density.

Bone mineral density (BMD) is assessed with dual-energy x-ray absorptiometry (DXA) at the spine (anteroposterior and lateral) and hip (total femur and femoral neck). To detect the secondary causes of osteoporosis, such as CD, individual's BMD is compared with that of an average person of the same sex and age. In this way, we obtain the number of standard deviations (SD) from age/sex-matched groups, ie, BMD Z-score (7). This study used Z-score as the main parameter to eliminate the effects of age, sex, and gonadal status on BMD.

The early outcome of pituitary surgery in patients with CD is affected by different factors: the surgeon's experience, tumor extent and size, an accurate identification of the tumor during the operation or preoperatively on the MRI scan, preoperative ACTH levels, and the clinical severity of the disease (8-10). In addition, in patients with CD, ACTH and the severity of hypercortisolism influence the degree of bone mineral loss (3,11-13). Since the severity of hypercortisolism is related to both the early outcome of pituitary surgery and the degree of BMD loss, we postulated that BMD might be inversely related to early postoperative remission. Therefore, we investigated whether BMD Z-scores were associated with the early-postoperative remission rate and clinical parameters in patients with CD.

PATIENTS AND METHODS

We retrospectively analyzed the electronic records (Nucleus and HUYE) of patients diagnosed with CD and operated on at the Pituitary Research Center of our institute between 2006 and 2020. CD was diagnosed with the following screening tests: midnight salivary cortisol, 24-hour urine free cortisol, a 2-mg low-dose dexamethasone suppression test, late-night serum cortisol, 8-mg high-dose dexamethasone suppression test, and corticotropin-releasing hormone (CRH) test or inferior petrosal sinus sampling,

when needed. Overall, 317 patients were available for the analysis. Exclusion criteria were a lack of preoperative DXA results, preoperative MRI scan images, insulin-like growth factor 1 (IGF-1), cortisol, or ACTH levels; impaired IGF-1 levels (below or above the age-adjusted reference range); age less than 18 years; severe renal or hepatic failure; untreated/uncontrolled hypogonadism or hypo/hyperthyroidism; history of a disease or drug use that may cause secondary osteoporosis; lower spinal and hip fractures; and history of instrumentation use for stabilization of the lumbar vertebrae and hip. After the exclusion of 230 patients, there remained 87 eligible CD patients.

BMD was determined by DXA at the lumbar spine 1-4 (L_{1-4}) and left femur (both femoral neck [F_N] and total femur [F_T]) using a GE Lunar Prodigy® 8743 densitometer (GE Healthcare, Machelen, Belgium). BMD values were expressed in grams per square centimeter. T-score and Z-score were expressed as standard deviation values, calculated through comparison with controls who were at peak BMD and persons matched for age and sex, respectively. BMD "below the expected range for age" was defined as a Z-score ≤ -2.00 SD (14).

Early-postoperative remission was defined as a morning cortisol concentration on the first day after surgery of less than 5 $\mu\text{g/dL}$ (15-18). Pituitary adenoma size was evaluated with a MRI scan. The maximum tumor diameter and the presence of cavernous sinus invasion were noted. We also recorded the results of DXA scan, postoperative morning serum cortisol levels, patient demographics, preoperative morning cortisol and ACTH concentrations, gonadal status, and if the operation was primary or a re-operation. ACTH/cortisol ratio was calculated. The study was approved by the ethics committee of Kocaeli University, Faculty of Medicine.

Statistical analysis

The normality of distribution was tested with the Kolmogorov-Smirnov and Shapiro-Wilk test. Continuous variables are expressed as mean \pm standard deviation (SD) or median (25th-75th percentile). Categorical variables are summarized as counts (percentages). The groups were compared with independent samples *t* test or Mann-Whitney U test and one-way analysis of variance (ANOVA), as appropriate. The Tukey test was used for multiple comparisons. Correlations between continuous variables were determined with the Pearson and Spearman correlation analysis, and associations between categorical variables with the χ^2 test. A

two-sided P value <0.05 was considered statistically significant. Statistical analyses were performed with SPSS for Windows, Version 20.0. (IBM Corp, Armonk, NY, USA)

RESULTS

Demographics and clinical characteristics are summarized in Table 1. The prevalence of BMD “below the expected range for age” was 36.8% ($n=32$) and that of normal BMD was 63.2% ($n=55$) (Table 1).

TABLE 1. Demographic and clinical characteristics ($n=87$)

| Characteristic | n (%) |
|----------------------------------|------------|
| Age (years), median (range) | 40 (18-73) |
| Sex | |
| male | 14 (16.1) |
| female | 73 (83.9) |
| Remission | |
| yes | 48 (55.2) |
| no | 39 (44.8) |
| Operation | |
| primary | 77 (88.5) |
| secondary | 10 (11.5) |
| Gonadal Status | |
| eugonadal | 68 (78.2) |
| menopausal | 19 (21.8) |
| Adenoma | |
| microadenoma | 49 (56.3) |
| macroadenoma | 38 (43.7) |
| Bone mineral density* | |
| below the expected range for age | 32 (36.8) |
| normal | 55 (63.2) |

*Below the expected range for age and normal groups defined as Z-score ≤ -2 and > -2 , respectively.

DXA results were not significantly associated with the early-postoperative remission (Table 2). They also did not significantly differ between eugonadal and menopausal groups ($P > 0.05$).

There was a weak significant negative correlation between preoperative morning cortisol level and the Z-score of F_T ($r = -0.276$, $P = 0.010$, Table 3), while there was a weak positive correlation between preoperative morning ACTH/cortisol ratio and DXA results (BMD, T-scores, and Z-scores of L_{1-4} : $r = 0.280$, $P = 0.009$; $r = 0.258$, $P = 0.016$; $r = 0.257$, $P = 0.016$, respectively). ACTH and adenoma size on MRI did not correlate with DXA results. Age did not correlate with preoperative morning cortisol, ACTH, or ACTH/cortisol ($P > 0.05$).

Z-scores of L_{1-4} were significantly lower than those of F_N and F_T (-1.33 ± 1.32 vs -0.49 ± 1.78 , $P < 0.001$ and -1.33 ± 1.32 vs -0.55 ± 1.18 , one-way ANOVA test with Tukey's test $P < 0.001$, respectively, Table 4). Z-scores of F_N and F_T did not significantly differ (-0.49 ± 1.78 vs -0.55 ± 1.18 , $P = 0.934$). Similarly, T-scores of L_{1-4} were significantly lower than T-scores of F_T (-1.03 ± 1.37 vs -0.52 ± 1.18 , $P = 0.021$); however, there was no significant difference between T-scores of L_{1-4} and F_N and those of F_N and F_T ($P = 0.158$ and $P = 0.672$, respectively).

DISCUSSION

In our study, BMD Z-score, even though a parameter known to be affected by the severity of GC excess, was not related to the early postoperative remission rate in patients with CD. This is the first study to evaluate the indirect pre-

TABLE 2. The association of dual-energy x-ray absorptiometry results with remission and gonadal status*

| | | Remission | | | Gonadal status | | |
|-----------|---------|--------------------------------|--------------------------------|----------------|------------------------|-------------------------|----------------|
| | | yes [†] | no [†] | p ^c | eugonadal [†] | menopausal [†] | p [§] |
| L_{1-4} | BMD | 1.05 ± 0.17 | 1.07 ± 0.15 | 0.516 | 1.07 ± 0.16 | 1.01 ± 0.14 | 0.202 |
| | T-score | -1.11 ± 1.45 | -0.92 ± 1.27 | 0.512 | -0.95 ± 1.40 | -1.31 ± 1.24 | 0.306 |
| | Z-score | -1.48 ± 1.39 | -1.14 ± 1.21 | 0.232 | -1.39 ± 1.40 | -1.12 ± 0.96 | 0.338 |
| F_N | BMD | 0.96 ± 0.18 | 0.96 ± 0.17 | 0.998 | 0.97 ± 0.18 | 0.90 ± 0.11 | 0.081 |
| | T-score | -0.68 ± 1.20 | -0.69 ± 1.12 | 0.970 | -0.58 ± 1.23 | -1.03 ± 0.79 | 0.135 |
| | Z-score | -0.55 ± 1.23 | -0.42 ± 1.12 | 0.604 | -0.51 ± 1.27 | -0.40 ± 0.78 | 0.713 |
| F_T | BMD | 0.97 ± 0.17 | 0.95 ± 0.14 | 0.560 | 0.97 ± 0.17 | 0.94 ± 0.12 | 0.511 |
| | T-score | $-0.45(-1.38-0.78)^{\ddagger}$ | $-0.30(-1.40-0.10)^{\ddagger}$ | 0.494* | -0.51 ± 1.25 | -0.55 ± 0.93 | 0.891 |
| | Z-score | -0.52 ± 1.31 | -0.60 ± 1.01 | 0.745 | -0.63 ± 1.23 | -0.28 ± 0.95 | 0.252 |

*Abbreviations: BMD – bone mineral density; F_N – femoral neck; F_T – total femur; L_{1-4} – lumbar vertebrae 1-4.

[†]Data are expressed as mean \pm standard deviation.

[‡]Data are expressed as median (25th-75th percentile).

§Independent-samples t test.

*Mann-Whitney U test.

dictivity of Z-score for the early-postoperative remission in CD patients. Chronic hypercortisolism associated with CD negatively affects bone turnover, which causes osteoporosis, osteopenia, and BMD "below the expected range for age." In our patients, osteoporosis prevalence was 16.1%, lower than the rate reported elsewhere (34% to 50%) (2,3). However, these studies generally had smaller sample sizes. Ohmori et al (2) reported a much higher prevalence of amenorrhea than we did (84.2% vs 21.8%). In the European Registry on Cushing's syndrome, with the largest sample size to date, of 136 CD patients with DXA records (4), 22% and 12% were diagnosed with spine and hip osteoporosis, respectively.

A study by McKiernan et al (7), which included 18674 patients, reported a significant inverse relationship between the presence of secondary causes of osteoporosis and Z-score. Similarly, Lewiecki et al (19) indicated that Z-score could be used for the diagnosis of osteoporosis caused by a secondary cause such as CD. Therefore, we decided to use Z-score as our primary variable, regardless of age, sex, and gonadal status. The prevalence of patients with Z-score below -2 SD was 36.8%, which was more than double the prevalence of osteoporosis. In a study by Sonino et al (10), patients with a severe clinical picture had significantly worse remission rates compared with patients without severe clinical picture. Additionally, in various studies DXA

results negatively correlated with biochemical parameters indicating the severity of hypercortisolism (3,12,13,20). A hypothetical indirect relationship between DXA results and the early-postoperative remission derived from these findings was not supported by our study.

In accordance with multiple studies (12,21), we found no significant difference between DXA results and the gonadal status of CD patients. This finding shows endogenous hypercortisolism to be a more important determinant of bone metabolism than the gonadal status.

Our study showed that preoperative morning serum cortisol concentration significantly inversely, but weakly, correlated with the Z-score of F_T . Similarly, another study demonstrated a significant negative association between serum cortisol and lumbar BMD (22). A recent study, which enrolled patients under long-term GC treatment, reported that cumulative GC dose negatively correlated with the Z-score of the total hip (23). Moreover, another study reported an inverse but non-significant correlation between presurgical morning cortisol level and BMD at both the lumbar spine and hip (12).

We also found a significant positive weak correlation between preoperative morning ACTH/cortisol ratio and DXA results of L_{1-4} . Interestingly, the effect of ACTH alone on

TABLE 3. Correlations between dual-energy x-ray absorptiometry results and preoperative morning cortisol, adrenocorticotrophic hormone (ACTH), ACTH/cortisol ratio, and adenoma size*

| | | L_{1-4} | | | F_N | | | F_T | | |
|---------------------|----------------|-----------|---------|---------|--------|---------|---------|--------|---------|---------|
| | | BMD | T-score | Z-score | BMD | T-score | Z-score | BMD | T-score | Z-score |
| Cortisol | r | -0.115 | -0.122 | -0.177 | -0.119 | -0.104 | -0.159 | -0.219 | -0.237 | -0.276 |
| | p [†] | 0.289 | 0.261 | 0.100 | 0.272 | 0.336 | 0.827 | 0.041 | 0.027 | 0.010 |
| ACTH | r | 0.181 | 0.163 | 0.123 | 0.116 | 0.078 | 0.024 | 0.105 | 0.039 | -0.21 |
| | p [†] | 0.094 | 0.130 | 0.255 | 0.285 | 0.473 | 0.827 | 0.335 | 0.720 | 0.845 |
| ACTH/cortisol | r | 0.280 | 0.258 | 0.257 | 0.199 | 0.156 | 0.126 | 0.249 | 0.206 | 0.160 |
| | p [†] | 0.009 | 0.016 | 0.016 | 0.065 | 0.150 | 0.244 | 0.020 | 0.055 | 0.139 |
| Size of the adenoma | r | 0.144 | 0.120 | 0.099 | 0.086 | 0.026 | -0.001 | 0.109 | 0.060 | 0.002 |
| | p [†] | 0.182 | 0.270 | 0.361 | 0.430 | 0.808 | 0.993 | 0.316 | 0.584 | 0.983 |

*Abbreviations: BMD – bone mineral density; r – correlation coefficient; F_N – femoral neck; F_T – total femur; L_{1-4} – lumbar vertebrae 1-4.

†Spearman's Rho correlation test.

TABLE 4. Multiple comparisons between dual-energy x-ray absorptiometry results of each anatomic location*

| | L_{1-4} mean±SD | F_N mean±SD | F_T mean±SD | F | P [†] |
|---------|----------------------|------------------|------------------|--------|----------------|
| Z-score | -1.33 ± 1.32 | -0.49 ± 1.78 | -0.55 ± 1.18 | 12.577 | <0.001 |
| T-score | -1.03 ± 1.37 | -0.68 ± 1.16 | -0.52 ± 1.18 | 3.788 | 0.024 |
| BMD | 1.06 ± 0.16 | 0.96 ± 0.17 | 0.96 ± 0.16 | 12.324 | <0.001 |

*Abbreviations: BMD – Bone mineral density; F_N – femoral neck; F_T – total femur; L_{1-4} – lumbar vertebrae 1-4, SD – standard deviation.

†one-way ANOVA test.

DXA results was non-significant, though the level of significance was higher for L_{1-4} than for F_N and F_T . ACTH stimulates osteoblastic cell proliferation and increases collagen synthesis in osteoblasts (11). Similarly, it significantly positively correlates with the lumbar BMD (22). We found that excess GC had a deteriorating effect mainly on the total hip, while ACTH exerted a protective effect on the lumbar spine.

As one of the richest sites in the trabecular bone, the lumbar spine was most vulnerable to the effect of GC excess. Its high sensitivity to hypercortisolism was explained by higher bone turnover rate due to a greater volume/surface ratio. Kawamata et al (24) showed preoperative lumbar spine T-scores to be significantly lower than those of the femoral neck. The lumbar spine also sustained more severe bone loss than the hip (12). In our study, Z-scores and T-scores of L_{1-4} were significantly lower than those of F_N and F_T .

The main limitation of our study was the fact that we did not assess body mass index and 25-hydroxyvitamin D levels, which were shown to be closely associated with bone structure and the osteoporosis development. Moreover, many patients were excluded from the study due to the absence of preoperative DXA scan.

In conclusion, our study found no significant relationship between the preoperative BMD Z-score and early-postoperative remission. To the best of our knowledge, this is the first study that investigated if the severity of bone loss predicted the failure of transsphenoidal surgery in CD. In addition, excess cortisol was negatively correlated with Z-score of F_T and ACTH/cortisol ratio was positively but weakly correlated with DXA results at the lumbar spine, which was the most adversely influenced bone site in CD. Bone properties were more affected by GC excess than by gonadal status.

Funding None.

Ethical approval granted by the Ethics Committee of Kocaeli University, Faculty of Medicine (KÜ GÖKAEK 2019/15.13 - 2019/256).

Declaration of authorship EG conceived and designed the study; EG, MS, OE, IA, SC acquired the data; ZC, AS, BC, IA, SC analyzed and interpreted the data; EG, MS, OE, IA drafted the manuscript; ZC, AS, BC, SC critically revised the manuscript for important intellectual content; all authors gave approval of the version to be submitted; all authors agree to be accountable for all aspects of the work.

Competing interests All authors have completed the Unified Competing Interest form at www.icmje.org/coi_disclosure.pdf (available on request from the corresponding author) and declare: no support from any organization for the submitted work; no financial relationships with any organizations that might have an interest in the submitted work in the previous 3 years; no other relationships or activities that could appear to have influenced the submitted work.

References

- 1 Cushing H. The basophil adenomas of the pituitary body and their clinical manifestations (pituitary basophilism). *Bull Johns Hopkins Hosp.* 1932;50:137-95.
- 2 Ohmori N, Nomura K, Ohmori K, Kato Y, Itoh T, Takano K. Osteoporosis is more prevalent in adrenal than in pituitary Cushing's syndrome. *Endocr J.* 2003;50:1-7. [Medline:12733704](#) [doi:10.1507/endocrj.50.1](#)
- 3 Trementino L, Appolloni G, Ceccoli L, Marcelli G, Concettoni C, Boscaro M, et al. Bone complications in patients with Cushing's syndrome: Looking for clinical, biochemical, and genetic determinants. *Osteoporos Int.* 2014;25:913-21. [Medline:24126765](#) [doi:10.1007/s00198-013-2520-5](#)
- 4 Valassi E, Santos A, Yaneva M, Tóth M, Strasburger CJ, Chanson P, et al. The European Registry on Cushing's syndrome: 2-year experience. Baseline demographic and clinical characteristics. *Eur J Endocrinol.* 2011;165:383-92. [Medline:21715416](#) [doi:10.1530/EJE-11-0272](#)
- 5 Weinstein RS, Jilka RL, Michael Parfitt A, Manolagas SC. Inhibition of osteoblastogenesis and promotion of apoptosis of osteoblasts and osteocytes by glucocorticoids: potential mechanisms of their deleterious effects on bone. *J Clin Invest.* 1998;102:274-82. [Medline:9664068](#) [doi:10.1172/JCI2799](#)
- 6 Gennari C. Glucocorticoid induced osteoporosis. *Clin Endocrinol (Oxf).* 1994;41:273-4. [Medline:7955431](#) [doi:10.1111/j.1365-2265.1994.tb02544.x](#)
- 7 McKiernan FE, Berg RL, Linneman JG. The utility of BMD Z-score diagnostic thresholds for secondary causes of osteoporosis. *Osteoporos Int.* 2011;22:1069-77. [Medline:20533026](#) [doi:10.1007/s00198-010-1307-1](#)
- 8 Tritos NA, Biller BMK, Swearingen B. Management of Cushing disease. *Nat Rev Endocrinol.* 2011;7:279-89. [Medline:21301487](#) [doi:10.1038/nrendo.2011.12](#)
- 9 Cannavò S, Almoto B, Dall'Asta C, Corsello S, Lovicu RM, De Menis E, et al. Long-term results of treatment in patients with ACTH-secreting pituitary macroadenomas. *Eur J Endocrinol.* 2003;149:195-200. [Medline:12943521](#) [doi:10.1530/eje.0.1490195](#)
- 10 Sonino N, Zielesny M, Fava GA, Fallo F, Boscaro M. Risk factors and long-term outcome in pituitary-dependent Cushing's disease. *J Clin Endocrinol Metab.* 1996;81:2647-52. [Medline:8675592](#)
- 11 Isales CM, Zaidi M, Blair HC. ACTH is a novel regulator of bone mass. *Ann NY Acad Sci.* 2010;1192:110-6. [Medline:20392225](#) [doi:10.1111/j.1749-6632.2009.05231.x](#)
- 12 Rahaman SH, Jyotsna VP, Kandasamy D, Shreenivas V, Gupta N, Tandon N. Bone health in patients with Cushing's syndrome. *Indian J Endocrinol Metab.* 2018;22:766-9. [Medline:30766815](#) [doi:10.4103/ijem.IJEM_160_18](#)
- 13 Braun LT, Fazel J, Zopp S, Benedix S, Osswald-Kopp A, Riester A, et al. The Effect of Biochemical remission on bone metabolism in Cushing's syndrome: a 2-year follow-up study. *J Bone Miner Res.*

2020. [Medline:32315096](#) doi:[10.1002/jbmr.4033](#)
- 14 Camacho PM, Petak SM, Binkley N, Diab DL, Eldeiry LS, Farooki A, et al. American Association of Clinical Endocrinologists/American College of Endocrinology Clinical Practice Guidelines for the Diagnosis and Treatment of Postmenopausal Osteoporosis-2020 Update. *Endocr Pract.* 2020;26 Supplement 1:1-46. [Medline:32427503](#) doi:[10.4158/GL-2020-0524SUPPL](#)
 - 15 Ironside N, Chatain G, Asuzu D, Benzo S, Lodish M, Sharma S, et al. Earlier post-operative hypocortisolemia may predict durable remission from Cushing's disease. *Eur J Endocrinol.* 2018;178:255-63. [Medline:29330227](#) doi:[10.1530/EJE-17-0873](#)
 - 16 Pendharkar AV, Sussman ES, Ho AL, Hayden Gephart MG, Katznelson L. Cushing's disease: Predicting long-term remission after surgical treatment. *Neurosurg Focus.* 2015;38. [Medline:25639315](#) doi:[10.3171/2014.10.FOCUS14682](#)
 - 17 Esposito F, Dusick JR, Cohan P, Moftakhar P, McArthur D, Wang C, et al. Clinical review: Early morning cortisol levels as a predictor of remission after transsphenoidal surgery for Cushing's disease. *J Clin Endocrinol Metab.* 2006;91:7-13. [Medline:16234305](#) doi:[10.1210/jc.2005-1204](#)
 - 18 Bansal P, Lila A, Goroshi M, Jadhav S, Lomte N, Thakkar K, et al. Duration of post-operative hypocortisolism predicts sustained remission after pituitary surgery for Cushing's disease. *Endocr Connect.* 2017;6:625-36. [Medline:28912338](#) doi:[10.1530/EC-17-0175](#)
 - 19 Lewiecki EM, Gordon CM, Baim S, Leonard MB, Bishop NJ, Bianchi ML, et al. International Society for Clinical Densitometry 2007 Adult and Pediatric Official Positions. *Bone.* 2008;43:1115-21. [Medline:18793764](#) doi:[10.1016/j.bone.2008.08.106](#)
 - 20 Tauchmanová L, Pivonello R, Di Somma C, Rossi R, De Martino MC, Camera L, et al. Bone demineralization and vertebral fractures in endogenous cortisol excess: Role of disease etiology and gonadal status. *J Clin Endocrinol Metab.* 2006;91:1779-84. [Medline:16522701](#) doi:[10.1210/jc.2005-0582](#)
 - 21 Van der Eerden AW, Den Heijer M, Oyen WJ, Hermus AR. Cushing's syndrome and bone mineral density: Lowest Z scores in young patients. *Neth J Med.* 2007;65:137-41. [Medline:17452762](#)
 - 22 Guo W, Li F, Zhu C, Wang B, Wang K, Dai C, et al. Effect of hypercortisolism on bone mineral density and bone metabolism: A potential protective effect of adrenocorticotrophic hormone in patients with Cushing's disease. *J Int Med Res.* 2018;46:492-503. [Medline:28851260](#) doi:[10.1177/0300060517725660](#)
 - 23 Iervolino LL, Ferraz-de-Souza B, Martin RM, Costa FC, Miranda MC, Mendonça BB, et al. Real-world impact of glucocorticoid replacement therapy on bone mineral density: retrospective experience of a large single-center CAH cohort spanning 24 years. *Osteoporos Int.* 2020;31:905-12. [Medline:31897546](#) doi:[10.1007/s00198-019-05268-0](#)
 - 24 Kawamata A, Iihara M, Okamoto T, Obara T. Bone mineral density before and after surgical cure of Cushing's syndrome due to adrenocortical adenoma: Prospective study. *World J Surg.* 2008;32:890-6. [Medline:18210182](#) doi:[10.1007/s00268-007-9394-7](#)