

PARTITIONS INTO TRIPLES WITH EQUAL PRODUCTS AND FAMILIES OF ELLIPTIC CURVES

AHMED EL AMINE YOUMBAL, ARMAN SHAMSI ZARGAR AND MAKSYM
VOZNYI

University of El Oued, Algeria, University of Mohaghegh Ardabili, Iran,
Toronto District School Board, Canada

ABSTRACT. Let $\mathcal{S}_\ell(M, N)$ denote a set of ℓ (distinct) triples of positive integers having the same sum M and the same product N . For each $2 \leq \ell \leq 4$ we establish a connection between a subset of $\mathcal{S}_\ell(M, N)$ with (integral) parametric elements and a family of elliptic curves. When $\ell = 2$ and 3 , we use certain known subsets of $\mathcal{S}_\ell(M, N)$ with parametric elements and respectively find families of elliptic curves of generic rank ≥ 5 and ≥ 6 , while for $\ell = 4$ we first obtain a subset of $\mathcal{S}_\ell(M, N)$ with parametric elements, then construct a family of elliptic curves of generic rank ≥ 8 . Finally, we perform a computer search within these families to find specific curves with rank ≥ 11 and in particular we found two curves of rank 14.

1. INTRODUCTION

For any positive integers ℓ and n , let $\mathcal{S}_\ell^{(n)}(M, N)$ denote a set of ℓ (distinct) triples $(x_{j1}^n, x_{j2}^n, x_{j3}^n)$ of positive integers having the same sum M and the same product N . For the j -th element of the set, we define

$$M_j^{(n)} = \sum_{i=1}^3 x_{ji}^n, N_j^{(n)} = \prod_{i=1}^3 x_{ji}^n, T_j^{(n)} = \sum_{1 \leq i < k \leq 3} x_{ji}^n x_{jk}^n \quad (1 \leq j \leq \ell).$$

We drop the notation “ (n) ” from the terminology when $n = 1$.

In this context, we are interested in the following partitioning problem, which, from the geometric point of view, is equivalent to examining the existence of ℓ rectangular boxes with integer sides having the same perimeter and the same volume.

2020 *Mathematics Subject Classification.* 14H52, 11D25, 11D09, 05A17.

Key words and phrases. Triples of integers, equal sum of integers, equal products, Diophantine equation, partition, parametric solution, elliptic curve.

PROBLEM 1.1. *For given positive integers $\ell \geq 2$ and n , construct a subset of $\mathcal{S}_\ell^{(n)}(M, N)$ with (integral) parametric elements.*

Note that solving this problem is equivalent to finding a parametric solution to the symmetric system of Diophantine equations

$$(1.1) \quad M = M_j^{(n)}, \quad N = N_j^{(n)},$$

for all $j = 1, \dots, \ell$.

Ever since Motzkin's conjecture, the aforesaid problem has been subjected to investigations by some authors (see [4, 23, 26]). In 1989, Kelly [24] showed the existence of an arbitrarily large number of such triples. In 1996, Schinzel [33] reproved the same result in a different way. Neither proofs was effective for constructing a numerical example until, in 2012, Choudhry [10] gave a constructive method. It is pertinent to note that, in 2013, Zhang and Cai [36] generalized the aforesaid result of Kelly from triples to m -tuples.

To the best knowledge of the authors, Problem 1.1 has been effectively solved only for the (ℓ, n) 's given in the following table:

TABLE 1. Known parametric solutions to (1.1)

(ℓ, n)	References in the chronological order of discovery
(2, 1)	[28, 22, 5, 9, 11]
(2, 2)	[2, 22, 35, 5, 25, 6, 18, 15]
(2, 3)	[22, 7]
(2, 4)	[8, 13]
(3, 1)	[31]
(3, 2)	[18]

In the other direction, making any connection between algebraic or geometric notions and elliptic curves has interested researchers. Specifically, symmetric Diophantine equations of certain forms have led to families of elliptic curves with higher ranks, see, for example, [1, 17, 20]. As far as we are aware, there are only a few works in literature that have studied some connections between elliptic curves of positive rank and triples satisfying (1.1). In 1989, Kelly [24] illustrated such a connection between positive rank elliptic curves of the form $y^2 - Mxy - Ny = x^3$ and triples satisfying (1.1) for $(\ell, n) = (1, 1)$. In 2015, Sadek and El-Sissi [31] extended this result by studying the same family assigned to triples satisfying (1.1) for $(\ell, n) = (2, 1)$ and $(3, 1)$ and showed that the generic ranks of the associated families are ≥ 2 and ≥ 3 , respectively. In 2019, Choudhry [14] constructed several families of elliptic

curves whose generic ranks range from ≥ 8 to ≥ 12 coming from the system $M = M_j$ and $T = T_j$ for some j 's.

In this work, we establish a new connection between each of certain solutions of (1.1) for $2 \leq \ell \leq 4$ with $n = 1$ and a family of elliptic curves, whose torsion subgroups are trivial in general. When $(\ell, n) = (2, 1)$ and $(3, 1)$, we use the solutions given in [11] and [31] and correspondingly introduce a family of generic rank ≥ 5 and ≥ 6 . For the case $(\ell, n) = (4, 1)$, we first obtain a parametric solution of (1.1) and then construct a family of elliptic curves of generic rank ≥ 8 . Finally, we perform a computer search within those families to find specific curves with high rank and in particular we found two curves of rank 14.

2. FAMILIES OF ELLIPTIC CURVES RELATED TO (1.1)

First we recall the following specialisation theorem of Néron [34, Theorem 20.1] that we need for our next results.

THEOREM 2.1. *Let K be a number field, and let E be an elliptic curve defined over the function field $K(\mathbb{P}^n)$. Then there are infinitely many points $t \in \mathbb{P}^n(K)$ such that the specialisation homomorphism*

$$\sigma_t : E(K(\mathbb{P}^n)) \rightarrow E_t(K)$$

is injective. The set of t for which σ_t is noninjective forms a thin set.

2.1. *A family with rank ≥ 5 .* According to [11, Subsection 3.1], Choudhry has fully determined the set $\mathcal{S}_2(M, N)$ by finding two complete solutions to the related equations (1.1). Without loss of generality, we consider his first solution, then

$$\mathcal{A} := \mathcal{A}(M, N) := \mathcal{S}_2(M, N) = \{(a_1, a_2, a_3), (b_1, b_2, b_3)\},$$

where

$$\begin{aligned} a_1 &= p(s + rt), & a_2 &= q(s + pt), & a_3 &= r(s + qt), \\ b_1 &= q(s + rt), & b_2 &= r(s + pt), & b_3 &= p(s + qt). \end{aligned}$$

To this set, we assign an elliptic curve described by the equation

$$y^2 = (x + a_1 a_2)(x + a_2 a_3)(x + a_1 a_3) + e_1^2 x^2,$$

where e_1 is a nonzero rational number which will be defined later.

The above elliptic curve can be rewritten as:

$$(2.1) \quad y^2 = x^3 + (T_1 + e_1^2)x^2 + MNx + N^2,$$

where $T_1 = a_1 a_2 + a_1 a_3 + a_2 a_3$.

By imposing

$$(2.2) \quad T_1 + e_1^2 = T_2 + e_2^2,$$

for some rational e_2 , the curve (2.1) will contain the seven rational points $P_1 = (0, N)$, $P_{ij} = (-a_i a_j, a_i a_j e_1)$ and $Q_{ij} = (-b_i b_j, b_i b_j e_2)$, $1 \leq i < j \leq 3$. Here, $T_2 = b_1 b_2 + b_1 b_3 + b_2 b_3$.

The quadratic (2.2) is easily accomplished by parametrising

$$e_1 = \frac{k^2 + T_2 - T_1}{2k}, \quad e_2 = \frac{k^2 + T_1 - T_2}{2k},$$

for any nonzero rational number k .

By this description, we have thus constructed a family of elliptic curves, defined over $\mathbb{Q}(p, q, r, s, t, k)$, coming from the elements of \mathcal{A} , that has the above rational points. We denote this family by $\mathcal{E}_{\mathcal{A}}$.

REMARK 2.2. Notice that the points P_{ij} are collinear (because their (x, y) -coordinates satisfy the linear equation $y + e_1 x = 0$), showing that at most two of them can be linearly independent. The same result holds for the points Q_{ij} . Therefore, at most five points of the seven points of $\mathcal{E}_{\mathcal{A}}$ can be linearly independent.

In the next theorem we show that five of seven points of $E_{\mathcal{A}}$ are linearly independent for infinitely many tuples (p, q, r, s, t, k) .

THEOREM 2.3. *For the set of all tuples (p, q, r, s, t, k) except for a thin subset, the rank of the family $\mathcal{E}_{\mathcal{A}}$ is at least five with the five linearly independent points $P_1, P_{12}, P_{13}, Q_{12}, Q_{13}$.*

PROOF. By Néron's specialisation theorem (Theorem 2.1), in order to prove that the family $\mathcal{E}_{\mathcal{A}}$ has rank ≥ 5 over $\mathbb{Q}(p, q, r, s, t, k)$, it suffices to find a specialisation $(p, q, r, s, t, k) = (p_0, q_0, r_0, s_0, t_0, k_0)$ such that the above points in the statement are linearly independent on the specialised curve over \mathbb{Q} . We take $(p, q, r, s, t, k) = (1, 4, 2, 4, 1, 1)$ for which we have

$$\mathcal{A}(42, 1920) = \{(6, 20, 16), (24, 10, 8)\}$$

and

$$T_1 = 536, \quad T_2 = 512, \quad e_1 = -\frac{23}{2}, \quad e_2 = \frac{25}{2}.$$

The five points are linearly independent and of infinite order on the specialised rank 5 elliptic curve

$$\mathcal{E}_{\mathcal{A}(42, 1920)} : y^2 = x^3 + \frac{2673}{4}x^2 + 80640x + 3686400.$$

Indeed, the determinant of the Néron–Tate height pairing matrix (a.k.a. regulator) of the specialised five points with x -coordinates

$$x(P_1) = 0, \quad x(P_{12}) = -120, \quad x(P_{13}) = -96, \quad x(Q_{12}) = -240, \quad x(Q_{13}) = -192,$$

is the nonzero value 23.4808049005680 as computed by SageMath [32]. This shows that the family of elliptic curves $\mathcal{E}_{\mathcal{A}}$ has rank ≥ 5 over $\mathbb{Q}(p, q, r, s, t, k)$ with independent points $P_1, P_{12}, P_{13}, Q_{12}, Q_{13}$ (except for a thin subset of (p, q, r, s, t, k) 's). \square

REMARK 2.4. Corresponding to each of the known subsets $\mathcal{S} \neq \mathcal{A}$ of $\mathcal{S}_\ell^{(n)}(M, N)$ with $\ell \geq 2$, being found in the references mentioned in Table 1 and in Subsection 2.3 (i.e., the set \mathcal{C}), one can assign a family of elliptic curves similar to $\mathcal{E}_\mathcal{A}$ and accordingly establish a result as that of Theorem 2.3.

2.2. *A family with rank ≥ 6 .* From [31, Section 4], we consider the subset $\mathcal{B} := \mathcal{B}(M, N)$ of $\mathcal{S}_3(M, N)$ with the elements

$$(a_1, a_2, a_3) = (pqrw, s, z), \quad (b_1, b_2, b_3) = (w, qrs, pz), \quad (c_1, c_2, c_3) = (pw, qs, rz),$$

where

$$s = pqr(r-p) + p^2 - p - r + 1, \quad w = q(r^2 - p - r + 1) + p - r, \\ \text{and} \quad z = pqr(qr - q - 1) + p + q - 1.$$

Then, we set

$$f_1(x) = \prod_{i=1}^3 (x + a_i), \quad f_2(x) = \prod_{i=1}^3 (x + b_i), \quad f_3(x) = \prod_{i=1}^3 (x + c_i),$$

so that

$$(2.3) \quad f_1(x) = f_i(x) + (T_1 - T_i)x, \quad i = 2, 3.$$

We now introduce the quartic elliptic curve

$$(2.4) \quad \begin{aligned} E : y^2 &= Ax f_1(x) + B^2 x^2 \\ &= Ax(f_2(x) + (T_1 - T_2)x) + B^2 x^2 \\ &= Ax(f_3(x) + (T_1 - T_3)x) + B^2 x^2, \end{aligned}$$

where

$$T_1 = \sum_{1 \leq i < j \leq 3} a_i a_j, \quad T_2 = \sum_{1 \leq i < j \leq 3} b_i b_j, \quad T_3 = \sum_{1 \leq i < j \leq 3} c_i c_j,$$

and the coefficients A and B will be given later. Note that the last two equalities in (2.4) come from (2.3). By the first equality of (2.4), the elliptic curve E clearly contains the three rational points with abscissas $-a_1$, $-a_2$, $-a_3$. Besides, by the second and third equality of (2.4), the curve E has the additional rational points with abscissas $-b_i$ and $-c_i$, $i = 1, 2, 3$, if and only if the system of equations

$$A(T_1 - T_{i+1}) = m_i^2 - B^2, \quad i = 1, 2,$$

is solvable. This system has the following parametric solution

$$(2.5) \quad \begin{aligned} A &= -4hk(h-k)((T_1 - T_2)h - (T_1 - T_3)k), \\ B &= (T_1 - T_2)h^2 - (T_1 - T_3)k^2, \\ m_i &= (T_1 - T_2)h^2 - 2(T_1 - T_{i+1})hk + (T_1 - T_3)k^2, \quad i = 1, 2, \end{aligned}$$

for any nonzero, unequal rationals h, k .

Henceforth, we have constructed a quartic family of elliptic curves, defined over $\mathbb{Q}(p, q, r, h, k)$, coming from the elements of \mathcal{B} , that has the nine points

$$P_i = (-a_i, a_i B), \quad Q_i = (-b_i, b_i m_1), \quad R_i = (-c_i, c_i m_2), \quad i = 1, 2, 3,$$

where A, B, m_1 and m_2 are given in (2.5).

The quartic elliptic curve E is birationally equivalent to the cubic elliptic curve $\mathcal{E}_{\mathcal{B}}$ given by the equation

$$Y^2 = X^3 + (AT_1 + B^2)X^2 + A^2 M_i N_i X + A^3 N_i^2, \quad i = 1, 2, 3,$$

via the rational transformation $X = AN_i/x, Y = AN_i y/x^2$.

In the view of this transformation, the nine points P_i, Q_i and $R_i, i = 1, 2, 3$, are mapped respectively to the points $\mathcal{P}_i, \mathcal{Q}_i$ and \mathcal{R}_i , given below,

$$\begin{aligned} \mathcal{P}_1 &= (-a_2 a_3 A, a_2 a_3 AB), \quad \mathcal{Q}_1 = (-b_2 b_3 A, b_2 b_3 A m_1), \quad \mathcal{R}_1 = (-c_2 c_3 A, c_2 c_3 A m_2), \\ \mathcal{P}_2 &= (-a_1 a_3 A, a_1 a_3 AB), \quad \mathcal{Q}_2 = (-b_1 b_3 A, b_1 b_3 A m_1), \quad \mathcal{R}_2 = (-c_1 c_3 A, c_1 c_3 A m_2), \\ \mathcal{P}_3 &= (-a_1 a_2 A, a_1 a_2 AB), \quad \mathcal{Q}_3 = (-b_1 b_2 A, b_1 b_2 A m_1), \quad \mathcal{R}_3 = (-c_1 c_2 A, c_1 c_2 A m_2). \end{aligned}$$

REMARK 2.5. Notice that the points \mathcal{P}_i are collinear (because their (x, y) -coordinates satisfy the linear equation $x + By = 0$), showing that at most two of them can be linearly independent. The same result holds for each of the points \mathcal{Q}_i and \mathcal{R}_i . Therefore, at most six of the nine points of $\mathcal{E}_{\mathcal{B}}$ can be linearly independent.

In the next theorem we show that six of the nine points of $\mathcal{E}_{\mathcal{B}}$ are linearly independent for infinitely many tuples (p, q, r, h, k) .

THEOREM 2.6. *For the set of all tuples (p, q, r, h, k) except for a thin subset, the rank of the family $\mathcal{E}_{\mathcal{B}}$ is at least six with the six linearly independent points $\mathcal{P}_i, \mathcal{Q}_i, \mathcal{R}_i, i = 1, 2$.*

PROOF. Take the specialisation $(p, q, r, h, k) = (2, 2, 4, 1, -1)$ which makes $s = 31, z = 83, w = 20$, and hence we have

$$\mathcal{B}(434, 823360) = \{(320, 31, 83), (20, 248, 166), (40, 62, 332)\}$$

and

$$\begin{aligned} T_1 &= 39053, \quad T_2 = 49448, \quad T_3 = 36344, \\ A &= -61488, \quad B = -13104, \quad m_1 = -28476, \quad m_2 = -2268. \end{aligned}$$

The regulator of the six specialised points with X -coordinates

$$\begin{aligned} X(\mathcal{P}_1) &= 158208624, \quad X(\mathcal{Q}_1) = 2531337984, \quad X(\mathcal{R}_1) = 1265668992, \\ X(\mathcal{P}_2) &= 1633121280, \quad X(\mathcal{Q}_2) = 204140160, \quad X(\mathcal{R}_2) = 816560640, \end{aligned}$$

on the specialised elliptic curve

$$\begin{aligned} \mathcal{E}_{\mathcal{B}(434, 823360)} : Y^2 &= X^3 - 2229576048X^2 + 1351015178454466560X \\ &\quad - 157597974109784775013171200 \end{aligned}$$

is the nonzero value 534.520794417629, carried out by SageMath. The result follows from Theorem 2.1. \square

REMARK 2.7. Corresponding to each of the known subsets $\mathcal{S} \neq \mathcal{B}$ of $\mathcal{S}_\ell^{(n)}(M, N)$ with $\ell \geq 3$, being found in [31] and [18] (cf. Table 1) and in Subsection 2.3, one can assign a family of elliptic curves similar to $\mathcal{E}_\mathcal{B}$ and accordingly establish a result as that of Theorem 2.6.

2.3. *A family with rank ≥ 8 .* For $(\ell, n) = (4, 1)$, there is no known parametric solution of (1.1). We first begin with presenting such a parametric solution.

PROPOSITION 2.8. *For arbitrary integers q and s with $q \neq 0, s^2, s \neq 0, 1$, let*

$$\begin{aligned} t_1 &= q^2 s^2 - 2q^2 s + qs + q - 1, \\ t_2 &= q^2 s^2 - 2q^2 s + qs^2 - qs + s^2 + q - s, \\ t_3 &= q^3 s^3 - q^2 s^4 - 4q^3 s^2 + 5q^2 s^3 - qs^4 + 4q^3 s - 6q^2 s^2 + 2qs^3 - s^4 + q^2 s \\ &\quad + 2s^3 - 2q^2 + qs - 2s^2 + q, \\ t_4 &= -q^2 s^5 + q^3 s^3 + 4q^2 s^4 - 4q^3 s^2 - 4q^2 s^3 - 2qs^4 + 4q^3 s + q^2 s^2 + 4qs^3 \\ &\quad - q^2 s - s^3 - 2q^2 - qs + s^2 + 2q - s, \\ t_5 &= q^2 s^7 + q^4 s^4 - 3q^3 s^5 - 4q^2 s^6 + qs^7 - 4q^4 s^3 + 15q^3 s^4 - qs^6 + 4q^4 s^2 \\ &\quad - 22q^3 s^3 + 12q^2 s^4 - 5qs^5 + 2s^6 + 10q^3 s^2 - 12q^2 s^3 + 7qs^4 - 4s^5 \\ &\quad - 4q^3 s + 12q^2 s^2 - 7qs^3 + 4s^4 - 4q^2 s + qs^2 - s^3 + q^2. \end{aligned}$$

Then,

$$\begin{aligned} (a_1, a_2, a_3) &= ((s-1)t_1 t_2 t_3, sq(1-s)t_1 t_5, (s^2-q)t_1^2 t_4), \\ (b_1, b_2, b_3) &= ((s-1)qt_1 t_2 t_3, -t_1^2 t_5, (s-1)s(s^2-q)t_1 t_4), \\ (c_1, c_2, c_3) &= ((1-s)t_1 t_5, (s-1)qt_1 t_2 t_4, s(s^2-q)t_1^2 t_3), \\ (d_1, d_2, d_3) &= ((1-s)qt_1 t_5, (s-1)st_1 t_2 t_4, (s^2-q)t_1^2 t_3) \end{aligned}$$

is a two-parameter solution to (1.1) with $(\ell, n) = (4, 1)$.

PROOF. Consider $(a_1, a_2, a_3) = (p, wqs, rz)$, $(b_1, b_2, b_3) = (pq, rw, sz)$, $(c_1, c_2, c_3) = (w, qr, spz)$ and $(d_1, d_2, d_3) = (wq, rs, pz)$. Then, clearly the products of the components of these triples are equal. Solving the equations $M_1 = M_2, M_2 = M_3, M_3 = M_4$ gives three formulas for w . By the equality between the first two and the second two of the resulting w 's, we get two

formulas for z . The equality between these z 's gives rise to

$$(q^2s^2 - 2q^2s + qs - rs + q + r - 1)p^2 + (-qrs^3 + q^2rs + r^2s^2 + q^2r - qr^2 - qr - qs + s)p - r(q^2s^2 - qs^3 + q^2r - q^2s - qr^2 + r^2s - rs + s^2) = 0.$$

By putting $q^2s^2 - 2q^2s + qs - rs + q + r - 1 = 0$ in the latter equation, it follows that

$$(2.6) \quad r = \frac{q^2s^2 - 2q^2s + qs + q - 1}{s - 1}$$

and

$$(2.7) \quad p = -r \frac{q^2s^2 - qs^3 + q^2r - q^2s - qr^2 + r^2s - rs + s^2}{qrs^3 - q^2rs - r^2s^2 - q^2r + qr^2 + qr + qs - s}.$$

Now, by substituting (2.6) in (2.7) and one of the obtained z 's and w 's, we get a rational solution of (1.1) for $(\ell, n) = (4, 1)$ in terms of q and s . By scaling we get the desired result as given in the statement of the theorem. \square

Now, we consider the subset $\mathcal{C} := \mathcal{C}(M, N)$ of $\mathcal{S}_4(M, N)$ with the elements introduced in Proposition 2.8. Then, we set

$$f_1(x) = \prod_{i=1}^3 \{(x + a_i)(x + b_i)\}, \quad f_2(x) = \prod_{i=1}^3 \{(x + c_i)(x + d_i)\},$$

so that

$$(2.8) \quad f_1(x) - f_2(x) = Tx^4 + MTx^3 + (T_1T_2 - T_3T_4)x^2 + NTx,$$

where

$$T_1 = \sum_{1 \leq i < j \leq 3} a_i a_j, \quad T_2 = \sum_{1 \leq i < j \leq 3} b_i b_j, \quad T_3 = \sum_{1 \leq i < j \leq 3} c_i c_j, \quad T_4 = \sum_{1 \leq i < j \leq 3} d_i d_j,$$

and

$$T = T_1 + T_2 - T_3 - T_4.$$

One can rewrite (2.8) as $\phi_1^2(x) - \phi_2^2(x)$ where

$$\phi_1(x) = x^3 + Mx^2 + \frac{4(T_1T_2 - T_3T_4) + T^2}{4T}x + N,$$

$$\phi_2(x) = x^3 + Mx^2 + \frac{4(T_1T_2 - T_3T_4) - T^2}{4T}x + N.$$

It follows that

$$\phi_i^2(x) - f_i(x) = Kx^4 + MKx^3 + Hx^2 + NKx, \quad i = 1, 2,$$

where

$$K = -\frac{T^2 + 2(T_3 + T_4)T - 4(T_1T_2 - T_3T_4)}{2T},$$

$$H = \frac{(T^2 + 4(T_1T_2 - T_3T_4))^2}{16T^2} - T_1T_2.$$

We now introduce the quartic elliptic curve

$$(2.9) \quad y^2 = Kx^4 + MKx^3 + Hx^2 + NKx,$$

which contains the rational points $P_{a_i}, P_{b_i}, P_{c_i}, P_{d_i}$ with abscissas $-a_i, -b_i, -c_i$ and $-d_i$ for $i = 1, 2, 3$, respectively.

The quartic curve (2.9) reduces to the following cubic elliptic curve

$$(2.10) \quad \mathcal{E}_C : Y^2 = X^3 + HX^2 + MNK^2X + N^2K^3,$$

by the rational transformation $X = NK/x, Y = NKy/x^2$. In the view of this transformation, the above twelve points are sent respectively to the points $\mathcal{P}_{a_i}, \mathcal{P}_{b_i}, \mathcal{P}_{c_i}, \mathcal{P}_{d_i}$ with X -coordinates

$$-\frac{NK}{a_i}, -\frac{NK}{b_i}, -\frac{NK}{c_i}, \text{ and } -\frac{NK}{d_i} \quad \text{for } i = 1, 2, 3.$$

We have thus constructed a family of elliptic curves, defined over $\mathbb{Q}(q, s)$, coming from the elements of \mathcal{C} , that has the above twelve points.

REMARK 2.9. Notice that the points \mathcal{P}_{a_i} are collinear, showing that at most two of them can be linearly independent. The same result holds for each of the points $\mathcal{P}_{b_i}, \mathcal{P}_{c_i}$ and \mathcal{P}_{d_i} . Therefore, at most eight points of the twelve points of \mathcal{E}_C can be linearly independent.

In the next theorem we show that eight of the twelve points of \mathcal{E}_C are linearly independent for infinitely many tuples (q, s) .

THEOREM 2.10. *For the set of all tuples (q, s) except for a thin subset, the rank of the family \mathcal{E}_C is at least eight with the eight linearly independent points $\mathcal{P}_{a_i}, \mathcal{P}_{b_i}, \mathcal{P}_{c_i}, \mathcal{P}_{d_i}, i = 1, 2$.*

PROOF. We specialise at $(q, s) = (3, 2)$ which makes

$$t_1 = 8, \quad t_2 = 11, \quad t_3 = 1, \quad t_4 = -6, \quad t_5 = 29,$$

and hence we have

$$\mathcal{C}(-1688, 47038464) = \{(88, -1392, -384), (264, -1856, -96), \\ (-232, -1584, 128), (-696, -1056, 64)\}$$

and

$$T_1 = 378240, \quad T_2 = -337152, \quad T_3 = 135040, \quad T_4 = 622848, \\ T = -716800, \quad K = 191008, \quad H = 140991510784.$$

The regulator of the eight points with X -coordinates

$$\begin{aligned} X(\mathcal{P}_{a_1}) &= -102099124224, & X(\mathcal{P}_{c_1}) &= 38727254016, \\ X(\mathcal{P}_{a_2}) &= 6454542336, & X(\mathcal{P}_{c_2}) &= 5672173568, \\ X(\mathcal{P}_{b_1}) &= -34033041408, & X(\mathcal{P}_{d_1}) &= 12909084672, \\ X(\mathcal{P}_{b_2}) &= 4840906752, & X(\mathcal{P}_{d_2}) &= 8508260352 \end{aligned}$$

on the specialised elliptic curve

$$\begin{aligned} \mathcal{E}_C : Y^2 &= X^3 + 140991510784X^2 - 2896867880665872334848X \\ &+ 15419167818458889008922652311552 \end{aligned}$$

is the nonzero value 15150.2483213544, carried out by SageMath. The result follows now from Theorem 2.1. \square

3. HEURISTICS

As all curves in the families \mathcal{E}_A , \mathcal{E}_B and \mathcal{E}_C have trivial torsion subgroups in general, to identify promising high-rank candidates we used heuristic arguments (motivated by the BSD conjecture) given by Mestre [27] and Nagao [30]. They suggest that for curves of high rank, certain sums should assume the largest values in the observed families. The first of these sums is

$$S_1(X) = \sum_{p \leq X} \frac{N_p + 1 - p}{N_p} \log p$$

where X is a prime bound for primes p , and $N_p = |E(\mathbb{F}_p)|$ is the number of points on elliptic curve E under the reduction modulo p . We followed the sieve phase of the general method for finding high-rank elliptic curves described in [21, pp. 64–68].

The Mestre–Nagao sum $S_1(X)$ with the prime bound $X = 10^6$ was calculated in Magma [3] for all curves with the absolute values of parameters p, q, r, s, t, k, h up to 40. For curves with $S_1(10^6) \geq 100$, `TwoSelmerGroup` function in Magma was used to deduce the upper bound R on the rank of the Mordell–Weil group of the curve. For all curves with $R \geq 11$, `DescentInformation` function in Magma was used to uncover the independent generators of the Mordell–Weil group of the curve.

Parameters for some high-rank curves are listed in the next tables. Note that all listed curves are non-isomorphic.

We were also able to find a number of curves of rank 15 in the form (2.10) where the set of four triples $\{(a_1, a_2, a_3), (b_1, b_2, b_3), (c_1, c_2, c_3), (d_1, d_2, d_3)\}$ was determined by a direct search given a fixed sum M rather than using Proposition 2.8. The first curve of rank 15 occurs for $M = 4907$ ($N = 1628394768$) whose respective set is

$$\{(632, 726, 3549), (312, 2054, 2541), (507, 924, 3476), (474, 1001, 3432)\};$$

the minimal model of the curve is

$$y^2 + xy = x^3 - 901882569760647935195484561648738x + 13932920298020241870290850727467604223423326273092.$$

Magma's special function `DescentInformation` was able to find 15 independent points on this curve in 184 core-hours on Intel Core i7-8700 CPU, although our knowledge of the 8 independent points on (2.10) reduced the search time to 1.6 core-hours (over a hundredfold calculation speedup).

In that light, Proposition 2.8 and a rank 13 curve obtained from rational (q, s) in the last table present significant value.

TABLE 2. High rank curves $\mathcal{E}_{\mathcal{A}}$

(p, q, r, s, t, k)	Rank	(p, q, r, s, t, k)	Rank
$(-71, -54, -36, 14, 36, 1)$	14	$(32, 41, 55, 60, 46, 1)$	14
$(-37, -36, 37, 25, 36, 37)$	13	$(-33, -29, 17, 6, 13, 9)$	13
$(-36, -33, 31, 6, 23, 7)$	13	$(-30, -27, 12, 16, 13, 17)$	13
$(-35, -34, 17, 29, 14, 13)$	13	$(-25, -15, 20, 8, 21, 22)$	13
$(-35, -31, 8, 12, 18, 11)$	13	$(-22, -11, 22, 13, 22, 2)$	13
$(-22, -18, 4, 20, 9, 19)$	12	$(-19, -18, -5, 6, 19, 1)$	12
$(-21, -13, 11, 14, 15, 19)$	12	$(-18, 6, 12, 16, 6, 11)$	12
$(-20, -16, 13, 3, 16, 7)$	12	$(-17, -10, 9, 1, 14, 9)$	12
$(-19, 8, 16, 7, 9, 16)$	12	$(-16, -5, 20, 2, 25, 1)$	12

4. A CONCLUDING REMARK

In this work, we considered the system of Diophantine equations

$$\sum_{i=1}^3 x_{1i} = \cdots = \sum_{i=1}^3 x_{\ell i}, \quad \prod_{i=1}^3 x_{1i} = \cdots = \prod_{i=1}^3 x_{\ell i}$$

for ℓ ranging from 2 to 4, which from the geometric point of view is equivalent to ℓ rectangular boxes with integer sides having the same perimeter and the same volume, and for each system we introduced a family of elliptic curves and then studied them closely. Here arises this natural question that whether there exists more than four rectangular boxes with integer sides that have the same perimeter and the same volume.

ACKNOWLEDGEMENTS.

We are grateful to the referees for the valuable comments that helped improve this article.

TABLE 3. High rank curves \mathcal{E}_B

(p, q, r, h, k)	Rank	(p, q, r, h, k)	Rank
$(-12, 7, -3, -1, 15)$	13	$(-1, -8, -6, 4, 5)$	13
$(-4, 5, -13, 1, -1)$	13	$(4, -3, -14, -14, 13)$	13
$(-2, -13, 11, 6, 1)$	13	$(11, -2, -12, -3, 4)$	13
$(-2, 8, 15, -4, 11)$	13	$(14, -2, -13, 8, 1)$	13
$(-11, 5, 3, 6, 5)$	12	$(-3, 10, 3, -5, 2)$	12
$(-8, -6, 9, -7, 5)$	12	$(-2, 10, 2, -7, 8)$	12
$(-7, 3, 2, -5, 8)$	12	$(-1, 7, -9, -6, 1)$	12
$(-5, -6, 5, 4, -3)$	12	$(3, -7, 9, 6, 7)$	12
$(-6, 3, -1, 6, -2)$	11	$(-2, 4, 3, 5, -4)$	11
$(-5, -5, 5, -3, -2)$	11	$(2, 3, 6, -2, -5)$	11
$(-4, -3, 4, 6, -2)$	11	$(2, 4, -5, -4, -3)$	11
$(-3, -4, 3, -5, -3)$	11	$(5, -2, -5, -1, 5)$	11

TABLE 4. High rank curves \mathcal{E}_C

(q, s)	Rank	(q, s)	Rank
$(7/11, -1)$	13		
$(-12/11, 2)$	12	$(-1/4, -1)$	12
$(-9/5, -2/5)$	12	$(3/4, 2)$	12
$(-8, -7/2)$	12	$(9/5, -1)$	12
$(-2/15, 2)$	12	$(12/7, 14/9)$	12
$(-11/7, -1/3)$	11	$(-4/13, 2)$	11
$(-9, -1)$	11	$(-2/3, 3)$	11
$(-5/7, -1/2)$	11	$(1/7, -1)$	11
$(-4/9, 2)$	11	$(11/8, 6/5)$	11

REFERENCES

- [1] J. Aguirre and J. C. Peral, *Sums of biquadrates and elliptic curves*, Glas. Mat. Ser. III **48(68)** (2013), 49–58.
- [2] U. Bini, *Sopra alcune proprietà delle forme binarie cubiche*, Interméd. Math. **16** (1909), 41–43.
- [3] W. Bosma, J. Cannon and C. Playoust, *The Magma algebra system. I. The user language*, J. Symbolic Comput. **24** (1997), 235–265.
- [4] B. Cha, A. Claman, J. Harrington, Z. Liu, B. Maldonado, A. Miller, A. Palma, T.W.H. Wong and H. Yi, *An investigation on partitions with equal products*, Int. J. Number Theory **15** (2019), 1731–1744.
- [5] A. Choudhry, *Symmetric Diophantine systems*, Acta Arith. **59** (1991), 291–307.

- [6] A. Choudhry, *On triads of squares with equal sums and equal products*, *Ganita* **49** (1998), 101–106.
- [7] A. Choudhry, *Triads of cubes with equal sums and equal products*, *Math. Student* **70** (2001), 137–143.
- [8] A. Choudhry, *Triads of biquadrates with equal sums and equal products*, *Math. Student* **70** (2001), 149–152.
- [9] A. Choudhry, *Some Diophantine problems concerning equal sums of integers and their cubes*, *Hardy–Ramanujan J.* **33** (2010), 59–70.
- [10] A. Choudhry, *Triads of integers with equal sums and equal products*, *Math. Student* **81** (2012), 185–188.
- [11] A. Choudhry, *Equal sums of like powers and equal products of integers*, *Rocky Mountain J. Math.* **43** (2013), 763–792.
- [12] A. Choudhry, *Equal sums of like powers with minimum number of terms*, *Integers* **16** (2016), Paper No. A77, 11.
- [13] A. Choudhry, *A diophantine problem on biquadrates revisited*, *Math. Student* **87** (2018), 129–132.
- [14] A. Choudhry, *Symmetric diophantine systems and families of elliptic curves of high rank*, *Rocky Mountain J. Math.* **49** (2019), 1419–1447.
- [15] A. Choudhry, *New solutions of the Tarry–Escott problem of degrees 2, 3 and 5*, *J. Integer Seq.* **24** (2021), Art. 21.8.1.
- [16] A. Choudhry, *Ideal solutions of the Tarry–Escott problem of degree seven*, *Integers* **22** (2022), Paper No. A115.
- [17] A. Choudhry and A. Shamsi Zargar, *An octic diophantine equation and related families of elliptic curves*, *Int. J. Number Theory* **19** (2023), 1967–1976.
- [18] A. Choudhry and J. Wróblewski, *Triads of integers with equal sums of squares and equal products and a related multigrade chain*, *Acta Arith.* **178** (2017), 87–100.
- [19] L. E. Dickson, *History of the theory of numbers II*, Chelsea Publishing Company, New York, 1920.
- [20] A. Dujella, *An example of elliptic curve over \mathbf{Q} with rank equal to 15*, *Proc. Japan Acad. Ser. A Math. Sci.* **78** (2002), 109–111.
- [21] A. Dujella, *Diophantine m -tuples and elliptic curves*, Springer, Cham, 2024.
- [22] A. Gloden, *Mehrgradige Gleichungen*, Noordhoff, Groningen, 1944.
- [23] R. K. Guy, *Unsolved problems in number theory*, Springer-Verlag, New York, 1994.
- [24] J. B. Kelly, *Partitions with equal products (II)*, *Proc. Amer. Math. Soc.* **107** (1989), 887–893.
- [25] J. B. Kelly, *Two equal sums of three squares with equal products*, *Amer. Math. Monthly* **98** (1991), 527–529.
- [26] J. G. Mauldon, *Problem E 2872*, *Amer. Math. Monthly* **88** (1981), 148.
- [27] J. F. Mestre, *Construction d’une courbe elliptique de rang ≥ 12* , *C. R. Acad. Sci. Paris Sér. I.* **295** (1982), 643–644.
- [28] A. Moessner, *Verschiedene Diophantische Probleme und numerische Identitäten*, *Tohoku Math. J.* **47** (1940), 188–200.
- [29] L. J. Mordell, *Diophantine equations*, Academic Press, London-New York, 1969.
- [30] K. Nagao, *An example of elliptic curve over Q with rank ≥ 20* , *Proc. Japan Acad. Ser. A Math. Sci.* **69** (1993), 291–293.
- [31] M. Sadek and N. El-Sissi, *Partitions with equal products and elliptic curves*, *Osaka J. Math.* **52** (2015), 515–525.
- [32] Sage Developers, *SageMath, Sage mathematics software system (ver. 9.3)*, 2021, available at <https://www.sagemath.org>
- [33] A. Schinzel, *Triples of positive integers with the same sum and the same product*, *Serdica Math. J.* **22** (1996), 587–588.

- [34] J. H. Silverman, *The arithmetic of elliptic curves*, Springer, Dordrecht, 2009.
- [35] D. Zeitlin, *Abstracts presented to American Mathematical Society*, **57** (1988), #843-11-50.
- [36] Y. Zhang and T. Cai, *n-tuples of positive integers with the same sum and the same product*, *Math. Comp.* **82** (2013), 617–623.

A. E. A. Youmbai
 LABTHOP Laboratory, Mathematics Department, Faculty of Exact Sciences
 University of El Oued
 PO Box 789, 39000 Echott El Oued
 Algeria
E-mail: youmbai-amine@univ-eloued.dz

A. Shamsi Zargar
 Department of Mathematics and Applications
 Faculty of Mathematical Sciences
 University of Mohaghegh Ardabili
 Ardabil
 Iran
E-mail: zargar@uma.ac.ir

M. Voznyy
 Department of Technology, Stephen Leacock CI
 Toronto District School Board
 Toronto
 Canada
E-mail: maksym.voznyy@tdsb.on.ca

Received: 18.7.2024.

Revised: 21.2.2025.

PARTICIJE U TROJKE S JEDNAKIM PRODUKTIMA I FAMILIJE ELIPTIČKIH KRIVULJA

A. E. A. YOUMBAL, A. SHAMSI ZARGAR I M. VOZNYI

SAŽETAK. Neka $S_\ell(M, N)$ označava skup ℓ (različitih) trojki pozitivnih cijelih brojeva koje imaju isti zbroj M i isti produkt N . Za svaki $2 \leq \ell \leq 4$ uspostavljamo vezu između podskupa $S_\ell(M, N)$ s (cjelobrojnim) parametarskim elementima i familijom eliptičkih krivulja. Kada je $\ell = 2$ i 3 , koristimo određene poznate podskupove $S_\ell(M, N)$ s parametarskim elementima i, respektivno, pronalazimo familije eliptičkih krivulja generičkog ranga ≥ 5 i ≥ 6 , dok za $\ell = 4$ prvo dobivamo podskup $S_\ell(M, N)$ s parametarskim elementima, a zatim konstruiramo familije eliptičkih krivulja generičkog ranga ≥ 8 . Na kraju, provodimo računalnu pretragu unutar tih familija kako bismo pronašli specifične krivulje s rangom ≥ 11 i posebno smo pronašli dvije krivulje ranga 14.