

EXPANSION OF THE NON-INDIGENOUS SPONGE
PARALEUCILLA MAGNA KLAUTAU, MONTEIRO
& BOROJEVIC, 2004 (PORIFERA: CALCAREA) ALONG THE
MEDITERRANEAN COAST OF MOROCCO

FOUZI ABDELKHALEQ TAYBI¹ , KAMAL GOURARI², ABDESSADEK ESSADEK³
& YOUNESS MABROUKI⁴ 

¹Multidisciplinary Faculty of Nador, Research Team in Applied Biology and Biotechnology, Mohamed Premier University, Nador, Morocco
(e-mail: a.taybi@ump.ac.ma)

²Faculty of Sciences, Laboratory for the Improvement of Agricultural Production, Biotechnologies and the Environment, Mohamed Premier University, Oujda, Morocco (e-mail: k.gourari@ump.ac.ma)

³Université Mohammed Premier, Laboratoire de Bioressources, Biotechnologie, Ethnopharmacologie et Santé. Équipe: Biochimie appliquée à la valorisation des bioressources Département de Biologie
(e-mail: a.essadek@ump.ac.ma)

⁴Faculty of Sciences of Dhar El Mehraz, Laboratory of Biotechnology, Conservation and Valorization of Bioresources, Sidi Mohamed Ben Abdellah University, Fez, Morocco
(e-mail: youness.mabrouki@usmba.ac.ma)

Taybi, F.A., Gourari, K., Essadek, A. & Mabrouki, Y.: Expansion of the non-indigenous sponge *Paraleucilla magna* Klautau, Monteiro & Borojevic, 2004 (Porifera: Calcarea) along the Mediterranean coast of Morocco. Nat. Croat., Vol. 34, No. 2, 327–335, 2025, Zagreb.

The Mediterranean Sea is experiencing an increased impact of biological invasions, including along the coastlines of North Africa. The non-indigenous sponge *Paraleucilla magna* Klautau, Monteiro & Borojevic, 2004 was detected recently in the western Mediterranean coast of Morocco. In this paper we document for the first time the occurrence of this non-indigenous species (NIS) in Marchica Lagoon, a Ramsar site and protected area situated on the eastern Mediterranean coast of Morocco. Additionally, we report it from the Saidia Marina, the easternmost location in the country where the species has been found, demonstrating its rapid expansion. The Marchica lagoon and Saidia marina have become hotspots for the proliferation of NIS, underscoring the pressing need for monitoring and management strategies to address the issue of biological invasions in this region.

Keywords: North Africa, protected area, Marchica lagoon, Saidia Marina, non-indigenous species, range-expanding species.

Taybi, F.A., Gourari, K., Essadek, A. & Mabrouki, Y.: Širenje alohtone spužve *Paraleucilla magna* Klautau, Monteiro & Borojevic, 2004 (Porifera: Calcarea) duž sredozemne obale Maroka. Nat. Croat., Vol. 34, No. 2, 327–335, 2025, Zagreb.

Sredozemno more doživljava sve veći utjecaj bioloških invazija, uključujući i sredozemne obale sjeverne Afrike. Alohtona spužva *Paraleucilla magna* Klautau, Monteiro & Borojevic, 2004 nedavno je zabilježena na zapadnoj sredozemnoj obali Maroka. U ovom radu prvi put dokumentiramo pojavu ove alohtone vrste (NIS) u laguni Marchica, Ramsarskom lokalitetu i zaštićenom području smještenom na istočnoj sredozemnoj obali Maroka. Osim toga donosimo i njen nalaz iz marine Saidia, najistočnijeg lokaliteta u zemlji na kojem je vrsta pronađena, što pokazuje njezino brzo širenje. Laguna Marchica i marina Saidia postale su žarišta za širenje alohtonih vrsta, što naglašava hitnu potrebu za monitoringom i strategijom upravljanja u cilju rješavanja problema bioloških invazija u ovoj regiji.

Ključne riječi: sjeverna Afrika, zaštićeno područje, laguna Marchica, Saidia Marina, alohtone vrste, proširenje rasprostranjenosti vrste

INTRODUCTION

The introduction and expansion of invasive and non-indigenous species represent a significant threat to biodiversity and ecosystem functioning on a global scale, resulting in substantial environmental and economic costs (DIAGNE *et al.*, 2021). In the Mediterranean region, the biological invasion of non-indigenous species has been on the rise since the 1970s, posing a serious threat to the region's native biodiversity (OJAVEER *et al.*, 2013; FRISTOE *et al.*, 2021). The number of reported invasive species in the Mediterranean is one of the highest worldwide (STREFTARIS *et al.*, 2005). Consequently, the study of invasive species has become a pressing research priority for the conservation of indigenous biodiversity (DUKES & MOONEY, 2004; TAYBI *et al.*, 2020a).

Paraleucilla magna Klautau, Monteiro & Borojevic, 2004 is a calcareous sponge of tubular or massive outer shape, presenting numerous flexible digitations of very variable size and terminated by an oscule of 5 to 20 mm in diameter. It was originally described from the Brazilian coast in the Atlantic Ocean, where it is the most abundant calcareous sponge (KLATAU *et al.*, 2004). The sponge is an important component of marine nutrient cycling and the filtration of large volumes of ambient water. However, when invasive it also poses a significant threat to native sponge communities. Previous studies have demonstrated that *P. magna* can displace native species through competition for space and resources, thus potentially changing the ecological balance in the invaded ecosystems (LANNA & KLAUTAU, 2010; GUZETTI *et al.*, 2019). The first record of *Paraleucilla magna* in the Mediterranean was in Mar Piccolo and Mar Grande in Taranto (north-western Ionian Sea) in 2001 (LONGO *et al.*, 2004). The species has certainly persisted in Mar Piccolo since then. However, local mussel farmers claim that the species colonised this environment 20–30 years earlier (LONGO *et al.*, 2004, 2007). Subsequently, the species has been found in various other locations in Italy, as well as in Spain, Malta, Croatia, Montenegro, Turkey, Cyprus, Greece, Tunisia and Algeria (GEROVASILEIOU *et al.*, 2017; BACHETARZI *et al.*, 2019; TIRALONGO *et al.*, 2022). In Morocco, the non-indigenous sponge was recorded recently from the western coast of the Mediterranean, where it was found at aquaculture facilities in Amsa and M'diq during February and March 2024 (EL MOUSSAOUI *et al.*, 2024). In this paper we provide the first record of *Paraleucilla magna* in the Marchica lagoon and Saidia Marina, representing the easternmost record of this species on the Mediterranean coast of Morocco.

MATERIAL & METHODS

Several sampling and prospecting trips were carried out along the Eastern Mediterranean coast of Morocco from June of 2019 (and ongoing) in order to monitor the spread of non-indigenous species (part of a structured survey), from the western side of Al Hoceima to the eastern side of Saidia, with a special interest to the protected areas of Morocco, such as the Marchica lagoon. The sampling was carried out by looking into the brackish and marine waters near the shore using a hawk net (mesh size: 1×1 mm / 0.5 mm) at a depth ranging between 0.5 and 2 m. Two voucher specimens of *Paraleucilla magna* were deposited at University Mohamed Premier of Oujda Morocco under the accession codes: F_A_T: spong.PM032023/1 and F_A_T: spong.PM032023/2. And two additional specimens from Marina Saïdia were catalogued under F_A_T: spong.PM062025/1 and /2. The species was identified using the spicule morphology and squelettic architecture reported by LONGO *et al.* (2007).

Specimens were kept in 70% ethanol. Each specimen has been analysed using an optical microscope to study its external morphology. Spicules were visible, fragments at oscula lateral and apical levels have been cut at its end. Samples were dehydrated through a graded ethanol series, rinsed with distilled water, and sectioned. The sections were then stained with eosin and photographed using a Leica Microsystems microscope fitted with a digital camera.

Duplicated profiles ($n = 3$ per site) were collected *in situ* at depths of 0.5–1 m in Marchica Lagoon and 1–3 m in Saïdia Marina using a WTW MPP 350 probe, which simultaneously logs temperature, pH, conductivity, salinity and dissolved oxygen.

RESULTS

Several individuals of *Paraleucilla magna* were found in the Marchica lagoon Ramsar site (35°11'09.2"N 2°55'25.9"W) on 15 March 2024 and in the Saïdia Marina (35°06'36.9"N 2°17'47.8"W) on 1 January 2025, at an average depth of 0.5–1 m, representing the easternmost records of the species on the Mediterranean coast of Morocco to date. The distance between the two sites is 70 km (Fig. 1). These results highlight the species' expansion across the Mediterranean. The mean values of physicochemical parameters of the water in the studied localities are presented in Tab. 1. Other non-indigenous species were found alongside *P. magna* in both Marchica lagoon and Saïdia Marina, such as the fanworm *Branchiomma luctuosum* (Grube, 1870), the Atlantic blue crab *Callinectes sapidus* Rathbun, 1896 and different non-indigenous cryptogenic tunicate species e.i. *Botrylloides niger* Herdman, 1886, *Microcosmus squamiger* Michaelsen, 1927, *Polyandrocarpa zorritensis* (Van Name, 1931) and *Styela plicata* (Lesueur, 1823).

Tab. 1. Mean values (\pm SD) values of the physico-chemical water parameters measured at each sampling site ($n = 3$ replicates per site).

Sampling site	Sampling depth (m)	Temperature (°C)	Dissolved oxygen (mg/l)	Electrical conductivity (ms/cm)	Salinity (g/l)	pH
Marchica lagoon	0.5 – 1	24.0 \pm 1.3	9.7 \pm 0.42	55.5 \pm 3.4	38.2 \pm 1.8	8.1 \pm 0.1
Saïdia Marina	1 – 3	22.3 \pm 2.2	10.3 \pm 0.37	43.5 \pm 2.7	4.0 \pm 1.3	8.6 \pm 1.7

The sponges were found attached to solid surfaces. The adult specimens were predominantly white to cream in colour and had a generally robust, massive body with an irregular, lobate outline. The surface bore numerous short, angular protrusions or digitiform branches of varying lengths and thicknesses, giving the sponge an uneven, dendritic appearance (Fig. 2). The measured specimens ranged in size from 0.5 to 8 cm in the Marchica lagoon and from 0.5 to 5 cm in the Saïdia Marina area. The body generally consisted of a broad basal mass or stalk, with dozens of cylindrical to finger-like branches arising from it at irregular intervals. Oscula (exhalant openings) were visible in many areas, particularly near the tips of the branches; these were relatively large, measuring 10–20 mm in diameter, and were positioned at the ends of the tubular branches.

In the cortical region, the spicules included numerous triactines (three-rayed spicules) and tetractines (four-rayed spicules). The triactines exhibited variable symmetry




Fig. 1. The updated distribution of *P. magna* in Morocco (new records red dots, old record black) in the Marchica Lagoon (A) and Saidia Marina (B).

(see Fig. 3). Many of them were sagittal, with one unpaired actine differing in length or angle from the other two. The remaining paired actines were roughly equiangular and typically pointed and conical. Relatively large cortical spicule actines had lengths ranging up to $\sim 400 \mu\text{m}$ (mean $\approx 287 \mu\text{m}$) and widths up to $\sim 40 \mu\text{m}$ (Tab. 2). The atrial skeleton consisted entirely of sagittal triactines.

The unpaired actines in the atrial triactines were consistently shorter than the paired actines. The paired triactines were long, slender and pointed, lying approximately parallel to the atrial surface. The unpaired (intratriatal) actine of these atrial triactines is markedly shorter than the paired actine (e.g. the mean unpaired actin length is $\sim 92 \mu\text{m}$, as against $\sim 196 \mu\text{m}$ for paired actine in one population). These short unpaired actines give the atrial triactines their distinctive T-shaped appearance (see Fig. 3). No tetractines were observed as part of the true atrial mesh. All spicule dimensions were consistent with those of *Paraleucilla magna* (Tab. 2).



Fig. 2. Photo of live specimens of *Paraleucilla magna* in the Marchica lagoon (left) and Saidia Marina (right).

Tab. 2. Measurements taken from *P. magna* spicules from eastern Morocco.

Spicules	Actine	Length (μm)	Thickness (μm)
Cortical triactine	Paired	90–320 (200 ± 60)	8–25 (16 ± 5)
	Unpaired	63–307 (180 ± 70)	7–22 (14 ± 6)
Cortical tetractine	Apical	170–550 (410 ± 100)	16–48 (30 ± 11)
	Basal	149–430 (315 ± 80)	14–40 (32 ± 9)
Subatrial triactine	Paired	65–272 (185 ± 55)	9–27 (18 ± 6)
	Unpaired	70–334 (225 ± 75)	8–29 (19 ± 7)
Subatrial tetractine	Paired	115–390 (255 ± 80)	12–45 (26 ± 10)
	Unpaired	95–380 (210 ± 70)	10–40 (22 ± 8)
Atrial triactine	Paired	122–315 (220 ± 60)	10–30 (20 ± 5)
	Unpaired	45–175 (104 ± 35)	5–18 (12 ± 4)




Fig. 3. The skeletal organisation and diagnostic spicules of *Paraleucilla magna*: Cortical tetractines with inward-pointing apical actines (A); Cortical triactines (B-C); Primary sub-auricular triactine (D); Primary sub-auricular tetractine (E); Atrial triactines (choanoskeletal) (G-F).

DISCUSSION

Morphological and microscopic examinations confirmed the identity of the non-indigenous calcareous sponges collected from the Mediterranean coast of eastern Morocco. *Paraleucilla magna* can be distinguished by a cortex that bares equiangular triactines and robust tetractines with very long apical actines. The sub-atrial layer contains smaller sagittal triactines and tetractines, and lacks diactines and pentactines. The atrial lining consists solely of T-shaped sagittal triactines (KLAUTAU *et al.*, 2004; LONGO *et al.*, 2007; MASTROTOTARO *et al.*, 2007; EL MOUSSAOUI *et al.*, 2024). This combination of characteristics clearly distinguishes *P. magna* from similar species such as *P. cucumis* (Haeckel, 1872) and *P. crosslandi* (Row, 1909) (BOROJEVIC *et al.*, 2002; LONGO *et al.*, 2007).

Paraleucilla magna has been present in the Mediterranean Sea for over several decades, including regions such as the Ionian Sea, Adriatic Sea, Tyrrhenian Sea, and the Sea of Marmara (MAČIĆ & PETOVIĆ, 2017). This non-indigenous calcareous sponge has also been reported on the southern shores of the Mediterranean in North Africa, specifically in Algeria and Tunisia (BACHETARZI *et al.*, 2019; BENSARI *et al.*, 2020; SGHAIER *et al.*, 2019) and recently from the western coast of Morocco (EL MANSOURI *et al.*, 2024). Here, we document the first record of *P. magna* on the eastern Mediterranean coast of Morocco. This finding highlights its well-established presence across much of the Mediterranean basin and underscores its ability to colonize diverse environments, including those on the southern coasts of the region. In the studied area, *P. magna* was not observed before June 2019, which means that the invasion process could be in its early stages. In invaded areas where the non-indigenous Porifera exists, the species has been reported to occur at shallow depths, in eutrophic environments with low water movement (LONGO *et al.*, 2007), as is case at the studied area of Marchica lagoon and Saidia Marina. The non-indigenous species has demonstrated its invasiveness by its ability to colonize different environments with high abundance in a short period of time (CVITKOVIĆ *et al.*, 2013; LONGO *et al.*, 2021). The same scenario is expected along the Moroccan coasts, where drastic measures are needed to increase monitoring and management of the invasion process.

The Mediterranean coast of eastern Morocco is located in a key area for the conservation of marine areas in the Alboran Sea, which is considered a very productive area and has an important socio-economic role in the region (ROBLES, 2010; EL HAMOUTI *et al.*, 2023). However, due to its geographical position and the concomitant intensification of maritime traffic, this area has become an important recipient of biological invasions in recent years, with a large number of NIS, including invertebrates and vertebrates, having been documented in recent years (MABROUKI *et al.*, 2021, 2025a, b; TAYBI & MABROUKI, 2020; TAYBI *et al.*, 2020a, b, 2024). Most of these biological invasions have been documented in semi-enclosed coastal environments such as lagoons and marinas (MABROUKI & TAYBI, 2024; TAYBI *et al.*, 2023, 2025; GOURARI *et al.*, 2025). The Marchica Lagoon and the Saidia Marina have emerged as prominent destinations of interest, attracting both national and international travellers. The accessibility of the Mediterranean coastlines of eastern Morocco, which can be reached by recreational boating and by maritime transport, makes this region vulnerable to the introduction of non-indigenous species. Furthermore, the phenomenon of biofouling has been demonstrated to increase the risk of such introductions and international travel. The identification and monitoring of NIS at regional and national levels have become an urgent requirement.

ACKNOWLEDGEMENTS

We would like to express our gratitude to the editor and reviewers for their helpful and sincere comments, which have greatly improved the manuscript.

Received January 30, 2025

REFERENCES

- BACHETARZI, R., DILMI, S., URIZ, M.J., VÁZQUEZ-LUIS, M., DEUDERO, S. & REBZANI-ZAHAF, C., 2019: The non-indigenous and invasive species *Paraleucilla magna* Klautau, Monteiro & Borojević, 2004 (Porifera: Calcarea) in the Algerian coast (Southwestern of Mediterranean Sea). *Acta Adriatica* **60**(1), 41–46. <https://doi.org/10.32582/aa.60.1.3>
- BENSARI, B., BAHBAH, L., LOUNAOUCI, A., EDDINA FAHCI, S., BOUDA, A. & ISLAM BACHARI, N., 2020: First records of non-indigenous species in port of Arzew (Algeria: southwestern Mediterranean). *Mediterranean Marine Science* 393–399. [doi:https://doi.org/10.12681/mms.21927](https://doi.org/10.12681/mms.21927)
- BOROJEVIC, R., BOURY-ESNAULT, N., MANUEL, M. & VACELET, J., 2002: Order Leucosolenida Hartman, 1958. *Systema Porifera* 1157–1184. https://doi.org/10.1007/978-1-4615-0747-5_120
- CVITKOVIĆ, I., DESPALATOVIĆ, M., GRUBELIĆ I., NIKOLIĆ V., PLEŠE B. & ŽULJEVIĆ, A., 2013: Occurrence of *Paraleucilla magna* (Porifera: Calcarea) in the eastern Adriatic Sea. *Acta Adriatica* **54**(1), 93–99.
- DIAGNE, C. LEROY, B., VAISSIÈRE, A.C., GOZLAN, R.E., ROIZ, D., JARIĆ, I., SALLES, J.M., BRADSHAW, C.J.A. & COURCHAMP, F., 2021: High and rising economic costs of biological invasions worldwide. *Nature* **592**, 571–576. <https://doi.org/10.1038/s41586-021-03405-6>
- DUKES, J. S. & MOONEY, H. A., 1999: Does global change increase the success of biological invaders? *Trends in Ecology and Evolution* **14**, 135–139. [https://doi.org/10.1016/S0169-5347\(98\)01554-7](https://doi.org/10.1016/S0169-5347(98)01554-7)
- EL MOUSSAOUI, A., EL AAMRI, F., CHIAAR, A., EL ASRI, F. & TAMSOURI, M.N., 2024: First record of the alien sponge *Paraleucilla magna* (Family Amphoriscidae) on the Moroccan Mediterranean coast. *Acta Adriatica* **65**, 223–228.
- EL HAMOUTI, C., CASTELLANO-HINOJOSA, A., MABROUKI, Y., CHAOUNI, B., GHAZAL, H., BOUKHATEM, N., CHAHBOUNE, R. & BEDMAR, E.J., 2023: Anthropogenic Nitrate Contamination Impacts Nitrous Oxide Emissions and Microbial Communities in the Marchica Lagoon (Morocco). *Sustainability* **15**, 4077. <https://doi.org/10.3390/su15054077>
- FRISTOE, T.S., CHYTRÝ, M., DAWSON, W., ESSL, F., HELENO, R., KREFT, H., MAUREL, N., PERGL, J., PYŠEK, P., SEEBENS, H., WEIGELT, P., VARGAS, P., YANG, Q., ATTORRE, F., BERGMEIER, E., BERNHARDT-RÖMERMANN, M., BIURRUN, I., BOCH, S., BONARI, G. & VAN KLEUNEN, M., 2021: Dimensions of invasiveness: Links between local abundance, geographic range size, and habitat breadth in Europe's alien and native floras. *Proceedings of the National Academy of Sciences of the United States of America* **118**(22), e2021173118. <https://doi.org/10.1073/pnas.2021173118>
- GEROVASILEIOU, V., AKEL, E.H.K., AKYOL, O., ALONGI, G., AZEVEDO, F., et al., 2017: New Mediterranean Biodiversity Records (July, 2017). *Mediterranean Marine Science* **18**(2), 355–384. <https://doi.org/10.12681/mms.13771>
- GOURARI, K., MABROUKI, Y., PARENTEAU-MAUFFETTE, É., LEGSSYER B., MANCINELLIV G. & TAYBI A.F., 2025: Assessing the Ecological and Socio-Economic Impacts of the Invasive Bleu Crab *Callinectes sapidus* Using Local Knowledge and Field Surveys in Eastern Morocco. *Thalassas* **41**, 118. <https://doi.org/10.1007/s41208-025-00876-6>
- GUZETTI, A., LONGO, C. & CORRIERO, G., 2019: Reproductive cycle of the invasive calcareous sponge *Paraleucilla magna* in the Mediterranean Sea. *Marine Ecology Progress Series* **609**, 113–123. <https://doi.org/10.3354/meps12819>
- KLAUTAU, M., MONTEIRO, L. & BOROJEVIC, R., 2004: First occurrence of the genus *Paraleucilla* (Calcarea, Porifera) in the Atlantic Ocean: *P. magna* sp. nov. *Zootaxa* **710**, 1–8.
- LANNA, E. & KLAUTAU, M., 2010: Life cycle of *Paraleucilla magna* Klautau, Monteiro, and Borojević, 2004 (Porifera: Calcarea). *Zoological Journal of the Linnean Society* **160**(1), 149–161. <https://doi.org/10.1111/j.1096-3642.2009.00598.x>

- LONGO, C., SCALERA -LIACI, L., MANUEL, M. & CORRIERO, G., 2004: Note sui poriferi del Mar Grande e del Mar Piccolo di Taranto (Mar Ionio). *Biologia Marina Mediterranea* **11**, 440–443.
- LONGO, C., MASTROTOTARO, F. & CORRIERO, G., 2007: Occurrence of *Paraleucilla magna* (Porifera: Calcareea) in the Mediterranean Sea. *Journal of the Marine Biological Association of the United Kingdom* **87**(6), 1749–1755. <https://doi.org/10.1017/S0025315407057748>
- LONGO, C., MASTROTOTARO, F. & CORRIERO, G., 2021: Ecological impact of the invasive calcareous sponge *Paraleucilla magna* in the Mediterranean Sea. *Biological Invasions* **23**(2), 457–472. <https://doi.org/10.1007/s10530-020-02388-9>
- MABROUKI, Y. & TAYBI, A.F., 2024: Range expansion of the alien polychaete *Branchiomma luctuosum* (Grube, 1870) (Sabellidae) in Morocco (the southwestern Mediterranean Sea). *Natura Croatica* **33**(1), 197–202. <https://doi.org/10.20302/NC.2024.1.17>
- MABROUKI, Y., CROCETTA F. & TAYBI, A.F. 2021: First Record of *Branchiomma luctuosum* (Grube, 1870) (Annelida: Polychaeta: Sabellidae) in Morocco. *Thalassas* **37**(5), 717–720. <https://doi.org/10.1007/S41208-021-00348-7>
- MABROUKI, Y., GOURARI, K., PARENTEAU-MAUFFETTE, É., SBAA, M., BOUZROUK, A. & TAYBI, A.F., 2025a: The Atlantic blue crab *Callinectes sapidus* (Rathbun, 1896) extends its southbound invasion along the Atlantic Coast of North Africa: first occurrence in the Atlantic Sahara shores of Morocco. *Thalassas* **41**(1), 12 <https://doi.org/10.1007/s41208-024-00767-2>
- MABROUKI, Y., GOURARI, K. & TAYBI F.A., 2025b: The Atlantic blue crab *Callinectes sapidus* Rathbun, 1896 (Portunidae, Malacostraca) could lead to the local extinction of endemic freshwater species in eastern Morocco. *Povolzhskii Ekologicheskii Zhurnal* **1**, 91–99. <https://doi.org/10.35885/1684-7318-2025-1-91-99>
- MAČIĆ, V. & PETOVIĆ, S., 2017: New data on the distribution of the alien sponge *Paraleucilla magna* Klautau, Monteiro & Borojević, 2004 in the Adriatic Sea. *Studia Marina* **29** (1), 63–68.
- OJAVEER, H., GALIL, B.S., MINCHIN, D., OLENIN, S., AMORIM, A., CANNING-CLODE, J. *et al.*, 2013: Ten recommendations for advancing the assessment and management of non-indigenous species in marine ecosystems. *Marine Policy* **44**, 1–6. <http://dx.doi.org/10.1016/j.marpol.2013.08.019>.
- ROBLES, R., 2010: Conservation and sustainable development of the Sea of Alborán. Gland, Switzerland and Malaga, Spain: IUCN. 83pp.
- SGHAIER, Y.R., ZAKHAMA-SRAIEB, R. & CHARFI, F., 2019: An inventory of non-indigenous species (NIS) inside and outside three tourist marinas from the southern Mediterranean coast outside three tourist marinas from the southern. *Journal of the Black Sea / Mediterranean Environment* **5**(1), 29–48.
- STREFTARIS, N., ZENETOS, A. & PAPATHANASSIOU, E., 2005: Globalisation in marine ecosystems: the story of non-indigenous marine species across European seas. *Oceanography and Marine Biology: An Annual Review* **43**, 419–453. <https://doi.org/10.1201/9781420037449.ch8>
- TAYBI, A.F. & MABROUKI, Y., 2020: The American blue crab *Callinectes sapidus* Rathbun, 1896 (Crustacea: Decapoda: Portunidae) is rapidly expanding through the Mediterranean coast of Morocco. *Thalassas* **36**(2), 267–271. <https://doi.org/10.1007/s41208-020-00204-0>
- TAYBI, A.F. MABROUKI, Y. & DOADRIO, I., 2020a: The occurrence, distribution and biology of invasive fish species in fresh and brackish water bodies of NE Morocco. *Arxius de Miscellania Zoologica* **18**, 59–73. <https://doi.org/10.32800/amz.2020.18.0059>
- TAYBI, A.F., MABROUKI, Y., CHAVANON, G. & MILLÁN, A., 2020b: The alien boatman *Trichocorixa verticalis verticalis* (Hemiptera: Corixidae) expanding in Morocco. *Limnetica* **39**(1), 49–59. <https://doi.org/10.23818/limn.39.04>
- TAYBI, A.F., MABROUKI Y. & PISCART, C., 2023: Distribution of Freshwater Alien Animal Species in Morocco: Current Knowledge and Management Issues. *Diversity* **15**(2), 169. <https://doi.org/10.3390/d15020169>
- TAYBI, A.F., MABROUKI, Y., GLÖER, P. & PISCART, C., 2024: Factors Explaining the Distribution of *Phylla acuta* (Draparnaud, 1805) in Freshwaters of Morocco. *Water* **16**, 803. <https://doi.org/10.3390/w16060803>
- TAYBI, F.A., LORENTI, M. & MABROUKI, Y., 2025: First Record of the Asiatic Isopod *Paranthurus japonica* Richardson, 1909 (Isopoda: Anthuroidea: Paranthuridae) in Morocco (Southern Mediterranean Sea). *Thalassas* **41**(2), 67 <https://doi.org/10.1007/s41208-025-00823-5>