Psychiatria Danubina, 2026 Vol. 38, No. 2, pp 111-135 https://doi.org/10.24869/psyd.2026.111 *Review Article*
© Medicinska naklada – Zagreb, Croatia

Cathexis and catharsis: An integrated neuropsychoanalytic model of serotonergic action

Arif de Mendelssohn[1] & Henriette Löffler-Stastka[2]

[1]Praxis Annagasse, Psychiatrische Kassenarztpraxis, Vienna, Austria
[2]Department of Psychoanalysis and Psychotherapy, Medical University Vienna, Vienna, Austria

received: 25 August 2025; revised: 23 February 2026; accepted: 20 May 2026

Summary

Background: *The history of mental science is marked by a divide between the subjective, energetic models of psychoanalysis and the objective models of computational neuroscience. Reductionist approaches often fail to bridge this gap or adequately explain the structural complexity of psychopathology. This conceptual synthesis aims to propose a unified, testable neurocomputational framework anchored in the Free Energy Principle (FEP).*

Subjects and Methods: *We conduct a theoretical synthesis mapping classical psychodynamic constructs onto the computational dynamics of Active Inference and neuromodulatory neurobiology. We formalize Freud's "Wish" as a high-precision allostatic prior, and drive as prediction error that compels the system to perform work. The Extended Reticular Thalamic Activating System (ERTAS) is analyzed as the physical operator of this economy.*

Results: *Catharsis (discharge/primary process) is mapped as the efficient minimization of free energy via established habits, mediated by routing dopaminergic drive into the dorsal striatum and stabilized by 5-HT1A signaling. Cathexis (binding/secondary process) involves inhibiting immediate discharge for generative model revision, triggered by the lateral habenula and mediated by 5-HT2A signaling. Consolidation occurs via the Gs pathway. We detail a serotonergic "precision switch" acting as a meta-regulator of the Ego, arbitrating between stability, reactivity, and growth.*

Conclusions: *The mind-brain functions as a hierarchical generative model striving to minimize entropy. This integrated model reframes psychopathology (e.g., depression, mania, psychosis) as "computational pathologies" of precision weighting where the switch between binding and discharge becomes stuck. This provides a rational basis for precision psychiatry, integrating pharmacological modulation of gain control with psychotherapeutic restructuring.*

Keywords: *Neuropsychoanalysis, Serotonin, Dopamine, Cathexis, Active Inference.*

* * * * *

THE ECONOMY OF THE MIND--BRAIN AND THE FAILURE OF REDUCTIONISM

The search for a unified model of the mind-brain

The history of mental science is characterized by a "Great Divide" between the subjective, energetic models of psychoanalysis and the objective, mechanistic descriptions of neuroscience. While psychoanalysis provides a rich phenomenology of motivation and internal conflict, its historical isolation from biology has limited its falsifiability. Conversely, neuroscience offers molecular precision but often descends into a reductionism that ignores the subjective agent. Reconciling these domains requires bridging experiential dynamics with biological mechanisms.

This article is a theoretical synthesis and hypothesis-generating paper: It integrates psychoanalytic concepts, the Free Energy Principle, and neuromodulatory pharmacology to propose a mechanistic account of cathexis and catharsis.

This work is not a systematic review: Rather, it selectively synthesizes molecular, systems-level, and clinical evidence to construct a coherent neurocomputational model. Where empirical foundations are robust, we cite meta-analytic or convergent evidence; where the synthesis relies on speculative inference, this is explicitly indicated and accompanied by testable predictions.

Throughout the text, we employ metaphorical terms as heuristic labels, in the spirit of Sigmund Freud's own hydraulic metaphors. These metaphors are not intended as explanatory substitutes but as conceptual guides; the underlying mechanisms are specified in biochemical and computational terms in the relevant sections.

Contemporary conceptual tools now enable a rigorous bio-psycho-social integration (Löffler-Stastka, 2023). We utilize **Neuropsychoanalysis**, which grounds psychodynamics in affective neuroscience (Panksepp, 1998; Solms, 2021) and the **Free Energy Principle (FEP)** (Friston, 2010), which views the brain as a predictive engine striving for homeostasis. This paper argues that Freud's concepts of **Cathexis** and **Catharsis** can be formally reinterpreted within Active Inference as operations of precision control over prediction error. We map these dynamics onto hierarchically organized neuromodulatory systems—specifically serotonergic and dopaminergic signaling acting through Gi, Gq, and Gs pathways—that regulate the fundamental stability-plasticity dilemma.

Addressing the brain-mind relation: a stance of dual-aspect monism

To navigate the "hard problem of consciousness" (Chalmers, 1996), we adopt the stance of **dual-aspect monism** (Solms, 2021), grounded in the spatiotemporal dynamics that bridge neural and mental states (Northoff, 2016). This perspective posits that mental and neural processes are subjective and objective aspects of a single reality, necessitating a **methodological dualism** that integrates the phenomenological language of psychoanalysis with the mechanistic metrics of neuroscience. We focus on the **neuromodulatory (precision) level** as the optimal organizational plane for this synthesis. This level reflects the **hierarchical dynamics of neuromodulation** (Deco & Kringelbach, 2020), connecting molecular receptor dynamics to behavioral phenomenology without reducing mental states to mere molecular events.

The failure of reductionism: beyond the "chemical imbalance"

Historically, reductive paradigms like the "chemical imbalance" theory (Coppen, 1967) have proven inadequate. Comprehensive reviews refute a consistent link between simple monoamine levels and depression (Delgado, 2000; Moncrieff et al., 2023), and such theories fail to account for the complex architecture of psychopathology. Modern systems-level views posit that neuromodulators do not create affect directly but act as **Gain Control** parameters (Stahl, 2013) within non-linear networks (Qi et al., 2013). Accordingly, we conceptualize monoamines as the **Extended Reticular–Thalamic Activating System (ERTAS; the integrated neuromodulatory control network)**—the effector arm of the Midbrain Self-System (Merker, 2007; Panksepp, 1998)— which acts as a **gain control mechanism** (Balasubramani et al., 2017; Stahl, 2013) to regulate the **precision** of cortical priors and prediction errors (Friston et al., 2012; Parr & Friston, 2017). This framework replaces the "chemical imbalance" with **"Computational Pathology,"** where the switch governing the transition between Cathexis (learning) and Catharsis (habit) becomes rigid.

Table 1: The Neuro-Computational Rosetta Stone

Psychoanalytic Concept	Computational Definition (Active Inference)	Neuroanatomical Substrate (Mechanism)
Drive Source *(Quelle)*	**Interoceptive Deviation:** The somatic error signal (e.g., hypoglycemia).	**PAG & Hypothalamus:** The „Vital Core" generating the raw, objectless affect (Need).
The Wish *(Wunsch)*	**The Fundamental Coupling:** The binding of Drive (Affect) to Object (Memory-Image). Acts as the **Allostatic Prior**.	**Basolateral Amygdala (BLA) & Ventral Striatum:** Binds Value (PAG) to Context (Hippocampus) to assign **Incentive Salience**.
The Drive *(Trieb)*	**Prediction Error Vector (Q):** A vector with **Magnitude** (Pressure) and **Direction** (Aim) demanding resolution.	**Ascending Arousal (ERTAS):** The energetic pressure arising from the discrepancy between Wish and Reality.
Cathexis *(Besetzung)*	**Minimizing Expected Free Energy (EFE):** Binding error to simulate future outcomes (Secondary Process).	**Layer V Pyramidal Neurons:** Sustained depolarization (Virtual Work) to search for a policy.
Catharsis *(Abfuhr)*	**Minimizing Variational Free Energy (VFE):** Immediate discharge via habit (Primary Process).	**Dorsal Striatum:** Efficient execution via the path of least resistance.
The Ego *(Ich)*	**Hierarchical Control System:** The Generator of Allostatic Priors (Self-Model) and the Arbitrator of shunting Drive to Habit.	**Systemic: DMN** (Seat) & **vmPFC** (Hand) steering the **Striatum** (Engine). **Cellular: Layer V Pyramidal Neuron** (Ca^{2+} Coupling of Prior and Error).

Defining the explanandum: cathexis and catharsis

To build an integrated model, we must first refine the Freudian economic concepts of Cathexis and Catharsis (Freud, 1950) through the lens of the **Hierarchical Generative Model (HGM)**. Throughout this model, we differentiate between **Content** (glutamatergic predictions and errors) and **Context** (neuromodulatory precision signals).

- **Cathexis** (translating *Besetzung*, meaning "investment"): The investment of psychic energy to "bind" a mental representation. In our framework, we operationalize **Active Cathexis** as the precision-weighted maintenance of prediction error to drive model revision. This state requires the inhibition of immediate discharge.
- **Catharsis** (*Abfuhr*, meaning "discharge"): The release of tension. Here, we operationalize **Passive Catharsis** as the efficient minimization of free energy through established, high-precision action policies. This corresponds to the **"path of least resistance"** (i.e., the model-free habitual policy).

Freud's original definitions: the hydraulic model

In his seminal *Project for a Scientific Psychology* (Freud, 1950) and later in *The Unconscious* (Freud, 1957b), Freud outlined these dynamics using a hydraulic metaphor. The **Primary Process (Catharsis)** is the default mode of the "Id," operating under the "Nirvana Principle" to immediately discharge excitation via the most direct motor pathway. The **Secondary Process (Cathexis)** is the mode of the "Ego," arising because immediate discharge is often maladaptive. To survive, the Ego must inhibit discharge, "binding" the energy to facilitate reality testing and thought (Freud, 1958).

The computational refinement: from hydraulics to information

While Freud's structural insights were profound, his hydraulic conceptualization is incompatible with the modern *thermodynamics* of a closed fluid system. However, the *topology of flow* (cascades and gating) remains a valid computational heuristic to describe cognitive processes, e.g. in the striatum. We therefore aim to refine Freudian Terminology within the FEP (Table 1), formalizing "Psychic Energy" (Q) as **Variational Free Energy** (prediction error) (Carhart-Harris & Friston, 2010).

The Full List of Abbreviations

5-HT	5-Hydroxytryptamine (Serotonin)
ACh	Acetylcholine
ATP	Adenosine Triphosphate
BDNF	Brain-Derived Neurotrophic Factor
BLA	Basolateral Amygdala
cAMP	Cyclic Adenosine Monophosphate
CNS	Central Nervous System
CREB	cAMP Response Element-Binding Protein
CSTC	Cortico-Striato-Thalamo-Cortical
DA	Dopamine
dACC	Dorsal Anterior Cingulate Cortex
DMN	Default Mode Network
EFE	Expected Free Energy
ERTAS	Extended Reticular Thalamic Activating System
FEP	Free Energy Principle
GIRK	G-protein-coupled Inwardly Rectifying Potassium
GPCRs	G-protein Coupled Receptors
HA	Histamine
HGM	Hierarchical Generative Model
LHb	Lateral Habenula
LSD	Lysergic Acid Diethylamide
LTP	Long-Term Potentiation
MDD	Major Depressive Disorder
mGluR2	Metabotropic Glutamate Receptor 2
mPFC	Medial Prefrontal Cortex
NA	Noradrenaline
NaSSA	Noradrenergic and Specific Serotonergic Antidepressant
OCD	Obsessive-Compulsive Disorder
OFC	Orbitofrontal Cortex
PAG	Periaqueductal Gray
PCC	Posterior Cingulate Cortex
PFC	Prefrontal Cortex
PKA	Protein Kinase A
PKC	Protein Kinase C
PLC	Phospholipase C
Q	Psychic Energy / Prediction Error Vector
REM	Rapid Eye Movement
RL	Reinforcement Learning
RMTg	Rostromedial Tegmental Nucleus
RNA	Ribonucleic Acid
SSRI	Selective Serotonin Reuptake Inhibitor
SWS	Slow-Wave Sleep
VFE	Variational Free Energy
vmPFC	Ventromedial Prefrontal Cortex
VTA	Ventral Tegmental Area
VTE	Vicarious Trial and Error

Difference 1: The "Wish" as a Structural Coupling vs. "Drive" as a Vector

Freud described the "**Wish**" as follows: *"An impulse of this kind is what we call a wish; the reappearance of the perception is the fulfillment of the wish, and the full cathexis of the perception, by the excitation springing from the need, constitutes the shortest path to the wish-fulfillment."* (Freud, 1953).

"Drive" (*Trieb*) is defined in *Instincts and their Vicissitudes* (Freud, 1957a) as the "measure of the demand for work" imposed upon the mind—thereby distinguishing the raw energetic pressure from the representational content of the Wish that guides its resolution.

To strictly satisfy Freud's definitions, we must distinguish between the structural representation (The Wish) and the quantitative pressure (The Drive).

- **The Wish (The Map):** The Wish is the **coupling** of a *Drive Trace* (affect from the PAG/Hypothalamus) with a *Memory-Image* (representation from the Hippocampus/Cortex). Anatomically, this intersection occurs in the **Basolateral Amygdala (BLA)**. This coupled state forms the **Allostatic Prior**—the top-down command sent to the cortex (e.g., "I am satisfied by this specific object").
- **The Drive (The Force):** The Drive is the **Prediction Error** (Q) generated when reality deviates from this Wish. Computationally, it acts as a **Vector**:
- **Magnitude (Pressure):** The intensity of the error (The "Demand for Work").
- Direction (Object): The coordinate pointing from the Current State toward the Wish.

Thus, the Wish provides the destination, and the Drive provides the force required to traverse the distance.

Difference 2: Cathexis as "Exploration" (The Binding of Drive).

We reframe Cathexis as the interruption of the drive's discharge. When the habit fails to minimize error, the drive energy (Dopamine) is "bound" or held in the cortex. This validates Freud's (1955) insight that "consciousness arises instead of a memory trace." When the "Zombie" agent (automatic model-free controller) hits a wall, the error cannot be discharged and must be "bound" in the cortex (Working Memory). This state is metabolically expensive (thermodynamic cost detailed in Section 3.5).

Difference 3: Catharsis as "Exploitation" (The Habit).

We redefine Catharsis as the successful channeling of Drive energy into a high-precision Action Policy. It is not merely the presence of Dopamine, but the specific routing of Mesolimbic drive (Ventral Striatum) into Nigrostriatal habit (Dorsal Striatum). This minimizes free energy efficiently, maintaining the organism in a low-energy attractor state.

We can integrate the energetic dynamics as the mathematical distinction between the Primary and Secondary processes:

- **Variational Free Energy (VFE) = The "Now" (Catharsis):** The measure of error in the current moment. It is the immediate **Drive Pressure**. If a habit exists to resolve this, VFE is minimized via Catharsis (Reactive Action).
- **Expected Free Energy (EFE) = The "Future" (Cathexis):** The predicted error associated with future policies. When habit fails, the brain cannot just react; it must minimize EFE. This requires **Cathexis** (Thinking)—binding the current Drive to simulate future outcomes via the Hippocampus and PFC to find a path to the Wish.

Clinical box A: The "autopilot" vs. The "detour"

To ground these computational dynamics in everyday experience, consider the act of driving a familiar route to work:

Catharsis (The Habit/Primary Process): When the route is clear, the driver navigates on "autopilot." The Generative Model perfectly predicts the environment (prediction error is near zero), and the biological drive to reach the destination flows seamlessly into the sensorimotor habit (Dorsal Striatum). The driver is in a state of efficient, low-energy *Passive Catharsis* and may arrive at work with no conscious memory of the journey.

Cathexis (The Binding/Secondary Process): Suddenly, the driver encounters a "Road Closed" sign. This generates a massive prediction error. The Lateral Habenula triggers the "brake," halting the automatic driving habit. The drive cannot be discharged. The error is shunted upward, initiating *Active Cathexis*. The driver grips the wheel, feels a surge of subjective tension (the metabolic cost of binding the error), and actively recruits Working Memory (Prefrontal Cortex/Hippocampus) to simulate potential detours. Immediate discharge is blocked to force a conscious revision of the model.

THE COMPUTATIONAL ARCHITECTURE—THE HIERARCHY OF INFERENCE

The central nervous system (CNS) functions as a deeply nested **Hierarchical Generative Model (HGM)** designed to minimize entropy (Friston, 2010).

The nested hierarchy: from core affect to cortical cognition

The HGM is organized evolutionarily:

- **Innermost Layer (Conscious Id):** Located in the upper brainstem centered on the Periaquaeductal Gray (PAG) and "Extended Reticular–Thalamic Activating System" (ERTAS) We here use this term as a conceptual aggregation rather than as a single anatomically unified system, referring collectively to heterogeneous ascending brainstem, hypothalamic, thalamic, and basal forebrain pathways that regulate cortical arousal, gain, and precision. This layer processes interoceptive signals and generates **Core Affect** (Panksepp, 1998; Solms, 2021).
- **Middle Layers (Limbic/Paralimbic):** Includes the amygdala, hippocampus, habenula, and cingulate. This level integrates affect with memory to determine **Salience**.
- **Outer Layers (Cortical Ego):** The neocortex, responsible for detailed modeling and executive control (Carhart-Harris & Friston, 2010).

The anatomical core and the "conscious Id"

Anatomically, the "Id" centers on the Midbrain Decision Triangle (Merker, 2007) and the SELF (Panksepp, 1998). This region performs the brain's fundamental calculation: Target Selection through integrating the current state of the body (PAG) with the environment (Superior Colliculi).

The ERTAS functions as the Effector Arm of this system (Alcaro et al., 2017). Upon detecting salient error (Drive), the ERTAS broadcasts neuromodulatory "Gain Control" signals to the cortex, commanding the Ego to "bind" attention to the problem.

Function: the "wish" and the generation of Q

We map Freud's tripartite definition directly onto specific neuroanatomical substrates:

1. **The Memory-Trace of Drive-Excitation ("The Need"):**
   - *Neuroanatomy:* **PAG & Hypothalamus**.
   - *Function:* These structures generate the raw, objectless affect (e.g., hunger, separation distress). This is the source of the "Demand for Work" (Q).
2. **The Memory-Image of Perception ("The Object"):**
   - *Neuroanatomy:* **Hippocampus & Sensory Association Cortices**.
   - *Function:* The Hippocampus stores the episodic context ("The Kitchen," "The Breast"), while sensory cortices store the specific perceptual modality. This provides the *content* of the solution.
3. **The "Fundamental Coupling" ("The Wish"):**
   - *Neuroanatomy:* **Basolateral Amygdala (BLA) & Ventral Striatum**.
   - *Function:* The BLA receives the Drive signal from the Hypothalamus and the Memory-Image from the Hippocampus. It **binds** them, assigning Valence to the memory. The Ventral Striatum then translates this valence into **Incentive Salience** ("Wanting"; Berridge & Robinson, 1998).

We map Freud's canonical components of the Drive (1957a) to the computational anatomy:

Note on the Ego's Hierarchy: While the BLA forms the immediate Affective Coupling (Need + Object), this signal is uploaded to the Default Mode Network (DMN). The DMN (The Ego) integrates this coupling into the persistent "Self-Model" and broadcasts it as the Top-Down

Table 2: Freud's Four Components of the Drive

Freudian Component	Definition	Computational Definition	Neuroanatomy
Source *(Quelle)*	Somatic process in an organ.	**Interoceptive Deviation:** Departure from homeostasis.	**PAG & Hypothalamus** (The Vital Core).
Pressure *(Drang)*	The „Demand for Work."	**Prediction Error (Q):** The magnitude of the Vector.	**Ascending Arousal** (ERTAS).
Object *(Objekt)*	The thing through which the aim is achieved.	**The Wish:** The coupled Memory-Image.	**Hippocampus / BLA.**
Aim *(Ziel)*	Satisfaction (Removal of Source).	**Free Energy Minimization:** Collapse of error ($Q \to 0$).	**Action / Update** (Striatum).

Allostatic Prior to the Layer V neurons. Thus, the BLA generates the Wish, but the Ego administers it.

The middle level: the "anti-reward" brake

The most critical regulatory mechanism at the limbic level—and the physical locus of the switch between Catharsis and Cathexis—is the **Lateral Habenula (LHb)**.

The "anti-reward" brake: the lateral habenula and RMTg

The LHb functions as the brain's "Anti-Reward" center (specifically the **LHb [8]$\rightarrow$ RMTg $\rightarrow$ VTA/Raphe inhibitory pathway** that signals negative prediction error).

Note: Although the lateral habenula plays a critical role in signaling negative prediction error and suppressing dopaminergic policy execution, it functions as a convergence and veto node within a distributed control architecture that also includes direct prefrontal–midbrain pathways and thalamocortical gating mechanisms. In parallel, prefrontal projections to midbrain dopaminergic nuclei permit top-down modulation of policy execution independently of habenular signaling, while thalamic nuclei provide additional gain control over cortical–striatal communication. The present model emphasizes the LHb because of its privileged access to aversive prediction error and its capacity to globally suppress dopaminergic output.

When the LHb detects that reality is "worse than predicted" (Frustration), it excites the **Rostromedial Tegmental Nucleus (RMTg)** (Hikosaka, 2010; Proulx et al., 2014). The RMTg is a GABAergic brake. When excited, it powerfully inhibits the Ventral Tegmental Area (VTA) and the Raphe Nuclei.

Computational Function: This blockade disables the mechanism of Catharsis (Habitual Discharge). By shutting down the Dopaminergic "Go" signal and reducing serotonergic stability signaling, the LHb/RMTg complex vetoes the "path of least resistance." This forces the system away from habitual Catharsis and the unprocessed prediction error to be shunted upward to the cortex, triggering Cathexis compelling the cortex to sustain error until a better model can be formed (binding).

The executive: the ego and the "waterfall" of action

Because Dopamine is implicated in both SEEKING/Exploration and habit/Exploitation we must distinguish between two distinct computational vectors based on anatomical pathways: **The Drive (Ventral)** vs. **The Discharge (Dorsal**) (Yin & Knowlton, 2006). We describe this as the 'Waterfall' dynamic, anatomically




Figure 1: The Precision Switch Network. The Lateral Habenula (LHb) acts as the Error Detector. Upon detecting negative prediction error, it activates the RMTg, which inhibits the VTA (Dopamine/Drive) and Raphe (Serotonin/Stability). This blockade of discharge forces the system into a high-energy binding state.

Table 3: The Neural Architecture (The Hardware)

Structure	Functional Role (Neuropsychology)	Computational Role (Active Inference)
PAG & Hypothalamus	**Affective Core** (The Conscious Id)	**Generator of Value:** Generates the Prior (Wish) and primary Prediction Error (Drive).
Midbrain Triangle	**Decision Generator** (Proto-Self)	**Action Selection:** Integrates Value and Sensory Data to initiate „Go“ signals.
ERTAS	**Precision Effector System**	**Gain Control:** Adjusts Synaptic Gain to enforce Stability or Plasticity.
Raphe Nuclei	**Precision Bias** (The Switch)	**Uncertainty Regulation:** Arbitrates between Exploitation and Exploration.
Lateral Habenula	**Error Detector** (The Brake)	**Inhibition:** Signals Negative Prediction Error. Blocks discharge to force *Binding*.
Ventral Striatum	**The Drive** (Incentive Salience)	**Input Vector (Q):** The „Demand for Work“ seeking a policy.
Dorsal Striatum	**The Automaton** (The Habit)	**Catharsis Engine:** Discharges energy via the „path of least resistance.“

corresponding to the **spiraling striato-nigro-striatal projections** described by Haber (2014), where information flows from the ventral (limbic) to the dorsal (motor) striatum.

Ventral striatum: the engine of SEEKING (the drive)

The **Ventral Striatum (Nucleus Accumbens)** receives the primary affective drive (Q) from the limbic system. It generates **Incentive Salience**—the urge to act or the “Precision of Opportunity” (Schultz, 1997). This is the input energy of the system.

The dorsal striatum and CATHARSIS (the habit)

The **Dorsal Striatum (Putamen/Caudate)** stores high-precision **Action Policies** (Habits).

- **Catharsis (The Flow):** If the Dorsal Striatum has a high-precision policy matching the context (“I know what to do”), the Drive energy (from the Ventral Striatum) flows seamlessly into the Habit (Dorsal). Action is executed, and energy is discharged.
- **Cathexis (The Dam):** If the policy fails, or if the “Anti-Reward” brake (LHb) is engaged, the transition from Ventral to Dorsal is blocked. The Drive energy cannot be discharged and is instead “bound” in the cortex. This is **Cathexis** (Bound SEEKING).

This account of catharsis as efficient discharge through habitual action is consistent with fronto–basal ganglia gating models, in which dopaminergic modulation biases Go/No-Go pathways to enable or suppress policy execution (Frank et al., 2004; Wiecki & Frank, 2013).

The anatomy of the Ego

The Ego is not localized to a single structure but instantiated across nested levels:

- cellular coincidence detection (Layer V pyramidal neurons),
- mesoscopic control (vmPFC/dACC) “The Hand”
- macroscopic self-modeling (DMN) “The Seat”

The Ego implements action selection. It can be defined as a Hierarchical Control System instantiated in the interaction between the Default Mode Network (DMN) and the Executive CSTC Loops.

The “seat” of the Ego: the self-model (DMN)

- **Anatomy: medial prefrontal cortex (mPFC), posterior cingulate (PCC), and precuneus** (The Default Mode Network).
- **Role:** This network encodes the **“Self-Model”** (The deepest High-Precision Priors). It maintains autobiographical continuity—the “I” that persists across time.
- **Function:** The DMN is the top of the hierarchy. It receives the “Wish” from the limbic system, integrates it into the self-narrative, and projects it downward as a stable **Allostatic Prior** to the sensory and motor cortices.

The “hand” of the ego: the controller of discharge

- **Anatomy: Ventromedial Prefrontal Cortex (vmPFC) / OFC** and **Dorsal Anterior Cingulate (dACC)**.
- **Role:** These structures act as the Interface between the Cortex and the Id.
- The **vmPFC/OFC** evaluates the *Value* of the Drive.

- The **dACC** monitors *Conflict* (Prediction Error) and recruits control.
- **The Mechanism:** Crucially, the mPFC sends glutamatergic projections to excite **GABAergic Interneurons** in the Amygdala and Ventral Striatum. This implements the active, top-down inhibition of the Id's immediate discharge to force Cathexis.

Steering the "Waterfall"

- **Anatomy:** The Spiraling **Cortico-Striatal Projections**.
- **Role:** The Ego "exists" in its capacity to act—specifically, to **steer the flow of Dopamine** from the **Ventral Striatum** (Drive/Exploration) to the **Dorsal Striatum** (Habit/Catharsis). The Ego uses the "Hand" (vmPFC) to open or close the gates of this waterfall.
- **Failure:** If the Cortex cannot steer this flow (e.g., in Mania), the Ego has failed, and the drive discharges blindly via the path of least resistance.

The biophysical ego: the layer v coincidence detector

The **Cortical Layer V Pyramidal Neuron** is described as the physical integrator of the HGM (Larkum, 2013). To be maximally precise, we must define the Ego at the cellular level, as this is where "Cathexis" physically occurs.

- **The Unit:** The **Layer V Pyramidal Neuron** in the Prefrontal Cortex can be conceptualized as the integrator that enforces the Wish.
- **The Mechanism:** The Ego is the **Integration** of:
  1. **The Apical Dendrites (Predictions/The Wish):** Receive the **"Coupled Wish"** (The high-precision prior) from the **DMN**. This prior is a synthesis of the Interoceptive Goal (BLA) and Exteroceptive Plan (Hippocampus/PFC) *(Top-Down Allostatic Prior)*.
  2. **The Basal Dendrite (The Reality):** Receives the **Bottom-Up Drive Vector** *(Prediction Error)* from the Thalamus/Id.

The Apical input instantiates the 'Wish' as a **Top-Down Allostatic Prior**. This prior essentially commands the lower hierarchies: 'Assume the state of being satisfied.' The mismatch between this Apical expectation and Basal reality generates the **Drive (Q)**—encoded as sustained somatic depolarization—which represents the metabolic cost of the discrepancy.




Figure 2: The Layer V Pyramidal Neuron. The Apical Tuft receives Top-Down Priors (Expectations) and is modulated by 5-HT2A (Plasticity). The Basal Dendrites receive Bottom-Up Prediction Error. Binding arises when apical excitability permits the maintenance of error in the apical-basal loop, preventing its resolution until the model is updated.

- **Definition:** The "Ego" is the **calcium-mediated coupling** (Backpropagation-activated Ca2+ spike firing) between these Apical and Basal compartments.
  0. Decoupling: When these compartments are decoupled (e.g., deep sleep, anesthesia), the Ego dissolves (loss of consciousness).
  1. Hyper-Coupling: When they are hyper-coupled (e.g., via 5-HT2A activation), the boundaries between the internal Wish and external Reality dissolve.

Function: "explaining away" and binding

The Ego's function is to minimize prediction errors via Active Inference.

- **Catharsis (Successful Inference):** If the Ego's model is accurate, the error is "explained away." The neuron remains stable, and the Id's energy is discharged efficiently via the CSTC motor loops (elaborated in Section 6.3.3).
- **Cathexis (Failed Inference/Binding):** If the model fails, the Bottom-Up error persists. The Ego must "bind" this error via sustained depolarization (thermodynamic cost detailed in Section 3.5). This high-energy state is the neural correlate of **Secondary Process Thinking** (Carhart-Harris & Friston, 2010).

The signaling architecture: content vs. Context

Throughout the model, we distinguish between **Content** (glutamatergic predictions and errors) and **Context** (neuromodulatory precision signals).

The substrate of content (the "what")

Content is mediated by **Glutamate** (excitation) and shaped by **GABA** (inhibition). These carry the specific informational content of predictions and errors.

Context: the neuromodulators (the "how much")

Mediated by the Neuromodulators (Monoamines/ACh). These regulate the **precision** (synaptic gain) of the content signals (Friston et al., 2017). By tuning precision, these molecules dictate the 'learning rate' and the balance between stability (exploitation) and plasticity (exploration) (Berger et al., 2009).

Crucially, we map this directly onto the biophysical models of serotonergic function described by Balasubramani et al. (2017). Where classical computational neuroscience describes serotonin as modulating the 'Signal-to-Noise Ratio' (SNR) via gain control, the Active Inference framework formalizes this as 'Precision Weighting.' These are not distinct operations but descriptions of the same mechanism at different levels of analysis: the biophysical modulation of gain (e.g., via 5-HT1A hyperpolarization or 5-HT2A depolarization) is the physical operator that adjusts the statistical confidence (precision) of the generative model.

The temporal hierarchy and development

The HGM operates across multiple timescales, from millisecond tactical processing (Glu/GABA) to strategic modulation (seconds/minutes via ERTAS). This architecture is constructed developmentally. Early caregiving shapes the baseline precision settings of the ERTAS (Schore, 2001). Adverse early life stress can dysregulate this system, encoding maladaptive priors (e.g., "internalizing a bad object") that function as rigid, pathogenic beliefs within the generative model (Fonagy et al., 2002).

MOLECULAR MECHANISMS — THE TOOLKIT OF STABILITY AND PLASTICITY

The computational imperatives of the HGM—balancing the efficiency of the Habit (Catharsis) against the necessity of Revision (Cathexis)—are physically implemented by the specific G-protein coupled receptors (GPCRs) engaged by the ERTAS. We identify a fundamental **Tripartite Logic** governing cellular computation, mapping the signal transduction pathways defined by Stahl (2013) onto the computational dynamics of precision: the three major G-protein families function as distinct 'Precision Operators'.

Gi/o signaling: the architecture of stability (catharsis)

The Stabilizer (Exploitation/Discharge): Receptors coupled to the **Gi/o family**—such as **5-HT1A** (postsynaptic), **D2** (dorsal striatum), and **α2-Adrenergic**—function as the physiological stabilizers of the generative model.

- **Mechanism:** Activation inhibits adenylyl cyclase (reducing cAMP) and, crucially, opens **GIRK channels**

(G-protein-coupled Inwardly Rectifying Potassium channels). This causes potassium efflux and rapid neuronal **hyperpolarization**.

- **Computational Function:** Hyperpolarization dampens neural noise and increases the threshold for firing. In Bayesian terms, this **increases the precision of the current Prior**. It "clamps" the system in a stable attractor, protecting the Generative Model from bottom-up perturbations. This is the biophysical basis of resilience and facilitates the noise-free execution of habits (Passive Catharsis) (Barnes & Sharp, 1999).

Gq/11 signaling: the architecture of reactivity (cathexis)

The Accelerator (Exploration/Binding): Receptors coupled to the **Gq/11 family** (e.g., **5-HT2A**, **5-HT2C**, **$\ alpha$1-Adrenergic**) function as precision modulators that increase systemic reactivity.

- **Mechanism:** Activation triggers Phospholipase C (PLC), mobilizing intracellular **Calcium (Ca^{2+})** and activating Protein Kinase C (PKC). This generally increases neuronal excitability.
- The Contextual Divergence (The Accelerator vs. The Brake):

Crucially, while Gq/11 signaling always "accelerates" cellular activity, the systemic outcome depends entirely on the target neuron. This resolves the apparent paradox where Gq signaling can drive both cognitive flexibility and behavioral inhibition:

1. **Cortical Expansion (5-HT2A):** Located on the apical dendrites of excitatory **Pyramidal Neurons**. Activation increases the gain on bottom-up sensory prediction errors, allowing them to override top-down priors. This "opens" the model for revision (Active Cathexis) (Carhart-Harris & Friston, 2019).
2. **Subcortical Braking (5-HT2C):** Located primarily on **GABAergic Interneurons** in the VTA and Striatum. Activation "accelerates" these inhibitory cells, which in turn suppress the dopaminergic drive. This "brakes" the motor output to prevent premature discharge (Di Giovanni et al., 2008).

- **Computational Function:** This synergistic push-pull mechanism defines **Active Cathexis**. The 5-HT2A receptor opens the "lens" of the mind (to process error), while the 5-HT2C receptor, acting though GABAergic Interneuron, functionally clamps the "motor" (to prevent discharge), forcing the system to bind the energy in the cortex.

Gs signaling: the architecture of growth (consolidation)

The Architect (Rewiring): Receptors coupled to the **Gs family**—specifically **5-HT4, 5-HT6, 5-HT7**, and **D1**—function as the structural architects of the HGM.

- **Computational Function:** This pathway is engaged when a prediction error is deemed sufficiently precise to warrant permanent encoding. It represents the "Save" function of the brain. While Gi/o and Gq manage the *dynamics* of the switch (seconds to minutes), Gs manages the *structure* of the model (hours to days), consolidating the transient solution of Cathexis into a new, efficient Habit (Catharsis) (Duman et al., 2016).
- **Mechanism:** Activation stimulates adenylyl cyclase, driving a robust increase in cAMP and PKA. Crucially, this initiates the **CREB** transcriptional cascade, leading to the synthesis of **BDNF** (Brain-Derived Neurotrophic Factor) and synaptic remodeling (LTP). This upregulation of neurotrophic factors is the critical step that translates transient signaling into permanent structural changes, effectively reversing neuronal atrophy .
- **Computational Function:** This pathway is engaged when a prediction error is deemed sufficiently precise to warrant permanent encoding. It represents the "Save" function of the brain. While Gi/o and Gq manage the *dynamics* of the switch (seconds to minutes), Gs manages the *structure* of the model (hours to days), consolidating the transient solution of Cathexis into a new, efficient Habit (Catharsis) (Duman et al., 2016).

Hierarchical dynamics of neuromodulation

Precision control is inherently hierarchical. Subcortical neuromodulatory nuclei regulate large-scale cortical gain, while cortical pyramidal neurons implement local binding dynamics (Deco & Kringelbach, 2020). This hierarchical neuromodulation ensures that Cathexis can occur at multiple levels—from localized perceptual uncertainty to global existential conflict—without collapsing the entire system into paralysis or noise.

Cathexis, catharsis, consolidation and energy.

Cathexis is metabolically costly. Sustained firing of deep-layer pyramidal neurons requires continuous ATP expenditure and calcium regulation (Attwell & Laughlin, 2001).

- **The Cost of Cathexis:** Binding energy (thinking) is metabolically expensive. Sustained firing of deep-layer pyramidal neurons requires continuous ATP expenditure and calcium regulation (Attwell & Laughlin, 2001). An organism cannot exist in perpetual Cathexis; it would succumb to metabolic exhaustion.
- **The Danger of Catharsis:** Discharging energy (habit) is metabolically cheap but rigid. An organism that never binds energy cannot learn. It remains trapped in the "automaton" state, vulnerable to environmental volatility.
- **The Optimum:** Mental health is the fluid oscillation between these states. We bind energy (Cathexis) only long enough to update the model, then consolidate that update into a new habit (Catharsis) to return to efficiency.

This sustained energetic burden is consistent with allostatic models, which emphasize the metabolic costs of maintaining stability through prolonged activation rather than through efficient discharge (McEwen, 2007; McEwen & Wingfield, 2003; Sterling & Eyer, 1988).

PRECISION MODULATION AND THE DIVISION OF ROLES WITHIN THE ERTAS

While the G-proteins provide the logic gates (Stability/Gi, Reactivity/Gq, Growth/Gs), the neuromodulators themselves—specifically **Dopamine** and **Serotonin**—orchestrate the timing and allocation of these resources. For conceptual clarity, it is imperative to differentiate between the **Source of Drive** (SEEKING) and the **Mechanism of Discharge** (Habit).

Specialized roles: modulating "setting" and "embodiment"

- **Histamine (HA):** Modulates "Embodied Precision," regulating the trade-off between interoceptive maintenance and exteroceptive arousal (Haas et al., 2008).
- **Noradrenaline (NA) & Acetylcholine (ACh):** Modulate the "Setting." NA signals **Unexpected Uncertainty** (Volatility), triggering a global reset (phasic) or vigilance (tonic). ACh signals **Expected Uncertainty**, enhancing the signal-to-noise ratio of specific sensory cues (Attentional Spotlight) (Yu & Dayan, 2005).

Dopamine (DA): the dual vectors of drive and discharge

Dopaminergic function is best described via a Dual-System Model based on the "Waterfall" dynamic of the striatum, distinguishing between the input energy **(Drive)** and the execution channel **(Discharge).**

Mesolimbic dopamine: precision of opportunity (the drive)

- **Anatomy:** Projections from the Ventral Tegmental Area (VTA) to the Ventral Striatum (Nucleus Accumbens).
- **Function (Q):** This system encodes **Incentive Salience** (Berridge & Robinson, 1998)—the "Precision of Opportunity." Computationally, this state is driven by **Reward Prediction Error** (Schultz, 1997), which signals that the environment contains a high-value means to reduce free energy. In our framework, this constitutes the **Drive Pressure (SEEKING;** Panksepp, 1998**):** a high-energy, exploratory state that acts as the input vector for the Ego.

Nigrostriatal dopamine: precision of execution (the habit)

- **Anatomy:** Projections from the Substantia Nigra to the **Dorsal Striatum (Putamen/Caudate)**.
- **Function (Catharsis):** This system encodes **Motor Confidence** or "Precision on Policy." It facilitates the automated execution of movement.
- **The Waterfall Dynamics:** In a healthy HGM, the energy from the Drive (Ventral) cascades into the Habit (Dorsal) to achieve **Catharsis** (Action). If this transfer is blocked (e.g., by the LHb/5-HT2C brake), the Drive energy is "shunted" to the cortex, becoming **Cathexis** (Bound Seeking).

Serotonin (5-HT): the meta-regulator of the "mindset"

While Dopamine provides the Drive (Quantity/Q), Serotonin regulates the **Global Gain** (Quality/Precision) applied to that drive. It functions as the "Meta-Precision Controller" (Deco & Kringelbach, 2020), determining *how* the system responds to the Drive:

1. **Stability Mode (5-HT1A/Gi):** "Ignore the error, trust the prior." Promotes passive Catharsis and resilience.
2. **Plasticity Mode (5-HT2A/Gq):** "Bind the error, revise the prior." Promotes Active Cathexis and learning.

3. **Inhibition Mode (5-HT2C/Gq):** "Stop the action, hold the drive." Promotes the necessary pause required for computation.

By arbitrating the interaction between the **Mesolimbic Drive** and the **Cortical Model**, Serotonin ultimately governs the stability-plasticity equilibrium of the Ego (Cools et al., 2011).

ARBITRATING CATHARSIS AND CATHEXIS: THE "PRECISION SWITCH"

Central to our model is the understanding that Serotonin (5-HT) acts as a **Meta-Precision Controller** (Deco & Kringelbach, 2020). By shifting the balance of G-protein signaling—**Gi/o (Stability)**, **Gq/11 (Reactivity)**, and **Gs (Growth)**—serotonin dictates the "Epistemic Mode" of the organism. We term this regulatory mechanism the **"Precision Switch."** It determines whether the **Mesolimbic Drive (Q)** is allowed to cascade into habitual discharge (Catharsis) or is intercepted for cortical revision (Cathexis).

The generator: dynamics of raphe secretory activity

The switch originates in the **Dorsal and Median Raphe Nuclei**, which function as the brain's central "uncertainty-ometer," where firing rates reflect the real-time estimation of the precision of high-level priors (Grossman et al., 2022).

Tonic firing: the signal of contextual stability (catharsis)

The baseline operation of the Raphe is **Tonic Firing** (0.5–3 Hz).

- **Computational Function:** This steady-state release maintains a "blanket" of Contextual Precision. It stabilizes global predictions, suppresses neural noise, and entrenches established schemata.
- **Pathway Modulation:** Tonic firing preferentially engages high-affinity receptors (e.g., **5-HT1A**) in the cortex and **5-HT1B** in the basal ganglia. This supports habitual, schema-driven behavior (**Passive Catharsis**), allowing the organism to operate on "autopilot" with high confidence in its priors.
- **Feedback Control:** This stability is strictly regulated by 5-HT1A autoreceptors (Gi/o) on the soma and dendrites of Raphe neurons (Barnes & Sharp, 1999). These receptors are activated by serotonin released from **local axon collaterals** and, primarily, via **somatodendritic volume transmission**, where serotonin diffuses through the extracellular space to influence neighboring neurons (Agnati et al., 1995). Upon activation, they drive potassium efflux to hyperpolarize the Raphe neuron, providing homeostatic negative feedback that curbs excessive release (Sprouse & Aghajanian, 1987). This autoregulatory loop acts as the "thermostat" of the Ego.

Phasic bursting: the signal of context updating (cathexis)

The Raphe shifts into **Phasic Bursting** (10–20 Hz) in response to "Unexpected Uncertainty" or high-precision prediction errors (e.g., punishment, surprise).

- **Computational Function:** These bursts flood the synapse, overcoming the high-affinity 1A buffer and recruiting lower-affinity receptors (e.g., **5-HT2A**, **5-HT4**). This signals a **Context Update** (Celada et al., 2013).
- **The Shift to Cathexis:** This phasic signal forces the "Binding" of energy. It disrupts the habitual flow, commanding the cortex to hold the error signal online for "cognitive processing."

Catharsis: the architecture of stability (Gi/o)

Catharsis—defined here as the efficient discharge of Drive via established habits—relies physically on the **Gi/o signaling pathway** (defined in Section 3.1).

The 5-HT1A receptor: the anchor of the Ego

As the primary Gi/o-coupled receptor in the cortex, the **5-HT1A receptor** functions as the anchor of the Ego. By recruiting the hyperpolarization mechanism, it stabilizes prior beliefs by filtering out bottom-up noise.

- **Mechanism:** It hyperpolarizes pyramidal neurons in the amygdala and hippocampus.
- **Outcome:** When 5-HT1A signaling dominates, the Ego is resilient. The integrity of the **Allostatic Prior (the 'Wish')** is protected from minor prediction errors (frustrations), allowing the Mesolimbic Drive to flow into the Dorsal Striatum for discharge.

The 5-HT1B receptor: the presynaptic filter

Located presynaptically in the Striatum, **5-HT1B** receptors inhibit the release of Glutamate and GABA. This effectively "cleans" the signal entering the habit system, ensuring that only high-salience motor commands actuate the dorsal striatum.

Cathexis: the architecture of plasticity (Gq/11)

Cathexis—defined here as the binding of prediction error to drive model revision—is mediated physically by the Gq/11 signaling pathway (defined in Section 3.2).

The brain achieves the state of Active Cathexis (Thinking) by coordinating two distinct Gq-mediated operations: Sensory Expansion (Cortical 5-HT2A) and Motor Inhibition (Subcortical 5-HT2C).

The cortical aperture: 5-HT2A (sensory expansion)

The **5-HT2A receptor** is the primary driver of cortical plasticity. Densely expressed on the apical dendrites of Layer V pyramidal neurons, its activation increases the gain on bottom-up sensory prediction errors.

- **Mechanism:** 5-HT2A activation depolarizes the apical trunk, lowering the threshold for incoming sensory data to override top-down priors. This underpins the **REBUS effect** (Relaxed Beliefs) (Carhart-Harris & Friston, 2019).
- **The Heterodimer Gate: Integrating Content and Context** The 5-HT2A receptor forms a functional heterodimer with the Metabotropic Glutamate Receptor 2 (mGluR2) (González-Maeso et al., 2008). This complex functions as the synaptic interface where **Context gates Content**: Serotonin binding (Context) destabilizes the mGluR2 unit, thereby releasing its inhibitory "brake" on Glutamate (Content). This disinhibition allows bottom-up sensory prediction errors to flood the cortex, physically enforcing the "opening of the aperture."
- **Computational Function:** This "opens the aperture" of the mind. It prevents the Ego from ignoring error signals, forcing the system to confront discrepancies between expectation and reality.

Crucially, this theoretical 'opening' is supported by empirical neuroimaging. Preller et al. (2019) demonstrated that 5-HT2A agonism (LSD) increases global functional connectivity, effectively 'flattening' the cortical hierarchy. Furthermore, Stoliker et al. (2024) recently showed that psilocybin disrupts the rigid top-down inhibitory control of the amygdala, providing a direct mechanism for how the 'aperture' allows previously suppressed affective prediction errors to surface.

The subcortical brake: 5-HT2C (trapping the drive)

While the cortex must be "opened" to process the error, the motor system must be "clamped" to prevent the premature discharge of that energy into habit. This is the role of the **5-HT2C receptor**.

- **Mechanism:** Unlike the cortical 2A, the 5-HT2C receptor is located primarily on GABAergic interneurons in the **Ventral Tegmental Area (VTA)** and **Striatum**.
- **The "Containment" Logic:** 5-HT2C activation excites these inhibitory interneurons, which brakes the **Ventral-to-Dorsal Transition**. We propose that 5-HT2C activation does not merely suppress dopamine; it **traps** the Dopaminergic Drive in the loop between the VTA and the Cortex (Mesocortical pathway), preventing it from leaking into the Motor system (Nigrostriatal).
- **A-to-I Editing:** This receptor undergoes **A-to-I RNA editing**, creating isoforms with varying G-protein coupling efficiencies (Burns et al., 1997). This allows the brain to fine-tune the "stiffness" of the brake.
- **Computational Function:** This acts as an "Anti-Discharge" mechanism. By blocking the flow to the dorsal striatum, 5-HT2C vetoes the path of least resistance. This creates a state of **Active Cathexis**. Subjectively, this binding of energy is felt as anxiety or tension—the metabolic cost of holding a Drive in check.

Crucially, this 'containment' logic is supported by direct in vivo recordings in the human striatum. Moran et al. (2018) demonstrated that serotonergic transients do not merely reflect negative valence, but specifically encode a **'Protective Action' signal**—a command to inhibit or withdraw to avoid punishment. This aligns precisely with our model of Active Cathexis: the serotonergic signal functions as a high-precision 'Emergency Brake' that opposes the dopaminergic 'Go' signal, forcing the organism to withhold discharge to preserve safety.

Synthesis: the "Stop and think" protocol

By integrating these two Gq pathways, we resolve the apparent paradox of Serotonin's role in "learning" vs. "inhibition." They are not contradictory; they are complementary operations of Cathexis. To revise a Generative Model, the organism must simultaneously:

1. **Stop Acting (5-HT2C):** Inhibit the habitual motor discharge that would simply repeat the error.

2. **Start Listening (5-HT2A):** Increase the sensitivity to the error signal to update the internal map.

Thus, Cathexis is the synergy of physical stillness and mental reactivity. Pathologies arise when this ratio is skewed, leading to **Stuck Binding** (All brake, no steering) or **Failed Binding** (All aperture, no grounding).

Formally, this 'Stop and Think' protocol corresponds to simultaneous

- increased cortical gain on prediction error and
- suppression of striatal motor policy execution.

This synergistic logic—where inhibition is the prerequisite for flexibility—is empirically supported by Matias et al. (2017). Using photometry in the dorsal raphe, they observed that serotonin neurons are intensely active during periods of 'active waiting' (binding the impulse to move) and that this activity is specifically required for **reversal learning** (updating the generative model when contingencies change). This confirms that the 'pause' mediated by serotonin is not a passive arrest, but a high-precision computational state designed to facilitate cognitive restructuring.

Consolidation: the architecture of growth (Gs)

The shift from the transient state of Cathexis to a new, stable Catharsis (Habit) requires **Consolidation**. This is the domain of **Gs signaling**, mediated by **5-HT4, 5-HT6, and 5-HT7 receptors** (defined in Section 3.3).

- **Mechanism:** Sustained Gs activation increases cAMP levels, activating the PKA-CREB cascade.
- **Function:** This initiates the transcription of **BDNF** and the structural remodeling of synapses (Castrén & Monteggia, 2021; Duman et al., 2021). The "thought" (Cathexis) is consolidated into a "memory trace" (Habit). The energy that was bound in the cortex is now released, as the new circuit can handle the prediction error efficiently.

THE DYNAMICS OF MEMORY AND ACTION: FROM "TRACE" TO "THOUGHT"

Having defined the molecular "Precision Switch," we now scale up to the systems level. The Hierarchical Generative Model (HGM) does not merely store "memories" as static archives; it utilizes them as active **Generative Policies** for "Wish-Fulfillment" (Priors). The brain faces a fundamental trade-off between efficiency and flexibility, a duality that maps precisely onto the distinct architectures of **Procedural (Habit)** and **Declarative (Explicit)** memory systems (Squire, 2004).

This systems-level analysis provides the structural validation for Freud's profound insight in *Beyond the Pleasure Principle* (1955) and *Letter 52* (1985): **"Consciousness arises instead of a memory trace."** In our neurocomputational framework, this is literal: Phenomenal consciousness (Cathexis) is mobilized only when the automatized policy (Catharsis/Memory Trace) fails to minimize prediction error.

Catharsis as procedural memory: the architecture of the "It" (Id/automaton)

Catharsis—defined here as the efficient routing of Drive into Habit—is the domain of **Procedural Memory**. It relies on the execution of established, high-precision action policies that reside primarily in the **Basal Ganglia**, specifically the **Sensorimotor Loop**.

The "waterfall": ventral to dorsal flow

In computational Reinforcement Learning (RL), this corresponds to **"Model-Free" control** (Daw et al., 2005). The brain does not simulate the outcome of an action; it simply executes the policy.

- **The Mechanism:** The **Mesolimbic Drive** (Q/Ventral Striatum) cascades into the **Nigrostriatal Habit** (Dorsal Striatum). This transition is gated by dopamine. When the dorsal striatum holds a high-precision policy for the current context, the gate opens, and the Drive is converted immediately into Action.
- **The "Zombie" Agent:** As noted by (Koch & Crick, 2001), these circuits mediate rapid, stereotyped behaviors that operate below the threshold of phenomenal consciousness. We refer to this as the 'Zombie' agent (Koch & Crick, 2001). In computational terms, this maps strictly onto **Model-Free control**, where action policies are executed based on cached values without access to a forward-looking model.. It corresponds to Freud's Primary Process in its adaptive aspect: the drive seeks immediate motor discharge via the most direct pathway.

The energetics of discharge

Why is this "Catharsis"? Because the prediction error is "explained away" through action without the need for upstream processing.

Neurochemical Stability: This state relies on **Tonic Dopamine** (acting on D2 receptors to inhibit competing

actions) and **Tonic Serotonin** (5-HT1A/1B). The 5-HT1B receptor acts as a presynaptic "filter," suppressing neural noise and preventing the cortex from interfering with the habit (Di Matteo et al., 2008).

- **Phenomenology:** Subjectively, this is the state of **"Flow"** or automatism. The Ego (as a conscious, deliberative agent) is transparent because the "It" (the habit) is functioning perfectly. Energy is discharged; tension is zero.

Cathexis as declarative and working memory: the architecture of the "I" (ego)

Cathexis—the binding of energy to drive model revision—is the domain of **Declarative and Working Memory**. It recruits the **Prefrontal Cortex (PFC)** and the **Hippocampus**.

The shift to model-based control: minimizing expected free energy

When the habit fails, the Lateral Habenula engages the "Anti-Reward" brake (LHb 5$\rightarrow$ RMTg; see Section 2.4.1), inhibiting the ventral-to-dorsal dopamine transfer. The Drive (Q) cannot be discharged and must be "bound" in the cortex.

This blockade forces a shift from **Model-Free** to **Model-Based** control (Daw et al., 2005). Computationally, this represents a transition from minimizing *Variational Free Energy* (VFE; immediate reflex) to minimizing ***Expected*** **Free Energy (EFE)**. The brain can no longer react; it must predict. Cathexis, therefore, is the active maintenance of the Drive Vector required to simulate future trajectories ("If I do X, will I match the Wish?"). This simulation of counterfactual outcomes—metabolically expensive and slow—is the computational definition of **Secondary Process Thinking**.

Mode A: episodic memory (the search for precedent)

The first line of defense is **Episodic Memory**. The PFC queries the **Hippocampus** to retrieve a specific trace that is *not* a habit but a record of a specific context.

- **Vicarious Trial and Error (VTE):** In rodents, this state is observable as VTE, where the animal pauses at a decision point and rapidly sweeps its head (Redish, 2016). Neurophysiologically, the Hippocampus "replays" potential future paths.
- **Thinking as Experimental Action:** This validates Freud (1958): "Thinking is an experimental action performed with small quantities of energy." The Hippocampus allows the organism to "act" in a virtual space, testing hypotheses without the risk of real-world discharge.

Mode B: working memory (the "Hold")

If no memory serves, the PFC must construct a *new* solution. This requires **Working Memory**.

- **The "Binding":** Working Memory is defined as the maintenance of information online in the absence of sensory input. This requires the **Persistent Firing** of Layer V pyramidal neurons in the PFC (Goldman-Rakic, 1995).
- **Metabolic Cost:** This persistent firing is metabolically demanding (the breakdown of ATP required to maintain membrane potentials against ion flux is the thermodynamic cost detailed in Section 3.5). This explains why "thinking is hard" and why Cathexis is felt as "tension." The energy (Error/Drive) is literally trapped in the reverberating circuits of the cortex.

Consolidation: the cycle from cathexis back to catharsis

The HGM is an economy; it cannot afford to maintain Cathexis indefinitely. The goal of Thinking (Cathexis) is to produce a new Habit (Catharsis).

The Two-Stage Model of Memory

This transition maps onto the standard Two-Stage Model of memory consolidation (Buzsáki, 1989; Frankland & Bontempi, 2005):

1. **Stage 1 (Cathexis/Encoding):** During the waking state of frustration/novelty, the cortex and hippocampus are plastic (high 5-HT2A/Gq). The solution is held precariously in the labile connections of the hippocampus.
2. **Stage 2 (Catharsis/Consolidation):** During sleep (specifically SWS and REM), the solution is transferred to the neocortex and striatum.

The "growth" mechanism (Gs)

The structural transfer of information relies on the **Gs pathway** (defined in Section 3.3). The energy "bound" during the waking state of Cathexis recruits the PKA-CREB cascade, initiating the transcription of **BDNF**. This consolidates the solution: the explicit thought ("I need to turn the handle left") becomes an implicit habit (Catharsis).

CLINICAL SYNTHESIS — THE COMPUTATIONAL PATHOLOGY OF THE HGM

If the healthy mind is defined by its metabolic agility—the fluid oscillation between the stability of the Habit (**Catharsis/Exploitation**) and the plasticity of the Open Mind (**Cathexis/Exploration**)—then psychopathology can be fundamentally reconceptualized as the seizure of this engine. Mental illness is not a deficiency of "chemical juice" but a **Computational Pathology of Precision Weighting** (Friston et al., 2017).

In this framework, the Hierarchical Generative Model (HGM) becomes trapped in a maladaptive attractor state—a **deep local minimum** (a rigid, low-probability configuration in the variational free-energy landscape that resists updating). The "Precision Switch," arbitrated by the integrated dynamics of the **Lateral Habenula (LHb)**, **Serotonin (5-HT)**, and **Dopamine (DA)**, becomes "stuck."

Depression (melancholia): pathological hyper-cathexis

Major Depressive Disorder (MDD) is classically described in psychoanalysis as "the shadow of the object falling upon the ego." In our computational framework, this is a **state of Stuck Binding** (also referred to as Toxic Binding). The brain is locked in the high-energy mode of "Error Processing" but lacks the plasticity to resolve it. This interpretation aligns with computational psychiatry accounts that conceptualize mood disorders as dysregulations of reinforcement learning, valuation, and policy selection rather than as isolated neurochemical deficits (Huys et al., 2016; Montague et al., 2012)

- **The Precision Failure:** The HGM assigns infinite precision to negative prediction errors while lacking the plasticity to update the model. The system creates a **High-Precision Negative Prior** (e.g., "Failure is inevitable") which becomes immune to positive sensory evidence (Badcock et al., 2017).
- **The Mechanism (The Locked Gate):** The physiological engine of this state is the hyperactivity of the **Lateral Habenula (LHb)**. As the "Anti-Reward" center (see Section 2.4.1), the LHb broadcasts a relentless "No-Go" signal (via the RMTg) that inhibits the VTA.
- **The Waterfall Blockade:** Crucially, this blocks the **Ventral-to-Dorsal waterfall** (detailed in Section 6.1.1). The Drive (Q) is generated but is prevented from cascading into the Dorsal Striatum for discharge. The "path of least resistance" (Catharsis) is barricaded.
- **The State:** Because discharge is blocked, the error is shunted to the cortex to be "bound." It reverberates in the Default Mode Network (rumination) without ever updating the underlying Generative Model and is subjectively experienced as exhaustion, reflecting the sustained metabolic cost of unresolved prediction error. This is the profound fatigue and "ego depletion" of melancholia: VTA/Cortical binding is high, but Striatal discharge cannot take place.

Clinical box B: Clinical translation

In the clinic, this manifests as the classic melancholic presentation: the patient is physically paralyzed in bed, entirely unable to initiate goal-directed action (blocked Catharsis), yet reports their mind is utterly exhausted from relentlessly racing with guilt and negative rumination (stuck cortical Cathexis). The energy is trapped in the cortex without a pathway for discharge.

Mania: unbound SEEKING

If depression is the blockage of flow, Mania is the loss of containment. It represents a **state of Unbound SEEKING**, where the Ego fails in its primary energetic function: regulating the dopaminergic “Waterfall”.

- **The Precision Failure (Broken Brake):** The physiological driver is a functional silencing or hypoactivity of the **Lateral Habenula (LHb)**. The "Anti-Reward" brake fails to engage (Grace, 2016; Proulx et al., 2014). Consequently, the brain stops detecting negative prediction errors (Risk/Social Cost), and the precision of "Caution" drops to zero (Huys et al., 2016).
- **The Mechanism (Unsteered Flow):** Without the LHb brake, the **vmPFC "Hand"** fails to recruit the necessary inhibition. The Drive (Q) generated in the Ventral Striatum is not routed into a specific stable Habit (Dorsal) nor bound in the DMN (Cathexis). Instead, it spills over into immediate, erratic motor discharge.
- **The State:** This results in **"Exploration without a Map"** (Daw et al., 2006; Moutoussis et al., 2021). It is a high-energy state where the **Precision of Opportunity** (Ventral DA) drives action directly, bypassing the "Secondary Process" of reality testing. Subjectively, this manifests as the "Flight of Ideas"—thoughts are not bound into coherent structures but flow rapidly along associative paths of least resistance.

Clinical box C: Clinical translation

Clinically, an example of this is observed when a patient abruptly liquidates their life savings to fund an impulsive, ill-planned business venture. The biological „precision of opportunity“ drives immediate motor execution (Primary Process Catharsis) without the necessary pause to simulate the long-term financial risks (failed Cathexis).

Addiction: artificial catharsis

Addiction represents a **state of Pathological Catharsis**, where the system is tricked into believing the Wish has been fulfilled.

- **The Precision Failure (Infinite Error):** Drugs of abuse bypass the generative model entirely to chemically spike Dopamine in the Ventral Striatum. Computationally, this mimics a "Better-than-Expected" prediction error of infinite magnitude (Redish, 2004).
- **The Mechanism (The Short Circuit):** This chemical spike signals to the HGM that the drug is the "Ultimate Object" of the Wish. The system registers a massive reduction in free energy (Satisfaction) without doing the actual work of survival.
- **The State:** This creates "Artificial Catharsis." Because the drug signals that the error has been perfectly resolved, the brain consolidates the drug-taking behavior into a rigid, high-precision Habit (Dorsal Striatum). The "Wish" is overwritten: the drug becomes the new Allostatic Prior, and the absence of the drug generates a catastrophic Drive Pressure(Q) that overrides all other biological needs.

Psychosis: failed binding

Psychosis represents a catastrophic failure of the hierarchy itself—a corruption of the Precision Weighting that distinguishes "Self" from "World." It is a **state of Failed Binding** (or Aberrant Salience).

- **The Precision Failure:** This state is driven by the hyperactivity of the **5-HT2A receptor** (Gq/11) and dysregulated **D2 signaling**.
- **The Mechanism (Aberrant Salience):**
  1. **Hyper-2A (The Broken Filter):** Excessive 5-HT2A activation pathologically relaxes high-level priors (REBUS effect). The "boundaries" that separate internal expectation from external reality dissolve adding to the presence of pre-existing glutamatergic, NMDA-receptor, or developmental vulnerabilities.
  2. **Dysregulated D2:** Without stabilizing priors, ordinary background noise is assigned the high precision of significant "News" (Kapur, 2003). This is **Aberrant Salience** (i.e., dopamine assigning high precision to irrelevant stimuli; see Section 4.2).
- **The State:** The "Stop and Think" protocol fails because the "Thinking" component (2A) is hyperactive while the grounding component is absent. The patient experiences a "porous" reality where internal thoughts are hallucinated as external objects. To stop the chaos (terror), the brain constructs a new, rigid High-Precision Prior (the **Delusion**) to forcibly bind the errors.

Anxiety and OCD: incomplete cathexis

Anxiety disorders represent specific corruptions of the "Precision Switch" where the system cannot determine if an error has been resolved.

- **Generalized Anxiety:** A **state of High Tonic Cathexis**. The system is perpetually "on guard." The **5-HT1A (Stability)** mechanism fails to dampen the "Noise" of the amygdala, leaving the HGM in a state of chronic vibration (persistent high-tonic NA/5-HT2A signaling).
- **OCD (The Broken Loop):** OCD is a specific "glitch" in the completion of Catharsis. The patient performs the ritual (Action), but the "Feeling of Knowing" or "Satisfaction" signal (OFC) never arrives (Pauls et al., 2014; Saxena & Rauch, 2000; Szechtman & Woody, 2004). The loop becomes broken: Action-without-Resolution.

Clinical box D: Clinical translation (OCD)

A patient washes their hands raw (the motor execution of Catharsis) but never experiences the subjective ‚click‘ of cleanliness. Because the expected neurobiological signal of completion fails to arrive, the prediction error remains high, forcing the patient to repeat the washing ritual endlessly in a futile attempt to discharge the tension.

THERAPEUTIC IMPLICATIONS: ENGINEERING THE STABILITY-PLASTICITY EQUILIBRIUM

If psychopathology is a “Computational Pathology,” then the goal of therapy is to restore the fluid oscillation of the HGM. We move from a “replacement model” (fixing chemical deficits) to a **“Tuning Model”**: Pharmacology adjusts the **Gain Parameters** (Hardware), while Psychotherapy edits the **Generative Model** (Software).

Pharmacological precision engineering

SSRIs: the stabilizers (enhancing resilience)

- **Target: 5-HT1A (Gi/o)**.
- **Mechanism:** SSRIs function by increasing the global tone of Serotonin. After initial autoreceptor desensitization (2–4 weeks), this tone preferentially engages high-affinity **5-HT1A receptors** in the limbic system.
- **Computational Effect:** This recruits the **Architecture of Stability**. It hyperpolarizes the amygdala, dampening the "noise" of bottom-up prediction errors. It effectively "thickens the skin" of the Ego, preventing "Toxic Binding" by raising the threshold for what counts as an error (Carhart-Harris & Nutt, 2017).

Clinical box E:
Clinical illustration – the stabilization of ms. D

The case of Ms. D illustrates how pharmacological enhancement of the 5-HT1A (Gi/o) pathway provides crucial psychic stabilization (passive coping).

Ms. D, a woman in her late 40s, presented with lifelong anxiety and recurrent depressive episodes since her early 20s, typically precipitated by significant life stressors. Her clinical course was characterized by a repeating pattern: treatment with Escitalopram (an SSRI) led to remission; discontinuation of medication led to stability for 6-12 months, followed by relapse necessitating the restart of therapy.

Analysis: Ms. D.'s history suggests a generative model characterized by relatively low resilience to stress. Her endogenous capacity for Catharsis (efficient tension management) is easily overwhelmed. Within the HGM framework, SSRI treatment chronically enhances 5-HT1A signaling (following autoreceptor desensitization). This increases the precision of her high-level priors, bolstering the stability of the HGM against perturbation and preventing a descent into a deep depressive attractor state.

NaSSAs (mirtazapine): the brake--cutters (restoring discharge)

- **Target: 5-HT2C Antagonism** (and α2-Adrenergic Blockade).
- **Mechanism:** Unlike SSRIs, NaSSAs specifically **cut the brake**. By blocking the **5-HT2C receptor**, they remove the inhibition on the VTA and Striatum.
- **Computational Effect:** This **Disinhibits the Waterfall**. It physically unblocks the transfer of Drive (Ventral) to Habit (Dorsal), directly treating Anhedonia by restoring the "Go" signal that was suppressed by the anxiety/inhibition system (Stahl, 2013).

Novel antipsychotics: the aperture closers (stabilizing boundaries)

- **Target: 5-HT2A Antagonism** (Gq/11).
- **Mechanism:** Atypical antipsychotics potently block the 5-HT2A receptor.
- **Computational Effect:** This **Closes the Aperture**. By opposing the "REBUS" effect (Relaxed Beliefs), they tighten the precision of high-level priors. This restores the "filter" that distinguishes internal noise from external reality, preventing the Sensory Flooding that drives delusion formation (Meltzer, 1999).

Clinical box F:
Clinical illustration – the aperture closer

The 5-HT2A (Gq/11) pathway modulates plasticity. Hyperactivity in this system is implicated in psychosis, where the excessive relaxation of priors leads to aberrant salience and distorted Cathexis. We illustrate this with a summary of a case reported by Sommer et al. (2018) concerning intractable visual hallucinations.

A 29-year-old woman with childhood-onset schizophrenia presented with severe, chronic, and treatment-resistant psychosis. She experienced persistent multimodal hallucinations, including profound visual hallucinations (seeing frightening people and objects morphing). These experiences were debilitating and these perceptions consumed her attentional resources. These symptoms persisted despite extensive treatment, including high doses of clozapine. The treatment team initiated ketanserin, a potent 5-HT2A receptor antagonist. This resulted in a significant reduction of her visual hallucinations but not of her auditory hallucinations (Sommer et al., 2018).

Analysis: This case illustrates psychosis as a disorder of aberrant precision. Excessive activity in the 5-HT2A pathway leads to a pathological decrease in the precision of high-level perceptual priors. This failure of top-down models allows unconstrained bottom-up errors (hallucinations) to dominate experience, resulting in an aberrant Cathexis. The successful treatment with a selective 5-HT2A antagonist demonstrates that blocking this pathway can increase the precision of priors, thereby stabilizing the HGM. The observed positive effect specifically on visual hallucinations corresponds to the high expression of 5-HT2A receptors in the visual cortex relative to the auditory cortex (Beliveau et al., 2017; Carhart-Harris & Nutt, 2017), supporting the anatomical specificity of this precision modulation system.

Psychotherapeutic synergy

Pharmacology opens the window (Plasticity), but Psychotherapy provides the context required to guide it.

The therapist as "auxiliary cortex"

Patients with trauma avoid Cathexis (thinking about the pain) because the prediction error is too high—it threatens to overwhelm the Ego.

- **The Safe Container:** The therapist acts as an **"Auxiliary Cortex."** Through empathy and non-judgmental attunement, the therapist provides a high-precision source of Safety by externally stabilizing precision signals.
- **Neurobiological Mechanism:** We hypothesize this social buffering recruits the **Oxytocin-5-HT1A** system. This enhances **Stability (Gi/o)**, creating a biological buffer that allows the patient to tolerate the high-arousal state of **Active Cathexis** (accessing the trauma) without triggering the defensive **5-HT2C** (Panic/Brake) reaction (Holmes & Nolte, 2019).

Reconsolidation: the new catharsis

The therapeutic aim of reconsolidation is to transform **Toxic Binding** into **Insight** by permanently updating the generative model. Although often described loosely, memory reconsolidation follows a highly specific computational and neurobiological sequence:

Prediction error: the triggering ingredient

A memory can only be revised when it is accompanied by **prediction error**—a discrepancy between what the old memory predicts and what the current context provides.

In therapy, this arises when the patient reactivates a traumatic schema ("I will be shamed," "I am to blame") in a relational environment that violates the expected outcome (e.g., the therapist is calm, non-judgmental, accepting).

This **"surprise" signal** gives the nervous system evidence that the old model is not fully accurate, opening the door to change.

Mismatch experience: the subjective entry point

Prediction error must take the form of a **felt mismatch** (Lane et al., 2015):

- The activated painful memory, *plus*
- A contradictory or disconfirming here-and-now experience.

Therapeutically, mismatch is created when:

- The patient expects rejection but receives attunement.
- The patient expects danger but encounters safety.
- The patient anticipates punishment but meets curiosity.

This mismatch is not merely cognitive — it must be **emotionally precise and embodied**, because the generative model being updated is itself affectively encoded.

Destabilization window: opening the trace

When prediction error is sufficiently strong and precisely targeted, the retrieved memory becomes **labile** — a state known as **destabilization**.

Neurobiologically, this involves:

- **5-HT2A/Gq-mediated relaxation of priors** (lowering the precision of the old model), and
- **Hippocampal reactivation of the trace**, which temporarily reopens the synaptic ensembles encoding it.

This destabilization forms a **time-limited window** (roughly minutes to hours) during which the memory can be rewritten at the molecular level. Without destabilization, the system merely appends new learning alongside the old model; with destabilization, the old model itself becomes modifiable.

Updating: integrating new evidence into the model

Within the destabilized state, the brain must receive **new affectively salient information** that resolves the original prediction error.

This is where the therapist's role as **contextual precision** becomes decisive:

- Their consistent safety provides a **high-precision new prior**,
- Allowing the cortex to bind the error without collapse into defenses or avoidance,
- And enabling the integration of contradictory information ("I am safe," "I am not to blame," "I can be seen without being harmed").

At the computational level, this corresponds to:

- Assigning high precision to the new relational evidence,
- Reducing precision on the old traumatic prior,
- Updating the prediction model to reduce future free energy.

Reconsolidation: the gs-mediated "save function"

Reconsolidation is the **molecular rewriting** of the memory trace.

Once the new solution is stabilized psychologically, the brain recruits **Gs-coupled pathways** (5-HT4/6/7, D1) that drive:

- cAMP → PKA → CREB activation → BDNF synthesis,
- Structural remodeling of synapses,
- Integration of the new prediction into the hierarchy.

Reconsolidation does not merely add new memories — it **replaces** the old, maladaptive expectation with a new prior.

This allows the patient to achieve a **New Catharsis**: the release of previously trapped energy as the updated model can now efficiently resolve prediction error without pathological Cathexis.

Clinical box G: Clinical illustration – psychotherapeutic reconsolidation and the auxiliary cortex

The case of Mr. E illustrates how psychotherapy utilizes the „Precision Switch" to rewrite a maladaptive Generative Model.

Case Presentation: Mr. E, a man with a history of an emotionally volatile, highly critical parent, presents with severe interpersonal anxiety. His generative model contains a rigid, high-precision prior: *"If I make a mistake, I will be viciously attacked."* His habitual policy (Catharsis) is to apologize profusely and withdraw to minimize this predicted error.

Therapeutic Intervention: During a session, Mr. E realizes he forgot to complete an agreed-upon therapeutic task. He immediately freezes, expects the therapist to become angry, and begins his habitual apologizing. Instead of validating this expectation, the therapist remains calm, warm, and curious, stating: *„I notice you are expecting me to be upset with you right now. Look at my face; what do you see?"*

Computational Analysis: This intervention generates a precise Prediction Error (a "Mismatch Experience"). Mr. E's activated traumatic prior predicts an attack, but external reality contradicts it. The therapist acts as an "Auxiliary Cortex," providing a high-precision signal of safety (hypothesized via oxytocinergic/5-HT1A buffering). This safety buffer allows Mr. E's system to tolerate the error without engaging the 5-HT2C panic brake (flight/freeze). Supported by the therapist, the 5-HT2A system relaxes the old prior, entering a transient state of *Active Cathexis* where the memory trace is destabilized. Through the Gs-pathway (BDNF/LTP), the new evidence (*"I can make a mistake and remain safe"*) is consolidated, structurally updating his generative model and allowing for a new, healthier Catharsis in future relationships.

SYNTHESIS OF DYNAMICS

Table 4. Hypothesized Neuromodulatory Profiles in States of Mind

State	Neuromodulatory Profile	Computational Characteristic
Stability (Catharsis)	**Balanced 5-HT1A/Gi + Nigrostriatal DA**	**Efficient Inference:** Low prediction error; „Flow" state; Drive cascades to Habit.
Cathexis (Binding)	**High Cortical 5-HT2A (Gq) + High VTA DA (Blocked)**	**Active Processing:** Drive is trapped (LHb brake); Priors relaxed (2A) to permit update.
Consolidation (Growth)	**High Gs (5-HT4/7, D1) + BDNF**	**Rewiring:** Structural encoding of the new model.

Table 5. Mental Disorders as Dysregulations of Precision

Disorder	Core Computational Pathology	Hypothesized Neuro-Mechanism
Depression	**Stuck Binding:** Drive trapped; failure of discharge.	**Hyperactive Habenula** blocks Ventral-to-Dorsal flow. Drive pools in cortex (Rumination).
Mania	**Unbound Seeking:** Pathological Discharge.	**Hypoactive Habenula** fails to brake Ventral DA. Drive floods motor system without binding.
Psychosis	**Failed Binding:** Dissolution of priors.	**5-HT2A Hyperactivity** dissolves boundaries; **D2** creates aberrant salience.
OCD	**Broken Catharsis:** Action without Resolution.	**CSTC Loop Glitch:** Satisfaction signal (OFC) missing; error persists despite action.

LIMITATIONS AND FALSIFIABILITY

Conceptual scope and epistemic status

The present work is a theoretical synthesis and hypothesis-generating model rather than a definitive mechanistic account. As outlined in Box 1, the framework integrates psychoanalytic constructs, the Free Energy Principle (FEP), and neuromodulatory neurobiology to propose a coherent neurocomputational interpretation of cathexis and catharsis. While many individual components of the model are grounded in established empirical findings, their integration into a unified "precision switch" architecture necessarily involves speculative inference.

Accordingly, the model should be evaluated not by whether every proposed mapping is already empirically confirmed, but by whether it is

- internally coherent

- consistent with known constraints from neuroscience and psychiatry
- capable of generating testable predictions that distinguish it from alternative accounts.

Throughout the manuscript, the term "prediction error" is used in a pragmatic computational sense to denote the dominant contributors to variational free energy under simplifying assumptions, rather than as a strict synonym for variational free energy itself. This usage follows pragmatic treatments of precision in active inference, where precision weighting is understood as a context-sensitive and hierarchically distributed quantity rather than a single, directly measurable variable (Friston et al., 2015; Parr & Friston, 2017).

Limitations of conceptual mapping

A first limitation concerns the translation of psychoanalytic concepts into computational and neurobiological terms. Constructs such as *cathexis*, *catharsis*, *drive*, and *wish* do not correspond one-to-one with discrete neural mechanisms. Our mappings are therefore functional and organizational rather than literal. For example, cathexis is operationalized as the sustained precision-weighted maintenance of prediction error, not as a unitary neural process. Alternative mappings—emphasizing different hierarchical levels or neuromodulatory systems—remain possible and may prove equally or more explanatory.

Relatedly, the model adopts a dual-aspect monist stance, treating subjective experience and neural dynamics as complementary descriptions of the same underlying process. This philosophical position is not empirically testable in itself and is offered as a pragmatic framework for integration rather than as a metaphysical claim.

Limitations of neurobiological specificity

A second limitation concerns neuroanatomical and neurochemical specificity. Although the model highlights particular structures (e.g., lateral habenula, raphe nuclei, striatum) and receptor families (Gi/o, Gq/11, Gs), it does not claim that these elements function in isolation or that they constitute the sole determinants of cathexis–catharsis dynamics. Neural control of precision is distributed, redundant, and context-dependent, involving cortical, thalamic, and neuromodulatory interactions that are simplified here for theoretical clarity.

Similarly, constructs such as the "Extended Reticular–Thalamic Activating System" are used as integrative labels rather than as claims about a single anatomically bounded system. The model abstracts across heterogeneous ascending neuromodulatory pathways to capture their shared computational role in gain control.

Limitations of clinical generalization

The clinical formulations presented (e.g., depression as "stuck binding," mania as "unbound seeking," psychosis as "failed binding") are intended as computational phenotypes rather than exhaustive pathophysiological explanations. Psychiatric disorders are heterogeneous, developmentally contingent, and multiply determined. The proposed mappings do not deny the roles of genetics, development, social context, or non-monoaminergic mechanisms, but instead aim to provide a unifying lens through which these factors may converge on shared failures of precision regulation.

As such, the framework should not be interpreted as a replacement for existing diagnostic or treatment models, but as a complementary perspective that may help explain why diverse interventions—pharmacological, psychotherapeutic, and contextual—can converge on similar clinical outcomes.

Falsifiability and testable predictions

Despite its integrative and theoretical nature, the model makes several falsifiable predictions that distinguish it from alternative accounts:

1. **Precision-Switch Manipulation** If cathexis corresponds to sustained precision-weighted error, then experimentally increasing cortical 5-HT2A signaling while simultaneously inhibiting striatal discharge (e.g., via 5-HT2C activation) should increase behavioral and neural markers of model-based control, working memory load, and cortical metabolic demand, even in the absence of overt action.
2. **Habenula-Dependent Gating** The model predicts that disruption of lateral habenula signaling should impair the transition from habitual (model-free) to deliberative (model-based) control following negative prediction error. Failure to observe such gating effects would directly challenge the proposed role of the LHb as a critical node in cathexis–catharsis arbitration.
3. **Metabolic Signature of Cathexis** If cathexis is metabolically costly, then states characterized by sustained binding (e.g., rumination, unresolved conflict) should show increased energy consumption in deep-layer prefrontal pyramidal neurons and associated networks,

relative to efficient habitual states. Absence of such metabolic differentiation would undermine the energetic dimension of the model.

4. **Therapeutic Precision Reset** The framework predicts that successful psychotherapy or pharmacotherapy should be associated not merely with symptom reduction, but with measurable normalization of precision weighting—manifesting as restored flexibility between exploratory and exploitative control modes. Interventions that improve symptoms without altering such flexibility would require revision of the model.

Scope for revision

Finally, the model is explicitly provisional. As empirical work in computational psychiatry, neuromodulation, and neuropsychoanalysis advances, individual components of the framework may require refinement, replacement, or rejection. The value of the present synthesis lies not in its finality, but in its capacity to organize disparate findings into a coherent explanatory architecture and to motivate targeted empirical investigation.

In this sense, the model should be regarded as a working hypothesis about the economy of the mind–brain, open to falsification, modification, and extension.

CONCLUSION: THE "PULSATION" OF THE MIND-BRAIN

This article endeavors towards a resolution of the century-long estrangement between the "Mental Apparatus" of Psychoanalysis and the "Neural Machinery" of Neuroscience. By adopting the **Free Energy Principle** as a translational framework, we conclude that the Mind-Brain is a homeostatic engine that pulses between two fundamental modes of being.

We posit that healthy mental functioning is defined by the fluid **Oscillation of Being**—the ability to bind energy to learn (**Cathexis/Exploration**) and discharge energy to live (**Catharsis/Exploitation**). This oscillation is not merely a psychological metaphor but a neurocomputational necessity, arbitrated by the integrated **ERTAS**.

- **Catharsis** is the efficient channeling of the **Mesolimbic Drive** into the **Striatal Habit**, stabilized by the "Keep Calm" signal of **5-HT1A** (Gi/o).
- **Cathexis** is the strategic interruption of this flow by the **Lateral Habenula** and **5-HT2C** (Brake), forcing the Drive to be "bound" in the cortex. Here, **5-HT2A** (Aperture) relaxes the model to permit revision.
- **Growth** is the structural consolidation of this revision via the **Gs-Pathway**, allowing the system to return to efficiency.

This oscillation is bounded by thermodynamics: Cathexis (binding) is metabolically expensive (ATP-demanding), whereas Catharsis (habit) is efficient. The pathology of depression is thus an 'energetic bankruptcy'—spending high energy on binding without achieving the release of discharge.

In this view, the human being is not a broken machine, but an active agent striving for equilibrium. Psychopathology is the "Stuck Switch" of this engine—a deep local minimum (a rigid maladaptive attractor) where the oscillation ceases. By tuning the gain on these circuits, we can restart the pulse, allowing the mind to once again move from the tension of the Question (Cathexis) to the relief of the Answer (Catharsis).

Conflicts of interest: No conflict of interest.
Ethical considerations: Not applicable.
Funding sources: None.
Authors' contributions: Arif de Mendelssohn: design of the review, literature searches and analyses, interpretation of data, manuscript writing. Henriette Löffler-Stastka: critical revision of the manuscript for important intellectual content.

References

Agnati, L. F., Zoli, M., Strömberg, I., & Fuxe, K. (1995). Intercellular communication in the brain: Wiring versus volume transmission. *Neuroscience, 69*(3), 711–726. https://doi.org/10.1016/0306-4522(95)00308-6

Alcaro, A., Carta, S., & Panksepp, J. (2017). The affective core of the self: A neuro-archetypical perspective on the foundations of human (and animal) subjectivity [Review]. *Frontiers in Psychology, 8*, Article 1424. https://doi.org/10.3389/fpsyg.2017.01424

Attwell, D., & Laughlin, S. B. (2001). An energy budget for signaling in the grey matter of the brain. *Journal of Cerebral Blood Flow & Metabolism, 21*(10), 1133–1145. https://doi.org/10.1097/00004647-200110000-00001

Badcock, P. B., Davey, C. G., Whittle, S., Allen, N. B., & Friston, K. J. (2017). The depressed brain: An evolutionary systems theory. *Trends in Cognitive Sciences, 21*(3), 182–194. https://doi.org/10.1016/j.tics.2017.01.005

Balasubramani, P. P., Chakravarthy, V. S., Wong-Lin, K., Wang, D. H., Cohen, J. Y., Nakamura, K., & Moustafa, A. A. (2017).

Neural circuit models of the serotonergic system: From microcircuits to cognition. In A. A. Moustafa (Ed.), *Computational models of brain and behavior* (pp. 389–400). Wiley-Blackwell. https://doi.org/10.1002/9781119159193.ch28

Barnes, N. M., & Sharp, T. (1999). A review of central 5-HT receptors and their function. *Neuropharmacology, 38*(8), 1083–1152. https://doi.org/10.1016/S0028-3908(99)00010-6

Beliveau, V., Ganz, M., Feng, L., Ozenne, B., Højgaard, L., Fisher, P. M., Svarer, C., Greve, D. N., & Knudsen, G. M. (2017). A high-resolution in vivo atlas of the human brain's serotonin system. *The Journal of Neuroscience, 37*(1), 120–128. https://doi.org/10.1523/JNEUROSCI.2830-16.2016

Berger, M., Gray, J. A., & Roth, B. L. (2009). The expanded biology of serotonin. *Annual Review of Medicine, 60*, 355–366. https://doi.org/10.1146/annurev.med.60.042307.110802

Berridge, K. C., & Robinson, T. E. (1998). What is the role of dopamine in reward: Hedonic impact, reward learning, or incentive salience? *Brain Research Reviews, 28*(3), 309–369. https://doi.org/10.1016/S0165-0173(98)00019-8

Burns, C. M., Chu, H., Rueter, S. M., Hutchinson, L. K., Canton, H., Sanders-Bush, E., & Emeson, R. B. (1997). Regulation of serotonin-2C receptor G-protein coupling by RNA editing. *Nature, 387*(6630), 303–308. https://doi.org/10.1038/387303a0

Buzsáki, G. (1989). Two-stage model of memory trace formation: A role for *noisy* brain states. *Neuroscience, 31*(3), 551–570. https://doi.org/10.1016/0306-4522(89)90423-5

Carhart-Harris, R. L., & Friston, K. J. (2010). The default-mode, ego-functions and free-energy: A neurobiological account of Freudian ideas. *Brain, 133*(Pt 4), 1265–1283. https://doi.org/10.1093/brain/awq010

Carhart-Harris, R. L., & Friston, K. J. (2019). REBUS and the anarchic brain: Toward a unified model of the brain action of psychedelics. *Pharmacological Reviews, 71*(3), 316–344. https://doi.org/10.1124/pr.118.017160

Carhart-Harris, R. L., & Nutt, D. J. (2017). Serotonin and brain function: A tale of two receptors. *Journal of Psychopharmacology, 31*(9), 1091–1120. https://doi.org/10.1177/0269881117725915

Castrén, E., & Monteggia, L. M. (2021). Brain-derived neurotrophic factor signaling in depression and antidepressant action. *Biological Psychiatry, 90*(2), 128–136. https://doi.org/10.1016/j.biopsych.2021.05.008

Celada, P., Puig, M. V., & Artigas, F. (2013). Serotonin modulation of cortical neurons and networks. *Frontiers in Integrative Neuroscience, 7*, Article 25. https://doi.org/10.3389/fnint.2013.00025

Chalmers, D. J. (1996). *The conscious mind: In search of a fundamental theory*. Oxford University Press.

Cools, R., Nakamura, K., & Daw, N. D. (2011). Serotonin and dopamine: Unifying affective, activational, and decision functions. *Neuropsychopharmacology, 36*(1), 98–113. https://doi.org/10.1038/npp.2010.121

Coppen, A. (1967). The biochemistry of affective disorders. *The British Journal of Psychiatry, 113*(504), 1237–1264. https://doi.org/10.1192/bjp.113.504.1237

Daw, N. D., Niv, Y., & Dayan, P. (2005). Uncertainty-based competition between prefrontal and dorsolateral striatal systems for behavioral control. *Nature Neuroscience, 8*(12), 1704–1711. https://doi.org/10.1038/nn1560

Daw, N. D., O'Doherty, J. P., Dayan, P., Seymour, B., & Dolan, R. J. (2006). Cortical substrates for exploratory decisions in humans. *Nature, 441*(7095), 876–879. https://doi.org/10.1038/nature04766

Deco, G., & Kringelbach, M. L. (2020). Turbulent-like dynamics in the human brain. *Cell Reports, 33*(10), Article 108471. https://doi.org/10.1016/j.celrep.2020.108471

Delgado, P. L. (2000). Depression: The case for a monoamine deficiency. *The Journal of Clinical Psychiatry, 61*(Suppl 6), 7–11.

Di Giovanni, G., Di Matteo, V., & Esposito, E. (2008). Serotonin-dopamine interaction: Experimental evidence and therapeutic relevance. Preface. In *Progress in Brain Research, 172*, ix. https://doi.org/10.1016/S0079-6123(08)00931-X

Di Matteo, V., Pierucci, M., Esposito, E., Crescimanno, G., Benigno, A., & Di Giovanni, G. (2008). Serotonin modulation of the basal ganglia circuitry: Therapeutic implication for Parkinson's disease and other motor disorders. In *Progress in Brain Research, 172*, 423–463. https://doi.org/10.1016/S0079-6123(08)00921-7

Duman, R. S., Aghajanian, G. K., Sanacora, G., & Krystal, J. H. (2016). Synaptic plasticity and depression: New insights from stress and rapid-acting antidepressants. *Nature Medicine, 22*(3), 238–249. https://doi.org/10.1038/nm.4050

Duman, R. S., Deyama, S., & Fogaça, M. V. (2021). Role of BDNF in the pathophysiology and treatment of depression: Activity-dependent effects distinguish rapid-acting antidepressants. *European Journal of Neuroscience, 53*(1), 126–139. https://doi.org/10.1111/ejn.14630

Fonagy, P., Gergely, G., Jurist, E. L., & Target, M. (2002). *Affect regulation, mentalization, and the development of the self*. Other Press.

Frank, M. J., Seeberger, L. C., & O'Reilly, R. C. (2004). By carrot or by stick: Cognitive reinforcement learning in parkinsonism. *Science, 306*(5703), 1940–1943. https://doi.org/10.1126/science.1102941

Frankland, P. W., & Bontempi, B. (2005). The organization of recent and remote memories. *Nature Reviews Neuroscience, 6*(2), 119–130. https://doi.org/10.1038/nrn1607

Freud, S. (1950). Project for a scientific psychology. In J. Strachey (Ed.), *The standard edition of the complete psychological works of Sigmund Freud* (Vol. 1, pp. 283–397). Hogarth Press.

Freud, S. (1953). *The interpretation of dreams*. In J. Strachey (Ed.), *The standard edition of the complete psychological works of Sigmund Freud* (Vols. 4–5). Hogarth Press.

Freud, S. (1955). *Beyond the pleasure principle*. In J. Strachey (Ed.), *The standard edition of the complete psychological works of Sigmund Freud* (Vol. 18, pp. 7–64). Hogarth Press.

Freud, S. (1957a). Instincts and their vicissitudes. In J. Strachey (Ed.), *The standard edition of the complete psychological works of Sigmund Freud* (Vol. 14, pp. 109–140). Hogarth Press.

Freud, S. (1957b). The unconscious. In J. Strachey (Ed.), *The standard edition of the complete psychological works of Sigmund Freud* (Vol. 14, pp. 159–215). Hogarth Press.

Freud, S. (1958). Formulations on the two principles of mental functioning. In J. Strachey (Ed.), *The standard edition of the complete psychological works of Sigmund Freud* (Vol. 12, pp. 213–226). Hogarth Press.

Freud, S. (1985). Letter 52 to Wilhelm Fliess (6 December 1896). In J. M. Masson (Ed.), *The complete letters of Sig-*

mund Freud to Wilhelm Fliess, 1887–1904. Harvard University Press.

Friston, K. (2010). The free-energy principle: A unified brain theory? *Nature Reviews Neuroscience, 11*(2), 127–138. https://doi.org/10.1038/nrn2787

Friston, K., FitzGerald, T., Rigoli, F., Schwartenbeck, P., & Pezzulo, G. (2017). Active inference: A process theory. *Neural Computation, 29*(1), 1–49. https://doi.org/10.1162/NECO_a_00912

Friston, K., Rigoli, F., Ognibene, D., Mathys, C., Fitzgerald, T., & Pezzulo, G. (2015). Active inference and epistemic value. *Cognitive Neuroscience, 6*(4), 187–214. https://doi.org/10.1080/17588928.2015.1020053

Friston, K. J., Shiner, T., FitzGerald, T., Galea, J. M., Adams, R., Brown, H., Dolan, R. J., Moran, R., Stephan, K. E., & Bestmann, S. (2012). Dopamine, affordance and active inference. *PLOS Computational Biology, 8*(1), e1002327. https://doi.org/10.1371/journal.pcbi.1002327

Goldman-Rakic, P. S. (1995). Cellular basis of working memory. *Neuron, 14*(3), 477–485. https://doi.org/10.1016/0896-6273(95)90304-6

González-Maeso, J., Ang, R. L., Yuen, T., Chan, P., Weisstaub, N. V., López-Giménez, J. F., Zhou, M., Okawa, Y., Callado, L. F., Milligan, G., Gingrich, J. A., Filizola, M., Meana, J. J., & Sealfon, S. C. (2008). Identification of a serotonin/glutamate receptor complex implicated in psychosis. *Nature, 452*(7183), 93–97. https://doi.org/10.1038/nature06612

Grace, A. A. (2016). Dysregulation of the dopamine system in the pathophysiology of schizophrenia and depression. *Nature Reviews Neuroscience, 17*(8), 524–532. https://doi.org/10.1038/nrn.2016.57

Grossman, C. D., Bari, B. A., & Cohen, J. Y. (2022). Serotonin neurons modulate learning rate through uncertainty. *Current Biology, 32*(3), 586–599.e587. https://doi.org/10.1016/j.cub.2021.12.006

Haas, H. L., Sergeeva, O. A., & Selbach, O. (2008). Histamine in the nervous system. *Physiological Reviews, 88*(3), 1183–1241. https://doi.org/10.1152/physrev.00043.2007

Haber, S. N. (2014). The place of dopamine in the cortico-basal ganglia circuit. *Neuroscience, 282*, 248–257. https://doi.org/10.1016/j.neuroscience.2014.10.008

Hikosaka, O. (2010). The habenula: From stress evasion to value-based decision-making. *Nature Reviews Neuroscience, 11*(7), 503–513. https://doi.org/10.1038/nrn2866

Holmes, J., & Nolte, T. (2019). "Surprise" and the Bayesian brain: Implications for psychotherapy theory and practice [Hypothesis and Theory]. *Frontiers in Psychology, 10*, Article 592. https://doi.org/10.3389/fpsyg.2019.00592

Huys, Q. J. M., Maia, T. V., & Frank, M. J. (2016). Computational psychiatry as a bridge from neuroscience to clinical applications. *Nature Neuroscience, 19*(3), 404–413. https://doi.org/10.1038/nn.4238

Kapur, S. (2003). Psychosis as a state of aberrant salience: A framework linking biology, phenomenology, and pharmacology in schizophrenia. *The American Journal of Psychiatry, 160*(1), 13–23. https://doi.org/10.1176/appi.ajp.160.1.13

Koch, C., & Crick, F. (2001). The zombie within. *Nature, 411*(6840), 893–893. https://doi.org/10.1038/35082161

Lane, R. D., Ryan, L., Nadel, L., & Greenberg, L. (2015). Memory reconsolidation, emotional arousal, and the process of change in psychotherapy: New insights from brain science. *Behavioral and Brain Sciences, 38*, e1. https://doi.org/10.1017/S0140525X14000041

Larkum, M. (2013). A cellular mechanism for cortical associations: An organizing principle for the cerebral cortex. *Trends in Neurosciences, 36*(3), 141–151. https://doi.org/10.1016/j.tins.2012.11.006

Löffler-Stastka, H. (2023). Psychiatry in the bio-psycho-social research paradigm as option for enduring and sustainable change in a time of chronic crisis. *Psychiatria Danubina, 35*(3), 295–306. https://doi.org/10.24869/psyd.2023.295

Matias, S., Lottem, E., Dugué, G. P., & Mainen, Z. F. (2017). Activity patterns of serotonin neurons underlying cognitive flexibility. *eLife, 6*, e20552. https://doi.org/10.7554/eLife.20552

McEwen, B. S. (2007). Physiology and neurobiology of stress and adaptation: Central role of the brain. *Physiological Reviews, 87*(3), 873–904. https://doi.org/10.1152/physrev.00041.2006

McEwen, B. S., & Wingfield, J. C. (2003). The concept of allostasis in biology and biomedicine. *Hormones and Behavior, 43*(1), 2–15. https://doi.org/10.1016/S0018-506X(02)00024-7

Meltzer, H. Y. (1999). The role of serotonin in antipsychotic drug action. *Neuropsychopharmacology, 21*(2 Suppl), 106S–115S. https://doi.org/10.1016/S0893-133X(99)00046-9

Merker, B. (2007). Consciousness without a cerebral cortex: A challenge for neuroscience and medicine. *Behavioral and Brain Sciences, 30*(1), 63–81; discussion 81–134. https://doi.org/10.1017/S0140525X07000891

Moncrieff, J., Cooper, R. E., Stockmann, T., Amendola, S., Hengartner, M. P., & Horowitz, M. A. (2023). The serotonin theory of depression: A systematic umbrella review of the evidence. *Molecular Psychiatry, 28*(8), 3243–3256. https://doi.org/10.1038/s41380-022-01661-0

Montague, P. R., Dolan, R. J., Friston, K. J., & Dayan, P. (2012). Computational psychiatry. *Trends in Cognitive Sciences, 16*(1), 72–80. https://doi.org/10.1016/j.tics.2011.11.018

Moran, R. J., Kishida, K. T., Lohrenz, T., Saez, I., Laxton, A. W., Witcher, M. R., Tatter, S. B., Ellis, T. L., Phillips, P. E. M., Dayan, P., & Montague, P. R. (2018). The protective action encoding of serotonin transients in the human brain. *Neuropsychopharmacology, 43*(6), 1425–1435. https://doi.org/10.1038/npp.2017.304

Moutoussis, M., Garzón, B., Neufeld, S., Bach, D. R., Rigoli, F., Goodyer, I., Bullmore, E., Guitart-Masip, M., & Dolan, R. J. (2021). Decision-making ability, psychopathology, and brain connectivity. *Neuron, 109*(12), 2025–2040.e2027. https://doi.org/10.1016/j.neuron.2021.04.019

Northoff, G. (2016). *Neuro-philosophy and the healthy mind: Learning from the unwell brain*. W. W. Norton & Company.

Panksepp, J. (1998). *Affective neuroscience: The foundations of human and animal emotions*. Oxford University Press.

Parr, T., & Friston, K. J. (2017). Uncertainty, epistemics and active inference. *Journal of The Royal Society Interface, 14*(136), 20170376. https://doi.org/10.1098/rsif.2017.0376

Pauls, D. L., Abramovitch, A., Rauch, S. L., & Geller, D. A. (2014). Obsessive-compulsive disorder: An integrative genetic and neurobiological perspective. *Nature Reviews Neuroscience, 15*(6), 410–424. https://doi.org/10.1038/nrn3746

Preller, K. H., Razi, A., Zeidman, P., Stämpfli, P., Friston, K. J., & Vollenweider, F. X. (2019). Effective connectivity changes in LSD-induced altered states of conscious-

ness in humans. *Proceedings of the National Academy of Sciences, 116*(7), 2743–2748. https://doi.org/10.1073/pnas.1815129116

Proulx, C. D., Hikosaka, O., & Malinow, R. (2014). Reward processing by the lateral habenula in normal and depressive behaviors. *Nature Neuroscience, 17*(9), 1146–1152. https://doi.org/10.1038/nn.3779

Qi, Z., Fieni, D., Tretter, F., & Voit, E. O. (2013). The neurochemical mobile with non-linear interaction matrix: An exploratory computational model. *Pharmacopsychiatry, 46*(Suppl 1), S53–S63. https://doi.org/10.1055/s-0033-1337920

Redish, A. D. (2004). Addiction as a computational process gone awry. *Science, 306*(5703), 1944–1947. https://doi.org/10.1126/science.1102384

Redish, A. D. (2016). Vicarious trial and error. *Nature Reviews Neuroscience, 17*(3), 147–159. https://doi.org/10.1038/nrn.2015.30

Saxena, S., & Rauch, S. L. (2000). Functional neuroimaging and the neuroanatomy of obsessive-compulsive disorder. *Psychiatric Clinics of North America, 23*(3), 563–586. https://doi.org/10.1016/S0193-953X(05)70181-7

Schore, A. N. (2001). Effects of a secure attachment relationship on right brain development, affect regulation, and infant mental health. *Infant Mental Health Journal: Infancy and Early Childhood, 22*(1-2), 7–66. https://doi.org/10.1002/1097-0355(200101/04)22:1<7::AID-IMHJ2>3.0.CO;2-N

Schultz, W. (1997). Dopamine neurons and their role in reward mechanisms. *Current Opinion in Neurobiology, 7*(2), 191–197. https://doi.org/10.1016/S0959-4388(97)80007-4

Solms, M. (2021). *The hidden spring: A journey to the source of consciousness*. W. W. Norton & Company.

Sommer, I. E. C., Kleijer, H., Visser, L., & van Laar, T. (2018). Successful treatment of intractable visual hallucinations with 5-HT2A antagonist ketanserin. *BMJ Case Reports, 2018*, bcr-2018-224340. https://doi.org/10.1136/bcr-2018-224340

Sprouse, J. S., & Aghajanian, G. K. (1987). Electrophysiological responses of serotoninergic dorsal raphe neurons to 5-HT1A and 5-HT1B agonists. *Synapse, 1*(1), 3–9. https://doi.org/10.1002/syn.890010103

Squire, L. R. (2004). Memory systems of the brain: A brief history and current perspective. *Neurobiology of Learning and Memory, 82*(3), 171–177. https://doi.org/10.1016/j.nlm.2004.06.005

Stahl, S. M. (2013). *Stahl's essential psychopharmacology: Neuroscientific basis and practical applications* (4th ed.). Cambridge University Press.

Sterling, P., & Eyer, J. (1988). Allostasis: A new paradigm to explain arousal pathology. In S. Fisher & J. Reason (Eds.), *Handbook of life stress, cognition and health* (pp. 629–649). Wiley.

Stoliker, D., Novelli, L., Vollenweider, F. X., Egan, G. F., Preller, K. H., & Razi, A. (2024). Neural mechanisms of resting-state networks and the amygdala underlying the cognitive and emotional effects of psilocybin. *Biological Psychiatry, 96*(1), 57–66. https://doi.org/10.1016/j.biopsych.2024.01.002

Szechtman, H., & Woody, E. (2004). Obsessive-compulsive disorder as a disturbance of security motivation. *Psychological Review, 111*(1), 111–127. https://doi.org/10.1037/0033-295X.111.1.111

Wiecki, T. V., & Frank, M. J. (2013). A computational model of inhibitory control in frontal cortex and basal ganglia. *Psychological Review, 120*(2), 329–355. https://doi.org/10.1037/a0031542

Yin, H. H., & Knowlton, B. J. (2006). The role of the basal ganglia in habit formation. *Nature Reviews Neuroscience, 7*(6), 464–476. https://doi.org/10.1038/nrn1919

Yu, A. J., & Dayan, P. (2005). Uncertainty, neuromodulation, and attention. *Neuron, 46*(4), 681–692. https://doi.org/10.1016/j.neuron.2005.04.026

Correspondence:
Henriette Löffler-Stastka
Department of Psychoanalysis and Psychotherapy,
Währinger Gürtel 18-20, Medical University Vienna,
A-1090 Vienna, Austria
henriette.loeffler-stastka@meduniwien.ac.at

Published under
https://creativecommons.org/licenses/by-nc-nd/4.0/