https://doi.org/10.5513/MSNE2250

Original Research Article

CODEN: PJZSAZ e-ISSN: 1331-7776

Effects of *Stevia rebaudiana* and *Hippophae rhamnoides* on dairy cow metabolism and milk quality

Radena Nenova*, Stanimir Enchev

Agricultural Academy - Sofia, Agricultural Institute - Shumen, Bulgaria

Correspondence: Radena Nenova (radena_nenova@abv.bg)

ARTICLE INFO

Received: 17 July 2026
Revised: 23 August 2026
Accepted: 25 August 2026
Available online: 25 August 2026

Academic Editor:
Prof. Marina Piria, University of Zagreb Faculty of Agriculture, Croatia

Keywords:
Stevia rebaudiana
Hippophae rhamnoides
Phytogenic additives
Dairy cows
Serum biochemistry

ABSTRACT

The present study aimed to evaluate the effects of the phytogenic feed additives *Stevia rebaudiana* Bertoni and *Hippophae rhamnoides* on the serum biochemical profile and physicochemical characteristics of milk in dairy cows. Thirty clinically healthy Black-and-white dairy cows were assigned to three experimental groups (Each group n = 10) using the analogue-group method: a control group (C), a *Stevia rebaudiana* group (SR) receiving 100 g/day of dried stevia leaves, and a *Hippophae rhamnoides* group (HR) receiving 100 g/day of dried sea buckthorn leaves over a period of 30 days. No formal randomisation procedure was applied. Serum concentrations of alanine aminotransferase (ALT), aspartate aminotransferase (AST), urea, glucose, total cholesterol, high-density lipoprotein (HDL) cholesterol, low-density lipoprotein (LDL) cholesterol, and triglycerides were determined on Days 0 and 30, while milk was analysed on Day 30. Supplementation had no statistically significant effect on milk fat, protein, lactose, total solids, solids-not-fat, density, pH, titratable acidity, ash content, or freezing point ($p > 0.05$). No significant changes were observed in serum AST activity, urea concentration, or glucose levels. Although ALT activity increased significantly in both supplemented groups, the recorded values remained within physiological reference limits. Both supplements had a positive effect by reducing total cholesterol, LDL cholesterol, and triglyceride concentrations while increasing HDL cholesterol levels. In this study, the tested supplements were not associated with adverse changes in the measured biochemical variables or physicochemical characteristics of milk and were associated with a more favourable serum lipid profile.

How to Cite

Nenova R., Enchev S. (2026). Effects of *Stevia rebaudiana* and *Hippophae rhamnoides* on dairy cow metabolism and milk quality. Agric Conspec Sci 91: e012.

INTRODUCTION

The production of high-quality milk while maintaining optimal animal health and reducing the use of antimicrobial agents has become a major priority in modern dairy farming. The intensification of dairy production, together with the increasing demands for high milk yield, imposes substantial metabolic stress on dairy cows, particularly during early and mid-lactation. As a consequence, increased generation of reactive oxygen species (ROS), disturbances in lipid and energy metabolism, alterations in hepatic function, and impairment of natural defence mechanisms may occur. These physiological changes can adversely affect animal health, productive performance, and milk quality (Sordillo & Aitken, 2009). Meanwhile, the growing prevalence of antimicrobial resistance (AMR) has emerged as one of the greatest challenges facing public health, veterinary medicine, and food production. The One Health concept recognises the close interconnection between human, animal, and environmental health, emphasising the need to reduce antimicrobial use through effective preventive strategies. In line with this approach, the European Union Veterinary Medicinal

© 2026 The Authors. Agriculturae Conspectus Scientificus by the University of Zagreb Faculty of Agriculture.
This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Products Regulation (Regulation (EU) 2019/6) promotes the development and application of innovative natural alternatives aimed at improving animal health while decreasing reliance on antimicrobial agents (European Parliament and Council, 2019). Within this framework, phytogenic feed additives have attracted increasing attention as safe and sustainable nutritional strategies for supporting animal health and productive performance.

Phytogenic additives comprise plant materials, extracts, or their bioactive constituents, which are rich in polyphenols, flavonoids, terpenes, organic acids, essential oils, and other secondary metabolites with well-documented biological activity. Numerous experimental and review studies have demonstrated that these compounds enhance antioxidant capacity, modulate immune responses, support liver function, improve lipid metabolism, and positively influence ruminal fermentation (Kholif & Olafadehan, 2021). Furthermore, phytogenic feed additives have been reported to improve nutrient utilization efficiency, metabolic performance, and certain indicators of milk production, highlighting their potential as valuable tools for sustainable ruminant nutrition (Windisch et al., 2008; Hashemi & Davoodi, 2011; Kholif & Olafadehan, 2021; Nastoh et al., 2024). Among the phytogenic plants with considerable potential for application in dairy cattle nutrition, *Stevia rebaudiana* (Bertoni) Bertoni (stevia) has received particular attention. In addition to its widespread use as a natural sweetener, *S. rebaudiana* represents a rich source of steviol glycosides, phenolic acids, flavonoids, vitamins, and other antioxidant compounds. These bioactive constituents have been shown to possess antioxidant, anti-inflammatory, hypolipidemic, hepatoprotective, and immunomodulatory properties by reducing lipid peroxidation, enhancing cellular antioxidant defence systems, and supporting normal hepatic function (Chatsudthipong & Muanprasat, 2009; Lemus-Mondaca et al., 2012; Gupta et al., 2013; Nenova et al., 2025).

In livestock species, dietary supplementation with stevia-derived products has been associated with improved antioxidant status, a more favourable lipid profile, and enhanced metabolic adaptation to physiological stress (Pirgozliev et al., 2021, 2022; Xiong et al., 2022; Nenova & Enchev, 2022). Another promising phytogenic source is *Hippophae rhamnoides* L. (sea buckthorn), which is characterised by an exceptionally rich phytochemical composition. Its leaves and berries contain substantial amounts of vitamin C, tocopherols, carotenoids, polyphenols, flavonoids, phytosterols, and essential fatty acids. Owing to this complex composition, *H. rhamnoides* exhibits pronounced antioxidant, anti-inflammatory, immunomodulatory, and hepatoprotective properties. Experimental studies have demonstrated that dietary supplementation with *H. rhamnoides* in farm animals improves antioxidant status, stabilises lipid metabolism, enhances resilience to physiological stress, and may positively influence productive performance (Beveridge et al., 1999; Suryakumar & Gupta, 2011).

Despite these promising findings, published in vivo studies evaluating the effects of *S. rebaudiana* and *H. rhamnoides* in dairy cows remain limited. Consequently, their influence on serum biochemical parameters and the physicochemical characteristics of milk has not yet been comprehensively established. Additional controlled studies under commercial production conditions are therefore required to evaluate the potential of these phytogenic plants as functional feed additives in sustainable dairy production systems.

The aim of the present study was to evaluate the effects of dietary supplementation with *S. rebaudiana* and *H. rhamnoides* on the serum biochemical profile and physicochemical characteristics of milk in Black-and-white dairy cows.

MATERIALS AND METHODS

Experimental animals and housing conditions

The experiment was conducted at the experimental dairy farm of the Agricultural Institute–Shumen in northeast Bulgaria (43.28° N, 26.93° E). Thirty clinically healthy Black-and-white dairy cows, comparable in lactation stage, age, body weight, and milk yield, were included in the study. All animals were housed under identical management and environmental conditions in a free-stall barn with unrestricted access to fresh drinking water (*ad libitum*). Cows were milked twice daily according to the routine management practices of the farm. Throughout the experimental period, all animals received the same basal diet consisting of 20 kg corn silage, 2 kg alfalfa hay, 2 kg wheat straw, and 5 kg concentrate per cow per day. The concentrate mixture consisted of ground corn (40%), barley (33%), wheat bran (15.9%), corn meal (9%), limestone (1.1%), salt (0.5%), and a vitamin–mineral premix (0.5%). The ration supplied 13.47 milk feed units (MFU) and 1,228 g crude protein per cow per day.

Experimental design

Cows were assigned to three experimental groups (n = 10 per group) using the analogue-group method. Animals were matched as closely as possible according to age, lactation stage, body weight, and milk production and were then allocated to the control, *S. rebaudiana*, or *H. rhamnoides* group to obtain comparable baseline groups. No formal randomisation procedure was applied.

- **Control group (C):** cows received only the basal diet.
- ***S. rebaudiana* group (SR):** cows received the basal diet supplemented with 100 g/day dried and ground *S. rebaudiana* leaves.
- ***H. rhamnoides* group (HR):** cows received the basal diet supplemented with 100 g/day dried and ground *H. rhamnoides* leaves.

The experimental period lasted 30 consecutive days.

Plant material

Dried leaves of *S. rebaudiana* and *H. rhamnoides* were prepared using a standardised processing protocol. Plant material was harvested at maximum physiological maturity and naturally dried under controlled environmental conditions, before milling to obtain a homogeneous particle size (0.25–3.0 mm). The residual moisture content was maintained below 10%, ensuring the stability of bioactive compounds throughout the experimental period. The selection of both plant species was based on their high concentrations of phenolic compounds, flavonoids, carotenoids, and other antioxidant constituents that have been reported to exert beneficial effects on the metabolic and physiological status of farm animals.

© 2026 The Authors. Agriculturae Conspectus Scientificus by the University of Zagreb Faculty of Agriculture.
This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Blood sampling and biochemical analysis

Blood samples were collected from the jugular vein of each animal on Day 0 (prior to dietary supplementation) and Day 30 (end of the experimental period). Samples were collected into vacuum tubes without anticoagulant, centrifuged at 3,000 rpm for 10 min to obtain serum, and stored at -20°C until analysis. Serum biochemical analyses were performed using an EXICO C200 automated biochemical analyser (EXICO Medical Systems, Sweden) with commercially available diagnostic reagent kits, according to the analyser and reagent manufacturers' instructions. The following parameters were determined: alanine aminotransferase (ALT), aspartate aminotransferase (AST), glucose, urea, total cholesterol, high-density lipoprotein (HDL) cholesterol, low-density lipoprotein (LDL) cholesterol, and triglycerides.

Milk sampling and analysis

Individual milk samples were collected on Day 30 during morning milking. After thorough homogenisation of the milk obtained from each cow, a 100 mL sample was collected, transported at 4°C, and analysed within 24 h. Milk fat, protein, lactose, total solids, solids-not-fat (SNF), density, pH, titratable acidity, ash content, and freezing point were determined using a calibrated MilkoScan infrared milk analyser (FOSS, Hillerød, Denmark), according to the manufacturer's operating and calibration procedures. Ash content and freezing point were recorded from the analyser outputs generated using its validated calibration equations.

Statistical analysis

Statistical analyses were performed using STATISTICA version 13.0 (TIBCO Software Inc., Palo Alto, CA, USA). Data are presented as mean ± standard deviation (SD). The normality of data distribution was assessed using the Shapiro-Wilk test. For serum biochemical variables measured on Days 0 and 30, the effects of treatment (Control, SR and HR groups), time, and their interaction (treatment × time) were evaluated using a linear mixed-effects model (LMM), with animal identity included as a random effect to account for within-animal correlation. When significant fixed effects were detected, pairwise comparisons were performed using Tukey-adjusted *post hoc* tests. Because milk was sampled only once, on Day 30, milk variables were analysed by one-way analysis of variance (ANOVA), with treatment as the fixed factor, followed by Tukey's *post hoc* test. Statistical significance was accepted at $p < 0.05$.

RESULTS

The effects of dietary supplementation with *S. rebaudiana* and *H. rhamnoides* on the serum biochemical profile and the physicochemical characteristics of milk in Black-and-white dairy cows were evaluated after a 30-day feeding period. Serum biochemical responses are presented in Tables 1 and 2, whereas the Day 30 milk characteristics are summarised in Table 3. Table 1 presents liver enzyme activities (ALT and AST) and indicators of energy and protein metabolism (glucose and urea) on Days 0 and 30. Table 2 summarises the serum lipid profile on Days 0 and 30, while Table 3 presents the physicochemical characteristics of milk measured on Day 30.

Table 1. Effects of dietary supplementation with *Stevia rebaudiana* Bertoni or *Hippophae rhamnoides* L. on liver enzymes and indicators of energy and protein metabolism in Black-and-white dairy cow serum

Variable	Control Day 0	Control Day 30	SR Day 0	SR Day 30	HR Day 0	HR Day 30	Treatment P	Time P	Treatment × Time P
ALT (U/L)	29.00 ± 1.30^{a}	29.50 ± 1.35^{a}	28.50 ± 1.20^{a}	39.60 ± 1.55^{b}	28.80 ± 1.22^{a}	40.20 ± 1.58^{b}	>0.05	>0.05	<0.001
AST (U/L)	133.00 ± 10.50	132.50 ± 10.20	132.00 ± 10.50	128.50 ± 9.80	132.80 ± 10.60	128.80 ± 9.60	>0.05	>0.05	>0.05
Glucose (mmol/L)	3.30 ± 0.30	3.32 ± 0.29	3.30 ± 0.30	3.45 ± 0.28	3.29 ± 0.30	3.48 ± 0.28	>0.05	>0.05	>0.05
Urea (mmol/L)	3.05 ± 0.30	3.10 ± 0.31	3.10 ± 0.32	3.05 ± 0.30	3.08 ± 0.30	3.02 ± 0.28	>0.05	>0.05	>0.05

Values are presented as mean ± SD (n = 10 cows per group). Statistical analysis was performed using a linear mixed-effects model with treatment, time and treatment × time interaction as fixed effects and animal as a random effect. a,b Values within a variable not sharing the same superscript differ significantly ($p < 0.05$). SR = *Stevia rebaudiana*; HR = *Hippophae rhamnoides*.

Table 2. Effects of dietary supplementation with *Stevia rebaudiana* Bertoni or *Hippophae rhamnoides* L. on the serum lipid profile of Black-and-white dairy cows

Variable	Control Day 0	Control Day 30	SR Day 0	SR Day 30	HR Day 0	HR Day 30	Treatment P	Time P	Treatment × Time P
Total cholesterol (mmol/L)	2.20 ± 0.19^{b}	2.18 ± 0.18^{b}	2.10 ± 0.18^{b}	1.80 ± 0.16^{a}	2.11 ± 0.18^{b}	1.82 ± 0.16^{a}	>0.05	>0.05	<0.001
Triglycerides (mmol/L)	0.24 ± 0.03^{b}	0.25 ± 0.03^{b}	0.25 ± 0.03^{b}	0.20 ± 0.02^{a}	0.24 ± 0.03^{b}	0.19 ± 0.02^{a}	>0.05	>0.05	<0.001
HDL cholesterol (mmol/L)	1.38 ± 0.12^{a}	1.37 ± 0.11^{a}	1.37 ± 0.11^{a}	1.52 ± 0.12^{b}	1.36 ± 0.11^{a}	1.54 ± 0.12^{b}	>0.05	>0.05	<0.001
LDL cholesterol (mmol/L)	0.66 ± 0.07^{b}	0.65 ± 0.07^{b}	0.64 ± 0.07^{b}	0.46 ± 0.05^{a}	0.65 ± 0.07^{b}	0.45 ± 0.05^{a}	>0.05	>0.05	<0.001

Values are presented as mean ± SD (n = 10 cows per group). Statistical analysis was performed using a linear mixed-effects model with treatment, time and treatment × time interaction as fixed effects and animal as a random effect. a,b Values within a variable not sharing the same superscript differ significantly ($p < 0.05$). SR = *Stevia rebaudiana*; HR = *Hippophae rhamnoides*.

© 2026 The Authors. Agriculturae Conspectus Scientificus by the University of Zagreb Faculty of Agriculture.
This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Table 3. Physicochemical characteristics of milk after 30-day dietary supplementation with *Stevia rebaudiana* Bertoni or *Hippophae rhamnoides* L. in Black-and-white dairy cows

Parameter	Control (n = 10)	SR (n = 10)	HR (n = 10)	*p*
Milk fat (%)	3.74 ± 0.23	3.78 ± 0.21	3.81 ± 0.20	0.74
Protein (%)	3.28 ± 0.11	3.31 ± 0.10	3.33 ± 0.12	0.69
Lactose (%)	4.71 ± 0.08	4.73 ± 0.07	4.74 ± 0.08	0.61
Total solids (%)	12.43 ± 0.32	12.49 ± 0.30	12.55 ± 0.29	0.72
Solids-not-fat (%)	8.69 ± 0.12	8.71 ± 0.11	8.73 ± 0.10	0.76
Density (g/cm^3)	1.0302 ± 0.0007	1.0304 ± 0.0006	1.0305 ± 0.0007	0.64
pH	6.67 ± 0.04	6.68 ± 0.03	6.68 ± 0.04	0.82
Titratable acidity (°T)	16.6 ± 0.6	16.5 ± 0.5	16.4 ± 0.5	0.57
Ash (%)	0.72 ± 0.02	0.72 ± 0.02	0.73 ± 0.02	0.81
Freezing point (°C)	−0.536 ± 0.005	−0.537 ± 0.004	−0.538 ± 0.005	0.70

Values are presented as mean ± SD (n = 10 cows per group) measured on Day 30. Data were analysed by one-way ANOVA followed by Tukey's *post hoc* test. No statistically significant differences were observed among groups ($p > 0.05$). SR = *Stevia rebaudiana* group; HR = *Hippophae rhamnoides* group.

Serum variables were evaluated using a linear mixed-effects model with treatment, time, and treatment × time interaction as fixed effects and animal as a random effect. Day 30 milk variables were evaluated by one-way ANOVA followed by Tukey's *post hoc* test.

DISCUSSION

The most pronounced metabolic response to dietary supplementation was observed in the serum lipid profile. Both *S. rebaudiana* and *H. rhamnoides* significantly reduced total cholesterol, LDL cholesterol and triglyceride concentrations while increasing HDL cholesterol. According to the linear mixed-effects analysis, significant treatment × time interactions were detected for all evaluated lipid variables, indicating that these changes were attributable to the dietary treatments rather than to temporal variation alone. The improvement in lipid metabolism occurred without concomitant disturbances in liver enzymes or protein metabolism, suggesting that the investigated phytogenic additives exerted targeted metabolic effects while maintaining overall physiological homeostasis.

The hypolipidemic effects observed in the present study are consistent with the known biological activities of stevia and sea buckthorn. Both plants are rich sources of polyphenols, flavonoids and other bioactive compounds that regulate lipid metabolism through multiple mechanisms, including suppression of hepatic cholesterol synthesis, stimulation of fatty acid β-oxidation, enhancement of reverse cholesterol transport and reduction of oxidative stress (Chatsudthipong & Muanprasat, 2009; Suryakumar & Gupta, 2011; Ramos-Tovar & Muriel, 2017; Ramos-Tovar et al., 2018). Experimental studies have also demonstrated that supplementation with stevia-derived products improves antioxidant capacity while reducing serum cholesterol and triglyceride concentrations in livestock species (Xiong et al., 2022; Nenova et al., 2025). Similarly, the lipid-modulating properties of *H. rhamnoides* have been attributed to its high content of flavonoids, carotenoids, tocopherols, phytosterols and unsaturated fatty acids, which collectively contribute to improved lipid metabolism and protection against oxidative damage (Wang et al., 2022).

Taken together, these findings indicate that both phytogenic feed additives improved lipid metabolism without adverse changes in the measured indicators of liver function or general metabolic status during the 30-day study. These responses support further investigation of plant-derived bioactive compounds as nutritional strategies for supporting metabolic health in dairy cattle. Dietary supplementation with *S. rebaudiana* and *H. rhamnoides* did not significantly affect the physicochemical characteristics of milk measured on Day 30. Milk fat, protein, lactose, total solids, solids-not-fat, density, pH, titratable acidity, ash content, and freezing point remained within the normal physiological ranges for clinically healthy dairy cows.

The absence of significant changes in milk composition is consistent with previous reports demonstrating that phytogenic feed additives primarily influence metabolic efficiency, antioxidant status and animal health rather than the basic chemical composition of milk (Windisch et al., 2008; Steiner, 2009). Similarly, Paglia et al. (2021) reported that supplementation with a phytogenic blend containing cardol, cardanol and ricinoleic acid did not significantly alter the principal physicochemical characteristics of milk. Kalaitsidis et al. (2025) observed that dietary supplementation with *S. rebaudiana* improved oxidative stability and modified the milk fatty acid profile without affecting milk fat, protein or lactose concentrations. These observations suggest that the biological activity of stevia is expressed mainly through modulation of metabolic and antioxidant pathways rather than changes in the gross composition of milk.

The present study demonstrated that 30-day dietary supplementation with *S. rebaudiana* and *H. rhamnoides* was not associated with adverse changes in the measured biochemical variables or the physicochemical characteristics of milk evaluated on Day 30. Significant treatment × time interactions

© 2026 The Authors. Agriculturae Conspectus Scientificus by the University of Zagreb Faculty of Agriculture.
This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

were observed for ALT and the serum lipid profile, whereas no significant effects were detected for AST, glucose, or urea. Although ALT activity increased in both supplemented groups, the values remained within physiological reference limits (Kaneko et al., 2008; Constable et al., 2017).

The maintenance of stable serum urea concentrations throughout the experimental period further suggests that dietary supplementation did not interfere with nitrogen metabolism or protein utilisation. Serum urea is widely used as an indicator of the dietary protein-to-energy balance and ruminal nitrogen utilisation efficiency, and values within the normal physiological range indicate adequate nutritional status and efficient microbial protein synthesis (National Academies of Sciences, Engineering, and Medicine, 2021). The slight increase in serum glucose observed in the supplemented groups was not statistically significant, indicating that the phytogenic additives did not markedly influence carbohydrate metabolism under the conditions of the present study. Similar observations have been reported for phytogenic feed additives rich in polyphenols and flavonoids, which improve metabolic efficiency and antioxidant status without causing clinically relevant alterations in routine serum biochemical variables (Windisch et al., 2008; Steiner, 2009).

Published information regarding the use of *H. rhamnoides* in dairy cattle remains limited. Nevertheless, the plant is recognised as an exceptionally rich source of flavonoids, carotenoids, tocopherols and other phenolic compounds with well-documented antioxidant and hypolipidemic properties (Suryakumar & Gupta, 2011; Wang et al., 2022). The present findings indicate that supplementation with sea buckthorn maintained normal milk composition while contributing to an improved serum lipid profile, suggesting that its primary effects are associated with metabolic regulation rather than direct modification of milk constituents.

The improvement in lipid metabolism observed in the present study, together with the absence of adverse changes in the measured liver-related variables and unchanged milk quality on Day 30 supports further evaluation of both plant species as functional phytogenic feed additives for dairy cattle. Longer studies with larger animal populations and additional health and performance endpoints are required before broader conclusions regarding their practical use can be drawn.

Several limitations of the present study should be acknowledged. The feeding trial was limited to 30 days and involved a relatively small number of animals, which may have reduced the ability to detect subtle physiological responses. Furthermore, only routine biochemical indicators and physicochemical characteristics of milk were evaluated. Future studies should include longer experimental periods, larger animal populations and additional biomarkers related to oxidative status, inflammatory response, rumen fermentation and milk fatty acid composition in order to further elucidate the mechanisms underlying the biological effects of *S. rebaudiana* and *H. rhamnoides*.

Overall, dietary supplementation with *S. rebaudiana* and *H. rhamnoides* for 30 days was associated with an improved serum lipid profile without adverse changes in the measured liver-related, protein metabolism, or milk physicochemical variables. These findings are limited to the variables and duration evaluated and warrant confirmation in longer and larger controlled studies.

CONCLUSIONS

The present study demonstrated that 30-day dietary supplementation with *S. rebaudiana* and *H. rhamnoides* was safe and well tolerated in Black-and-white dairy cows. Supplementation did not adversely affect liver function, protein metabolism or the physicochemical characteristics of milk, and all evaluated variables remained within physiological reference ranges. According to the linear mixed-effects analysis, significant treatment × time interactions were observed for the serum lipid profile, whereas no significant effects were detected for AST, glucose or urea.

The most important finding was the improvement in the serum lipid profile, characterised by reduced concentrations of total cholesterol, LDL cholesterol, and triglycerides, together with increased HDL cholesterol levels. These changes occurred without adverse alterations in other measured biochemical variables or Day 30 milk characteristics.

The present findings expand the limited knowledge regarding the use of *S. rebaudiana* and *H. rhamnoides* in dairy cattle nutrition. Nevertheless, the conclusions are restricted to the measured variables and the 30-day experimental period. Studies involving larger animal populations, longer supplementation periods, different stages of lactation, and additional indicators of oxidative status, inflammation, animal performance, and milk composition are required to confirm these observations.

ACKNOWLEDGEMENTS

The study is part of ongoing work within the framework of COST (European Cooperation in Science and Technology) project CA22109 "MEDICINAL PLANTS FOR ANIMAL HEALTH CARE: TRANSFORMING TRADITION INTO MODERN VETERINARY MEDICINE (MEDPLANTS4VET)", and also this document is part of the project KP-06-COST/23/16.12.2024. "Development of a new protective and therapeutic system based on natural functional components/ stevia and sea buckthorn/with antioxidant and prebiotic activity, as an alternative to antibiotic preparations" funded by FNI - Ministry of Education and Science, Republic of Bulgaria.

CRediT AUTHORSHIP CONTRIBUTION STATEMENT

Radena Nenova: Conceptualization, Methodology, Investigation, Data curation, Formal analysis, Visualization, Writing – original draft, Writing – review & editing; **Stanimir Enchev:** Supervision, Methodology, Validation, Resources, Writing – review & editing; All authors have read and approved the final version of the manuscript.

DECLARATION OF COMPETING INTEREST

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

ETHICS AND PERMIT APPROVALS

All procedures involving animals were conducted in accordance with the applicable Bulgarian and European Union legislation on animal welfare and with the institutional animal-care guidelines of the Agricultural Institute - Shumen. The trial was conducted under standard farm management conditions. Jugular blood samples were collected by a qualified veterinarian

© 2026 The Authors. Agriculturae Conspectus Scientificus by the University of Zagreb Faculty of Agriculture.
This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

using routine aseptic venipuncture and appropriate restraint, and milk samples were obtained during routine morning milking.

DATA AVAILABILITY STATEMENT

The data supporting the findings of this study are available from the corresponding author upon reasonable request.

FUNDING STATEMENT

This research was supported by the COST (European Cooperation in Science and Technology) Action CA22109 – *Medicinal Plants for Animal Health Care: Transforming Tradition into Modern Veterinary Medicine (MEDPLANTS4VET)*. This work was also supported by the Bulgarian National Science Fund (NSF), Ministry of Education and Science, Republic of Bulgaria, under Project KP-06-COST/23/16.12.2024, entitled *"Development of a novel protective and therapeutic system based on natural functional components (stevia and sea buckthorn) with antioxidant and prebiotic activity as an alternative to antibiotic preparations"*.

AI USE STATEMENT

None.

SAŽETAK

Učinci stevije (*Stevia rebaudiana* Bertoni) i pasjeg trna (*Hippophae rhamnoides* L.) na metabolizam mliječnih krava i kvalitetu mlijeka

Cilj ovog istraživanja bio je procijeniti učinke biljnih hranidbenih dodataka, *Stevia rebaudiana* Bertoni i *Hippophae rhamnoides* L. na biokemijski profil seruma i fizikalno-kemijska svojstva mlijeka mliječnih krava. Trideset klinički zdravih crno-bijelih mliječnih krava raspoređeno je metodom analognih skupina u tri pokusne skupine (svaka grupa n = 10): kontrolnu skupinu (C), skupinu *Stevia rebaudiana* (SR), koja je primala 100 g osušenih listova stevije dnevno, te skupinu *Hippophae rhamnoides* (HR), koja je primala 100 g osušenih listova pasjeg trna dnevno. Formalni postupak randomizacije nije primijenjen, a pokus je trajao 30 dana. Aktivnosti alanin-aminotransferaze (ALT) i aspartat-aminotransferaze (AST) te koncentracije ureje, glukoze, ukupnog kolesterola, HDL-kolesterola, LDL-kolesterola i triglicerida u serumu određene su na početku pokusa (0. dan) i 30. dana pokusa, dok su svojstva mlijeka analizirana 30. dana.

Hranidbeni dodatci nisu značajno utjecali na sadržaj mliječne masti, proteina, laktoze, ukupne i bezmasne suhe tvari, gustoću, pH, titracijsku kiselost, sadržaj pepela ni točku smrzavanja mlijeka ($p > 0,05$). Također, nisu utvrđene značajne promjene aktivnosti AST-a ni koncentracija ureje i glukoze u serumu.

Iako se aktivnost ALT-a značajno povećala u objema skupinama koje su primale dodatke, sve su zabilježene vrijednosti ostale unutar fizioloških referentnih granica. Oba dodatka imala su pozitivan učinak, smanjujući koncentracije ukupnog kolesterola, LDL-kolesterola i triglicerida te povećavajući koncentraciju HDL-kolesterola ($p < 0,05–0,01$). U uvjetima ovog 30-dnevnog istraživanja, primjena dodataka nije bila povezana s nepovoljnim promjenama u izmjerenim biokemijskim pokazateljima ni fizikalno-kemijskim svojstvima mlijeka, dok je bila povezana s povoljnijim lipidnim profilom seruma.

Ključne riječi: *Stevia rebaudiana*, *Hippophae rhamnoides*, biljni hranidbeni dodaci, mliječne krave, biokemija seruma

REFERENCES

Beveridge T., Li T.S.C., Oomah B.D., Smith A. (1999). Sea buckthorn products: Manufacture and composition. J Agric Food Chem 47(9): 3480–3488. https://doi.org/10.1021/jf981331m

Chatsudthipong V., Muanprasat C. (2009). Stevioside and related compounds: Therapeutic benefits beyond sweetness. Pharmacol Ther. 121(1): 41–54. https://doi.org/10.1016/j.pharmthera.2008.09.007

Constable P.D., Hinchcliff K.W., Done S.H., Grünberg W. (2017). Veterinary Medicine: A Textbook of the Diseases of Cattle, Horses, Sheep, Pigs and Goats. 11th Edition. Elsevier, St. Louis, MO, USA.

European Parliament and Council (2019). Regulation (EU) 2019/6 of the European Parliament and of the Council of 11 December 2018 on veterinary medicinal products and repealing Directive 2001/82/EC. Off J Eur Union L4: 43–167.

Gupta E., Purwar S., Sundaram S., Rai G.K. (2013). Nutritional and therapeutic values of *Stevia rebaudiana*: A review. J Med Plants Res 7(46): 3343–3353. Doi: 10.5897/JMPR2013.5276

Hashemi S.R., Davoodi H. (2011). Herbal plants and their derivatives as growth and health promoters in animal nutrition. Vet Res Commun 35(3): 169–180. https://doi.org/10.1007/s11259-010-9458-2

Kalaitsidis K., Parissi Z., Theodoridis A., Savvidou S., Kotsampasi B., Vasilopoulou K., Dokou S., Chantzi P., Tsaliki E., Lazari D., Valergakis G. E., Giannenas I. (2025). Effects of *Stevia rebaudiana* and hemp by-products (*Cannabis sativa*) on milk yield, fatty acid profile, and oxidative stability in dairy cows. J Dairy Res 92(1), 26–31. https://doi.org/10.1017/S0022029925000275

Kaneko J.J., Harvey J.W., Bruss M.L. (2008). Clinical Biochemistry of Domestic Animals. 6th Edition. Academic Press, Burlington, MA, USA.

Kholif A. E., Olafadehan O. A. (2021). Essential oils and phytogenic feed additives in ruminant diets: Chemistry, ruminal microbiota and fermentation, feed utilization and productive performance. Phytochem Rev 20, 1087–1108. https://doi.org/10.1007/s11101-021-09739-3

Lemus-Mondaca R., Vega-Gálvez A., Zura-Bravo L., Ah-Hen K. (2012). *Stevia rebaudiana* Bertoni, source of a high-potency natural sweetener: A comprehensive review on the biochemical, nutritional and functional aspects. Food Chem 132(3): 1121–1132. https://doi.org/10.1016/j.foodchem.2011.11.140

National Academies of Sciences, Engineering, and Medicine. (2021). Nutrient Requirements of Dairy Cattle. 8th rev. ed. National Academies Press, Washington, DC, USA. https://doi.org/10.17226/25806

Nastoh N. A., Waqas M., Çınar A. A., Salman M. (2024). The impact of phytogenic feed additives on ruminant production: A review. J Anim Feed Sci 33(4): 431–453. https://doi.org/10.22358/jafs/191479/2024

Nenova R., Enchev S. (2022). Influence of the inclusion of Stevia (*Stevia rebaudiana* Bertoni) in the compound feed of lactating sows on their haematological parameters. J Mt Agric Balk 25(1): 1–14.

Nenova R., Penchev P., Enchev S., Ilieva Y., Schreiner M. (2025). Stevia as a regulator of antioxidative and biochemical status in the buffalo and its effect on lactic fatty acids. Himera Ltd., Shumen, Bulgaria. ISBN 978-619-7745-23-8.

Paglia D. K., Collet S. G., Camillo G., Prestes A. M., Cony A. V., Pazinato F. M., Girardini L. K. (2021). Effect of a phytogenic feed additives mixture on milk physico-chemical properties and biochemical parameters of Holstein cows. Cienc Rural 51(12): e20200682. https://doi.org/10.1590/0103-8478cr20200682

Pirgozliev V., Kljak K., Whiting I. M., Rose S. P., Mansbridge S. C., Enchev S., Atanasov A., Stringhini J. H. (2021). Feeding dry stevia leaf (*Stevia rebaudiana*) or xylanase improves the hepatic antioxidative status of broiler chickens. Res Vet Sci 136: 227–229. https://doi.org/10.1016/j.rvsc.2021.03.001

© 2026 The Authors. Agriculturae Conspectus Scientificus by the University of Zagreb Faculty of Agriculture.
This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Pirgozliev V. R., Whiting I. M., Kljak K., Mansbridge S. C., Atanasov A. G., Rose S. P., Enchev S. B. (2022). Stevia (*Stevia rebaudiana*) improves carotenoid content in eggs when fed to laying hens. Foods 11(10): 1418. https://doi.org/10.3390/foods11101418

Ramos-Tovar E., Muriel P. (2017). Stevia as a putative hepatoprotector. In: Liver Pathophysiology: Therapies and Antioxidants (Muriel P., ed.), Academic Press, London, United Kingdom, pp. 715–727. https://doi.org/10.1016/B978-0-12-804274-8.00051-5

Ramos-Tovar E., Hernández-Aquino E., Casas-Grajales S., Buendia-Montaño L. D., Galindo-Gómez S., Camacho J., Tsutsumi V., Muriel P. (2018). Stevia prevents acute and chronic liver injury induced by carbon tetrachloride by blocking oxidative stress through Nrf2 upregulation. Oxid Med Cell Longev 2018: 3823426. https://doi.org/10.1155/2018/3823426

Sordillo L. M., Aitken S. L. (2009). Impact of oxidative stress on the health and immune function of dairy cattle. Vet Immunol Immunopathol 128(1–3): 104–109. https://doi.org/10.1016/j.vetimm.2008.10.305

Steiner T. (ed.) (2009). Phytogenics in Animal Nutrition: Natural Concepts to Optimize Gut Health and Performance. Nottingham University Press, Nottingham, United Kingdom.

Suryakumar G., Gupta A. (2011). Medicinal and therapeutic potential of sea buckthorn (*Hippophae rhamnoides* L.). J Ethnopharm 138(2): 268–278. https://doi.org/10.1016/j.jep.2011.09.024

Wang Z., Zhao F., Wei P., Chai X., Hou G., Meng Q. (2022). Phytochemistry, health benefits, and food applications of sea buckthorn (*Hippophae rhamnoides* L.): A comprehensive review. Front Nutr 9: 1036295. https://doi.org/10.3389/fnut.2022.1036295

Windisch W., Schedle K., Plitzner C., Kroismayr A. (2008). Use of phytogenic products as feed additives for swine and poultry. J Anim Sci 86 (Suppl. 14): E140–E148. https://doi.org/10.2527/jas.2007-0459

Xiong Y., Liu S., Xiao H., Wu Q., Chi L., Zhu L., Fang L., Li Y., Jiang Z., Wang L. (2022). Dietary stevia residue extract supplementation improves the performance and antioxidative capacity of growing-finishing pigs. J Sci Food Agric 102(11): 4724–4735. https://doi.org/10.1002/jsfa.11833

© 2026 The Authors. Agriculturae Conspectus Scientificus by the University of Zagreb Faculty of Agriculture.
This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.