

**ON THE ORDER STRUCTURE ON THE SET OF
COMPLETELY MULTI-POSITIVE LINEAR MAPS ON
 C^* -ALGEBRAS**

MARIA JOIȚA, TANIA-LUMINIȚA COSTACHE AND MARIANA ZAMFIR

University of Bucharest, University "Politehnica" of Bucharest and
Technical University of Civil Engineering Bucharest, Romania

ABSTRACT. In this paper we characterize the order relation on the set of all completely n -positive linear maps on C^* -algebras in terms of the representation associated to each completely n -positive linear map given by Suen's construction.

1. INTRODUCTION AND PRELIMINARIES

Completely positive linear maps are an often used tool in operator algebras theory and quantum information theory [1, 3, 5, 7, 10].

In the mathematical framework of quantum information theory, all admissible devices are modelled by the so-called quantum operations (that is, completely positive linear maps on the algebra of observables (C^* -algebra) of the physical system under consideration). A good analysis of completely multi-positive maps between C^* -algebras involves understanding and solving certain problems in quantum information theory and understanding the infinite dimensional non-commutative structure of topological $*$ -algebras [2, 5, 7, 10]. The theorems on the structure of completely linear maps and Radon-Nikodym type theorems for completely positive linear maps are an extremely powerful and veritable tool for problems involving characterization and comparison of quantum operations.

2000 *Mathematics Subject Classification.* 46L05, 47A20, 47L90.

Key words and phrases. C^* -algebra, completely multi-positive linear map, Radon-Nikodym type theorem, pure completely multi-positive linear map, extremal completely multi-positive linear map.

Given a C^* -algebra A and a positive integer n , we denote by $M_n(A)$ the C^* -algebra of all $n \times n$ matrices over A with the algebraic operations and the topology obtained by regarding it as a direct sum of n^2 copies of A .

DEFINITION 1.1. A linear map $\rho: A \rightarrow B$ between two C^* -algebras is completely positive if the linear maps $\rho^{(n)}: M_n(A) \rightarrow M_n(B)$ defined by

$$\rho^{(n)}([a_{ij}]_{i,j=1}^n) = [\rho(a_{ij})]_{i,j=1}^n$$

are positive for any positive integer n .

DEFINITION 1.2. Let A and B be two C^* -algebras. An $n \times n$ matrix $[\rho_{ij}]_{i,j=1}^n$ of linear maps from A to B can be regarded as a linear map ρ from $M_n(A)$ to $M_n(B)$ defined by

$$\rho([a_{ij}]_{i,j=1}^n) = [\rho_{ij}(a_{ij})]_{i,j=1}^n.$$

We say that $[\rho_{ij}]_{i,j=1}^n$ is a completely n -positive linear map from A to B if ρ is a completely positive linear map from $M_n(A)$ to $M_n(B)$.

We shall denote by $CP_\infty(A, B)$ the set of all completely positive linear maps from A to B and by $CP_\infty^n(A, B)$ the set of all completely n -positive linear maps from A to B .

In [9], Suen showed that any completely n -positive linear map from a C^* -algebra A to $L(H)$, the C^* -algebra of all bounded linear operators on a Hilbert space H , is of the form $[V^*T_{ij}\Phi(\cdot)V]_{i,j=1}^n$, where Φ is a representation of A on a Hilbert space K , $V \in L(H, K)$ and $[T_{ij}]_{i,j=1}^n$ is a positive element in $M_n(\Phi(A)')$ ($\Phi(A)'$ denotes the commutant of $\Phi(A)$ in $L(K)$).

THEOREM 1.3 ([9, 4]). Let A be a C^* -algebra, let H be a Hilbert space and let $\rho = [\rho_{ij}]_{i,j=1}^n$ be a completely n -positive linear map from A to $L(H)$. Then there is a representation Φ_ρ of A on a Hilbert space H_ρ , $V_\rho \in L(H, H_\rho)$ and a positive element $T^\rho = [T_{ij}^\rho]_{i,j=1}^n$ in $M_n(\Phi_\rho(A)')$ with $\sum_{k=1}^n T_{kk}^\rho = nid_{L(H_\rho)}$ such that:

- i. $\{\Phi_\rho(a)V_\rho\xi; a \in A, \xi \in H\}$ spans a dense subspace in H_ρ ;
- ii. $\rho_{ij}(a) = V_\rho^*T_{ij}^\rho\Phi_\rho(a)V_\rho$, for all $a \in A$ and for all $i, j = 1, \dots, n$.

The quadruple $(\Phi_\rho, H_\rho, V_\rho, T^\rho)$ will be called the Suen's construction associated with ρ and it is unique up to unitary equivalence [4, Theorem 2.3].

REMARK 1.4. The triple $(\Phi_\rho, H_\rho, V_\rho)$ is the Stinespring representation associated with $\tilde{\rho} = \frac{1}{n} \sum_{k=1}^n \rho_{kk}$ (see, [4, the proof of Theorem 2.3]).

In this paper we characterize the order relation on the set of all completely n -positive linear maps on C^* -algebras in terms of the representation associated to each completely n -positive linear map given by Suen's construction [9].

We also give sufficient conditions for that a completely n -positive linear map from a unital C^* -algebra A to $L(H)$ to be an extreme point in the set of all completely n -positive linear maps $[\rho_{ij}]_{i,j=1}^n$ from A to $L(H)$ such that $[\rho_{ij}(1_A)]_{i,j=1}^n = T^0$ for some $T^0 \in M_n(L(H))$.

2. THE MAIN RESULTS

Let $\rho = [\rho_{ij}]_{i,j=1}^n$ be an element in $CP_\infty^n(A, L(H))$ and let $(\Phi_\rho, H_\rho, V_\rho, T^\rho)$ be the construction associated to ρ given by Theorem 1.3.

LEMMA 2.1. *Let $S = [S_{ij}]_{i,j=1}^n$ be a positive element in $M_n(\Phi_\rho(A)')$. The map $\rho_S = [\rho_{S_{ij}}]_{i,j=1}^n$ from $M_n(A)$ to $M_n(L(H))$ defined by*

$$\rho_S([a_{ij}]_{i,j=1}^n) = [V_\rho^* S_{ij} \Phi_\rho(a_{ij}) V_\rho]_{i,j=1}^n$$

is a completely n -positive linear map from A to $L(H)$.

PROOF. It is not difficult to see that ρ_S is an $n \times n$ matrix of linear maps from A to $L(H)$ whose (i, j) -entry is the linear map $\rho_{S_{ij}}$ from A to $L(H)$ defined by $\rho_{S_{ij}}(a) = V_\rho^* S_{ij} \Phi_\rho(a) V_\rho$ for all $a \in A$ and for all $i, j = 1, \dots, n$.

To show that ρ_S is a completely n -positive linear map from A to $L(H)$ it is sufficient to show that $\Gamma(\rho_S) \in CP_\infty(A, M_n(L(H)))$, where Γ is the map from $CP_\infty^n(A, B)$ onto $CP_\infty(A, M_n(B))$ defined by $\Gamma([\rho_{ij}]_{i,j=1}^n)(a) = [\rho_{ij}(a)]_{i,j=1}^n$ for all $a \in A$ [2, Theorem 1.4]. For this, let m be a positive integer, $a_1, \dots, a_m \in A, \xi_1 = (\xi_1^i)_{i=1}^n, \dots, \xi_m = (\xi_m^i)_{i=1}^n \in H^n$. Then we have

$$\begin{aligned} \sum_{k,l=1}^m \langle \Gamma(\rho_S)(a_l^* a_k) \xi_k, \xi_l \rangle &= \sum_{k,l=1}^m \langle [V_\rho^* S_{ij} \Phi_\rho(a_l^* a_k) V_\rho]_{i,j=1}^n (\xi_k^i)_{i=1}^n, (\xi_l^i)_{i=1}^n \rangle \\ &= \sum_{k,l=1}^m \sum_{i,j=1}^n \langle V_\rho^* S_{ij} \Phi_\rho(a_l^* a_k) V_\rho \xi_k^j, \xi_l^i \rangle \\ &= \sum_{i,j=1}^n \left\langle S_{ij} \sum_{k=1}^m \Phi_\rho(a_k) V_\rho \xi_k^j, \sum_{k=1}^m \Phi_\rho(a_k) V_\rho \xi_k^i \right\rangle \\ &= \left\langle [S_{ij}]_{i,j=1}^n \left(\sum_{k=1}^m \Phi_\rho(a_k) V_\rho \xi_k^i \right)_{i=1}^n, \left(\sum_{k=1}^m \Phi_\rho(a_k) V_\rho \xi_k^i \right)_{i=1}^n \right\rangle \geq 0 \end{aligned}$$

since $[S_{ij}]_{i,j=1}^n$ is a positive element in $M_n(\Phi_\rho(A)')$. From this fact we conclude that $\Gamma(\rho_S) \in CP_\infty(A, M_n(L(H)))$ and the lemma is proved. \square

REMARK 2.2. It is not difficult to check that:

1. $\rho_{T^\rho} = \rho$;
2. $\rho_{\alpha S} = \alpha \rho_S$, for all positive numbers α and for all positive elements S in $M_n(\Phi_\rho(A)')$;
3. $\rho_{S_1+S_2} = \rho_{S_1} + \rho_{S_2}$, for all positive elements S_1, S_2 in $M_n(\Phi_\rho(A)')$;

4. $\rho_{S_1} \leq \rho_{S_2}$ if and only if $S_1 \leq S_2$, where S_1, S_2 are positive elements in $M_n(\Phi_\rho(A)')$.

Let $\rho, \theta \in CP_\infty^n(A, L(H))$. We say that ρ *dominates* θ , and we write $\theta \leq \rho$, if $\rho - \theta \in CP_\infty^n(A, L(H))$.

For $\rho \in CP_\infty^n(A, L(H))$, we put:

$$[0, \rho] = \{\theta = [\theta_{ij}]_{i,j=1}^n \in CP_\infty^n(A, L(H)); \theta \leq \rho\}$$

and

$$[0, T^\rho] = \{S = [S_{ij}]_{i,j=1}^n \in M_n(\Phi_\rho(A)'); 0 \leq S \leq T^\rho\}.$$

THEOREM 2.3. *The map $S \longrightarrow \rho_S$ is an affine order isomorphism from $[0, T^\rho]$ to $[0, \rho]$.*

PROOF. By Lemma 2.1 and Remark 2.2, the map $S \longrightarrow \rho_S$ from $[0, T^\rho]$ to $[0, \rho]$ is well-defined and moreover, it is affine.

To show that the map is injective, let $S = [S_{ij}]_{i,j=1}^n$ be an element in $[0, T^\rho]$ such that $\rho_S = 0$. Then $[\rho_{S_{ij}}]_{i,j=1}^n = 0$, that is $V_\rho^* S_{ij} \Phi_\rho(a) V_\rho = 0$, for all $a \in A$ and for all $i, j = 1, \dots, n$.

For each $a, b \in A, \xi, \eta \in H$ and $i, j = 1, \dots, n$, we have

$$\begin{aligned} \langle S_{ij} \Phi_\rho(a) V_\rho \xi, \Phi_\rho(b) V_\rho \eta \rangle &= \langle V_\rho^* \Phi_\rho(b)^* S_{ij} \Phi_\rho(a) V_\rho \xi, \eta \rangle \\ &= \langle V_\rho^* \Phi_\rho(b^*) S_{ij} \Phi_\rho(a) V_\rho \xi, \eta \rangle \\ &= \langle V_\rho^* S_{ij} \Phi_\rho(b^* a) V_\rho \xi, \eta \rangle = 0. \end{aligned}$$

From this fact and taking into account that $\{\Phi_\rho(a) V_\rho \xi; a \in A, \xi \in H\}$ spans a dense subspace of H_ρ , we conclude that $S_{ij} = 0$. Hence $S = 0$ and the map $S \longrightarrow \rho_S$ is injective.

It remains to show that the map $S \longrightarrow \rho_S$ from $[0, T^\rho]$ to $[0, \rho]$ is surjective.

Let $\sigma = [\sigma_{kl}]_{k,l=1}^n$ be an element in $[0, \rho]$. By [4, the proof of Theorem 2.3] (see also [6]), $\frac{1}{n} \sigma_{kk}, \frac{1}{2} \tilde{\sigma} \pm \frac{1}{n} \operatorname{Re} \sigma_{kl}, \frac{1}{2} \tilde{\sigma} \pm \frac{1}{n} \operatorname{Im} \sigma_{kl} \in [0, \tilde{\rho}]$, where $\tilde{\rho} = \frac{1}{n} \sum_{j=1}^n \rho_{jj}$

and $\tilde{\sigma} = \frac{1}{n} \sum_{j=1}^n \sigma_{jj}$, for all $k, l = 1, \dots, n$ with $k \neq l$. Let $(\Phi_\rho, H_\rho, V_\rho, T^\rho)$ be

the Suen's construction associated with ρ . By Remark 1.4, $(\Phi_\rho, H_\rho, V_\rho)$ is the Stinespring representation of A associated with $\tilde{\rho}$. Then by [1, Theorem 1.4.6], for each $j = 1, \dots, n$, there is a positive element $S_{jj} \in \Phi_\rho(A)'$ such that

$$\sigma_{jj}(a) = V_\rho^* S_{jj} \Phi_\rho(a) V_\rho$$

for all $a \in A$ and for all $k, l = 1, \dots, n$ with $k \neq l$, there are two positive elements $S_{kl}^1, S_{kl}^2 \in \Phi_\rho(A)'$ such that

$$\frac{n}{2} \tilde{\sigma}(a) + (\operatorname{Re} \sigma_{kl})(a) = V_\rho^* S_{kl}^1 \Phi_\rho(a) V_\rho$$

and

$$\frac{n}{2}\tilde{\sigma}(a) + (\text{Im}\sigma_{kl})(a) = V_\rho^* S_{kl}^2 \Phi_\rho(a) V_\rho$$

for all $a \in A$.

From these relations, we deduce that $\sigma_{kl}(a) = V_\rho^* S_{kl} \Phi_\rho(a) V_\rho$ for all $a \in A$, where

$$S_{kl} = S_{kl}^1 + iS_{kl}^2 - \frac{1+i}{2} \sum_{j=1}^n S_{jj}.$$

Clearly $S = [S_{ij}]_{i,j=1}^n \in M_n(\Phi_\rho(A)')$. Moreover, S is positive (see, for example, [4, the proof of Theorem 2.3]) and $\sigma = \rho_S$. Since $\sigma \leq \rho$, by Remark 2.2, $S \in [0, T^\rho]$ and the theorem is proved. \square

DEFINITION 2.4. Let A be a C^* -algebra and let H be a Hilbert space. A completely n -positive linear map $\rho = [\rho_{ij}]_{i,j=1}^n$ from A to $L(H)$ is said to be pure if for every completely n -positive linear map $\theta = [\theta_{ij}]_{i,j=1}^n \in [0, \rho]$, there is a positive number α such that $\theta = \alpha\rho$.

PROPOSITION 2.5. Let $\rho = [\rho_{ij}]_{i,j=1}^n$ be an element in $CP_\infty^n(A, L(H))$. Then ρ is pure if and only if $[0, T^\rho] = \{\alpha T^\rho; 0 \leq \alpha \leq 1\}$.

PROOF. First we suppose that ρ is pure. Let $S = [S_{ij}]_{i,j=1}^n$ be an element in $[0, T^\rho]$. By Theorem 2.3, $\rho_S \in [0, \rho]$ and since ρ is pure, there is a positive number α such that $\rho_S = \alpha\rho$. From this fact, Remark 2.2 and Theorem 2.3, we deduce that $S = \alpha T^\rho$ for some $0 \leq \alpha \leq 1$.

Conversely, suppose that $[0, T^\rho] = \{\alpha T^\rho; 0 \leq \alpha \leq 1\}$. Let $\theta = [\theta_{ij}]_{i,j=1}^n$ be an element in $[0, \rho]$. By Theorem 2.3, there is $S \in [0, T^\rho]$ such that $\rho_S = \theta$ and since $S = \alpha T^\rho$ for some positive number α , $\theta = \alpha\rho$ and the proposition is proved. \square

Let A be a unital C^* -algebra, let H be a Hilbert space and $\rho = [\rho_{ij}]_{i,j=1}^n \in CP_\infty^n(A, L(H))$. We denote by $CP_\infty^n(A, L(H), T^0)$, where

$$T^0 = \text{diag}(V_\rho^*, \dots, V_\rho^*) T^\rho \text{diag}(V_\rho, \dots, V_\rho),$$

the set of all completely n -positive linear maps $\sigma = [\sigma_{ij}]_{i,j=1}^n$ from A to $L(H)$ such that $\sigma_{ij}(1_A) = (T^0)_{ij}$, for all $i, j = 1, \dots, n$. Clearly, $CP_\infty^n(A, L(H), T^0)$ is a convex set.

PROPOSITION 2.6. Let $\rho = [\rho_{ij}]_{i,j=1}^n$ be an element in $CP_\infty^n(A, L(H), T^0)$ and let P_{H_0} be the projection on the closed subspace H_0 of H_ρ generated by $\{V_\rho \xi; \xi \in H\}$. If the map $S \rightarrow \text{diag}(P_{H_0}, \dots, P_{H_0}) S \text{diag}(P_{H_0}, \dots, P_{H_0})$ from $M_n(\Phi_\rho(A)')$ to $M_n(L(H_\rho))$ is injective then ρ is an extreme point in $CP_\infty^n(A, L(H), T^0)$.

PROOF. Let θ, σ be elements in $CP_\infty^n(A, L(H), T^0)$ and $\alpha \in (0, 1)$ such that $\alpha\theta + (1 - \alpha)\sigma = \rho$. Since $\alpha\theta \in [0, \rho]$, by Theorem 2.3 there is a positive element S in $M_n(\Phi_\rho(A)')$ such that $\alpha\theta = \rho_S$. Then

$$\begin{aligned} \langle P_{H_0}(S_{ij} - \alpha T_{ij}^\rho) P_{H_0} V_\rho \xi, V_\rho \eta \rangle &= \langle S_{ij} V_\rho \xi, V_\rho \eta \rangle - \alpha \langle T_{ij}^\rho V_\rho \xi, V_\rho \eta \rangle \\ &= \alpha \langle \theta_{ij}(1_A) \xi, \eta \rangle - \alpha \langle \rho_{ij}(1_A) \xi, \eta \rangle = 0, \end{aligned}$$

for all $\xi, \eta \in H$ and for all $i, j = 1, \dots, n$.

From this fact we deduce that $P_{H_0}(S_{ij} - \alpha T_{ij}^\rho) P_{H_0} = 0$ for all $i, j = 1, \dots, n$ and since the map $S \rightarrow \text{diag}(P_{H_0}, \dots, P_{H_0}) S \text{diag}(P_{H_0}, \dots, P_{H_0})$ from $M_n(\Phi_\rho(A)')$ to $M_n(L(H_\rho))$ is injective, $S = \alpha T^\rho$. Thus we showed that $\theta = \rho$ and so ρ is an extreme point in $CP_\infty^n(A, L(H), T^0)$. \square

By Remark 1.4, $(\Phi_\rho, H_\rho, V_\rho)$ is the Sinespring representation of A associated to $\tilde{\rho}$. If $\rho = [\rho_{ij}]_{i,j=1}^n \in CP_\infty^n(A, L(H), T^0)$, then

$$\tilde{\rho}(1_A) = \frac{1}{n} \sum_{k=1}^n \rho_{kk}(1_A) = \frac{1}{n} \sum_{k=1}^n V_\rho^* T_{kk} V_\rho = V_\rho^* V_\rho,$$

and by [1, Theorem 1.4.6], $\tilde{\rho}$ is an extreme point in $CP_\infty(A, L(H), V_\rho^* V_\rho)$ if and only if the map $S \rightarrow P_{H_0} S P_{H_0}$ from $\Phi_\rho(A)'$ to $L(H_\rho)$ is injective.

COROLLARY 2.7. *Let $\rho = [\rho_{ij}]_{i,j=1}^n$ be an element in $CP_\infty^n(A, L(H), T^0)$.*

If $\tilde{\rho} = \frac{1}{n} \sum_{k=1}^n \rho_{kk}$ is an extreme point in $CP_\infty(A, L(H), V_\rho^ V_\rho)$, then ρ is an extreme point in $CP_\infty^n(A, L(H), T^0)$.*

PROOF. Since $\tilde{\rho}$ is an extreme point in the set $CP_\infty(A, L(H), V_\rho^* V_\rho)$, the map $S_0 \rightarrow P_{H_0} S_0 P_{H_0}$ from $\Phi_\rho(A)'$ to $L(H_\rho)$ is injective [1, Theorem 1.4.6], and so the map $S \rightarrow \text{diag}(P_{H_0}, \dots, P_{H_0}) S \text{diag}(P_{H_0}, \dots, P_{H_0})$ is injective. From this fact and Proposition 2.6, we deduce that ρ is an extreme point in $CP_\infty^n(A, L(H), T^0)$. \square

ACKNOWLEDGEMENTS.

This research was supported by CNCSIS grant code A 1065/2006.

REFERENCES

- [1] W. Arveson, *Subalgebras of C^* -algebras*, Acta Math. **123** (1969), 141-224.
- [2] J. Heo, *Completely multi-positive linear maps and representation on Hilbert C^* -modules*, J. Operator Theory **41** (1999), 3-22.
- [3] A. S. Holevo, *Radon-Nikodym derivatives of quantum instruments*, J. Math. Phys. **39** (1998), 1373-1387.
- [4] M. Joița, *On representations associated with completely n -positive linear maps on pro- C^* -algebras*, Chin. Ann. Math. Ser. B **29** (2008), 55-64.
- [5] P. Jorgensen, *Some connection between operator algebras and quantum information theory*, AIMS "Fifty International Conference on Dynamical Systems and Differential Equations", California State Polytechnic University, June 16-19, 2004.

- [6] V. I. Paulsen and C.Y. Suen, *Commutant representations of completely bounded maps*, J. Operator Theory **13** (1985), 87-101.
- [7] M. Ragsky, *Radon-Nikodym derivatives of quantum operations*, J. Math. Phys. **44** (2003), 5003-5020.
- [8] W. Stinespring, *Positive functions on C^* -algebras*, Proc. Amer. Math. Soc. **6** (1955), 211-216.
- [9] C. Y. Suen, *An $n \times n$ matrix of linear maps of a C^* -algebra*, Proc. Amer. Math. Soc. **112** (1991), 709-712.
- [10] R. F. Werner, *Quantum Information Theory-An Invitation*, in: Alber G., Beth T., Horodecki M. et al. (eds), Quantum Information: An Introduction to Basic Theoretical Concepts and Experiments, Springer Tracts in Modern Physics **173**, Springer-Verlag, Berlin, 2001, 14-57.

M. Joița
Department of Mathematics
Faculty of Chemistry
University of Bucharest
Bd. Regina Elisabeta nr. 4-12
Romania
E-mail: mjoita@fmi.unibuc.ro

T.-L. Costache
Faculty of Applied Sciences
University "Politehnica" of Bucharest
Romania
E-mail: lumycos@yahoo.com

M. Zamfir
Department of Mathematics and Informatic
Technical University of Civil Engineering Bucharest
Romania
E-mail: zamfirvmariana@yahoo.com

Received: 9.10.2007.

Revised: 13.4.2008.