

HYPERSENSITIVITY TO THE POLLEN OF *OLEA EUROPEA* IN THE MEDITERRANEAN AREA

Nataša SKITARELIĆ and Neven SKITARELIĆ

Zadar General Hospital, Zadar, Croatia

Received in February 2009

Accepted in February 2009

Olive pollen is one of the most important causes of inhalant allergy in the Mediterranean countries (1, 2). We investigated the frequency of hypersensitivity to the pollen of *Olea europea* in pollen allergic patients in Dalmatia, Croatia (3). A total of 810 patients with pollen allergy, were examined: 469 (58 %) male and 341 (42 %) female, age range from 4 to 53 years.

The patients were assessed on the basis of medical history, clinical examination, measurement of pulmonary function (adults and children older than 7 years), intradermal prick test, and immuno-enzyme UniCAP test (UniCAP, Fluoroenzymeimmunoassay; Pharmacia, Upjohn, Sweden) for specific IgE antibodies. We used standard allergen prick tests produced by the Institute of Immunology in Zagreb, Croatia.

Our study, like two earlier (4, 5), confirmed that grasses, especially *Parietaria officinalis*, are the main pollen allergens. Trees rarely cause allergies in the Croatian part of the Adriatic coast (4). Hypersensitivity to *Olea europea* occurred in 66 of 810 (8.15 %) assessed patients with pollen allergy. *Olea europea* and *Fraxinus ornus* belong to the family of Oleaceae. Both trees are part of the local flora. We confirmed the hypersensitivity to *Olea europea* allergen and excluded cross-reactivity between these plants by specific IgE antibodies measurements.

The most prevalent clinical manifestations were rhinitis in 39 (59 %) of 66 patients allergic to *Olea*

europea pollen and bronchial asthma in 20 (30.3 %) patients. It is interesting to note that 51 patients (77 %) lived in towns and the rest in villages; only 3 patients (4 %) lived on the islands. The prevalence of pollen allergies in patients living in the coastal region is recognised in other Mediterranean countries (6).

The hypersensitivity to *Olea europea* allergen in our patients was significantly lower than in other Mediterranean countries (Figure 1); ($p < 0.01$), where varied from 21 % to 31.8 % (7-10). The reason for this difference may be sought in greater industrial pollution (11, 12) and greater area cultivated with olives (13) in other Mediterranean countries.




Figure 1 Hypersensitivity to the pollen of *Olea europea* in five Mediterranean countries, * $p < 0.01$.

REFERENCES

1. Asturias JA, Arilla MC, Gomez-Bayon N, Martinez J, Martinez A, Palacios R. Cloning and expression of the panallergen profilin and the major allergen (Ole e 1) from olive tree pollen. *J Allergy Clin Immunol* 1997;100:365-72.
2. Tejera ML, Villalba M, Batanero E, Rodriguez R. Identification, isolation and characterization of Ole e 7, a new allergen of olive tree pollen. *J Allergy Clin Immunol* 1999;104:797-802.
3. Skitarelić N, Sindik N, Skitarelić N, Mazzi A, Vuletić A, Mišulić J. Učestalost senzibilizacije na maslinu u bolesnika s peludnom alergijom na području Zadra i Dubrovnika [Hypersensitivity to pollen of *Olea europea* in patients with pollen allergy in the areas of Zadar and Dubrovnik, in Croatian]. *Liječ Vjesn* 2004;126: 65-70.
4. Sindik N, Restović-Sirotković M, Skitarelić B, Balić J, Kavurić-Hafner K, Peharda-Sabolić V, Čvorišćec B. Polinoze na području Dalmacije, Istre i Hrvatskog primorja s posebnim osvrtom na pelud *Parietariae* [Pollinosis in the area of the Dalmatian, Istrian and Croatian coast with special emphasis on *Parietariae* pollen, in Croatian]. *Liječ Vjesn* 1989;111:263-5.
5. Cvitanović S, Marušić M, Juričić M, Vrdoljak E, Petrovečki M, Rozga M, Stavljenić-Rukavina A. Hypersensitivity to *Parietaria officinalis* pollen in newcomers to the area with the plant. *Allergy* 1993;48:592-7.
6. Ariano R, Passalacqua G, Panzani R, Scordamaglia A, Venturi S, Zoccali P, Canonica GW. Airborne pollens and prevalence of pollenosis in western Liguria: a 10-year study. *J Invest Allergol Clin Immunol* 1999;9:229-34.
7. Verini M, Rossi N, Verotti A, Pelaccia G, Nicodemo A, Chiarelli F. Sensitization to environmental antigens in asthmatic children from a central Italian area. *Sci Total Environ* 2001;270:63-9.
8. Geller-Bernstein C, Lahoz C, Cardaba B, Hassoun G, Iancovic-Kidon M, Kenett R et. al. Is it "bad hygiene" to inhale pollen in early life? *Allergy* 2002;57(Suppl. 71):37-40.
9. Ledesma A, Rodriguez R, Villalba M. Olive-pollen profilin. Molecular and immunologic properties. *Allergy* 1998;53:520-6.
10. Gioulekas D, Papakosta D, Damialis A, Spiexsma F, Giouleka P, Patakas D. Allergenic pollen records (15 years) and sensitization in patients with respiratory allergy in Thessaloniki, Greece. *Allergy* 2004;59:174-84.
11. Zhang XD, Lotvall J, Arakawa H, Welinder H, Skerfving S. Relationship between IgG1 levels and airway responses in guinea pigs actively and passively sensitized to hexahydrophthalic anhydride. *Allergy* 1998;53:20-7.
12. Salvi S. Pollution and allergic airways disease. *Curr Opin Allergy Clin Immunol* 2001;1:35-41.
13. Macchia L, Aliani M, Caiaffa MF, Carbonara AM, Gatti E, Iacobelli A, Strada S, Casella G, Tursi A. Monitoring of atmospheric conditions and forecast of olive pollen season. *Experientia Suppl* 1987;51:95-9.

CORRESPONDING AUTHOR:

Nataša Skitarelić, MD
Put Murvice 33
23000 Zadar, Croatia
E-mail: neven.skitarelic@zd.t-com.hr