

GEOL. CROAT.	52/1	77 - 81	2 Figs.		ZAGREB 1999
--------------	------	---------	---------	--	-------------

Stable Isotope Event Markers Near the Permo-Triassic Boundary in the Karavanke Mountains (Slovenia)

Tadej DOLENEC^{1,2}, Sonja LOJEN², Stanko BUSER¹ and Matej DOLENEC³

Key words: Permo-Triassic boundary, Karavanke Mountains, Slovenia, Oxygen, Carbonate carbon, Organic carbon stable isotopes.

Abstract

Stable isotope analyses of carbonates and organic matter from the Permo-Triassic boundary section in the Karavanke Mountains, Slovenia, indicate a further example of the "light carbon" event across the boundary. In this section the changes in carbon isotope values were a direct result of the culmination of the marine regression and associated events at the end of the Permian, which caused a drop in primary productivity, as well as related local environmental changes, with no evidence of any considerable diagenetic overprint.

over the world. One of the most remarkable anomalies is the worldwide negative shift of ¹³C of inorganic and organic carbon across the P/Tr boundary (MAGARITZ et al., 1992; WANG et al., 1994; WOLBACH et al., 1994; FAURE et al., 1995). A corresponding oxygen isotopic shift is more or less parallel, but less pronounced. Furthermore, significant shifts in sulphur (KAJIWARA et al., 1994) and strontium isotopes (KRAMM & WEDEPOHL, 1991) have also been recorded.

In this study we present the results of stable isotope analysis of the P/Tr boundary section in the Karavanke Mountains (Fig. 1), and discuss their implications with respect to a better understanding of the nature and causes of the P/Tr boundary events in this part of Western Palaeotethys.

1. INTRODUCTION

Permo-Triassic (P/Tr) boundary events, which took place approximately 250 Ma ago, led to the most extensive mass extinction in the history of life. A number of possible explanations for this profound break in the evolution of life have been proposed, such as volcanic activity, sea-level fluctuation, changes in sea-water chemistry, an extra-terrestrial impact event and various related factors (YOICHI, 1994). The most recently proposed cause of the mass extinction at the end of the Permian is a combination of these more or less co-occurring events operating in three phases (ERVIN, 1996). The first began with the marine regression during the Late Permian and resulted in the destruction of many marine basins, reduction in the habitat area of many organisms and increased climatic instability. The second phase involved the eruption of the Siberian Traps and further environmental degradation. The final phase may have started immediately prior to the boundary when the Late Permian regression ended and the earliest Triassic transgression begun. The global events outlined above coincide with isotope and elemental anomalies recorded in several P/Tr boundary sections all

2. GEOLOGICAL SETTING AND STRATIGRAPHY

In the southern Karavanke Mountains, the Middle Permian Val Gardena Formation of mostly fluvial origin, is overlain by a 270 m thick Upper Permian carbonate sequence, which was named the Karavanke Formation (BUSER, 1974). The boundary between the two formations is transitional and is characterized by thin sandy red dolomite layers alternating with the topmost Val Gardena shales and sandstones. The thickness of the transitional unit which grades upward into the Karavanke Formation is about 5 m (DOLENEC et al., 1981). The evaporitic sequence, up to 70 m thick, represents the basal unit of the Karavanke Formation composed of cellular dolomite, which alternates with rare black bituminous shales, and grey vuggy dolomites. The evaporitic sequence is overlain by a 200 m thick succession of fossiliferous biomicritic dolomites. The Upper Permian age of these beds is indicated by calcareous algal assemblages, as well as by foraminifera (RAMOVŠ, 1986). The lithostratigraphic boundary between the Upper Permian Karavanke Formation and the Lower Triassic (Scythian) beds is placed at the end of the sedimentation of the well-bedded grey dolomicrite. It is followed by a red coloured partly terrigenous sequence predominantly composed of thin-bedded siltstones, mudstones and sandstones alternating with micritic dolomites, that contain no characteristic fossils. These earliest Triassic beds were deposited in an extremely shallow sea, which

¹ Department of Geology, University of Ljubljana, Aškerčeva 12, SLO-1000 Ljubljana, Slovenia.

² Department of Environmental Sciences, Jožef Stefan Institute, Jamova 39, SLO-1000 Ljubljana, Slovenia.

³ Geoxp d.o.o., Slap 21, SLO-4290 Tržič, Slovenia.




Fig. 2 Stable isotope composition of carbonates ($^{13}\text{C}_{\text{carb.}}$, ^{18}O), total organic carbon ($^{13}\text{C}_{\text{org.}}$) and carbon isotope fractionation between carbonate and total organic carbon ($^{13}\text{C}_{\text{carb.-org.}}$) across the Permo-Triassic boundary in the Karavanke Mountains. a) Dark grey and brown micritic and sparitic limestone intercalated with oolitic limestone, marls and shales; b) a red partly terrigenous sequence composed of siltstones, mudstones and sandstones alternating with micritic dolomites; c) light grey fossiliferous biomicritic dolomite; d) an evaporitic sequence composed of cellular dolomite intercalated with black bituminous shales and grey vuggy dolomites; e) a transitional unit composed of sandy red dolomite alternating with shales and sandstones; 1-3) carbonate and organic carbon anomalies.

The dolomites of the basal evaporitic sequence show a variation of $^{13}\text{C}_{\text{carb.}}$ in the range of +0.69 to +3.83‰, and of ^{18}O between -5.88 to -3.34‰. The oxygen isotopic composition of the evaporitic sequence is not as high as expected from recent evaporitic environments (TUCKER, 1990). The observed reduction in ^{18}O indicate an influx of fresh water into the evaporitic basin, less evaporation and/or a variety of different post-depositional processes, that may affect either oxygen or carbon or both stable isotopic records (BRAND & VEIZER, 1981; MAGARITZ & HOLSER, 1991). By selecting the least visibly weathered and recrystallized samples from the investigated sections we attempted to minimise the possible post-depositional effects. A weak positive correlation ($r = 0.40$) between ^{18}O and ^{13}C of the dolomite samples and a weak negative correlation ($r = -0.22$) between ^{18}O and ^{13}C of the limestone samples most probably suggests that the isotopically light meteoric water and dolomitizing fluids during diagenesis and burial to some extent reset the original whole rock oxygen and carbon isotopic composition. The low positive correlation ($r = 0.51$) between $^{13}\text{C}_{\text{carb.}}$ and $^{13}\text{C}_{\text{org.}}$ indicates that the whole rock isotopic composition was not modified by the oxidation of organic matter during diagenesis. Based on these observations, we tend to believe that the isotopic composition of the investigated Karavanke Mountain carbonate

rocks has not been seriously altered after their formation and that the primary palaeoceanographic signal was not completely overprinted.

The transition from Permian to Triassic is characterized by a prominent negative shift of carbonate ^{13}C as well as total organic carbon and a similar, but less pronounced ^{18}O decrease. The major drop of $^{13}\text{C}_{\text{carb.}}$ and $^{13}\text{C}_{\text{org.}}$, similar to that seen globally, begins approximately 15 m below the boundary. The $^{13}\text{C}_{\text{carb.}}$ curve reaches a minimum peak value of -1.86‰ about 8 m below the boundary and after that there is another minimum of -1.57‰ at the end of the Permian and a positive excursion of +0.55‰ in the lowermost Scythian, before settling to values which are 1 to 2‰ lower relative to those in the Upper Permian. The position of the negative $^{13}\text{C}_{\text{carb.}}$ peak anomaly may indicate that the P/Tr boundary in the Karavanke Mountains should be placed a little further downsection, since in the Carnic Alps a dramatic $^{13}\text{C}_{\text{carb.}}$ drop occurs right after the stratigraphic P/Tr boundary which is placed within the lowermost 0.5 m of the 4 m thick oolitic Tesserò Horizon or at its base (HOLSER et al., 1991). In the Karavanke Mountains this oolitic unit at the base of the Scythian was not recognized. The corresponding total organic carbon isotope curve is essentially parallel. The first (-29.89‰) and the second (-26.65‰) negative anomaly for $^{13}\text{C}_{\text{org.}}$ are coeval with the first two nega-

tive shifts observed for ^{13}C of carbonate carbon, while the third decrease of $^{13}\text{C}_{\text{org.}}$ (-26.92‰) occurs in the Triassic, 7 m below the third negative shift of $^{13}\text{C}_{\text{carb.}}$ values. During the Permian-Triassic transition the Karavanke Mountains constituted a semi-restricted marine basin, surrounded by vast areas of an extremely shallow epicontinental sea into which a considerable amount of terrigenous material was transported (DOLENEC et al., 1981).

The environmental stress conditions caused decreased bioproductivity in the upper water column and an associated decrease in dissolved carbonate (ZHENG et al., 1993). A drop of ^{13}C values in carbonate precipitates reflects this decrease. The incorporation of light carbon from eroded and oxidized organic matter into carbonates and photosynthetic marine organisms would also cause both to become isotopically lighter (WOLBACH et al., 1994). The variation between $^{13}\text{C}_{\text{carb.}}$ and $^{13}\text{C}_{\text{org.}}$ values, which can be used to evaluate major changes in the carbon cycle by removing the effect of changes in the ^{13}C of the surface water dissolved inorganic carbon reservoir (HOLLANDER et al., 1993), also indicates changes in bioproduction. For the P/Tr transition in the Karavanke Mountains the decreasing isotopic composition of total organic carbon of 6.5‰ (from -23.19 to -29.69‰) and increasing $^{13}\text{C}_{\text{carb.-org.}}$ of 3.78‰ (from 24.66 to 28.44‰) about 8 m below the boundary can be explained by a dramatic decrease in primary production. This breakdown in the photosynthesis/respiration cycle appears to coincide with events related to the accelerated fall in sea level and/or widespread volcanic activity in Siberia. These events began substantially before the end of the Permian and affected the carbon cycle long enough to cause a clear perturbation in both ^{13}C records. The ^{13}C shift of 3.5‰ (from -8.89 to -25.39‰) in total organic carbon toward higher values and the reduction in $^{13}\text{C}_{\text{carb.-org.}}$ by 4.1‰ (from 28.23 to 24.13‰) at the P/Tr boundary could be the consequence of a subsequent recovery period in the biological system before the sea gradually became a wide, extensive mud flat. This situation was changed after the sedimentation of the red terrigenous sequence, due to the earliest Triassic marine transgression which spread anoxic bottom water over the entire region of the Karavanke Mountains, and marked the end of the period of extreme variability in ^{13}C of both carbonate and organic carbon. During this phase deposition of the Lower Scythian dark grey and brown limestones began.

The variability of ^{18}O in the interval straddling the P/Tr boundary shows slightly different trends with respect to those of ^{13}C in carbonate and organic carbon, roughly indicating environmental changes from marine to desultory evaporitic conditions affected by an excessive input of terrigenous material, and by local freshening of waters, due to the influx of more or less isotopically modified meteoric waters into the sedimentary environment. A significant drop of ^{18}O from -4.19‰ to -9.51‰ approximately 25 m above the boundary is stratigraphically coincident with abrupt chan-

ges in the lithology, as well as carbonate mineralogy of the rocks. The Scythian limestones are considerably depleted in ^{18}O (from 3 to 8‰) relative to both the underlying lowermost Scythian dolomite and the Upper Permian dolomite of the Karavanke Formation. Such depletion cannot be interpreted only in terms of dolomite-calcite fractionation which is thought to be between +3 and +4‰ (DICKSON, 1990). It may also be caused by a change in the ^{18}O of seawater, a decrease of salinity, changes in the depositional environment, diagenetic processes and burial, or some combination of all of these factors. Due to the muddled oxygen isotopic pattern, it is difficult to give a definite explanation of the observed ^{18}O variations at the P/Tr transition. However, it is important to note that the ^{13}C signals of inorganic and organic carbon are independent of lithology and are preserved in both Scythian dolomite and limestone.

5. CONCLUSIONS

The results we have presented in this study indicate that the transition from the Permian to the Triassic in the Karavanke Mountains is characterized by a strongly negative complex pattern of carbonate and organic carbon anomalies, accompanied by changes in the ^{18}O values of the boundary carbonate rocks. These anomalies are supposed to record extremely adverse changes of the environment due to global events which marked the terminal Permian productivity crash and the P/Tr boundary. We suggest a causal connection between the global carbon isotopic anomalies and the marine regression at the end of the Permian, but the exact controls remain complex and enigmatic.

Acknowledgements

The Ministry of Science and Technology, Republic of Slovenia, and Geoxp d.o.o., Tržič, Slovenia financially supported this study. To both these institutions we express our sincere thanks.

6. REFERENCES

- ASSERETO, R., BOSSELINI, A., FANTINI SESTINI, N. & SWEET, W.C. (1973): The Permian-Triassic boundary in the Southern Alps (Italy).- In: LOGAN, A. & HILLS, L.V. (eds.): The Permian and Triassic Systems and Their Mutual Boundary. *Albertal. Soc. Petrol. Geol. Mem.*, 2, 176-199.
- BEUKES, N.J., KLEIN, C., KAUFMAN, A.J. & HAYES, J.M. (1990): Carbonate petrography, kerogen distribution, and carbon and oxygen isotope variations in an Early Proterozoic transition from limestone to iron-formation deposition, Traansvaal Super-group, South Africa.- *Econ. Geology*, 85, 663-690.

- BRAND, U. & VEIZER, J. (1981): Chemical diagenesis of a multicomponent carbonate system - 2: stable isotopes.- *J. Sediment. Petrol.*, 51, 987-997.
- BUSER, S. (1974): Neue Feststellungen im Perm der westlichen Karavanken.- *Carinthia II*, 164/84, 27-37.
- COMPTON, J.S., SNYDER, S.W. & HODELL, D.A. (1990): Phosphogenesis and weathering of shelf sediments from the southern United States: Implications for Miocene ^{13}C excursions and global cooling.- *Geology*, 18, 1227-1230.
- CRAIG, H. (1957): Isotopic standards for carbon and oxygen and correction factors for mass spectrometric analysis of carbon dioxide.- *Geochim. Cosmochim. Acta*, 12, 133-149.
- DICKSON, T. (1990): Carbonate mineralogy and chemistry.- In: TUCKER, M.E. & WRIGHT, V.P.: *Carbonate Sedimentology*. Blackwell Scientific Publications, Oxford, 482 p.
- DOLENEC, T. (1984): The origin of the uranium mine Žirovski vrh.- Unpublished PhD Thesis, University of Ljubljana, 187 p.
- DOLENEC, T., OGORELEC, B. & PEZDIČ, J. (1981): Upper Permian and Scythian beds in the Tržič area.- *Geologija*, 24/2, 217-238, Ljubljana.
- ERVIN, D.H. (1996): Permian global bio-events.- In: WALISER, O.H. (ed.): *Global Events and Event Stratigraphy in the Phanerozoic*. Springer-Verlag, Berlin, 251-264.
- FAURE, K., DE WIT, M.J. & WILLIS, J.P. (1995): Late Permian global coal hiatus linked to ^{13}C -depleted CO_2 flux into the atmosphere during the final consolidation of Pangea.- *Geology*, 23, 507-510.
- HOLLANDER, D.J., MCKENZIE, J.A. & KENNETH, J.H. (1993): Carbon isotope evidence for unusual plankton blooms and fluctuation of surface water CO_2 in "Strangelove Ocean" after terminal Cretaceous event.- *Palaeogeogr. Palaeoclim. Palaeoecol.*, 104, 229-237.
- HOLSER, W.T., SCHÖNLAUB, H.P., BOECKELMANN, K. & MAGARITZ, M. (1991): The Permian-Triassic of the Gartnerkofel-1 core (Carnic Alps, Austria): synthesis and conclusions.- *Abh. Geol. B.*, A 45, 213-232.
- KAJIWARA, Y., YAKAMITA, S., ISHIDA, K., ISHIGA, H. & IMAI, A. (1994): Development of a largely anoxic stratified ocean and its temporary massive mixing at the Permian/Triassic boundary supported by the sulfur isotopic record.- *Palaeogeogr. Palaeoclim. Palaeoecol.*, 111, 367-379.
- KRAMM, U. & WEDEPOHL, K.H. (1991): The isotopic composition of strontium and sulfur in seawater of Late Permian (Zechstein) age.- *Chem. Geol.*, 90, 253-262.
- MAGARITZ, M. & HOLSER, W.T. (1991): The Permian-Triassic of the Gartnerkofel-1 Core (Carnic Alps, Austria): carbon and oxygen isotope variation.- *Abh. Geol. B.*, A 45, 149-163.
- MAGARITZ, M., KRISHNAMURTHY, R.V. & HOLSER, W.T. (1992): Parallel trends in organic and inorganic carbon isotopes across the Permian/Triassic boundary.- *Am. Jour. Sci.*, 292, 727-739.
- RAMOVŠ, A. (1986): Marine development of the uppermost Žažar beds and the lowermost Scythian beds.- In: *Permian and Permian-Triassic Boundary in the South Alpine Segment of the Western Tethys. Excursion Guidebook* (ed. by the Italian IGCP 203 Group), 39-42.
- SUN, S., LI, J., CHEN, H., PENG, W., HSU, K.J. & SHELTON, J.W. (1989): Mesozoic and Cenozoic sedimentary history of South China.- *Am. Assoc. Pet. Geol. Bull.*, 73, 1247-1269.
- TUCKER, M.E. (1990): Geological background to carbonate sedimentation.- In: TUCKER, M.E. & WRIGHT, V.P.: *Carbonate Sedimentology*. Blackwell Scientific Publications, Oxford, 28-69.
- WANG, E., GELDSETZE, H.H.J. & KRUISE, H.R. (1994): Permian-Triassic extinction: Organic ^{13}C evidence from British Columbia, Canada.- *Geology*, 22, 580-584.
- WOLBACH, W.S., ROEGGE, D.R. & GILMOUR, I. (1994): The Permian-Triassic of the Gartnerkofel-1 core (Carnic Alps, Austria): Organic carbon isotope variation.- In: *New Developments Regarding the KT Event and Other Catastrophes in Earth History*. LP, Contribution No. 825, Lunar and Planetary Institute, Houston, 138 p.
- YOICHI, E. (1994): Patterns and paleoenvironmental implications of end-Permian extinction of Rugosa in South China.- *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 107, 165-177.
- ZHENG, Y., HONG-FEI, H. & LIAN-FENG, Y. (1993): Carbon and oxygen isotope event markers near the Frasnian/Famian boundary, Luxin section, South China.- *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 104, 97-104.

Manuscript received January 18, 1999.

Revised manuscript accepted May 28, 1999.

