

Amino acid derivatives. Part I. Synthesis, antiviral and antitumor evaluation of new α -amino acid esters bearing coumarin side chain

NAJIM A. AL-MASOUDI^{1*}
IMAN A. AL-MASOUDI²
IBRAHIM A. I. ALI³
YASEEN A. AL-SOUDI⁴
BAHJAT SAEED²
PAOLO LA COLLA⁵

¹ *Fachbereich Chemie, Universität
Konstanz, Postfach 5560
D-78457 Konstanz, Germany*

² *College of Veterinary Medicine
University of Basrah, Basrah, Iraq*

³ *Department of Chemistry
Faculty of Science, Suez Canal
University, Ismailia, Egypt*

⁴ *Department of Chemistry
College of Science, University
of Al al-Bayt, Jordan*

⁵ *Universita di Cagliari, Dipartimento
di Biologia Sperimentale Sezione di
Microbiologia, 09042 Cagliari, Italy*

A series of amino acid esters bearing coumarin (3–15) were synthesized and evaluated, *in vitro*, against HIV-1, and bovine viral diarrhoea virus (BVDV). The *in vitro* cytotoxicity of 3–10 and 12 was assayed against a panel of tumor cell lines consisting of CD4 human T-cells. Compound 14 showed inhibition of HIV-1 with $EC_{50} > 1.6 \mu\text{g mL}^{-1}$, meanwhile compound 9 exhibited activity against leukaemia (MT4) with $CC_{50} = 24 \mu\text{mol L}^{-1}$.

Keywords: amino acids, antitumor activity, antiviral activity, coumarin

Accepted January 16, 2006

Several compounds containing coumarin residue are known to possess useful biological activity (1). Geiparvarin, a naturally occurring product, bearing a coumarin nucleus, has been shown to possess significant inhibitory activity against a variety of cell Walker 256 carcinoma sarcoma (2). Warfarin and some bis-hydroxycoumarins have been used as oral anticoagulants (3), β -adrenergic blocking agents (4) and vasorelaxant agents (5). Furumi *et al.* (6) have prepared new furanocoumarin ethers of faltarindol, named japonagelol, as novel antiproliferative agents. On the other hand, a new target for the

* Correspondence, e-mail: najim.al-masoudi@gmx.de

development of anti-HIV and antitumor therapies has been reported by the use, *in vivo* and *in vitro*, of amino acid derived heterocycles. Such compounds are the lysyl amide prodrug of 2-(4-amino-3-methylphenyl)-5-fluorobenzothiazole (7), amino acid derivatives of paclitaxel (8), cysteine-modifying agents (9) and isoquinoline carboxylic acid derivatives as building blocks for HIV protease inhibitors (10).

The virus family *Flaviviridae* includes the genera *Flavivirus*, *Pestivirus* and *Hepacivirus* (hepatitis C viruses, HCVs). BVDV (Bovine Viral Diarrhea Virus) is one of the classical examples of *Flaviviridae*. Only a few examples have been reported for the treatment of BVDV, such as ribavirin (11) and mycophenolate acid (12). Pharmacological properties of coumarins and amino acid derivatives prompted us to prepare new coumarin bearing amino acid derivatives to study, *in vitro*, their activity against tumor models, along with inhibitory activity for HIV-1 and BVDV.

EXPERIMENTAL

Melting points are uncorrected and were measured on a Büchi melting point apparatus B-545 (BÜCHI Labortechnik AG, Switzerland). Microanalytical data were obtained with a Vario, Elementar apparatus (Shimadzu, Japan). NMR spectra were recorded on

Table I. Physical data of the newly prepared compounds

| Compd. No. | Yield (%) | M.p. (°C) | Mol. Formula (M_r) | Found/calcd. (%) | | |
|------------|-----------|-----------|---------------------------------------------------------------------------|------------------|------|------|
| | | | | C | H | N |
| 3 | 81 | 98–99 | C ₁₄ H ₁₃ NO ₆ (291.26) | 57.60 | 4.42 | 4.67 |
| | | | | 57.73 | 4.40 | 4.81 |
| 4 | 76 | oil | C ₁₈ H ₂₁ NO ₆ (347.14) | 62.02 | 6.00 | 3.89 |
| | | | | 62.24 | 6.09 | 4.03 |
| 5 | 85 | 78–80 | C ₁₅ H ₁₅ NO ₆ (305.28) | 58.86 | 4.82 | 4.38 |
| | | | | 59.01 | 4.95 | 4.59 |
| 6 | 71 | oil | C ₁₇ H ₁₉ NO ₆ | – | – | – |
| 7 | 56 | 86–88 | C ₂₁ H ₁₇ NO ₆ (367.35) | 65.39 | 4.66 | 3.81 |
| | | | | 65.13 | 4.46 | 3.67 |
| 8 | 62 | 105–106 | C ₂₃ H ₂₁ NO ₆ (407.42) | 67.54 | 5.09 | 3.18 |
| | | | | 67.81 | 5.20 | 3.44 |
| 9 | 66 | 133–134 | C ₂₃ H ₂₀ N ₂ O ₆ (420.41) | 65.50 | 4.69 | 6.45 |
| | | | | 65.71 | 4.79 | 6.66 |
| 10 | 73 | 145–146 | C ₂₁ H ₁₉ NO ₇ (397.38) | 63.22 | 4.69 | 3.27 |
| | | | | 63.47 | 4.82 | 3.52 |
| 11 | 59 | oil | C ₃₆ H ₃₉ NO ₁₇ | – | – | – |
| 12 | 62 | oil | C ₃₄ H ₃₄ NO ₁₂ | – | – | – |
| 13 | 71 | 121–124 | C ₁₄ H ₁₃ NO ₅ (275.08) | 60.75 | 4.66 | 4.89 |
| | | | | 61.09 | 4.76 | 5.09 |
| 14 | 65 | 89–94 | C ₂₁ H ₁₉ NO ₇ (397.38) | 63.22 | 4.69 | 3.27 |
| | | | | 63.47 | 4.82 | 3.52 |

250 and 600 MHz (^1H) and 150.91 MHz (^{13}C) spectrometers (Bruker, Germany) with TMS as internal standard and on δ scale in ppm. Signal assignments for protons were identified by selective proton decoupling or by COSY spectra. Heteronuclear assignments were verified by ^1H - ^{13}C COSY, or HMQC experiments. Mass spectra were recorded on 70 eV EI and FAB MAT 8200 spectrometers (Finnigana MAT, USA), using nitrobenzyl alcohol (NBOH) or glycerol as matrixes.


Physicochemical and spectral data for the synthesized compounds are given in Tables I and II. Synthetic routes are presented in Schemes 1–3.

Table II. Spectral data of the newly prepared compounds


| Compd. | Mass (m/z) ^a | ^1H NMR (δ , ppm) |
|--------|------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 3 | 292 $[\text{M}+\text{H}]^+$ | 7.62 (d, 1H, $J = 8.9$ Hz, H-4), 7.40 (d, 1H, $J = 8.9$ Hz, H-5), 7.19 (br s, 1H, NH), 6.83 (dd, 1H, $J = 2.3$ Hz, 9.5 Hz, H-6), 6.81 (d, 1H, $J = 2.3$ Hz, H-8), 6.26 (d, 1H, $J = 9.3$ Hz, H-3), 4.56 (s, 2H, CH_2 -10), 4.12 (d, 2H, $J = 2.2$ Hz, CH_2 -13), 3.73 (s, 3H, OAc), 1.45 (br s, 1H, CH_2 -15), 1.09–0.96 (m, 1H, H-16), 0.71, 0.67 (2xs, 6H, Me_2) |
| 4 | 370 $[\text{M}+\text{Na}]^+$ | 7.53 (d, 1H, $J = 9.6$ Hz, H-4), 7.28 (d, 1H, $J = 8.4$ Hz, 2.6 Hz, H-6), 6.68 (d, 1H, $J = 2.6$ Hz, H-8), 6.18 (d, 1H, $J = 9.5$ Hz, H-3), 4.47 (s, 2H, CH_2 -10), 4.42 (m, 1H, H-13), 3.57 (s, 3H, OAc), 2.06 (m, 2H, CH_2 -15), 1.06 (m, 1H, H-16), 0.78, 0.76 (2s, 6H, 2xMe) |
| 5 | 306 $[\text{M}+\text{H}]^+$ | 7.53 (d, 1H, $J = 9.3$ Hz, H-4), 7.30 (d, 1H, $J = 8.5$ Hz, H-5), 7.05 (d, 1H, $J = 8.5$ Hz, NH), 6.87 (dd, 1H, $J = 8.6$ Hz, 2.5 Hz, H-6), 6.80 (d, 1H, $J = 2.3$ Hz, H-8), 6.17 (d, 1H, $J = 9.3$ Hz, H-3), 4.60 (m, 1H, H-13), 4.56 (s, 2H, CH_2 -10), 3.67 (s, 3H, OAc), 1.38, 1.34 (2xd, 3H, $J = 7.2$ Hz, C_{13} -Me, two rotameres) |
| 6 | 356 $[\text{M}+\text{Na}]^+$ | 7.53 (d, 1H, $J = 9.6$ Hz, H-4), 7.28 (d, 1H, $J = 9.6$ Hz, H-5), 7.03 (d, 1H, $J = 8.8$ Hz, NH), 6.78 (dd, 1H, $J = 8.7$ Hz, 2.6 Hz, H-6), 6.68 (d, 1H, $J = 2.6$ Hz, H-8), 6.18 (d, 1H, $J = 9.5$ Hz, H-3), 4.47 (s, 2H, CH_2 -10), 4.42 (dd, 1H, $J = 9.5$ Hz, 5.1 Hz, H-13), 3.57 (s, 3H, OAc), 2.06 (m, 2H, CH_2 -15), 0.78, 0.76 (2s, 6H, 2xMe) |
| 7 | 390 $[\text{M}+\text{Na}]^+$ | 7.67 (d, 1H, $J = 9.2$ Hz, H-4), 7.55 (d, 1H, $J = 9.5$ Hz, H-5), 7.33–7.18 (m, 6H, NH, Ar-H, H-5), 6.80 (dd, 1H, $J = 8.6$ Hz, 2.5 Hz, H-6), 6.72 (d, 1H, $J = 2.0$ Hz, H-8), 6.14 (d, 1H, $J = 9.5$ Hz, H-3), 5.58 (d, 1H, $J = 7.2$ Hz, H-13), 4.49 (s, 2H, CH_2 -10), 3.62 (s, 3H, OAc) |
| 8 | 408 $[\text{M}+\text{H}]^+$ | 7.54 (d, 1H, $J = 9.4$ Hz, H-4), 7.29–7.20 (m, 7H, H-5, NH, CH_2Ph), 6.82 (dd, 1H, $J = 8.7$ Hz, 2.4 Hz, H-6), 6.72 (d, 1H, $J = 2.1$ Hz, H-8), 6.14 (d, 1H, $J = 9.5$ Hz, H-3), 5.10 (s, 2H, CH_2Ph), 4.69 (s, 2H, CH_2 -10), 4.54 (m, 1H, H-16), 3.58 (m, 2H, CH_2 -13), 2.29–1.81 (m, 4H, CH_2 -14, CH_2 -15) |
| 9 | 398 $[\text{M}+\text{H}]^+$ | 9.03 (br s, 1H, NH-indole), 7.48 (m, 2H, H-4, H-5), 7.42–6.51 (m, 5H, NH, indole-5H), 6.54 (m, 2H, H-6, H-8), 6.16 (d, 1H, $J = 9.4$ Hz, H-3), 4.92 (m, 1H, H-13), 4.38 (s, 2H, CH_2 -10), 3.67 (s, 3H, OAc), 3.30 (br s, 2H, CH_2 -15) |
| 10 | 398 $[\text{M}+\text{H}]^+$ | 7.54 (d, 1H, $J = 9.4$ Hz, H-4), 7.29–7.20 (m, 7H, H-5, NH, CH_2Ph), 6.82 (dd, 1H, $J = 8.7$ Hz, 2.4 Hz, H-6), 6.72 (d, 1H, $J = 2.1$ Hz, H-8), 6.14 (d, 1H, $J = 9.5$ Hz, H-3), 5.10 (s, 2H, CH_2Ph), 4.69 (s, 2H, CH_2 -10), 4.54 (m, 1H, H-16), 3.58 (m, 2H, CH_2 -13), 2.29–1.81 (m, 4H, CH_2 -14, CH_2 -15) |

Table II. continued


| Compd. | Mass (m/z) ^a | ¹ H NMR (δ , ppm) |
|--------|-----------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 11 | 780 [M+Na] ⁺ | 7.65 (d, 1H, $J = 9.6$ Hz, H-4), 7.50 (d, 1H, $J = 8.1$ Hz, NH), 7.43 (d, 1H, $J = 8.8$ Hz, H-5), 7.38–7.27 (m, 5H, CH ₂ Ph), 6.96 (dd, 1H, $J = 8.8$ Hz, $J = 2.0$ Hz, H-6), 6.91 (d, 1H, $J = 2.0$ Hz, H-8), 6.14 (d, 1H, $J = 9.5$ Hz, H-3), 5.26 (d, 2H, $J = 1.2$ Hz, CH ₂ Ph), 5.24 (m, 2H, H-3', H-4'), 5.15 (br s, 1H, H-2'), 4.93 (m, 1H, H-13), 4.74 (d, 1H, $J_{1',2'} = 2.8$ Hz, H-1'), 4.62 (s, 2H, CH ₂ -10), 4.19 (dd, 1H, $J_{5',6'a} = 5.5$ Hz, H-6'a), 4.09 (dd, 1H, $J_{6'a,6'b} = 12.0$ Hz, $J_{5',6'b} = 3.7$ Hz, H-6'b), 4.06, 4.00 (d, 2H, $J = 4.0$ Hz, CH ₂ -17), 3.88 (dt, 1H, $J_{4',5'} = 9.5$ Hz, H-5'), 2.16, 2.15, 2.07, 1.98 84xs, 12H, 4xOAc) |
| 12 | 680 [M+Na] ⁺ | 7.60 (d, 1H, $J = 9.6$ Hz, H-4), 7.80 (d, 1H, $J = 8.0$ Hz, NH), 7.37 (d, 1H, $J = 8.6$ Hz, H-5), 7.35–7.24 (m, 5H, CH ₂ Ph), 6.85 (d, 1H, $J = 8.5$ Hz, H-6), 6.80 (d, 1H, $J = 2.4$ Hz, H-8), 6.24 (d, 1H, $J = 9.5$ Hz, H-3), 5.23 (d, 2H, $J = 5.7$ Hz, H-1'), 5.16 (d, 2H, $J = 7.7$ Hz, CH ₂ Ph), 4.82 (m, 2H, H-2', H-13), 4.67–4.51 (m, 3H, CH ₂ -10, H-4'), 4.74 (d, 1H, $J_{1',2'} = 2.8$ Hz, H-1'), 4.19 (dd, 1H, $J_{5',6'a} = 5.5$ Hz, H-6'a), 4.28 (d, 1H, $J_{3',4'} = 6.0$ Hz, H-3'), 4.27 (dd, 1H, $J_{4',5'} = 5.5$ Hz, $J_{5',6'a'} = 7.6$ Hz, H-5'), 4.05 (dd, 1H, $J_{5',6'a} = 3.7$ Hz, H-6'a), 4.01 (m, 1H, H-17a), 3.98 (m, 1H, H-17b), 3.92 (dd, 1H, $J_{6'a,6'b} = 12.0$ Hz, H-6'b), 1.39, 1.27, 1.25, 1.22 (4xs, 6H, 2xCM ₂) |
| 13 | 680 [M+H] ⁺ | 7.32 (d, 1H, $J = 8.7$ Hz, H-5), 7.11 (br s., 1H, NH), 7.19 (dd, 1H, $J = 2.4$ Hz, 8.7 Hz, H-7), 7.08 (m, 2H, H-6, H-8), 6.17 (s, 1H, H-3), 4.50 (s, 2H, CH ₂ -12), 3.71 (s, 3H, OAc), 2.90 (s, 1H, CH ₂ -9) |
| 14 | 398 [M+H] ⁺ | 7.30 (d, 1H, $J = 9.4$ Hz, H-5), 7.08 (br s., 1H, NH), 7.17 (dd, 1H, $J = 2.6$ Hz, 9.4 Hz, H-7), 7.07 (m, 2H, H-6, H-8), 6.18 (s, 1H, H-3), 4.45 (s, 2H, CH ₂ -12), 3.60 (s, 3H, OAc), 2.94 (m, 1H, CHMe ₂), 2.89 (s, 2H, CH ₂ -9), 1.00, 0.97 (2s, 6H, CHMe ₂) |



Scheme 1



Scheme 2



Scheme 3

Methyl 2-(2H-1-benzopyran-2-one-7-yloxy)acetate (1)(13)

A suspension of 7-hydroxy coumarin (1.0 g, 6.17 mmol) in acetone (20 mL) was refluxed with methyl bromoacetate (1.40 g, 9.15 mmol) and K_2CO_3 (4.69 g, 33.91 mmol) for 12 h. After cooling, the mixture was evaporated to dryness and the residue was partitioned between $CHCl_3$ (50 mL) and water (50 mL). The organic phase was dried (Na_2SO_4), filtered and evaporated to dryness. The residue was recrystallized from acetone to give **1** (1.00 g, 70%, m.p. 150–152 °C; ref. 13: 61%, m.p. 148 °C).

2-(2H-1-benzopyran-2-one-7-yloxy)acetic acid (2)(14)

A solution of **1** (1.50 g, 6.46 mmol) in EtOH (35 mL) and 5% NaOH (6 mL) was heated under reflux for 2 h. After cooling, the solution was evaporated to dryness and the residue was dissolved in water and acidified with 6 mol L⁻¹ HCl. The white precipitate was filtered, dried and crystallized from EtOH to give **2** (1.36 g, 96%), m.p. 218–220 °C, (ref. 14: m.p. 219–220 °C).

Synthesis of coumarin-bearing amino acid esters (3–10)

General procedure. – To a cold solution of the amino acid ester hydrochloride (10 mmol), at –5 °C, in MeCN (20 mL), 7-coumarin acetic acid (10 mmol), hydroxybenzotriazole (HOBt) (1.35 g, 10 mmol) and *N,N'*-dicyclohexyl-carbodiimide (DCC) (10 mmol) were added successively. The reaction mixture was stirred at 0 °C for 1 h, at 5 °C for 1 h, and at 23 °C for 16 h. Dicyclohexylurea (DCU) was filtered, and the filtrate was evaporated to dryness and the residue was dissolved in ethyl acetate, filtered, washed successively with saturated NaCl solution, 5% NaHCO₃ solution, 1.0 mol L⁻¹ HCl, followed by washing with saturated NaCl solution and finally with water. The residue was dried (Na₂SO₄), filtered, evaporated to dryness and recrystallized from the appropriate solvent.

Synthesis of benzyl 2-[2-(2-(2H-benzopyran-2-on-7-yloxy)acetylamino)-3-(2,3,4,6-tetra-O-acetyl- α -D-glucopyranos-1-yloxy)]propanoate (11) and benzyl 2-[2-(2-(2H-benzopyran-2-on-7-oxyl)acetylamino)-3-(2,3,4,6-di-O-isopropylidene- α -D-mannofuranos-1-yloxy)]propanoate (12)

General procedure. – A solution of 2,3,4,6-tetra-O-acetyl- α -D-mannopyranosyl trichloroacetimidate or 2,3,4,6-di-O-isopropylidene- α -D-mannofuranosyl trichloroacetimidate (0.90 mmol) and **10** (358 mg, 0.90 mmol) in dry CH₂Cl₂ (25 mL) was stirred under nitrogen at room temperature for 5 min, followed by the addition of trimethylsilyl trifluoromethanesulphonate (TMSOTf) (19.5 μ L, 0.09 mmol). After stirring for 2 h, solid NaHCO₃ was added slowly, filtered and the filtrate was washed with water (30 mL), the organic layer was dried (Na₂SO₄), filtered and evaporated to dryness. The residue was purified on SiO₂ column (20 g), using ethyl acetate/petroleum ether as eluent to give **11** and **12**, respectively.

Synthesis of 4-substituted coumarin with amino acid esters 13 and 14

General procedure. – The title compounds were prepared by the coupling method applied for preparation of compounds **3–10**, using the amino acid ester hydrochloride (10 mmol), 4-coumarin-acetic acid (10 mmol), hydroxybenzotriazole (HOBt) (1.35 g, 10 mmol) and DCC (10 mmol).

Antitumor screening

Compounds **2–10** and **12** were tested, *in vitro* against a panel of tumor cell lines consisting of CD4 human T-cells containing an integrated human T-leukaemia virus type 1

(HTLV-1), CD4 human acute T-lymphoblastic leukaemia, human splenic B-lymphoblastoid cells, human acute B-lymphoblastic leukaemia, human skin melanoma, human breast adenocarcinoma, human lung squamous carcinoma, human hepatocellular carcinoma, human prostate carcinoma, human foreskin fibroblasts, and human lung fibroblasts. The Microculture Tetrazolium Assay (MTT) method (15) was used for estimation of the *in vitro* tumor-inhibiting activity of the tested compounds. The cell lines of tumor subpanels were incubated within five concentrations (0.01–100 $\mu\text{g mL}^{-1}$) of each tested compound for 48 h.

RESULTS AND DISCUSSION

The starting material 2-(2H-1-benzopyran-2-one-7-yloxy)acetic acid (**2**) was prepared by saponification of the corresponding ester **1** (**13**) in 96% yield. Compound **2** was prepared previously by direct condensation of 7-hydroxy coumarin with chloroacetic acid, in 61% yield (**14**).

A suitable coupling method (16) was employed for the formation of peptides by reaction of the carboxylic acid group with acylated amino acid, using 1-hydroxybenzotriazole (HOBt) (**17**, **18**) and *N,N'*-dicyclohexylcarbodiimide (DCC) (**19**) as coupling reagents. HOBt (**1**) is currently the most frequently used activating agent for the carboxyl group of amino acids. The procedure is fast and suppresses racemization, especially in the presence of DCC (**20**).

Amides **3–10** were prepared by coupling **2** with the appropriate acylated amino acids (L-glycine acetate hydrochloride, L-leucine acetate hydrochloride, L-alanine acetate hydrochloride, L-valine acetate hydrochloride, L-phenylglycine acetate hydrochloride, benzyl-L-proline carboxylate hydrochloride, D,L-tryptophane acetate hydrochloride and benzyl-L-serine carboxylate hydrochloride) in the presence of HOBt and DCC as coupling reagents to give **3–10** in 56–85% yield (Scheme 1).

Recently, Schmidt *et al.* (21, 22) reported an efficient anomeric stereocontrolled glycosylation method, with high yield of α -anomer, by using *O*-glycosopyranosyl trichloroacetimidate derivative as donor and alcohol as acceptor precursors in the presence of catalytic amounts of trimethylsilyl trifluoromethanesulfonate (TMSOTf) as Lewis acid. By applying Schmidts procedure, **10** was selected in our present work as alcohol acceptor for coupling with the protected *O*-mannopyranosyl- and *O*-mannofuranosyl-1-trichloroacetimidate derivatives as donor precursors in the synthesis of new glycoside derivatives **11** and **12** in 59% and 62% yield, respectively (Scheme 2). Trichloroacetimidates, characterized by their stabilities at room temperature for long periods of time without any decomposition, were prepared from the reaction of their corresponding 1-hydroxy-protected glycosyl precursors (21, 22) with trichloroacetonitrile in the presence of 1,5-diazabicyclo[5,4,0]undec-5-ene (DBU) as catalyst.

The structures of **3–10** were determined by their ^1H , ^{13}C NMR and by mass spectra (Tables II and III). The coumarin protons showed a similar pattern. H-3, H-4, H-5 and H-8 protons appeared as doublets in region δ 6.14–6.24 ($J = 9.3\text{--}9.5$ Hz), 7.48–7.67 ($J = 8.9\text{--}9.6$ Hz), 7.28–7.55 ($J = 8.4\text{--}9.6$ Hz) and 6.68–6.81 ppm ($J = 2.0\text{--}2.6$ Hz), respectively. H-6 protons were oriented as a doublet of doublets in the region δ 6.78–6.87 ppm ($J =$

8.6–9.5 and $J = 2.3$ – $2.2.6$ Hz), except for **9** which showed multiplets at δ 7.54, attributed to H-3 and H-4 and at δ 6.54 ppm, belonging to H-3 and H-4. The singlets that appeared in the region δ 4.38–4.69 ppm were characterized as CH₂-10 protons. H-13 protons appeared as doublets, a doublet of doublets or multiplets at δ 3.58–4.92 ppm, depending on the functional group adjacent to C-13. The other protons of the amino acid ester were fully analyzed. The ¹³C NMR spectra of **3**–**10** contained similar resonance signals of the coumarin carbons ring C-2–C-8a. The chemical shifts between δ_C 161.5 and 161.8 ppm were assigned to the carbonyl group of the benzopyran ring (C-2), while the resonances in the range of δ_C 66.8–67.4 ppm were assigned to CH₂-10 protons. Compounds **11** and **12** were identified from the ¹H NMR and ¹³C NMR spectra, which showed almost similar resonance of benzopyran ring atoms as those of **10**. The sugar proton H-1' of **12** appeared as a doublet at δ 5.23 ppm, while H-2' was oriented with H-13 as a multiplet at δ 4.82 ppm. H-4' appeared together with CH₂-10 as a multiplet at δ 4.67–4.51 ppm. The small $J_{1',2'}$ coupling (5.23 Hz) confirmed the α -anomer of glycoside **12**. Assignment of proton resonances of **11** was confirmed by homo decoupling and by COSY spectra. ¹H-¹³C COSY, DFQ-COSY, HSQC, HMQC and HMBC were used for assignments of the ¹³C resonances.

The ¹³C NMR spectrum of **11** showed signals at higher field: δ 170.6–169.6 ppm were attributed to the acetoxy groups of the sugar moiety while resonances at δ 160.6, 167.0, 168.0 ppm were assigned to the carbonyl groups C-2, C-11 and C-14, respectively. C-4a, C-7 and C-8a appeared at δ 113.8, 159.8 and 155.6 ppm, respectively. The gradient selected ¹H,¹³C-HSQC spectrum in CDCl₃ showed couplings between the benzopyran ring H-3–H-8 as well as CH₂-10, H-13 and CH₂-17 at δ 6.14 ppm (d, $J = 9.5$ Hz), 7.65 ppm (d, $J = 9.6$ Hz), 7.43 ppm (d, $J = 8.8$ Hz), 6.96 ppm (d, $J = 8.8$ Hz), 6.91 ppm (d, $J = 2.0$ Hz), 4.62 ppm (s), 4.93 ppm (m), 4.00 ppm (d, $J = 4.0$ Hz) and C-3–C-8, C-10, C-13 and C-17 at δ 114.2, 143.0, 129.2, 112.3, 102.5, 52.4, 52.2, 68.5 ppm, respectively. The small $J_{1',2'}$ coupling (2.8 Hz) is typical of the α -configured mannopyranoside. The DFQ-COSY spectrum showed a linear correlation between the sugar protons H-1' (δ 4.74 ppm), H-3' (δ 4.28 ppm) and the amino acid protons CH₂-10 (δ 4.62 ppm), H-13 (δ 4.93 ppm) and CH₂-17 (δ 4.00 ppm), indicating α -configuration. Additional proof for α -configuration came from the ROESY spectrum, which suggested a large distance between H-1' and H-5'. The large $J_{4',5'}$ coupling (9.5 Hz) of **11** is indicative of the ⁴C₁ conformation of the pyranose ring. Furthermore, C-17 at δ 68.5 ppm was inferred from the HMBC and HMQC spectra by showing ³J_{C,H} coupling with H-1' at δ 5.23 ppm.

Further, our work was modified by selecting the 4-coumarin acetic acid (**23**) as precursor for the synthesis of new derivatives to examine their antiviral activity in comparison to the 7-coumarin analogues **3**–**10**. Compounds **13** and **14** were prepared in 71 and 65% yields from 4-coumarin acetic acid by applying the coupling method used previously in the presence of HOBt and DCC as coupling reagents (Scheme 3). The structures of **13** and **14** were determined from their ¹H NMR and mass spectra.

Antiviral assay

Compounds **3**–**10** were tested for their anti-HIV-1 activity *in vitro*, using III_B strain in human T-lymphocyte (MT-4) cells, based on a Microculture Tetrazolium (MTT) assay and the results are reported in Table IV. Cytotoxicity was also measured on MT-4 cells.

Table III. ^{13}C NMR data of the newly prepared compounds

| Compd. | ^1H NMR (δ , ppm) |
|--------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 3 | 172.4 (C-14), 166.7 (C-11), 160.1 (C-2), 159.8 (C-7), 155.1 (8a), 142.8 (C-4), 128.8 (C-5), 113.4 (C-3), 113.2 (C-4a), 112.0 (C-6), 102.0 (C-8), 67.1 (C-10), 51.8 (C-13), 50.0 (C-14), 40.7 (C-15), 24.4 (C-16), 22.3, 21.3 ($\text{C}_{16}\text{-Me}_2$) |
| 4 | 171.8 (C-14), 167.1 (C-11), 160.6 (C-2), 160.1 (C-7), 155.4 (8a), 143.2 (C-4), 129.2 (C-5), 113.7 (C-3), 113.5 (C-4a), 112.4 (C-6), 102.1 (C-8), 67.3 (C-10), 56.8 (C-13), 52.1 (C-14), 30.9 (C-15), 18.8, 17.7 (2xMe) |
| 5 | 172.8 (C-14), 166.6 (C-11), 160.7 (C-2), 160.1 (C-7), 155.6 (8a), 143.1 (C-4), 129.2 (C-5), 114.0 (C-3), 113.6 (C-4a), 112.3 (C-6), 102.4 (C-8), 67.4 (C-10), 52.6 (C-13), 47.7 (C-14), 18.2 ($\text{C}_{13}\text{-Me}$) |
| 6 | 171.8 (C-14), 167.1 (C-11), 160.6 (C-2), 160.1 (C-7), 155.4 (8a), 143.2 (C-4), 129.2 (C-5), 113.7 (C-3), 113.5 (C-4a), 112.4 (C-6), 102.1 (C-8), 67.3 (C-10), 56.8 (C-13), 52.1 (C-14), 30.9 (C-15), 18.8, 17.7 (2xMe) |
| 7 | 170.8 (C-14), 166.7 (C-11), 160.7 (C-2), 160.1 (C-7), 155.5 (8a), 143.2 (C-4), 135.9 (Ar-C _a), 129.2 (Ar-C _f), 128.9 (Ar-C _g), 128.6 (Ar-C _d), 127.3 (C-5), 113.7 (C-3), 113.5 (C-4a), 112.5 (C-6), 102.2 (C-8), 67.3 (C-10), 56.1 (C-13), 52.8 (C-13), 20.9 (C-14) |
| 8 | 183.1 (C-17), 171.5 (C-14), 165.8 (C-11), 161.1 (C-2), 160.9 (C-7), 155.5 (8a), 143.4 (C-4), 135.9, 129.0, 128.7, 128.5, 128.3, 128.2 ($\text{CH}_2\text{Ar-C}$), 127.9 (C-5), 113.4 (C-3), 113.2 (C-4a), 112.6 (C-6), 102.0 (C-8), 67.0 (C-16), 66.8 (C-10), 59.2 ($\text{CH}_2\text{Ph-13}$), 28.6 (C-15), 24.9 (C-14) |
| 9 | 170.8 (C-14), 166.7 (C-11), 160.7 (C-2), 160.1 (C-7), 155.5 (8a), 143.2 (C-4), 136.3 (indol-C _i), 129.0 (C-5), 127.4 (indol-C _d), 123.3 (indol-C _b), 122.0 (indol-C _f), 119.6 (indole-C _g), 118.2 (indole-C _e), 113.6 (C-3), 113.5 (C-4a), 112.2 (C-6), 111.7 (indole-C _h), 109.0 (indole-C _c), 102.2 (C-8), 67.2 (C-10), 52.8 (C-13), 52.5 (C-14), 27.3 ($\text{CH}_2\text{-15}$) |
| 10 | 169.8 (C-14), 167.6 (C-11), 161.0 (C-2), 160.0 (C-7), 155.5 (8a), 143.3 (C-4), 129.2 (C-5), 135.1, 128.6, 128.5, 128.1, 128.2 ($\text{CH}_2\text{Ar-C}$), 127.9 (C-5), 113.9 (C-3), 113.6 (C-4a), 112.4 (C-6), 102.5 (C-8), 67.6 (C-16), 67.4 (C-10), 62.7 (CH_2Ph), 54.6 (C-13) |
| 11 | 170.6, 169.8, 169.7, 169.6 (COMe), 168.8 (C-14), 167.0 (C-11), 160.6 (C-2), 159.8 (C-7), 155.6 (8a), 143.0 (C-4), 129.2 (C-5), 134.8, 128.7, 128.6, 128.3 ($\text{CH}_2\text{Ph-C}$), 129.2 (C-5), 114.2 (C-3), 113.8 (C-4a), 112.3 (C-6), 102.5 (C-8), 98.2 (C-1'-sugar), 69.3 (C-2'-sugar), 69.2 (C-5'-sugar), 68.6 (C-3'-sugar), 68.5 (C-17), 67.9 [C-16 (CH_2Ph)], 67.4 (C-10), 62.3 (C-6'-sugar), 52.2 (C-13), 20.8, 20.7, 20.6, 20.5 (4xCOMe) |
| 12 | 170.1 (C-14), 168.2 (C-11), 160.9 (C-2), 160.2 (C-7), 154.2 (8a), 143.3 (C-4), 135.9 ($\text{CH}_2\text{Ph-C}$), 129.2 (C-5), 128.9, 128.8, 128.2 ($\text{CH}_2\text{Ph-C}$), 114.7 (C-3), 114.2 (C-4a), 112.6 (C-6), 103.6 (C-8), 105.1 (C-1'-sugar), 84.0 (C-2'-sugar), 80.5 (C-3'-sugar), 78.4 (C-5'-sugar), 76.3 (C-4'-sugar), 72.1 (C-6'-sugar), 68.1 (C-16 CH_2Ph), 67.5 (C-10), 63.8 (C-17, $\text{CH}_2\text{-glycoside}$), 59.2, 57.1 (Me, acetal groups), 52.7 (C-13) |

The gp41 subunit of the human immunodeficiency virus type 1 (HIV-1) envelope glycoprotein plays an important role in HIV-1 entry and serves as an attractive target for the development of HIV-1 entry inhibitors, a new class of anti-HIV drugs (24). Triggered by gp120 binding to CD4 and a coreceptor, gp41 undergoes a conformation shift from a native prefusogenic state to a fusogenic state, in which the N-terminal heptad repeat (NHR) and C-terminal heptad repeat (CHR) associate to form a six-helix bundle, representing the fusion-active gp41 core. Any compound that disrupts the gp41 sixhelix bun-

dle formation may inhibit the gp41-mediated membrane fusion, thereby blocking HIV-1 entry into target cells (25).

Our target was the disruption of the gp41 sixhelix bundle formation by the newly synthesized amino acids, leading to inhibition of HIV. None of the *in vitro* tested compounds **3–10** and **12** were found to inhibit HIV-replication, at EC_{50} lower than the CC_{50} compared to the antiviral agents efavirenz (EFV) (26) and azidothymidine (AZT) (27), whereas compound **11** was not tested against HIV. On the other hand, **13** and **14** were tested against HIV-1 (III_B strain) with $EC_{50} > 14$ and $1.6 \mu\text{g mL}^{-1}$, respectively, and against HIV-2 (ROD strain) with $EC_{50} > 19$ and $3.2 \mu\text{g mL}^{-1}$, respectively, at non-toxic concentrations.

The above data showed no selective anti-HIV activity, except for compound **14**, which showed an encouraging result by inhibiting of HIV with EC_{50} value $> 1.6 \mu\text{g mL}^{-1}$.

Compounds **3–10** and **12** were evaluated against BVDV, and showed no inhibition at non-toxic concentrations, in comparison to the iminosugar derivatives (28), since the minimum inhibitory concentration required to reduce the virus-induced cytopathogenicity by 50% was higher than $100 \mu\text{g mL}^{-1}$ as shown in Table IV. Compounds **11**, **13** and **14** were not tested against BVDV.

Table IV. *In vitro* anti-HIV-1^a and anti-BVDV activity of some coumarin compounds

| Compd. | CC_{50} MT-4 ($\mu\text{g mL}^{-1}$) ^b | EC_{50} HIV-1 ($\mu\text{g mL}^{-1}$) ^c | CC_{50} BVDV ($\mu\text{g mL}^{-1}$) | EC_{50} BVDV ($\mu\text{g mL}^{-1}$) |
|-----------------------|----------------------------------------------------------|-----------------------------------------------------------|---------------------------------------------|---------------------------------------------|
| 3 | > 100 | > 100 | > 100 | 91 |
| 4 | > 100 | > 100 | > 100 | > 100 |
| 5 | > 100 | > 100 | > 100 | > 100 |
| 6 | > 100 | > 100 | > 100 | 66 |
| 7 | > 100 | > 100 | > 100 | 100 |
| 8 | > 100 | > 100 | > 100 | > 100 |
| 9 | 24 | > 24 | > 100 | 100 |
| 10 | > 100 | > 100 | > 100 | > 100 |
| 12 | 49 | > 49 | > 100 | > 100 |
| 13 | > 14 | > 14 | | |
| 13^d | > 19 | > 19 | | |
| 14 | > 1.6 | > 1.6 | | |
| 14^d | > 3.2 | > 3.2 | | |
| EFV | 40 | 0.003 | | |
| AZT | 63 | 0.02 | | |

^a Anti-HIV-1 activity measured with strain III_B.

^b CC_{50} concentration required to reduce the viability of mock-infected MT-4 cells by 50%, as determined by the MTT (Microculture Tetrazolium Assay) method.

^c EC_{50} concentration required to achieve 50% protection of MT-4 cells from the HIV-1 induced cytopathogenicity, as determined by the MTT method.

^d Anti-HIV-2 activity measured with strain ROD.

EFV – efavirenz

AZT – azidothymidine

Anticancer assay

The Microculture Tetrazolium Assay (MTT) method (15), which is based on metabolic reduction of 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide, was used for preliminary estimation of the *in vitro* tumor-inhibiting activity of the coumarin derivatives 3–10 and 12 against a panel of tumor cell lines. The results are summarized in Table V.

Table V. Antitumor activity in most sensitive tumor cell lines

| Compd. | Tumor | Cell line | CC ₅₀ ($\mu\text{mol L}^{-1}$) ^a |
|--------|------------------------------------|------------------------|----------------------------------------------------------|
| 9 | Leukaemia/ lymphoma | MT4 ^b | 24 |
| | | CCRF-CEM ^c | 36 |
| | | WIL-2NS ^d | 45 |
| | | CCRF-SB ^e | 48 |
| | MT-4 | | 24 |
| | MDBK | | > 100 |
| | Solid tumor- derived cell lines | SK-MEL-28 ^f | 60 |
| | | MCF7 ^g | > 100 |
| | | SKMES-1 ^h | 80 |
| | | HepG2 ⁱ | > 100 |
| | | DU145 ^j | 90 |
| | | CRL 7065 ^k | > 100 |
| | Normal-cell lines | MRC-5 ^l | > 100 |
| | | | > 100 |
| 12 | Leukaemia/ lymphoma | MT4 ^b | 49 |
| | | CCRF-CEM ^c | 53 |
| | | WIL-2NS ^d | 68 |
| | | CCRF-SB ^e | 59 |
| | MT-4 | | > 100 |
| | MDBK | | > 100 |
| | Solid tumor- derived cell lines | SK-MEL-28 ^f | 60 |
| | | MCF7 ^g | > 100 |
| | | SKMES-1 ^h | 80 |
| | | HepG2 ⁱ | > 100 |
| | | DU145 ^j | 90 |
| | | CRL 7065 ^k | > 100 |
| | Normal-cell lines | MRC-5 ^l | > 100 |
| | | | > 100 |

^a Compound concentration required to reduce cell proliferation by 50%, as determined by the MTT method, under conditions allowing untreated controls to undergo at least three consecutive rounds of multiplication.

^b CD4 human T-cells containing an integrated HTLV-1.

^c CD4 human acute T-lymphoblastic leukaemia.

^d Human splenic B-lymphoblastoid cells.

^e Human acute B-lymphoblastic leukaemia.

^f Human skin melanoma.

^g Human breast adenocarcinoma.

^h Human lung squamous carcinoma.

ⁱ Human hepatocellular carcinoma.

^j Human prostate carcinoma.

^k Human foreskin fibroblasts.

^l Human lung fibroblasts.

All the new compounds were inactive against all tumor cell lines ($CC_{50} > 100 \mu\text{mol L}^{-1}$, except for compounds **9** and **12**, which showed marked activity against leukaemia/lymphoma. Comparing the activity of **9** to the other coumarin bearing amino acid derivatives showed that the inclusion of the indole moiety shifted the threshold of potency from the inactive side toward activity, particularly against the leukaemia cell lines MT4 ($CC_{50} = 24 \mu\text{mol L}^{-1}$). It is noticed that the substances with aliphatic substituents are inactive whereas the tryptophane derivative **9** shows significant activity. This may be due to the indole residue, which performs more intermolecular interactions.

The introduction of a furanose sugar moiety into the OH group of **12** by glycoside linkage may be responsible for a slight change in the antitumor activity.

CONCLUSIONS

The structure activity relationship suggested that the substituted carbon-coumarin linkage showed higher HIV inhibition than the corresponding analogues having oxygen linkage, and this result should lead us to modify our new target molecules by introducing more potent groups with carbon linkage.

Acknowledgement. – We thank Dr. Gosselin of Laboratoire de Chimie Organique, Université de Montpellier II (France), for helping with the antiviral and antitumor screening. Miss Friemel and Mr. Galtesky of Konstanz University (Germany) are gratefully acknowledged for the 2D-NMR experiments and mass measurements, respectively.

REFERENCES

1. G. Feuer, *The Metabolism and Biological Actions of Coumarins*, in *Progress in Medicinal Chemistry* (Eds. G. P. Ellis and G. B. West), Vol. 10, North-Holland and Publishing Company, New York 1974, pp. 85.
2. K. Padmawinata, Isolation and identification of cancer delaying compounds from the leaves of *Giejera salicifolia*, *Acta Pharmacol.* **4** (1973) 1–9.
3. K. G. Mann, The challenge of regulating anticoagulant drugs: focus on warfarin, *Am. Heart J.* **149** (2005) 495–536.
4. Y. Sato, Y. Kobayashi, T. Nagasaki, T. Oshima, S. Kumakura, K. Nakayama, H. Koike and H. Takagi, Studies on new β -adrenergic blocking agents I. Synthesis and pharmacology of coumarin derivatives, *Chem. Pharm. Bull.* **20** (1972) 905–917.
5. Y.-L. Chen, T.-C. Wang, N.-C. Chang, Y.-L. Chang, C.-M. Teng and C.-C. Tzeng, α -Methylene- γ -butyrolactones: synthesis and vasorelaxing activity assay of coumarin, naphthalene and quinoline derivatives, *Chem. Pharm. Bull.* **46** (1998) 962–965.
6. K. Furumi, T. Fujioka, H. Fujii, H. Okab, Y. Nakano, H. Matsunaga, M. Katano, M. Mori and K. Mihashi, Novel antiproliferative falcarindiol furanocoumarin ethers from the root of *Angelica japonica*, *Bioorg. Med. Chem. Lett.* **8** (1998) 93–96.
7. T. D. Bradshaw, M. C. Bibby, J. A. Double, I. Fichtner, P. A. Cooper, M. C. Alley, S. Donohue, S. F. Stinson, J. E. Tomaszewski, E. A. Sausville and M. F. G. Stevens, Preclinical evaluation of amino acid prodrugs of novel antitumor 2-(4-amino-3-methylphenyl)benzothiazoles, *Mol. Cancer Ther.* **1** (2002) 239–246.

8. M. D. Wittman and J. F. Kadow, *Amino acid derivatives of paclitaxel*, U.S. Pat. 5,489,589, 6 Feb. 1996; ref. *Chem. Abstr.* **124** (1996) 289947a.
9. A. Casini, A. Scozzafava and C. T. Supuran, Cystein-modifying agents: A possible approach for effective anticancer and antiviral drugs, *Environ. Health Perspect.* **110** (2002) 801–806.
10. K.-L. Yu, W. E. Harte, P. Spinazze, J. C. Martin and M. M. Mansuri, Synthesis of 1,3-diamino-2-hydroxypropane derivatives as pseudosymmetric HIV protease inhibitors, *Bioorg. Med. Chem. Lett.* **3** (1993) 535–538.
11. V. Escuret, P. Parvaz, O. Hantz, M. A. Petit, C. Trepo and F. Zoulim, Study of the antiviral mechanism of action of ribavirin in the bovine viral diarrhea virus model, *Gastroenterol. Clin. Biol.* **26** (2002) 584–590.
12. J. R. Stangl, K. L. Carroll, M. Illichmann and R. Striker, Effect of antimetabolite immunosuppressants on Flaviviridae, including hepatitis C virus, *Transplantation* **77** (2004) 562–567.
13. B. S. Joshi, N. Viswanathan, C. L. Kaul and R. S. Grewal, Evaluation of some naturally occurring & synthetic coumarins for hypotensive activity, *Indian J. Chem.* **19B** (1980) 495–499.
14. M. Birau and Z. Y. Wang, A dual-mode molecular switch based on a chiral binaphthol-coumarin compound, *Tetrahedron Lett.* **41** (2000) 4025–4028.
15. M. C. Alley, D. A. Scudiero, A. Monks, M. L. Hursey, M. J. Czerwinski, D. L. Fine, B. J. Abbott, J. G. Mayo, R. H. Shoemaker and M. R. Boyd, Feasibility of drug screening with panels of human tumor cell lines using a microculture tetrazolium assay, *Cancer Res.* **48** (1988) 589–601.
16. A. Asagarasu, T. Uchiyama and K. Achiwa, Design and synthesis of HIV protease inhibitor linked with binding unit to gp 120, *Chem. Pharm. Bull.* **46** (1986) 697–703.
17. S. Davis and A. K. Mohammed, Assessment of racemization *n*-alkylated amino acid derivatives during peptide coupling in a model dipeptide system, *J. Chem. Soc. Perkin Trans. 1* **1981**, 2982–2990.
18. W. König and R. Geiger, Eine neue Methode zur Synthese von Peptiden: Hydroxybenzotriazol-en, *Chem. Ber.* **103** (1970) 788–798.
19. R. M. Pennington and R. R. Fischer, Dicyclohexylcarboimide modification of bovine heart mitochondrial transhydrogenase, *J. Biol. Chem.* **256** (1981) 8963–8969.
20. A. Katritzky, K. Suzuki and S. K. Singh, N-Acylation in combinatorial chemistry *ARKIVOC* **2004**, 12–35.
21. I. A. I. Ali, E. S. H. El-Ashry and R. R. Schmidt, Protection of hydroxyl groups with diphenyl and 9-fluorenyl trichloroacetimidates – effect on anomeric stereocontrol, *Eur. J. Org. Chem.* **2003**, 4121–4131.
22. A. I. Ali, E. S. H. El-Ashry and R. R. Schmidt, Imidomethylation of C-nucleophiles using *O*-phthalimidomethyl trichloroacetimidate and catalytic mounts of TMSOTf, *Tetrahedron* **60** (2004) 4773–4780.
23. A. N. Brubaker, J. DeRuiter and W. L. Whitmer, Synthesis and rat lens aldose reductase inhibitory activity of some benzopyranones, *J. Med. Chem.* **29** (1986) 1094–1099.
24. M. J. Root and H. K. Steger, HIV-1 gp41 as a target for viral entry inhibition, *Curr. Pharm. Design*, **10** (2004) 1805–1825.
25. L. Ni, G. F. Gao and P. Tien, Rational design of highly potent HIV-1 fusion inhibitory proteins: Implication for developing antiviral therapeutics, *Biochem. Biophys. Res. Commun.* **332** (2005) 831–836.
26. S. D. Young, S. F. Britcher, L. O. Tran, L. S. Payne, W. C. Lumma, T. A. Lyle, J. R. Huff, P. S. Anderson, D. B. Olsen, S. S. Carroll, D. J. Pettibone, J. A. O'Brien, R. G. Ball, S. K. Balani, J. H. Lin, I. W. Chen, W. A. Schleif, V. V. Sardana, W. J. Long, V. W. Byrnes and E. A. Emimi, Novel, highly potent non-nucleoside inhibitor of the human immunodeficiency-virus type-1 reverse transcriptase, *Antimicrob. Agents Chemother.* **39** (1995) 2602–2605.

27. H. Mitsuya, K. J. Weinhold, P. A. Furman, M. H. St. Clair, S. N. Lehrmann, R. C. Gallo, D. Bolognesi, D. W. Barry and S. Broder, 3'-Azido-3'-deoxythymidine (BW A509U), an antiviral agent that inhibits the ineffectivity and cytopathic effect of human T-lymphotropic virus type III/lymphadenopathy-associated virus *in vitro*, *Proc. Natl. Acad. Sci. USA* **82** (1985) 7096–7100.
28. D. Durantel, N. Branza-Nichita, S. Carrouee-Durantel, T. D. Butters, R. A. Dwek and N. Zitzmann, Study of the mechanism of antiviral action of iminosugar derivatives against bovine viral diarrhoea virus, *J. Virol.* **75** (2001) 8987–8998.

S A Ž E T A K

Derivati aminokiselina. Dio I. Sinteza, antivirusno i antitumorsko djelovanje novih estera α -aminokiselina s kumarinskim supstituentom

NAJIM A. AL-MASOUDI, IMAN A. AL-MASOUDI, IBRAHIM A. I. ALI,
YASEEN A. AL-SOUD, BAHJAT SAEED i PAOLO LA COLLA

U radu je opisana sinteza estera aminokiselina s kumarinskim ostatkom **3–15**. Ispitano je antivirusno djelovanje sintetiziranih spojeva na HIV-1 i goveđi virus diareje (BVDV), te *in vitro* citotoksičnost spojeva **3–10** i **12** na tumorskim linijama CD4 humanih T-stanica. Spoj **14** pokazao je inhibiciju HIV-1 s $EC_{50} > 1,6 \mu\text{g mL}^{-1}$, dok je spoj **9** djelotvoran na leukemiju (MT4) s $CC_{50} = 24 \mu\text{mol L}^{-1}$.

Ključne riječi: aminokiseline, antitumorsko djelovanje, antivirusno djelovanje, kumarin

Fachbereich Chemie, Universität Konstanz, Postfach 5560, D-78457 Konstanz, Germany

College of Veterinary Medicine, University of Basrah, Basrah, Iraq

Department of Chemistry, Faculty of Science, Suez Canal University, Ismailia, Egypt

Department of Chemistry, College of Science, University of Al al-Bayt, Jordan

Universita di Cagliari, Dipartimento di Biologia Sperimentale Sezione di Microbiologia, Cagliari, Italy